

~~BEHAVIOR OF THE TOTAL ELECTRON CONTENT DURING GEOMAGNETIC STORM 17 MARCH 2015 OVER THE ARCTIC REGION~~

THE IONOSPHERIC TOTAL ELECTRON CONTENT RESPONSE TO THE 17TH -18TH MARCH 2015 GEOMAGNETIC STORM OVER THE ARCTIC REGION

Abstract

We describe the ionospheric response to the St. Patrick's Day storm, which was the biggest geomagnetic storm of the 24th solar cycle up until that point. The storm occurred on March 17–18, 2015. TEC measurements were obtained using Global Positioning System (GPS) receivers of the Canadian High Arctic Ionospheric Network (CHAIN) from three stations: Cambridge Bay, Eureka, and Rabbit Lake. Based on the results, we observed that during stormy conditions, TEC was significantly lower on Cambridge Bay than it was on Eureka and Rabbit Lake.

- Give a general introduction of introduction of the geomagnetic storm and a brief summary of the methodology, main results and conclusion.

Introduction:

The ionospheric reaction to geomagnetic storms has been one of the essential subjects in space climate research for quite a long time (Gonzalez et al., 1994; Richmond et al., 2000). During geomagnetic storms, the improved infusion of energy from the sun oriented breeze causes unsettling influences in the science and elements of the coupled thermosphere and ionosphere framework (Prölss, et al., 1995; Buonsanto, et al., 1999). Many examinations have been given to considering ionospheric storm impacts close to the F district top (Fuller-Rowell et al., 1994; Burns et al., 1995; Mendillo, 2006), yet the ionospheric reaction to geomagnetic storms is as yet not completely comprehended.

Examinations of ionosphere elements and aggravations during geomagnetic storms are still of high significance because of impact of the ionosphere on the space climate. The ionosphere, as a functioning layer, assumes a significant part in the space climate because of its affectability to the sun powered action. Thusly, the space climate influences satellite frameworks, energy transport, air traffic and, in outcome, the economy (Schrijver et al., 2015). Subsequently, the investigations of the ionosphere conduct assume a vital part in comprehension and estimating of the space climate. The ionosphere is a layer of the air comprising for the most part of ionized particles, which causes the satisfy light electromagnetic signs to be deferred, changes in signal strength and other unfriendly impacts. The extent of these ionospheric impacts generally relies upon the condition of the shifting ionosphere, its complete electron content (TEC) and the recurrence of electromagnetic waves. The unique changes of the ionosphere are most effectively separated into normal and abrupt ones. Standard changes are straightforwardly identified with the periodicity of the variables impacting them, like the sun powered cycle. These ordinary varieties are to some degree simple to display (Bilitza et al., 2008; Maruyama et al. 2009). The ionospheric storms (aggravations in the earthly ionosphere delivered by geomagnetic storms)

cause huge unsettling influences for innovative frameworks, for example, the static or dynamical situating with GNSS satellites, and others, which rely upon the transionospheric correspondences. Essentially, in correlation with under calm conditions, the electron thickness can increment or diminishing during geomagnetic storm periods. These changes, which have been called positive ionospheric tempest or positive tempest impact and negative ionospheric tempest or negative tempest impact individually, happen in light of the fact that there is critical energy input (from the sun based breeze) into the polar ionosphere, for the most part over a time of a few hours to a day. A few driver powers have been utilized to clarify the ionospheric impacts during storms at various scopes. For instance, it is accepted that level convection overwhelms in the polar covers, and structure changes, molecule precipitation and electric fields rule in the auroral zones, while electric fields, meridional breezes and creation changes rule at tropical and low scopes. In the auroral and polar ionosphere, those abnormalities at an alternate scale have a typical element during geomagnetic storms, which causes vacillations in the Total Electron Content (TEC). Thusly, the limited scale ionospheric plasma thickness inconsistencies produce quick change in the abundance and period of transionospheric radio signs, which is known as sparkle. The huge scope inconsistencies and related TEC vacillations an muddle stage vagueness goal, increment the quantity of uncorrected cycle slips and misfortunes of sign lock in GNSS. Since there are moderately inadequate outcomes for TEC. The large-scale irregularities and associated TEC fluctuations can complicate phase ambiguity resolution, increase the number of uncorrected cycle slips and losses of signal lock in GNSS. Because there are relatively sparse results for TEC at sub-auroral, auroral and polar latitudes in this paper we analyze the behavior of TEC at three stations located in the Arctic sectors during intense geomagnetic storms in the 2015.

- Check on the flow of your introduction.
- Mention a few recent related studies on geomagnetic storms (on ionospheric TEC) in other regions and their findings. Identify notable gaps in these studies and bring in your study, highlighting the importance of the study.

Data and methodology

The GNSS perception information of Crustal Movement Observation Network CHAIN in three stations in particular given in table 1, which incorporates three GPS beneficiaries, are utilized to reproduce the ionospheric complete electron content (TEC) circulation during 17 March, 2015 geomagnetic storm periods. The scope and longitude of these three GPS recipients are displayed in Table 1. Fig. 1 presents the topographical area of GPS beneficiaries over High Latitude stations. The CHAIN GPS recipients are GPS Ionospheric Scintillation and TEC Monitors (GISTMs) model GSV4000B. In outline, a GISTM comprises of a NovAtel OEM4 double recurrence beneficiary with uncommon firmware explicitly arranged to gauge adequacy and stage shine got from the L1 recurrence GPS signals and ionospheric absolute electron content (TEC) got from the L1 and L2 recurrence GPS signals. This collector is fit for following and

revealing glimmer and TEC estimations from up to ten GPS satellites in see. Stage and plentifulness information are examined and logged, either in crude structure or detrended, at a pace of 50 Hz. Nine of the ten collectors are as of now took care of by a NovAtel GPS-702 receiving wire, the exemption being the Qikiqtarjuaq recipient what shares an Ashtech ASH701945E_M radio wire with a pre-existing Natural Resources Canada GPS beneficiary through a splitter. At most occasions, 8 to 10 GPS satellites are apparent to a solitary ground recipient. From GPS information we figure the complete electron content (TEC), a worth coordinated in a 1 m² section along the satellite to collector raypath. Detective is determined in TEC units (TECu), where 1 TECu = 10¹⁶ electrons for each square meter.




Figure 1: CHAIN Ionspheric Network over Arctic Region

Table 1. Geographic and Geomagnetic Coordinates of Stations Used in the Study

| Name | Geo Lat | Geo Long | Instrument | Model |
|---------------------|-------------|--------------|------------|----------|
| Cambridge Bay (CBB) | 69.101929 N | 254.884829 E | GISTM/GPS | GSV4004B |
| Eureka (EUR) | 79.990089 N | 274.097557 E | GISTM/GPS | GSV4004B |
| Rabbit Lake (RAB) | 58.226935 N | 256.322945 E | GISTM/GPS | PolaRxS |

In this paper, we are researching subtly the overall effect of the St. Patrick's Day tempests of March 2015 on the ionosphere over the northern polar cap locale. Exceptionally compelling are the positive ionospheric storms saw at the northern polar stations Cambridge Bay, Eureka and Rabbit Lake in the attractive tempests, which propose that the plasma thickness develops along these lines during comparable geophysical conditions. While the commitments of different tempest time wonders toward the arrangement of TOI during these tempests have been broadly contemplated, the job of outer driving instruments in altering the reaction of the polar ionosphere has not been focused. Consequently, the principle objective of the investigation is to break down

exhaustively the systems that lead to the age of positive ionospheric storms over the northern polar cap area. In the accompanying, we depict our outcomes in subtleties and talk about their importance in creating comprehension of the reaction of the Earth's polar ionospheric framework to geomagnetic storms.

- For data and methodology, identify data sources for TEC and solar wind parameters and geomagnetic indices (SYM-H and AE). Consider using the following format:
 2. Data and methodology
 - 2.1 Data acquisition
 - Give out the methodology i.e. how is the TEC data retrieved (website) and processed?, how are the solar wind parameters and geomagnetic indices retrieved(website)?
 - 2.2 Methodology
 - How are the global TEC maps generated/plotted from TEC data?, how are the solar wind parameters and geomagnetic indices plotted?

Result

Observation of geomagnetic condition we are describing in figure 2. The severe geomagnetic storm occurred on 17 March 2015 and caused the dramatic response in the ionosphere–plasmasphere–magnetosphere system. Figure 1 shows the variations of interplanetary and geomagnetic parameters during 15–20 March 2015.




Fig 2.Observation of geomagnetic condition

The sudden storm commencement (SSC) was registered at ~0445 UT and then there was a quick drop of the SYM-H index to the value of -226 nT, observed at ~2300 UT, with a couple of local minima of -93 and -164 nT at ~0940 and ~1740 UT respectively (Fig. 1). The planetary index of the geomagnetic activity Kp reached the maximum value of 8 after ~12 UT on 17 March 2015, qualifying it as a severe geomagnetic storm. During the main phase of the storm (17 March), the interplanetary magnetic field (IMF) orientation displayed a highly complex behavior. Three IMF components (top panels of Fig. 1) switched several times from positive to negative values and vice versa. Right after the shock arrival, the northward IMF Bz component reached the value of about 25 nT. At ~0530 UT the IMF Bz turned southward and reached the first minimal value of -18 nT at 0615 UT. Then the IMF Bz sharply turned northward and varied significantly between north and south during ~8 h. After ~1340 UT the Bz turned southward again and remained south till the end of this day. From ~06 till 11 UT, there are observed dominating positive Bx and negative By with peak values of 16.5 and -16.8 nT for Bx and By, respectively. During 11–15 UT with the new southward turning of Bz, the opposite situation with Bx/By domination occurred—Bx became negative with the minimal values of -14 nT while By component became positive with the peak of 30 nT. After 15 UT, IMF By turned sharply to negative values, reaching -8 nT, and then again to the positive ones with the new peak of 20 nT around 18 UT. Kamide and Kusano (2015) reported that this severe geomagnetic storm (G4 level) was a result from the superposition of two successive, moderate storms, driven by two successive, southward IMF structures. The intense geomagnetic storm on 17–18 March 2015 leads to the auroral particle precipitation and an enhancement of the substorm activity.

Response of the High latitude northern ionosphere

To understand the plasma density variations in the southern polar ionosphere during 17 March 2015, we examined the variation of TEC at the Arctic stations, is shown in Figures 3 respectively




Figure 3. TEC variation over Arctic Region ionosphere during 15-20, March 2015

Figure 3 is composed of four panel including Dst fluctuation during storm from 15 march to 20 march over arctic region. TEC data are plotted with time and latitude scale during the storm period, from the first panel showing the TEC fluctuation over Cambridge Bay, during the observation we notice the TEC is decreasing (<8 TECU) during the storm time and after 24 hrs it's taking regular pattern, whole period of storm there is very less TEC observed. Second panel of figure showing Eureka station, there is also TEC activity is decreasing (>10 TECU) but little bit high as compare with Cambridge way we can see clearly in graph. Penal third is showing Rabbit Lake, there is TEC activity is high (>12 TECU) as compere to both stations.

- Combine the two sub-headings i.e. Results and discussion.
- Present your results using the plots and discuss them comprehensively. Defend your results by citing works from other researchers.
- Try to compare your results with results from other recent similar studies i.e. results from same or other storms in other regions. Do they agree with yours or not? Give possible reasons.

Discussion

Above segment we depicted exhaustively the perceptions. In this part, we will examine point by point the outcomes all together toward decipher them. In Table 2, we have set up together the fundamental attributes of the tempest relying upon the longitudinal. There are a few actual systems for the arrangement of ionospheric anomalies in the polar ionosphere during TOI advancement, specifically Kelvin–Helmholtz and slope flow instabilities (e.g., van der Meeren et al. 2014). Sojka et al. (1998) examined one such component. For the March 17, 2015 tempest, the two elements of the quick unbiased and plasma streams, affirmed by the Millstone Hill ISR estimations along with outdoors plasma upgrade (got from Swarm LP and outdoors GPS TEC estimations), support that ideal conditions for GDI advancement caused the event of the plasma thickness inconsistencies in the outdoors ionosphere. Expanded amplitudes of TEC varieties with expanding coupling rate probably mirror an expanded power of vivacious molecule precipitation, expanded thickness of polar patches comparative with the ionospheric foundation thickness, and, as a rule, a more unique and tempestuous polar cap with expanded magnetospheric convergence of sun oriented breeze energy. Biggest sufficiency TEC varieties were seen across early afternoon at scopes of $74.0\text{--}78.0^\circ$ MLat. Huge postnoon amplitudes were noticed for lower coupling rates (<5000) and keeping in mind that prenoon and early afternoon amplitudes were biggest for higher coupling rates (>5000). These appropriations demonstrate expanded precipitation power in the postnoon area with expanded coupling rate and extraordinary precipitation around early afternoon and prenoon for high coupling rates >5000 . Nilsson et al. [1998] noticed E district ionization because of particle precipitation at areas planning to the low scope cusp, in spite of the fact that it is hazy whether expanded coupling rates would strengthen this precipitation. The attractive field that interfaces the high-latitude ionosphere to the magnetosphere permits the immediate passage of particles of sunlight based breeze and magnetospheric beginning into the polar ionosphere during calm/upset occasions. The power of these cycles improves during times of toward the south interplanetary attractive field (IMF) which is an important condition for the magnetopause disintegration (Aubry et al., 1970; Meng, 1970). Dayside reconnection brings about the reallocation of attractive motion and fortifying of the magnetospheric convection electric fields and field-aligned flows, prompting polar cap development and equatorward development of the auroral oval (Le et al., 2016). The high-latitude ionosphere is likewise influenced by the attractive substorms (Elphinstone et al., 1996)

CONCLUSION

17 March 2015 was considerably bigger in size when contrasted with different tempests and shifted in accordance with the force of the geomagnetic storm. The St. Patrick's Day tempest of 2015 was an exceptionally upset period with solid coupling of the SWMI framework during the whole principle period of the tempest. The presence of improved plasma thickness in the TOI

shaped in the southern polar cap on 17 March 2015 is a consequence of the mind boggling coupling of the sunlight based wind-magnetosphere-ionosphere framework through electric fields and nonpartisan breezes for long lengths. The solid and supported magnetopause disintegration on 17 March 2015 prompted the pervasiveness of more grounded storm time electric fields beginning from the early principle period of the tempest. This examination shows that the length and degree of magnetopause disintegration assume a significant part in the spatiotemporal development of the plasma thickness dissemination in the high-multiplicity ionosphere. This investigation features the way that the conduct of the polar ionosphere is unequivocally affected by the outside drivers of geomagnetic storms and gives understanding into the idea of association of the sun oriented breeze with the earthly IT framework during the St. Patrick's Day of 2015. In this paper, ionospheric reactions to the geomagnetic storms on 17 March 2015 are explored by utilizing the GPS perceptions. Our principle discoveries are summed up as follows.

Total electron content fluctuation was very low observed over Cambridge Bay during storm time, 15 March to 20 March 2015 over arctic region and after 24 hrs it's taking regular pattern. Over Eureka station, there is also TEC activity is decreasing but little bit high as compare with Cambridge way. For the Rabbit Lake station, there is TEC activity is high as compare to both stations during the storm period.

- **CONCLUSION:** Give the overall findings if your study.
- **ACKNOWLEDGEMENTS:** Include acknowledgements section after the conclusion section.
- **REFERENCES:** Include recent references i.e. from 2020, 2021, 2022, 2023 and 2024 to make research unique and more conclusive.
- Work on the flow of the whole paper and read through it keenly to remove the grammatical errors in the manuscript.

Reference

van der Meeren C, Oksavik K, Lorentzen D, Moen JI, Romano V (2014) GPS scintillation and irregularities at the front of an ionization tongue in the nightside polar ionosphere. *J Geophys Res Space Phys* 119:8624–8636. doi:10.1002/2014JA020114

Aubry, M. P., Russel, C. T., & Kivelson, M. G. (1970). Inward motion of the magnetopause before a substorm. *Journal of Geophysical Research*, 75, 7018–7031. <https://doi.org/10.1029/JA075i034p07018>.

Buonsanto, M. J. (1999). Ionospheric storms-Review. *Space Science Reviews*, 88, 563–601.

Elphinstone, R. D., Murphree, J. S., & Cogger, L. L. (1996). What is a global auroral substorm? *Reviews of Geophysics*, 34, 169–232.

Gonzalez, W. D., Joselyn, J. A., Kamide, Y., Kroehl, H. W., Rostoker, G., Tsurutani, B. T., & Vasyliunas, V. M. (1994). What is a geomagnetic storm. *Journal of Geophysical Research*, 99(A4), 5771–5792.

Le, G., Luhr, H., Anderson, B. J., Strangeway, R. J., Russell, C. T., Singer, H., et al. (2016). Magnetopause erosion during the 17 March 2015 magnetic storm: Combined field-aligned currents, auroral oval, and magnetopause observations. *Geophysical Research Letters*, 43, 2396–2404. <https://doi.org/10.1002/2016GL068257>.

Meng, C. I. (1970). Variation of the magnetopause position with substorm activity. *Journal of Geophysical Research*, 75, 3252.

Pröls, G. W. (1995). Ionospheric F-region storms. In H. Volland (Ed.), *Handbook of atmospheric electrodynamics* (Vol. 2, pp. 195–248).

Mendillo, M., 2006. Storms in the ionosphere: patterns and processes for total electron content. *Rev. Geophys.* 44, RG4001. <https://doi.org/10.1029/2005RG000193>.

Fuller-Rowell, T.J., Codrescu, M.V., Rishbeth, H., Moffett, R.J., Quegan, S., 1996. On the seasonal response of the thermosphere and ionosphere to geomagnetic storms. *J. Geophys. Res.* 101, 2343–2353.

Bilitza, D., and B. W. Reinisch (2008), *International Reference Ionosphere 2007: Improvements and new parameters*, *Adv. Space Res.*, 42, 599–609, doi:10.1016/j.asr.2007.07.048.

Jayachandran, P. T., et al. (2009), Canadian High Arctic Ionospheric Network (CHAIN), *Radio Sci.*, 44, RS0A03, doi:10.1029/2008RS004046.

Jayachandran, P. T., C. Watson, I. J. Rae, J. W. MacDougall, D. W. Danskin, R. Chadwick, T. D. Kelly, P. Prikryl, K. Meziane, and K. Shiokawa (2011), High-latitude GPS TEC changes associated with a sudden magnetospheric compression, *Geophys. Res. Lett.*, 38, L23104, doi:10.1029/2011GL050041.

Jayachandran, P. T., K. Hosokawa, K. Shiokawa, Y. Otsuka, C. Watson, S. C. Mushini, J. W. MacDougall, P. Prikryl, R. Chadwick, and T. D. Kelly (2012), GPS total electron content variations associated with poleward moving Sun-aligned arcs, *J. Geophys. Res.*, 117, A05310, doi:10.1029/2011JA017423.

Nilsson, H., S. Kirkwood, and T. Moretto (1998), Incoherent scatter radar observations of the cusp acceleration region and cusp field-aligned currents, *J. Geophys. Res.*, 103, 26,721–26,730, doi:10.1029/98JA02269.

Prikryl, P., P. T. Jayachandran, R. Chadwick, and T. D. Kelly (2015), Climatology of GPS phase scintillation at northern high latitudes for the period from 2008 to 2013, *Ann. Geophys.*, 33, 531–545, doi:10.5194/angeo-33-531-2015.

Burns, A. G., T. L. Killeen, W. Deng, G. R. Carignan, and R. G. Roble (1995), Geomagnetic storm effects in the low-latitude to middle-latitude upper thermosphere, *J. Geophys. Res.*, 100, 14,673–14,691, doi:10.1029/94JA03232.

Richmond, A. D., and G. Lu (2000), Upper-atmospheric effects of magnetic storms: A brief tutorial, *J. Atmos. Sol. Terr. Phys.*, 62, 1115–1127, doi:10.1016/S1364-6826(00)00094-8.

Prölss, G. W. (1995), Ionospheric F-region storms, in *Handbook of Atmospheric Electrodynamics*, edited by H. Volland, pp. 195–248, CRC Press, Boca Raton, Fla.

UNDER PEER REVIEW