

Original Research Article

SCREENING INDICATIONS, SOCIO-DEMOGRAPHIC, VIROLOGIC AND RADIOLOGIC CHARACTERISTICS OF VIRAL HEPATITIS B PATIENTS IN A HIGHLY ENDEMIC POPULATION IN SOUTHERN NIGERIA

Abstract

This study aims to identify indications for HBV screening among patients visiting a tertiary outpatient clinic, virologic characteristics as well as the sociodemographic variations of this populace. There are multiple barriers/challenges to screening and access to care for Hepatitis B in Nigeria and these screening challenges include inadequate National strategies/ guidelines on viral hepatitis infection in Nigeria especially a lack of screening programs such as the PROLIFICA program in Gambia. This was a cross-sectional, descriptive, observational study conducted by reviewing the database of patients with viral hepatitis B seen from July 2017 to January 2019. Radiological assessment was carried out on all patients irrespective of their group as well as laboratory evaluation. The study observed a varied array of reasons for HBV screening ranging from non-health issues which were tagged as incidental indications to hepatic and non-hepatic related diseases. HBV prevention and care should be integrated into all health sectors be it public or private. Blood for donation should be screened with the triple panel screening kit rather than the regular RDTs and ELISA tests as these have a low sensitivity and are unable to screen HBV mutant strains.

Keywords: HBV screening, ELISA tests, HBV prevention, Hepatitis B

Comment [P1]: Please mention the names of the clinics from which the samples were collected) Rivers State University Teaching Hospital) or replace the tertiary with specialized outpatient clinic.

Comment [P2]: Delete

Comment [P3]: Hepatitis B virus

Comment [P4]: Hepatitis B virus

Introduction

Viral hepatitis B, a potentially life-threatening and global dilemma with a disparate epidemiology, is a highly endemic infection in Africa affecting approximately 81 million people with viral hepatitis-related liver cirrhosis as one of the leading causes of mortality in Africa.(1,2) The implication of chronic hepatitis B (HBV) infection is progressively being acknowledged with a 2019 WHO report of over 820,000 annual deaths and 1.5million new infections yearly.(1) Furthermore, Lozano et al submitted that HBV now ranks 15th among all causes of human mortality. (3)

Despite the high burden of childhood HBV infection in Africa, horizontal transmission remains the major route of HBV transmission (4) compared with the perinatal route of transmission and in adults, male sex is a major risk factor for chronic HBV infection. (5)

In Sub-Saharan Africa, HBV genotypes E, A and D respectively are the dominant genotypes(6) resulting in a predominance of HBeAg-negative hepatitis B infection. (7) Thus, HBeAg-negative/anti-HBe-positive chronic hepatitis B (CHB) are 7-9 times more common in Africa due to the molecular characteristics of the prevailing HBV genotype. (7)Also, it has been observed that patients infected with sub genotype A1 which is prevalent in Africademonstrate an earlier loss of HBeAg and seroconversion to anti-HBe during the natural course of HBV infection compared to those infected with sub genotype A2. (7)

The 2023 CDC guideline for screening and testing for Hepatitis B virus infection recommends screening for all adults aged 18 years and above at least once in a lifetime using the triple panel test which includes the Hepatitis B surface antigen, antibody to Hepatitis B surface antigen and antibody to core antigen.(8) The CDC also recommends testing for everyone born in regions with HBV infection prevalence of $\geq 2\%$ which includes Nigeria and most Sub-Saharan African countries.(9)A 2021 meta-analysis reported a pooled prevalence of 9.5% for Hepatitis B in Nigeria (10) with the North-Western region having the highest prevalence of 12.1%. this places Nigeria in a region with high endemicity (>8%) for HBV using the WHO criteria. (11)

Comment [P5]: Replace vertical transmission ,

Comment [P6]: How did the researcher determine that?

Comment [P7]: Write Centers for Disease Control and Prevention (CDC)

The Nigerian national guideline for viral Hepatitis in Nigeria (12) published in 2016 recommends routine viral hepatitis screening for high-risk populations such as female sex workers, male sex workers, healthcare workers, and PWID to mention but a few. There, however, were no screening strategies defined for the general population despite an estimated 19million Nigerians living with HBV or HCV, and a high prevalence of 11-13% and 2.1% for HBV and HCV respectively. (13)

Moreover, there are multiple barriers/challenges to screening and access to care for Hepatitis B in Nigeria and these screening challenges include inadequate National strategies/ guidelines on viral hepatitis infection in Nigeria especially a lack of screening programs such as the PROLIFICA program in Gambia(14) that will access in-depth level of HBV screening and awareness in both urban and rural centers. Additional challenges comprise Poor funding for Viral Hepatitis program, lack of prioritisation of viral Hepatitis infection by donor agencies in Nigeria, poor impact of ongoing screening strategies for viral Hepatitis, inadequate supply of test kits, limitation of testing to urban regions of the country, and underdiagnosis of viral Hepatitis due to the use of testing kits that are low in sensitivity and specificity.

Comment [P8]: delete

Olakunde et al (15) in their study on Barriers to hepatitis B virus screening of pregnant women in primary healthcare centers in Nigeria identified that "perceived barriers exist at patient, provider, and health system levels and they include the lack of test kits, unaffordability of HBV test, shortage of trained personnel, poor awareness and knowledge of HBV, and unavailability of treatment and prevention interventions for HBV."

Comment [P9]: What proves the truth of this statement? Are there statements by the ministry or the government? Or a previous source

Comment [P10]: When reading the conclusions of the researcher referred to, I did not find these details. Please, please adhere to the researcher's conclusions

There are also identifiable barriers to access to treatment currently in Nigeria which may consist of the cost of DNA assays, absence of easy, cheap and noninvasive screening tools for assessment of liver diseases, lack of widespread educational programs on the prevention and treatment of HBV, and restriction of the free generic Tenofovir to HIV-infected patients only thereby discriminating against HBV monoinfected individuals. Furthermore, there is no national access program nor subsidized care for HBV-infected persons and the cost of treatment is borne individually which impacts and impoverishes the patients as reported in a recently published study in southern Nigeria. (16)

Comment [P11]: Reference? who said that ?

Lastly, another noticeable barrier is the HBV knowledge gap among healthcare workers and the general population (15,17) which may affect the demand and uptake of HBV screening test as well as its treatment. This study aims to identify indications for HBV screening among patients visiting a tertiary outpatient clinic, virologic characteristics as well as the sociodemographic variations of this populace.

Comment [P12]: What do you mean?
Replace (Rivers State University Teaching Hospital)

Methodology

Rivers State University Teaching Hospital is a 500-bed specialist tertiary health center with 2 medical schools and residency training programs that serves Rivers State and neighboring states in the south-south region of Nigeria.

Comment [P13]: Delete this part , not necessary

This was a cross-sectional, descriptive, observational study conducted by reviewing the database of patients with viral hepatitis B seen from July 2017 to January 2019.

Comment [P14]: Added the place (in Rivers State University Teaching Hospital)

Ethical approval was obtained from the hospital ethical committee with approval number RSUTH/REC/2022213.

Inclusion criteria: Data of Adult HbsAg positive patients (>18 years) who remained in treatment for more than 6 months in the outpatient Gastroenterology clinic was obtained from the viral hepatitis B register and included in the study.

Exclusion criteria: Co-infected patients (Hepatitis C and HIV) and patients who have a recent history of vaccination within the past 30 days were excluded from the study.

Classification of the study population

Patients were classified into two main groups according to the initial HBeAg status: HBeAg positive (HBeAg+) and negative (HBeAg-) groups. Radiological assessment was carried out on all patients irrespective of their group as well as laboratory evaluation. The limits of normal /detection were dependent on the laboratory kit available during the study period and were established by our laboratory.

Comment [P15]: How was it classified? What is the approved method?

Hepatitis B viral DNA obtained at first visit (initial HBV DNA) also known as Viral load is classified into 3 depending on its levels- undetectable, low and high.

Comment [P16]: How was the presence of DNA determined?!

Undetectable level is HBV DNA ≤ 10 iu/ml as determined by the manufacturers of COBAS® TaqMan HBV test (Roche Diagnostics).(18)

Comment [P17]: Please, mention the method or device?

Low levelsof HBV DNA was adopted as PCR quantification levels < 2000 iu/ml and high levelsas HBV DNA ≥ 2000 iu/ml. (12)

Data analysis

Data was entered into Microsoft Excel and was exported and analyzed using Statistical Package for Social Sciences (SPSS) version 26. Categorical variables were reported as frequencies and percentages and compared with the Chi-square test. Continuous variables were reported as means \pm SD, and the student t-test was used to compare means. A p-value of ≤ 0.05 was considered to be statistically significant.

Results

Data of Four hundred and fifty-four (454) hepatitis B positive patients with complete data and no contraindication were extracted from the Viral hepatitis register comprising 198(43.8%) females and 255(56.2%) males. The mean age of the respondents was 38.33 ± 11.16 years with a range of 18 to 76 years with a predominance of young people (N=355, 78.2%) table 1. Notably, over 50% of our study population are in a legally and socially sanctioned union we however did not explore the type of union (table 2).

Comment [P18]: The total of females and males is 453, while you are writing 454. please edit.

Comment [P19]: Must added to table below.

Comment [P20]: Pleases ad

Table 1: Age distribution of the study population

Age group	Frequency (%)
Young (< X)	355(78.2)
Middle aged (X – Z)	84(18.5)
Elderly (> Z)	15(3.3)

Comment [P21]: What do the symbols mean?

Comment [P22]: What do the symbols X and Z mean?


Table 2: marital status of the study population

Marital status	Frequency (%)
Single	124(27.3)
Married	314(69.2)
Divorced	5(1.1)
Widowed	11(2.4)

Geopolitical variation of the study population

Geopolitically, Nigeria consists of 6 regions as shown in Picture 1. All regions of the country are represented in the study population showing the cosmopolitan nature of Rivers State as our tertiary Hospital serves the south-south region of Nigeria.


The South-South region being the host, has the highest number at 220 patients while the North-East region has the lowest representation. The distribution of the study population by state is shown in table 3.



Picture 1: Geopolitical distribution of the study population

Table 3: Distribution of the study population by State and Nationality.

State	Frequency	Percent
Abia	71	15.6
Akwa Ibom	12	2.6
Anambra	14	3.1
Bauchi	3	0.7
Bayelsa	8	1.8
Benue	2	0.4
Cross River	4	0.9
Delta	18	4.0
Ebonyi	4	0.9
Edo	20	4.4
Ekiti	6	1.3
Enugu	12	2.6
Imo	61	13.4
Jigawa	2	0.4
Kaduna	4	0.9
Kogi	2	0.4
Kwara	9	2.0
Niger	3	0.7
Ogun	4	0.9
Ondo	7	1.5
Osun	15	3.3
Oyo	2	0.4
Plateau	11	2.4
Rivers	158	34.8
Other countries -Ghana	2	0.4
Total	454	100.0



Picture 2: map of Nigeria showing the distribution of the state of origin of the study population

Indications for screening for viral Hepatitis B

The indications for screening are shown in table 4. Incidental finding which makes up 64% of the indications include pre-employment screening, pre-marriage screening, screening before blood donation, pre-surgery screening for non-Hepatic related diagnosis and public health awareness campaigns by non-governmental organizations and individuals. 51(11.3%) of the study population presented late with suspected Liver disease and 20 (4.4%) were diagnosed from the compulsory antenatal screening done in our institution.

Comment [P23]: Must added to the table below

Indication	Frequency (%)
Abdominal pain	58(12.8)
Abdominal swelling	2(0.4)
Antenatal screening	20(4.4)
Febrile illness	18(4.0)
Malaise	2(0.4)
Incidental finding	291(64.0)
Suspected liver disease	51(11.3)
Others	12(2.6)

Comment [P24]: It was not mentioned in the above section?

Table 4: Indications for screening for Hepatitis B

Comment [P25]: above the table please

Laboratory (virologic) parameters

We observed a predominance of HBeAg negative category compared to HBeAg positive {430(94.7%) vs 24(5.3%)} in our cohort while the HBV DNA (Viral load) ranged from <20 to 170500000iu/ml with a mean value of 4509723.97 ± 26264722.07 and a median value of 1130.50.

Comment [P26]: Why wasn't it put in a table? The results are not clear. The results do not match the tables !

Increased AST levels were observed in 256(56.4%) persons while ALT was increased in 193(42.5%) persons. Comparing the mean of AST to ALT, the mean of the AST levels was notably higher than that of the ALT though their ranges were comparable. (AST mean 53.905±66.10, range 4.0 to 544.0: ALT mean 42.88±49.57, range 3.0 to 544.0).

Comment [P27]: He did not mention in methods

Comment [P28]: also

Table 5. HBV DNA levels and E antigen comparison

	Undetectable	Low	High
	<i>HBV DNA</i>		
Number	38	224	192
Frequency %	8.4	49.3	42.3
	<i>E antigen comparison</i>		
E antigen negative	35	207	188
E antigen positive	3	17	4
Total	38	224	192

$\chi^2=6.823$, $p=0.033$


low HBV DNA= <2000iu/ml, high HBV DNA= ≥2000iu/ml, undetectable HBV DNA = < 20iu/ml

Comment [P29]: Why were the results of this table not presented in detail above? What is the purpose of the table then?

Radiological features of the study population

An abdominal ultrasound scan was done routinely at first visit and the results are shown in picture 3. While more than half of the study population had normal findings, we detected hepatobiliary disorders in a significant number of the patients ranging from Hepatic steatosis 42(9.3%), Hepatomegaly 41(9.0%), Chronic liver disease 39(8.6%), Cholelithiasis 11(2.4%), Chronic hepatitis 2(0.4%) and Primary Liver cell carcinoma in 8 (1.8%). Other reported findings on abdominal USS include bilateral renal parenchymal disease, Hydronephrosis, isolated Splenomegaly and Cystitis and these were described in 2.6% (12) of the study population.

Comment [P30]: It is preferable to put the results in a table



Picture 3: Abdominal ultrasound scan findings. PLCC= Primary Liver cell carcinoma. CLD= Chronic Liver Disease.

UNDER PEER REVIEW

Discussions

This study aimed at an in-depth analysis of the indications for screening, the sociodemographic, virologic and radiologic characteristics of HBV population.

The study revealed a gender disparity in the prevalence of hepatitis B, with more males being affected than females among adults. This observation aligns with previous studies that have also reported a higher prevalence of hepatitis B among males compared to females. (19,20) Multiple studies have also reported a male predominance in the distribution of hepatitis B infection especially in the adult population. (19–22) A Zambian study reported a male-to-female carrier state of 14.6% to 4.6% for HbsAg and 82.6% of the male population were positive for at least one marker compared to 69.4% of females. (23) The mean age of the study population was 38.33 years with >78% being young adults (table 1) similar to figures reported by other African investigators. (24,25)

Several factors could contribute to this gender disparity. One possible explanation is differences in behaviors and risk factors between males and females. For example, males may be more likely to engage in high-risk behaviors such as intravenous drug use or unprotected sex, which can increase the risk of hepatitis B transmission. Additionally, cultural and societal factors may also play a role, influencing healthcare-seeking behaviors and access to screening and vaccination services.

Comment [P31]: Reference ?

Understanding these gender disparities is crucial for developing targeted prevention and intervention strategies to reduce the burden of hepatitis B, particularly among males who are disproportionately affected. Further research is needed to explore the underlying reasons for this gender disparity and to develop effective public health interventions aimed at reducing the prevalence of hepatitis B among males.

Geopolitically, all six geopolitical zones of the country and >2/3rd of the 36 states of Nigeria are represented in this study demonstrating the metropolitan nature of the study site. The higher representation from the south-south zone can be explained by the high population of Locals as well as persons from nearby states residing in Rivers state. This picture is expectedly in contrast with the 2021 review and meta-analysis on HBV in Nigeria that recorded the North-West zone as having the highest prevalence of viral Hepatitis B (26) due to multiple factors including

Comment [P32]: Can you replace

low immunization coverage, geographical isolation, and birthing outside healthcare facilities.

The study observed a varied array of reasons for HBV screening ranging from non-health issues which were tagged as incidental indications to hepatic and non-hepatic related diseases. Sixty-four percent (64%) of the study population were diagnosed with viral hepatitis B incidentally while a considerable percentage were diagnosed with HBV during an illness with $\geq 11\%$ exhibiting clinical features of liver disease already. Persons who presented to the clinic for pre-employment medical examination, pre-marriage screening, screening before blood donation, pre-surgery screening for non-Hepatic related diagnosis and public health awareness campaigns by non-governmental organizations, were screened for viral hepatitis B with some turning out positive. This finding underscores the importance of routine screening among this population. This strengthens the need for the prioritization of viral hepatitis B by the government and aid agencies and the need for the development and implementation of an effective public health strategy towards the 2030 WHO elimination goals for viral Hepatitis. (26,27)

Assessment of the characteristics of the Hepatitis B virus in the study population revealed that 94.7% were HBeAg negative compared to HBeAg positive with a few numbers (n=38) of the patients having undetectable HBV DNA. Furthermore, among the 194 patients with high HBV DNA levels, 97.9% (n= 188) were HBeAg negative ($p= 0.033$). Studies have however shown that HBeAg negative chronic HBV is the most predominant form of the disease worldwide (27,28,29) and HBeAg-negative/anti-HBe-positive chronic hepatitis B (CHB) are more common in Africa due to the molecular characteristics of the prevailing HBV genotype. (7)

Notably, 42.5% of the patients have elevated ALT levels indicating some degree of necroinflammation. ALT could provide a sensitive alternative to liver biopsy in the assessment of hepatic inflammation and a cheaper alternative to HBV DNA is lacking hence determining when to initiate therapy for chronic HBV patients.

Limitations

Comment [P33]: This must be put in the result part not here .

This study has some limitations. It is a single-center hospital-based study and thus may not capture the enormity of the burden of chronic hepatitis B in the general population in Nigeria.

Recommendations

Despite the high prevalence of viral hepatitis B in Nigeria, we are yet to have a comprehensive and effective care package for patients living with viral Hepatitis. Thus, the recommendation to the federal government of Nigeria and other relevant bodies is to make HBV a priority by improving screening and increasing access to care. There is a need for a global and locally domiciled plan for HBV care including education, population-based surveys as against sporadic hospital/healthcare facility-based reports and unitary, unabridged and full funding for HBV care in Nigeria.

Pre-employment medical examination, pre-marital screening should be made compulsory by law. This will improve the screening yield of viral hepatitis B infection.

The relevant health authorities need to adopt 'point of care' (POC) tests for HbsAg and HCV antibodies at the community level as well as POC for cheaper and easy HBV DNA quantification as a means of treatment evaluation.

Other important measures include improving and implementing the test-and-treat strategy for HBV-infected pregnant women, and improving access to birth dose immunoglobulin and neonatal vaccination.

Furthermore, HBV prevention and care should be integrated into all health sectors be it public or private. Blood for donation should be screened with the triple panel screening kit rather than the regular RDTs and ELISA tests as these have a low sensitivity and are unable to screen HBV mutant strains.

References

1. 91 million Africans infected with Hepatitis B or C | WHO | Regional Office for Africa [Internet]. [cited 2023 Feb 12]. Available from: <https://www.afro.who.int/news/91-million-africans-infected-hepatitis-b-or-c>
2. UNAIDS_2021_FactSheet_en_em.pdf [Internet]. [cited 2022 Feb 26]. Available from: https://embargo.unaids.org/static/files/uploaded_files/UNAIDS_2021_FactSheet_en_em.pdf
3. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet Lond Engl*. 2012 Dec 15;380(9859):2095–128.
4. Kiire CF. The epidemiology and prophylaxis of hepatitis B in sub-Saharan Africa: a view from tropical and subtropical Africa. *Gut*. 1996;38(Suppl 2):S5-12.
5. Ruggieri A, Gagliardi MC, Anticoli S. Sex-Dependent Outcome of Hepatitis B and C Viruses Infections: Synergy of Sex Hormones and Immune Responses? *Front Immunol*. 2018 Oct 8;9:2302.
6. Ahmad AE, Bakari AG, Musa BOP, Mustapha SK, Jamoh BY, Abdullahi IN, et al. Pattern of Prevalent Hepatitis B Virus Genotypes in Zaria, Nigeria. *Niger Postgrad Med J*. 2019;26(2):80–6.
7. Hadziyannis SJ. Natural history of chronic hepatitis B in Euro-Mediterranean and African countries. *J Hepatol*. 2011 Jul;55(1):183-91. doi: 10.1016/j.jhep.2010.12.030. Epub 2011 Jan 14. Erratum in: *J Hepatol*. 2013 May;58(5):1062. PMID: 21238520.
8. Screening and Testing Recommendations for Chronic Hepatitis B Virus Infection (HBV) | CDC [Internet]. 2023 [cited 2023 Jun 30]. Available from: <https://www.cdc.gov/hepatitis/hbv/testingchronic.htm>
9. Connors EE. Screening and Testing for Hepatitis B Virus Infection: CDC Recommendations — United States, 2023. *MMWR Recomm Rep* [Internet].

2023 [cited 2023 Jun 30];72. Available from: <https://www.cdc.gov/mmwr/volumes/72/rr/rr7201a1.htm>

10. Ajuwon BI, Yujuico I, Roper K, Richardson A, Sheel M, Lidbury BA. Hepatitis B virus infection in Nigeria: a systematic review and meta-analysis of data published between 2010 and 2019. *BMC Infect Dis.* 2021 Oct 30;21(1):1120.
11. Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2010 (GBD 2010) Results by Risk Factor 1990-2010 [Internet]. Institute for Health Metrics and Evaluation (IHME); 2012 [cited 2023 Jun 30]. Available from: <http://ghdx.healthdata.org/record/ihme-data/gbd-2010-results-risk-factor-1990-2010>
12. Nigeria-Hepatitis-Guidelines-TX-guidelines.pdf [Internet]. [cited 2022 Feb 19]. Available from: <https://www.hepb.org/assets/Uploads/Nigeria-Hepatitis-Guidelines-TX-guidelines.pdf>
13. Odukoya O, Odeyemi K, Odubanjo O, Isikekpei B, Igwilo U, Disu Y, et al. Hepatitis B and C seroprevalence among residents in Lagos State, Nigeria: A population-based survey [Internet]. Vol. 29, *Nigerian Postgraduate Medical Journal.* 2022. p. 75–81. Available from: <https://www.npmj.org/article.asp?issn=1117-1936;year=2022;volume=29;issue=2;spage=75;epage=81;aualast=Odukoya>
14. Howell J, Ladep NG, Nayagam S, Lemoine M, Garside DA, Crossey MME, et al. PROLIFICA: a story of West African clinical and research collaborations to target hepatitis B-related hepatocellular carcinoma in West Africa. *QJM Int J Med.* 2016 Jun;109(6):373–5.
15. Olakunde BO, Adeyinka DA, Olakunde OA, Raji HB, Yahaya HB, Ijaodola OA, et al. Barriers to hepatitis B virus screening of pregnant women in primary healthcare centers in Nigeria: health workers' perspective. *BMC Prim Care.* 2023 Oct 17;24(1):209.
16. Alali A, Osaro B, Jaja I, Abere S, Fajola A. Catastrophic Health Expenditure and Impoverishment Amongst Hepatitis B Patients in a Resource-Poor Setting. *Asian J Med Health.* 2024 Feb 5;22(2):23–32.

17. Abere S, Oyan B, Alali DJ, Omunakwe H, Ejikem M. Viral Hepatitis B and C: Knowledge Gaps and Patterns of Preventive Practices Among Medical Doctors in Rivers State, Nigeria. *Cureus*. 2022 Mar;14(3):e22928.
18. Belopolskaya M, Avrutin V, Firsov S, Yakovlev A. HBsAg level and hepatitis B viral load correlation with focus on pregnancy. *Ann Gastroenterol Q Publ Hell Soc Gastroenterol*. 2015;28(3):379–84.
19. Baig S. Gender disparity in infections of Hepatitis B virus. *J Coll Physicians Surg-Pak JCPSP*. 2009 Sep;19(9):598–600.
20. Tsega E, Mengesha B, Hansson BG, Lindberg J, Nordenfelt E. Hepatitis A, B, and delta infection in Ethiopia: a serologic survey with demographic data. *Am J Epidemiol*. 1986 Feb;123(2):344–51.
21. Forbi JC, Iperepolu OH, Zungwe T, Agwale SM. Prevalence of Hepatitis B e Antigen in Chronic HBV Carriers in North-central Nigeria. *J Health Popul Nutr*. 2012;30(4):377–82.
22. Bowry TR. The pattern of transmission of hepatitis B virus in an urban population of Kenya. *East Afr Med J*. 1984 May;61(5):385–92.
23. Tabor E, Bayley AC, Cairns J, Pelleu L, Gerety RJ. Horizontal transmission of hepatitis B virus among children and adults in five rural villages in Zambia. *J Med Virol*. 1985 Feb;15(2):113–20.
24. Hamza M, Samaila AA, Yakasai AM, Babashani M, Borodo MM, Habib AG. Prevalence of hepatitis B and C virus infections among HIV-infected patients in a tertiary hospital in North-Western Nigeria. *Nigerian Journal of Basic and Clinical Sciences*. 2013 Jul 1;10(2):76-81.
25. Kolou M, Katawa G, Salou M, Gozo-Akakpo KS, Dossim S, Kwarteng A, Prince-David M. High Prevalence of Hepatitis B Virus Infection in the Age Range of 20-39 Years Old Individuals in Lome. *Open Virol J*. 2017 Jan 12;11:1-7
26. Ajuwon, B.I., Yujuico, I., Roper, K. et al. Hepatitis B virus infection in Nigeria: a systematic review and meta-analysis of data published between 2010 and 2019. *BMC Infect Dis* 21, 1120 (2021).

27. World Health Organisation. Hepatitis B fact sheet. 2019. <https://www.who.int/newsroom/factsheets/detail/hepatitis-b>. accessed 16 April 2024
28. Forbi JC, Iperepolu OH, Zungwe T, Agwale SM. Prevalence of hepatitis B e antigen in chronic HBV carriers in North-central Nigeria. *J Health Popul Nutr*. 2012 Dec;30(4):377-82
29. Bonino F, Colombatto P, Brunetto MR. HBeAg-Negative/Anti-HBe-Positive Chronic Hepatitis B: A 40-Year-Old History. *Viruses*. 2022; 14(8):1691

UNDER PEER REVIEW