

**Influence of Plant Growth Regulators on Yield and Qualitative Parameters in Strawberry (*Fragaria x anannasa*) cv. Winter Dawn under Protected Cultivation**

**ABSTRACT:** The present investigation was carried out during 2021-2022 to find out the effect of plant growth regulators on yield and qualitative parameters in strawberry (*Fragaria x anannasa*) cv. Winter Dawn under protected cultivation at Precision Farming Development Centre (PFDC), Department of Horticulture, Chaudhary Charan Singh Haryana Agricultural University, Hisar. The experiment was conducted with ten treatments using randomized block design. Each treatment had three replication. The treatment consisted of three different concentration of plant growth regulators viz. T<sub>1</sub>: NAA 25 ppm, T<sub>2</sub>: NAA 50 ppm, T<sub>3</sub>: NAA 75 ppm, T<sub>4</sub>: GA<sub>3</sub> 25 ppm, T<sub>5</sub>: GA<sub>3</sub> 50 ppm, T<sub>6</sub>: GA<sub>3</sub> 75 ppm, T<sub>7</sub>: CCC 500 ppm, T<sub>8</sub>: CCC 750 ppm, T<sub>9</sub>: CCC 1000 ppm and T<sub>10</sub>: Control. The yield and quality of strawberry were significantly influenced by plant growth regulators. The highest quality of fruit in term of berry weight (16.57g), berry breadth (2.87cm), berry length (3.97cm), TSS (9.38°Brix), TSS/acid ratio (14.89) and ascorbic acid (60.13mg/100g) were observed in T<sub>9</sub> treatment which was consist cycocel@1000 ppm while GA<sub>3</sub>@75 ppm gave maximum results in terms of marketable yield (88.27%), yield per plant (304.26g) and anthocyanin (5.67mg/100g) in strawberry fruit.

**Key words:** Plant growth regulators, quality, strawberry, Winter Dawn, yield.

**INTRODUCTION:**

“The cultivated strawberry (*Fragaria x ananassa* Duch.) is one of the luscious and soft fruits of the world. It is a hybrid of two Native American species *Fragaria chilonensis* and *Fragaria virginiana*. *Fragaria* species belongs to the family Rosaceae, with basic chromosome number  $x=7$ ” (Choudhary *et al.*, 2024). “All cultivated varieties are octoploid ( $2n=56$ ). Botanically it is an aggregate fruit which is highly perishable in nature. Strawberry usually grows in temperate zones, and it is classified as a short-day plant based on its behaviour and eco system” (Gautam and Mishra, 2024). “In temperate climate condition, its plants behave like a small perennial herb with shallow root system whereas in sub-tropical climate it behaves as annuals. It has short stem known as crown. The crown produces leaves at very close interval along the stem axis and flowers at terminal position on stem axis. The edible portion of strawberry includes the ripened receptacle and achenes (true fruit and seed). Strawberry is usually propagated through runners. Strawberry is rich source of vitamins and minerals and coupled with delicate flavor, the red color of the fruit is mainly due to the presence of the anthocyanin, pelargonidin, 3-monoglucoside and traces of cyaniding” (Srivastav, A., *et al.*, 2018).

It is commercially grown in Maharashtra, Punjab, Haryana, Karnataka, Himachal Pradesh, Madhya Pradesh, Jammu & Kashmir and Uttarakhand in India (Thakur *et al.*, 2017). Strawberry cultivation area in India is nearly 1000 ha, with an annual production of 20,000 MT (pib.govt.in, 2020-21). Strawberry covers approximately 200 acres of land in Haryana (Hortharyana 2018-19), and the Hisar district has emerged as a hub. Strawberry cultivation is gaining popularity throughout India, with Haryana leading the way in terms of both area and production. “However, its marketing and profit are suffering as a result of a lack of desired fruit size and quality. Application of growth regulators has been practiced commercially to increase the production and quality of crops. Plant hormones are a small group of signaling molecules that plays a vital role in various morphological,

photosynthetic, biochemical and developmental processes in crop plants” (Nazir, *et. al.*, 2019). The growth and quality of fruits depends on different attributes which are closely associated with nutrient uptake by the plant and also with PGR’s. **Plant growth regulators have become very important in growing larger and better crops.**

GA<sub>3</sub> has been found to increase the vegetative growth of strawberry; Cycocel and NAA improved the yield and quality of strawberry. Although, the use of Cycocel (CCC) was reported to reduce plant height by which fruit yield is positively increased (Tiwari, A.K., *et.al.* 2017). Use of GA<sub>3</sub> in strawberry has been reported in early flowering, increased duration of flowering, harvesting and yield. It increases yield and quality of fruits, helps in cell elongation and cell enlargement, increases vegetative growth and minimizes time of maturity and increases fruit set. Application of NAA increases fruit size and delays ripening and increases anthocyanin accumulation in strawberry fruits. It also increases duration of flowering, improves yield and quality of fruits (Yadav, I., *et.al.* 2017). **Yield and quality are both pursued aiming at food security as well as better life.** With the goal of increasing strawberry production, marketability, and profit margins in mind, the current study was designed to determine the optimal concentration of plant growth regulators and their effect on yield and quality of strawberry.

## **MATERIALS AND METHODS:**

The present study was carried out at Precision Farming Development Centre (PFDC), Department of Horticulture, Chaudhary Charan Singh Haryana Agricultural University (CCS HAU), Hisar during 2021-2022. Hisar is located at an altitude of 215m (705ft) above MSL with coordinates of 29° 09'N latitude and 75° 42'E longitudes in western Haryana. This region is part of alluvial Ghaggar – Yamuna

plain and its southern and western portion mark a gradual transition to the desert. An experiment was conducted using foliar spray of PGRs viz. Naphthalene Acetic acid (NAA), Gibberellic acid (GA<sub>3</sub>) and Cycocel (CCC) with different concentration (Table 1) where planting material (runners) of Winter Dawn variety were obtained from Saharwa village of Hisar district (Haryana). The strawberry runners with uniform size were planted in month of October at a distance of 25 x 30 cm on raised bed. The experiment was laid out in randomized block design (RBD) with ten treatments where each treatment replicated thrice. The various combination of PGRs were: T<sub>1</sub>-Naphthalene Acetic acid (NAA) 25 ppm, T<sub>2</sub>-Naphthalene Acetic acid (NAA) 50 ppm, T<sub>3</sub>-Naphthalene Acetic acid (NAA) 75 ppm, T<sub>4</sub>-Gibberellic acid (GA<sub>3</sub>) 25 ppm, T<sub>5</sub>- Gibberellic acid (GA<sub>3</sub>) 50 ppm, T<sub>6</sub>-Gibberellic acid (GA<sub>3</sub>) 75 ppm, T<sub>7</sub>-Cycocel (CCC) 500 ppm, T<sub>8</sub>-Cycocel (CCC) 750 ppm, T<sub>9</sub>-Cycocel (CCC) 1000 ppm, T<sub>10</sub>-Control.

Foliar spray of PGRs was done at 30 and 45 days after planting. The required dose of PGRs was measured by digital analytical electronic balance and dissolved in a solvent and then the final volume was made up to one liter using distilled water. Growth regulators was sprayed with the help of knap sack sprayer. Observations were recorded by using standard method. The data of yield and qualitative parameters was tabulated and subjected to statically analysis using method of analysis of variance (ANOVA) for Randomized Block Design (RBD) by Fisher and Yates (1963). Whenever 'F' test was found significant for comparing the means of two treatment, critical difference (C.D. at 5%) were worked out.

## **RESULT AND DISCUSSION:**

There were significant differences among the treatments of naphthalene acetic acid, gibberellic acid, and cycocel on yield and qualitative traits in strawberry cv. Winter Dawn under protected cultivation. The

table 1 revealed that the application of gibberellins and cycocel significantly influenced the number of fruit per plant, berry length, berry breadth, berry weight, marketable yield, unmarketable yield and yield per plant. Different combinations of plant growth regulators exhibit significant influence on yield and qualitative traits of strawberry fruit. The maximum number of fruit per plant (21.10), yield per plant (304.26g), and marketable yield (88.27%) were found in GA<sub>3</sub> 75ppm. The increase in the number of fruits per plant can be attributed to the vigorous growth of plants treated with gibberellins, which accumulate more starch, carbohydrates, and photosynthates, resulting in a greater number of berries per plant. This result is in line with the findings of Khalid *et al.* (2013) and Saima *et al.* (2014) in strawberry crop. The increase in fruit yield per plant in gibberellin-treated plants could be attributed to increased vegetative growth, which allows for greater fruit set and weight. The yield attributes on the sink capacity of crop is determined by its vegetative growth throughout the life cycle of plants. Vigorous growth is associated with higher sink capacity of a crop. This result is in conformity with the findings of Asadi *et al.* (2013), Saima *et al.* (2014), and Tanuja *et al.* (2019).

The highest marketable yield with GA<sub>3</sub> 75ppm because gibberellic acid regulates the movement of metabolites from the sink to the source (Iqbal *et al.*, 2011). By stimulating enzymes needed during the post-fertilization stage, the exogenous injection of GA<sub>3</sub> may control this balance in favour of fruit-forming metabolic processes, improving fruit setting and ultimately leading to the greatest number of fruits/plant (Sharma and Sharma, 2004). A further benefit of using GA<sub>3</sub> on strawberry blooms is reported to be improved pollen germination (Paroussi *et al.*, 2002). GA<sub>3</sub> at all concentrations produced significantly longer fruits over the control. The earlier cell elongation and cell division at apical region by GA<sub>3</sub> might have contributed to increased fruit length which was also explained by Moore *et al.* (1970). The maximum berry breadth (2.87cm), berry length (3.97cm) and berry weight (16.57g) were obtained in cycocel 1000ppm. The increase in breadth, length and weight of berries with the application

of CCC might be due to greater supply of photosynthates to the berries. This may also be due to lower percentage of fruit set and lower number of fruits which enables fruit growth to its maximum extent. The current results are in correspondence with the findings of Saima *et al.* (2014).

**Table 1. Effect of plant growth regulators on yield parameters in strawberry cv. Winter Dawn**

<b>Treatments</b>	<b>No. of fruit/plant</b>	<b>Berry length (cm)</b>	<b>Berry breadth (cm)</b>	<b>Berry weight (gm)</b>	<b>Yield/plant (g)</b>	<b>Marketable fruit (%)</b>	<b>Unmarketable fruit (%)</b>
<b>T<sub>1</sub> (NAA 25 ppm)</b>	16.27	3.32	2.60	12.73	207.11	79.89	20.11
<b>T<sub>2</sub> ( NAA 50 ppm)</b>	17.19	3.40	2.52	12.97	222.95	80.54	19.46
<b>T<sub>3</sub> (NAA 75 ppm)</b>	18.24	3.51	2.45	13.27	242.04	82.43	17.57
<b>T<sub>4</sub> (GA<sub>3</sub> 25 ppm)</b>	18.90	3.59	2.37	13.57	256.47	83.64	16.36
<b>T<sub>5</sub> (GA<sub>3</sub> 50 ppm)</b>	19.50	3.71	2.29	13.95	272.02	86.38	13.62
<b>T<sub>6</sub> (GA<sub>3</sub> 75 ppm)</b>	21.10	3.79	2.21	14.42	304.26	88.27	11.73
<b>T<sub>7</sub> (CCC 500 ppm)</b>	15.40	3.84	2.72	16.07	247.47	76.14	23.86
<b>T<sub>8</sub> (CCC 750 ppm)</b>	14.70	3.91	2.79	16.30	239.61	77.96	22.04
<b>T<sub>9</sub> (CCC 1000 ppm)</b>	13.40	3.97	2.87	16.57	222.02	79.36	20.64

<b>T<sub>10</sub> ( Control)</b>	13.10	2.90	2.08	12.32	161.39	74.37	25.63
<b>C.D. @ 5%</b>	0.53	0.17	0.09	0.45	12.13	3.38	0.96
<b>S.Em±</b>	0.21	0.05	0.03	0.15	4.05	1.13	0.32

The data presented in table 2 showed that there was significant difference between different plant growth regulators but there were no significant differences between different treatments. The maximum and minimum titratable acidity (0.89% and 0.63%) were recorded in control and CCC@1000ppm respectively. The reduction in titratable acidity could be attributed to the conversion of more organic acids and photosynthates into sugars during the fruit ripening stage (Saima *et al.*, 2014, and Rajesh *et al.*, 2012a).

**Table 2. Effect of plant growth regulators on qualitative parameters in strawberry cv. Winter Dawn**

<b>Treatments</b>	<b>TSS (°Brix)</b>	<b>Titratable acidity (%)</b>	<b>TSS/acid ratio</b>	<b>Ascorbic acid (mg/100gm)</b>	<b>Anthocyanin (mg/100g)</b>
<b>T<sub>1</sub> (NAA 25 ppm)</b>	8.81	0.80	11.00	52.02	59.61
<b>T<sub>2</sub> ( NAA 50 ppm)</b>	8.60	0.79	10.84	53.96	62.23
<b>T<sub>3</sub> (NAA75 ppm)</b>	8.42	0.78	10.75	55.15	65.02
<b>T<sub>4</sub> (GA<sub>3</sub> 25 ppm)</b>	7.91	0.82	9.84	53.73	63.86

<b>T<sub>5</sub> (GA<sub>3</sub> 50 ppm)</b>	7.70	0.83	9.24	55.29	65.15
<b>T<sub>6</sub> (GA<sub>3</sub> 75 ppm)</b>	7.51	0.85	8.57	56.99	67.33
<b>T<sub>7</sub> (CCC 500 ppm)</b>	8.57	0.66	12.98	58.21	56.55
<b>T<sub>8</sub> (CCC 750 ppm)</b>	8.78	0.64	13.71	59.8	57.17
<b>T<sub>9</sub> (CCC 1000 ppm)</b>	9.38	0.63	14.89	60.13	58.17
<b>T<sub>10</sub> ( Control)</b>	7.40	0.89	8.27	50.17	54.95
<b>C.D. @ 5%</b>	0.36	0.04	0.62	2.56	2.45
<b>S.Em±</b>	0.01	0.12	-	0.85	0.81

The highest ascorbic acid (60.13mg/100g), TSS (9.38° Brix) and TSS/acid ratio (14.89) was found in CCC 1000ppm. The higher ascorbic acid content could be attributed to the effect of growth regulators on biosynthesis from its precursor glucose-6-phosphate, which may increase ascorbic acid content (Tripathi and Shukla, 2007). While increase in sugar and TSS content of fruits with plant bio-regulators treatments, particularly CCC, could be attributed to the rapid metabolic transformation of starch and pectin into soluble sugars and the rapid translocation of sugars from leaves to developing fruits by reducing plant vegetative growth. The increase in total sugar content resulted in a higher TSS/acid ratio (Sevan *et al.*, 2014). In strawberry fruits, the maximum anthocyanin content (67.33mg/100gm) was recorded in treatment T<sub>6</sub> (GA<sub>3</sub> 75 ppm), whereas the minimum anthocyanin (54.95mg/100g) was observed in treatment T<sub>10</sub> (control). Montero *et al.* (1998) discovered increased anthocyanin content with increasing GA<sub>3</sub> concentrations, which could be due to either direct or indirect involvement of GA<sub>3</sub> in the synthesis

of anthocyanin pigment or its precursor, or involvement in the movement of its precursor. Furthermore, the higher anthocyanin levels observed in this study could be attributed to increased carbohydrate accumulation as a result of increased photosynthesis under the influence of plant growth regulators.

## **CONCLUSION:**

From the present study, it is concluded that foliar application of GA<sub>3</sub>@75 ppm resulted in improved yield and anthocyanin while CCC@1000ppm improve the quality of strawberry. These results could be utilized to assess the impact of plant growth regulators on various strawberry cultivars grown under protected agriculture in the future.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## **REFERENCES:**

- Asadi, Z., Jafarpour, M., Golparvar, A. R. And Mohammadkhani, A. (2013). Effect of GA<sub>3</sub> application on fruit yield, flowering and vegetative characteristics on early yield of Strawberry cv. Gaviota. *Int. J. Agri. CropSci.*, **5**(15): 1716-1718.
- Choudhary, Sakshi., Topno, Samir E. and Joseph, Annjoe V. (2024). “Effect of PGR and Micronutrients on Growth, Yield and Quality of Strawberry (*Fragaria × ananassa*) Cv. Winter Dawn”. *Journal of Advances in Biology & Biotechnology*, **27**(6):968-73.
- Gautam, Manish Kumar and Mishra, Saket. (2024). “Impact of Plant Growth Regulators on Growth, Fruit Yield and Quality of Strawberry (*Fragaria Ananassa* Duch.) Cv. Winter Dawn”. *Journal of Experimental Agriculture International*, **46**(7):1205-12.
- Iqbal, N., Nazar, R., Khan, M.I.R., Masood, A. and Khan, N.A. (2011). Role of gibberellins in regulation of source-sink relations under optimal and limiting environmental conditions. *Current Science*. **100**(7): 998-1007.
- Khalid, M. Q., Saman, C., Usman, S. Q. and Nadeem, A. A. (2013), Impact of exogenous application of salt and growth regulators on growth and yield of strawberry. *Pakistan J. Bot.*, **45**(4): 1179-1185.
- Montero T., Molla E., Martin Cabresas M. A. and Lopez Andreu F.J. (1998). Effect of GA<sub>3</sub> on strawberry, Phenylalanine ammonia lyase and Tyrosine ammonia lyase enzymes activities. *Journal of Science Food and Agriculture* **77** (2): 230- 234.
- Moore, J.N., G.R. Brown and E.D. Brown (1970). Comparison of factors influencing fruit size in large-fruited and small-fruited clones of strawberry. *J. Amer. Soc. Hort. Sci.* **95**:827-831.
- Nazir F, Hussain A, Fariduddin (2019). Interactive role of epibrassinolide and hydrogen peroxide in regulating stomatal physiology, root morphology, photosynthetic and growth traits in *Solanum lycopersicum* L. under nickel stress. *Environ Exp Bot* 162:479–495.

- Paroussi, G., Voyiatzis, D.G., Paroussis, P. and Drogoudi, P.D. (2002). Growth, flowering, and yield response to GA<sub>3</sub> of strawberry grown under different environmental conditions. *Sci. Hortic.*, **96**: 103-114.
- Rajesh, K., Manish, B. and Singh, D. B. (2012a). Influence of plant growth regulators on growth, yield and quality of Strawberry (*Fragaria X ananassa* Duch.) under U. P. sub tropics. *The Asian J. Hort.*, **7**(2): 434-436.
- Saima, Z., Sharma, A., Umar, I. and Wali, V. K. (2014). Effect of plant bio-regulators on vegetative growth, yield, and quality of strawberry cv. Chandler. *African Journal of Agricultural Research*, **9**(22): 1694-99.
- Sevan, D. K., Anil, K., Vijay, K., Shambu, S. and Saravanam, S. (2014). Effect of growth regulators and organic manure on physico-chemical properties of Strawberry (*Fragaria X ananassa* Duch.) cv. Chandler. *International Journal for Scientific Research and Education*. **2**(7): 1424-1435.
- Sharma, V. P., and Sharma, R. R. (2004). The Strawberry, ICAR, New Delhi, India. pp. 106-109.
- Srivastav, A., Singh, B.K., Pandey, R., Singh, K. and Singh, V. (2018). Effect of organic manures and bio-fertilizers on vegetative growth and yield of strawberry cv. Chandler. *Journal of Pharmacognosy and Phytochemistry*. **7**(5): 2841-2844.
- Tanuja and Rana, D.K. (2019). Assessment of planting dates, GA<sub>3</sub> concentrations and Azotobacter on Chandler strawberry. *Indian J. Hort.* **76**(4): 641-645.
- Tiwari, A.K., Saravanan, S. and Lall, D. (2017). Influence of different plant growth regulators on vegetative growth and physico-chemical properties of strawberry (*fragaria x ananassa* duch.) cv. Chandler. *Plant Archives*. **17** (1); 367-370.

Tripathi, V, K. and Shukla, P. K. (2007). Effect of plant bio-regulators, zinc sulphate and boric acid on yield and quality of Strawberry cv. Chandler. *J. Asian Hort.*, **4**(1): 15-18.

Yadav, I., Singh, J., Meena, B., Singh, P., Meena, S., Neware, S and Patidar, D.K. (2017). Strawberry Yield and Yield Attributes after Application of Plant Growth Regulators and Micronutrients on Cv. Winter Dawn. *Chem Sci Rev Lett*, **6**(21), 589-594.