

Original Research Article

“Interaction effect of various establishment methods and solid and liquid organic manures on growth of finger millet (*Eleusine coracana* L.)”

Abstract

A field experiment was conducted in certified organic field SMOF, SHUATS during *Kharif* season (2021 and 2022) to study the impact of establishment methods, solid and liquid organic manures on growth and yield of finger millet. The experiment was carried out in split-split plot design replicated thrice with 27 treatments. Treatments comprised of three establishment methods (Line sowing, Broadcasting and Transplanting), three solid organic supplements (100% FYM, 100% poultry manure and 100% vermicompost) each in combination with three different liquid organic supplements (3% panchagavya, 3% jeevamrutham and 3% vermiwash, respectively). All the individual factors had significantly influence on the growth characters of finger millet. The higher growth parameters at 80 DAS was significantly recorded among establishment methods in Transplanting, among solid organic manures in Vermicompost and among liquid organic manures in Vermiwash. All three parameters recorded no significant effect in the first year. plant height 97.85 cm in second year, number of tillers first year 3.87 and pooled 3.83, dry weight 49.54, 46.90 and 48.22 g/plant in first, second year and pooled respectively.

Keywords: *Finger millet, organic manure, poultry manure, panchagavya, growth*

Introduction

Finger millet (*Eleusine coracana* L.) is one among the foremost important millet grown for both grain and fodder purpose in India. Finger millet also called as ragi contains higher calcium content which was 10 times more than rice or wheat (Michaelraj and Shanmugam 2013). Finger millet was the third most millet next to sorghum and pearl millet. In India, finger millet constitutes an area of 1.19 m ha with annual production of 1.98 m t and productivity of 1662 kg/ha. In Tamil Nadu, finger millet could be a prominent crop among small millets growing in a part of 0.86 L ha with 3.21 L t production and 3714 kg/ha productivity (INDIASTAT, 2018).

The main reasons for low productivity is due to an imbalance in nutrients coupled with adverse climatic conditions, late transplanting, faulty methods of cultivation and little or no use of fertilizers. The secret of boosting its yields mainly lies in suitable planting method and properly fertilizing the crop. Method of sowing is important agronomic factor affecting the productivity of crop. Proper sowing method is the important non-monetary input in crop production, which affects the crop growth, yield and quality to greater extent. Method of establishment play important role to fully exploit all available resources for growth as it provides optimum growing condition [Sarawale *et al.*, 2022].

Intensive cultivation, unbalanced and inadequate fertilizers with restricted use of organic manures have made soil deficient in nutrients and health. Therefore, organic farming is gaining importance which mainly involves the use of on-farm resources largely avoiding the utilization of chemical fertilizers. Liquid and solid manures having higher amount of beneficial microbes, macro and micro nutrients, essential amino acids, growth promoting substance like IAA, GA may greatly help in increasing soil microbial population and soil fertility further increasing the crop growth, yield and quality (Sreenivasa *et al.*, 2012). Organic farming practices are gaining importance as farmers realized benefits in terms of soil fertility, soil health and sustainable productivity. Most of the research on organic production of finger millet was applied with utilization of FYM, green manures, compost, neem cake, etc. Less number of researches was done on the effect of liquid organic manures like panchagavya, jeevamrutham, vermiwash alone or together with solid organic manures in finger millet Organic liquid formulations like jeevamrutha and panchgavya helps for quick build-up of soil fertility through enhanced activity of microflora and fauna. These have the properties of both fertilizer and biopesticide and play a key role in promoting growth and immunity to the plant system. Any combination that reduce the dependence on chemical fertilizers and other resources can go an extended way in maintain the soil fertility as well as the financial conditions of the farming

community. Hence, the experiment was carried out with an objective to find out the effect of solid and liquid organic supplements on growth and yield of transplanted finger millet.

Material and Methods

A research trial was conducted at Crop Research Farm, Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) during *kharif* season of 2021 and 2022 finger millet sown by broadcasting, line sowing and transplanting with spacing of 30 cm x 10 cm. The location is situated at 25.57° N latitude, 87.19° E longitude and at an altitude of 98 m above mean sea level. In broadcasting method of establishment, seeds and fertilizers were dispersed randomly in the experimental plot. In line sowing method, seeds were sown directly and application of fertilizers were done in rows with a definite spacing of 30 cm x 10 cm. Whereas, in transplanting method, 18 days old seedlings were transplanted with a definite row to row and plant to plant pattern of 30 cm x 10 cm with 2 seedlings each. For this, one raised nursery beds were prepared and seeds were sown on beds in a row, so that seedlings can be uprooted easily at the time of transplanting. As it is a rainfed crop, no irrigation is needed but after transplanting two irrigations with alternate days were given for better crop establishment. Manually two hand weedings were done at 25 and 50 DAS/DAT with '*khurpi*'. At every 20 days interval, observations such as plant height, number of total tillers/plant, Leaf Area Index (LAI) and dry weight. Solid organic manures *viz.* Framyard manure, Poultry manure and Vermicompust were applied to fulfil the nitrogen requirement. Liquid manures- Vermiwash, Jeevamrutha and Panchgavya were applied to Finger millet crop at an interval of every 20 days after sowing. Necessary aftercare operations were followed as per the recommendations. No major pest and disease incidences were noticed during crop growth. Experimental data collected was subjected to statistical analysis by adopting Fisher's method of Analysis of Variance (ANOVA) as outlined by Gomez and Gomez (1984). Critical Difference (CD) values were calculated whenever the 'F' test was found significant at 5 per cent level.

Result and Discussion

Plant height

There was no significant effect in number of tillers at 40 DAS and pooled. AT 80 DAS significantly higher plant height was recorded in Transplanting + VC + Panchgavya (97.85 cm) was recorded in second year. However, treatments Direct sowing + VC + Vermiwash, Transplanting + FYM + Panchgavya, Transplanting + FYM + Vermiwash, Transplanting + PM + Jeevamrut, Transplanting + PM + Vermiwash, Transplanting + PM + Panchgavya and Transplanting + VC + Vermiwash were

statistically at par with Transplanting + VC + Panchgavya. This might be due to higher macro and micro nutrient content of the poultry manure which enables continuous slow and steady release of nutrients coupled with panchagavya foliar spray increased the nutrient uptake which might have helped in better growth (Priya and Sathyamoorthi, 2019).

Number of tillers

There was no significant effect in number of tillers at 40 DAS in both the years and pooled. There was significant effect of interaction effect at 80 DAS in first year and pooled. In first year significantly highest number of tillers was recorded in Transplanting + VC + Panchgavya (3.87). Direct sowing + FYM + Panchgavya, Direct sowing + PM + Panchgavya, Direct sowing + PM + Vermiwash, Direct sowing + VC + Panchgavya, Direct sowing + VC + Vermiwash, Transplanting + FYM + Panchgavya, Transplanting + PM + Jeevamrut, Transplanting + PM + Vermiwash, Transplanting + VC + Vermiwash were statistically at par with Transplanting + VC + Panchgavya. Further, in pooled significantly highest number of tillers was recorded in Transplanting + VC + Panchgavya (3.83). Treatments Direct sowing + PM + Panchgavya, Direct sowing + VC + Panchgavya, Transplanting + FYM + Panchgavya, Transplanting + PM + Vermiwash and Transplanting + VC + Vermiwash were statistically at par with Transplanting + VC + Panchgavya. Transplanted plants would have utilized the available sources such as spacing, forage area for root system, light utilization further enhanced the tiller development (Vanukuri *et al.*, 2022). This ensured continuous availability of nutrients throughout the crop growth stages due to steady transformation, mineralization, solubilisation, decomposition of minerals and nutrients that might helped in ensuring superior yield attributing characters by organics. Similar findings were observed with Gawade *et al.*, (2013) and Ananda *et al.*, (2017).

Dry weight

However, in the first year of 80 DAS significantly higher dry weight was recorded in Transplanting + Vermicompost + Panchgavya (49.54 g/plant). Treatments Transplanting + Farmyard Manure + Panchgavya, Transplanting + Farmyard Manure + Jeevamrut, Transplanting + Farmyard Manure + Vermiwash, Transplanting + Poultry Manure + Panchgavya, Transplanting + Poultry Manure + Vermiwash and Transplanting + Vermicompost + Vermiwash were statistically at par with Transplanting + Vermicompost + Panchgavya. Similarly, in the second year Transplanting + Vermicompost + Panchgavya (46.90 g/plant) recorded significantly higher dry weight. However, treatments Transplanting + Farmyard Manure + Panchgavya, Transplanting + Farmyard Manure + Vermiwash, Transplanting + Poultry Manure + Panchgavya, Transplanting + Poultry Manure +

Jeevamrut, Transplanting + Poultry Manure + Vermiwash, Transplanting + Vermicompost + Jeevamrut and Transplanting + Vermicompost + Vermiwash were statistically at par with Transplanting + Vermicompost + Panchgavya. Also, in pooled significantly higher dry weight was recorded in Transplanting + Vermicompost + Panchgavya (48.22 g/plant). Treatments Transplanting + Farmyard Manure + Panchgavya, Transplanting + Farmyard Manure + Vermiwash, Transplanting + Poultry Manure + Panchgavya, Transplanting + Poultry Manure + Vermiwash and Transplanting + Vermicompost + Vermiwash were statistically at par with Transplanting + Vermicompost + Panchgavya. Transplanting and organic nutrients increase of dry weight is due to the fact that the crop absorbed proportionately higher amount of N, P and K due to their higher availability under lower plant population and less competition among the plants for growth resources. The increased growth attributing characters in respect to the application of organic supplements might be due to enhanced nutrient availability (Yogananda *et al.*, 2019). Since poultry manure and panchagavya contains high nitrogen, macro and micro nutrients and growth promoting substance which helped in increased yield attributes and yield (Kumar *et al.*, 2018).

Conclusion

Transplanting finger millet seedlings along with application of solid organic manure of Vermicompost and Panchgavya recorded significantly higher growth attributes.

References

Ananda MR, Sharanappa, Kalyana Murthy KN. Response of Finger Millet under Organic Nutrient Management in Groundnut (*Arachis hypogaea* L.) – Finger Millet (*Eleusine coracana* L.) Cropping System. *International Journal of Pure & Applied Bioscience*. 2017; 5(5):200- 206.

Gawade MB, Mahadkar UV, Jagtap DN. Effects of organic manures, sources and levels of fertilizers on yield attributes and yield of finger millet (*Eleusine coracana* G.). *International Journal of Agricultural Sciences* 2013; 9(2):795-798.

Kumar, M.V., Velayutham, A., Kumar, N.S., Vasanthi, D. 2018. Influence of Different Organic Manures on the Growth and Yield of Baby Corn. *International Journal of Advances in Agricultural Science and Technology*, 5(7):167-174.

Michaelraj PSJ, Shanmugam A. 2013. A study on millets based cultivation and consumption in India *Int J Market Financ Serv Manag Res*. 2(4):49-58.

Priya, G. and Sathyamoorthi, K. 2019. Influence of Organic Manures on the Growth and Yield of Foxtail Millet [*Setaria Italica* (L.) Beauv]. *Chemical Science Review and Letters*, 8(29):114-117

Sarawale, P.P., Rajemahadik, V.A., Shendage, G.B. and Mane, S.V. 2016. Effect of different varieties and establishment methods on growth and yield of finger millet (*Eleusine coracana* (L.) Gaertn.) under konkan condition. *Journal of the Indian Society of Coastal Agricultural Research*, 34(2):22- 26.

Sreenivasa, M.N., Nagaraj, N. and Bhat, S.N. 2012. Nutrient status and microbial load of different organic liquid manures. *Karnataka Journal of Agricultural Sciences*, 24(4).

Vanukuri, B.R, Singh, V., George, S.G. and Vivek, 2022. Effect of Establishment Methods and Nutrient Levels on Growth and Yield of Finger Millet (*Eleusine coracana* L.). *International Journal of Environment and Climate Change*, 12(9): 201-205.

Yogananda S.B., Devkumar N., Gowda P.T., Shruthi, G.K. 2019. Influence of bio-digester liquid and farmyard manure on growth and yield of rice (*Oryza sativa*) in Cauvery Command Area of Karnataka. *Indian Journal of Agronomy* 64(1): 54-57

Table 1 Interaction effect of establishment methods, solid and liquid organic manures on plant height

| Plant height (cm) | | | | | | | | | | | | | |
|-------------------|----|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|-------|-------|
| 40 DAS | | | | | | | | | | | | | |
| treatment | | 2021 | | | | 2022 | | | | POOLED | | | |
| | | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN |
| DR | O1 | 36.71 | 36.56 | 35.66 | 36.31 | 33.00 | 31.85 | 33.95 | 32.93 | 34.85 | 34.20 | 34.80 | 34.62 |
| | O2 | 39.12 | 35.46 | 39.83 | 38.13 | 41.07 | 34.75 | 37.45 | 37.76 | 40.09 | 35.11 | 38.64 | 37.95 |
| | O3 | 38.15 | 35.99 | 36.87 | 37.00 | 37.44 | 31.61 | 34.83 | 34.63 | 37.80 | 33.80 | 35.85 | 35.81 |
| BR | O1 | 33.13 | 28.17 | 32.47 | 31.26 | 29.76 | 27.46 | 28.76 | 28.66 | 31.45 | 27.81 | 30.62 | 29.96 |
| | O2 | 35.83 | 34.46 | 32.20 | 34.16 | 35.79 | 34.09 | 34.49 | 34.79 | 35.81 | 34.28 | 33.35 | 34.48 |
| | O3 | 32.70 | 31.12 | 32.25 | 32.02 | 34.99 | 35.41 | 33.54 | 34.65 | 33.84 | 33.27 | 32.90 | 33.33 |
| TR | O1 | 42.00 | 40.82 | 42.35 | 41.72 | 40.95 | 37.44 | 39.93 | 39.44 | 41.48 | 39.13 | 41.14 | 40.58 |
| | O2 | 48.07 | 41.84 | 46.02 | 45.31 | 48.69 | 41.13 | 44.31 | 44.71 | 48.38 | 41.49 | 45.17 | 45.01 |
| | O3 | 43.95 | 41.48 | 44.28 | 43.24 | 48.90 | 41.44 | 44.24 | 44.86 | 46.43 | 41.46 | 44.26 | 44.05 |
| Mean | | 38.85 | 36.21 | 37.99 | 37.68 | 38.95 | 35.02 | 36.83 | 36.94 | 38.90 | 35.62 | 37.41 | 37.31 |
| F-test | | NS | | | | NS | | | | NS | | | |
| SEm± | | 1.96 | | | | 1.60 | | | | 1.23 | | | |
| CD (P=0.05) | | - | | | | - | | | | - | | | |
| Plant height (cm) | | | | | | | | | | | | | |
| 80 DAS | | | | | | | | | | | | | |
| treatment | | 2021 | | | | 2022 | | | | POOLED | | | |
| | | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN |
| DR | O1 | 79.89 | 76.58 | 79.42 | 78.63 | 79.94 | 75.30 | 77.78 | 77.67 | 79.92 | 75.94 | 78.60 | 78.15 |
| | O2 | 83.65 | 82.07 | 83.03 | 82.92 | 85.01 | 80.23 | 85.23 | 83.49 | 84.33 | 81.15 | 82.63 | 82.70 |
| | O3 | 80.10 | 79.97 | 82.53 | 80.87 | 82.37 | 80.17 | 87.73 | 83.42 | 81.24 | 80.07 | 82.13 | 81.15 |
| BR | O1 | 74.62 | 74.58 | 74.32 | 74.51 | 74.42 | 70.38 | 73.78 | 72.86 | 74.52 | 72.48 | 74.05 | 73.69 |
| | O2 | 79.38 | 73.33 | 78.14 | 76.95 | 75.18 | 73.13 | 74.61 | 74.31 | 77.28 | 73.23 | 76.37 | 75.63 |
| | O3 | 76.59 | 76.66 | 75.48 | 76.24 | 80.72 | 79.46 | 81.95 | 80.71 | 78.65 | 78.06 | 78.72 | 78.48 |
| TR | O1 | 92.42 | 85.71 | 90.92 | 89.68 | 89.89 | 86.51 | 87.72 | 88.04 | 90.66 | 86.11 | 89.32 | 88.70 |
| | O2 | 98.38 | 88.17 | 91.45 | 92.67 | 94.29 | 91.30 | 94.58 | 94.58 | 98.11 | 89.74 | 93.01 | 93.62 |
| | O3 | 96.16 | 89.66 | 93.60 | 93.14 | 97.85 | 86.12 | 93.40 | 91.27 | 95.22 | 87.89 | 93.50 | 92.20 |
| Mean | | 84.58 | 80.75 | 83.21 | 82.84 | 84.30 | 84.41 | 80.29 | 84.09 | 82.93 | 80.52 | 83.15 | 82.70 |
| F-test | | NS | | | | S | | | | NS | | | |
| SEm± | | 2.65 | | | | 2.29 | | | | 1.74 | | | |
| CD (P=0.05) | | - | | | | 10.24 | | | | - | | | |

Table 2 Interaction effect of establishment methods, solid and liquid organic manures on number of tillers

| No. of tillers | | | | | | | | | | | | | | |
|----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|------|--|
| 40 DAS | | | | | | | | | | | | | | |
| treatment | 2021 | | | | | 2022 | | | | | POOLED | | | |
| | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN | | |
| DR | O1 | 1.60 | 1.40 | 1.47 | 1.49 | 1.33 | 1.20 | 1.40 | 1.31 | 1.47 | 1.30 | 1.43 | 1.40 | |
| | O2 | 1.80 | 1.67 | 1.53 | 1.67 | 1.47 | 1.27 | 1.67 | 1.47 | 1.63 | 1.47 | 1.60 | 1.57 | |
| | O3 | 1.80 | 1.60 | 1.53 | 1.64 | 1.53 | 1.47 | 1.53 | 1.51 | 1.67 | 1.53 | 1.53 | 1.58 | |
| BR | O1 | 1.33 | 1.13 | 1.33 | 1.27 | 1.40 | 1.27 | 1.53 | 1.40 | 1.37 | 1.20 | 1.43 | 1.33 | |
| | O2 | 1.60 | 1.40 | 1.33 | 1.44 | 1.53 | 1.40 | 1.33 | 1.42 | 1.57 | 1.40 | 1.33 | 1.43 | |
| | O3 | 1.80 | 1.33 | 1.53 | 1.56 | 1.73 | 1.33 | 1.67 | 1.58 | 1.77 | 1.33 | 1.60 | 1.57 | |
| TR | O1 | 1.73 | 1.53 | 1.93 | 1.73 | 1.73 | 1.53 | 1.80 | 1.69 | 1.73 | 1.53 | 1.87 | 1.71 | |
| | O2 | 2.07 | 1.33 | 1.47 | 1.62 | 1.93 | 1.40 | 1.60 | 1.64 | 2.00 | 1.37 | 1.53 | 1.63 | |
| | O3 | 2.00 | 1.73 | 1.60 | 1.78 | 1.67 | 1.67 | 2.00 | 1.78 | 1.83 | 1.70 | 1.80 | 1.78 | |
| Mean | 90.23 | 86.54 | 89.51 | 88.76 | 89.22 | 83.96 | 88.01 | 87.06 | 89.73 | 85.25 | 88.76 | 87.91 | | |
| F-test | NS | | | | | NS | | | | | NS | | | |
| SEm± | 2.79 | | | | | 2.94 | | | | | 1.92 | | | |
| CD (P=0.05) | - | | | | | - | | | | | - | | | |
| No. of tillers | | | | | | | | | | | | | | |
| 80 DAS | | | | | | | | | | | | | | |
| treatment | 2021 | | | | | 2022 | | | | | POOLED | | | |
| | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN | | |
| DR | O1 | 3.33 | 3.07 | 3.20 | 3.29 | 3.40 | 3.33 | 3.20 | 3.22 | 3.37 | 3.20 | 3.20 | 3.26 | |
| | O2 | 3.47 | 3.20 | 3.27 | 3.38 | 3.60 | 3.40 | 3.33 | 3.38 | 3.53 | 3.30 | 3.30 | 3.38 | |
| | O3 | 3.73 | 3.00 | 3.40 | 3.49 | 3.40 | 3.33 | 3.27 | 3.22 | 3.57 | 3.17 | 3.33 | 3.36 | |
| BR | O1 | 3.10 | 2.60 | 3.13 | 2.93 | 2.87 | 2.47 | 2.67 | 2.71 | 3.03 | 2.53 | 2.90 | 2.82 | |
| | O2 | 2.93 | 2.87 | 2.47 | 3.09 | 2.80 | 2.87 | 3.00 | 2.87 | 2.87 | 2.83 | 3.23 | 2.98 | |
| | O3 | 2.73 | 2.67 | 2.47 | 3.02 | 2.67 | 2.67 | 2.93 | 2.80 | 2.80 | 2.73 | 3.20 | 2.91 | |
| TR | O1 | 3.60 | 3.20 | 3.07 | 3.31 | 3.65 | 3.27 | 3.40 | 3.40 | 3.60 | 3.23 | 3.23 | 3.36 | |
| | O2 | 2.87 | 3.27 | 3.73 | 3.36 | 3.60 | 3.47 | 3.60 | 3.56 | 3.23 | 3.37 | 3.77 | 3.46 | |
| | O3 | 3.87 | 3.07 | 3.53 | 3.49 | 3.80 | 3.20 | 3.53 | 3.51 | 3.83 | 3.13 | 3.53 | 3.50 | |
| Mean | 3.32 | 3.10 | 3.36 | 3.26 | 3.30 | 3.01 | 3.24 | 3.19 | 3.31 | 3.06 | 3.30 | 3.22 | | |
| F-test | S | | | | | NS | | | | | S | | | |
| SEm± | 0.20 | | | | | 0.16 | | | | | 0.11 | | | |
| CD (P=0.05) | 0.64 | | | | | - | | | | | 0.37 | | | |

Table 3 Interaction effect of establishment methods, solid and liquid organic manures on dry weight

| Dry weight | | | | | | | | | | | | | |
|------------|-------------|-------|-------|-------|-------|-------|-------|-------|-------|--------|-------|-------|-------|
| 40 DAS | | | | | | | | | | | | | |
| treatment | | 2021 | | | | 2022 | | | | POOLED | | | |
| | | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN |
| DR | O1 | 6.20 | 5.15 | 6.48 | 5.94 | 5.71 | 4.82 | 5.38 | 5.31 | 5.96 | 4.99 | 5.93 | 5.62 |
| | O2 | 8.41 | 6.64 | 7.35 | 7.47 | 6.28 | 5.14 | 6.14 | 5.86 | 7.34 | 5.89 | 6.75 | 6.66 |
| | O3 | 7.38 | 5.74 | 6.45 | 6.52 | 8.41 | 6.59 | 7.93 | 7.64 | 7.90 | 6.17 | 7.19 | 7.08 |
| BR | O1 | 5.11 | 4.71 | 4.77 | 4.86 | 5.15 | 4.97 | 4.98 | 5.03 | 5.13 | 4.84 | 4.87 | 4.95 |
| | O2 | 5.14 | 4.24 | 5.03 | 4.80 | 5.38 | 5.24 | 5.87 | 5.50 | 5.26 | 4.74 | 5.45 | 5.15 |
| | O3 | 5.48 | 5.21 | 4.58 | 5.09 | 7.99 | 9.48 | 7.75 | 8.41 | 6.73 | 7.34 | 6.17 | 6.75 |
| TR | O1 | 7.28 | 7.99 | 8.86 | 8.04 | 6.48 | 5.95 | 5.52 | 5.99 | 6.88 | 6.97 | 7.19 | 7.01 |
| | O2 | 9.49 | 8.14 | 9.10 | 8.91 | 9.10 | 7.36 | 6.58 | 7.68 | 9.30 | 7.75 | 7.84 | 8.30 |
| | O3 | 10.16 | 8.42 | 9.05 | 9.21 | 9.86 | 9.10 | 8.38 | 9.12 | 10.01 | 8.76 | 8.72 | 9.16 |
| | Mean | 1.59 | 1.46 | 1.75 | 1.60 | 1.42 | 1.27 | 1.56 | 1.42 | 1.51 | 1.36 | 1.66 | 1.51 |
| | F-test | NS | | | | NS | | | | NS | | | |
| | SEm± | 0.24 | | | | 0.23 | | | | 0.12 | | | |
| | CD (P=0.05) | - | | | | - | | | | - | | | |
| Dry weight | | | | | | | | | | | | | |
| 80 DAS | | | | | | | | | | | | | |
| treatment | | 2021 | | | | 2022 | | | | POOLED | | | |
| | | L1 | L2 | L3 | Mean | L1 | L2 | L3 | Mean | L1 | L2 | L3 | MEAN |
| DR | O1 | 35.14 | 34.70 | 38.08 | 33.94 | 33.12 | 35.57 | 33.12 | 33.94 | 34.13 | 35.14 | 35.60 | 34.96 |
| | O2 | 41.57 | 36.96 | 42.31 | 38.72 | 40.03 | 38.09 | 38.05 | 38.72 | 40.80 | 37.53 | 40.18 | 39.50 |
| | O3 | 44.45 | 37.95 | 39.72 | 40.49 | 38.53 | 40.20 | 42.73 | 44.82 | 39.99 | 39.08 | 41.23 | 40.10 |
| BR | O1 | 34.70 | 34.85 | 41.57 | 34.20 | 35.70 | 33.96 | 32.95 | 33.87 | 35.20 | 34.41 | 37.26 | 35.62 |
| | O2 | 36.09 | 34.98 | 38.95 | 38.05 | 39.85 | 38.98 | 35.31 | 36.05 | 37.97 | 36.98 | 37.13 | 37.36 |
| | O3 | 36.05 | 33.98 | 39.54 | 34.24 | 34.70 | 34.20 | 33.83 | 38.91 | 35.38 | 34.09 | 36.69 | 35.38 |
| TR | O1 | 46.73 | 45.49 | 47.73 | 47.24 | 46.42 | 34.97 | 43.06 | 36.15 | 46.98 | 39.34 | 45.77 | 44.03 |
| | O2 | 47.53 | 44.20 | 46.44 | 46.06 | 44.82 | 41.95 | 43.20 | 44.35 | 45.78 | 44.64 | 47.40 | 45.94 |
| | O3 | 49.54 | 43.70 | 48.48 | 45.22 | 46.90 | 43.78 | 45.06 | 45.24 | 48.22 | 43.08 | 45.82 | 45.37 |
| | Mean | 40.01 | 37.97 | 38.81 | 40.83 | 40.90 | 37.74 | 38.92 | 39.19 | 40.49 | 38.25 | 40.67 | 39.81 |
| | F-test | S | | | | S | | | | S | | | |
| | SEm± | 1.46 | | | | 1.70 | | | | 0.99 | | | |
| | CD (P=0.05) | 4.77 | | | | 5.54 | | | | 3.24 | | | |