

IMMUNOGENICITY OF BACTERIAL PROTOPLASMIC SONICATE PROTEINS

Abstract

Bacterial Protoplasmic Sonicate Proteins (BPSP) were recovered from skin burn *Pseudomonas aeruginosa* and *Klebsiella oxytoca* infections. These proteins were considered as immunogens in rabbits. Mucosal and systemic, humoral and cellular immune responses were matched. The *P. aeruginosa* protoplasmic sonicate proteins induce an increase in; NBT % neutrophil phagocytosis, significant leukocyte migration inhibition cytokines, IL6 cytokine elevation and rise up of humoral agglutinins responses. While, *K. oxytoca* protoplasmic sonicate proteins induce increase of; humoral agglutinins and IL6 cytokine as well as non-significant increase in leukocyte inhibitory cytokines. Both at mucosal and systemic compartments. The study highlights the potentials of these proteins as candidates for developing vaccines against multidrug resistance infections, particularly in burn patients. These findings underscore the importance of immunogenicity in advancing bacterin development in clinical use.

Key Words

Antigens, Bacterin, Cellular, Humoral, immunogen, Protein, vaccine.

Introduction

Bacterial antigens [BAGs] were being investigated to be of use in, bacterial diagnosis, bacterial infection diagnosis, vaccine development and vaccine production. Immunogenicity of BAGs holds a backbone position in vaccine development strategies [1 – 7]. The objective of the present work was the investigation of immunogenicity of protoplasmic sonicate protein antigens of skin burn infections with *P. aeruginosa* and *K. oxytoca*.

Materials and Methods

Protoplasmic Sonicate Protein Antigens (PSPA);

The PSPA separation was in direct way for *P. aeruginosa* [8], while, by indirect way as it needs removal of the capsule [9, 10] before processing for obtaining PSPs. The method for separation, identification, purification and quantitation of these PSPs were as in the method described by Bjorn et al. [8]. The details of the processing method was as in the followings; Six ml of tris buffer 0.01 N and PH 8 was added to the surface of 24 hrs growth of *P. aeruginosa* and *K. Oxytoca* (after suspension of bacteria then removal of capsule [9, 10]) on the nutrient agar plates. Growth were vortexed in vortex test tubes for three minutes. Suspensions were centrifuged at 5000 rpm for ten minutes. Supernatant were discarded and pellets Ps were kept. Ps suspensions were tubed and washed three times with tris buffer 0.01N. Ps were reconstituted with 6 ml, and were tubed in the cell disintegrator tubes. Then jacketed with cooled ice. The best sonication conditions were five times for five minutes at 20 oscillation amplitude. The sonicated cell suspensions were centrifuged at 5000 rpm for ten minutes. Supernats were collected and ultrafiltered with 0.22 um millipore filter. Filterates were collected in sterile plastic tubes, then proteins were separated with PEG 6000, 6% as in Shnawa and AlSadi [11]. The protein concentration determination was

measured by Biurt test[12].To this end the preparations can be designated as PAPSP and KOPSP both *P.aeruginosa* and *K.oxytoca* respectively.These PSPs were distributed into aliquotes of 0.5 mls. In an appendroff plastic tubes and kept at -20C,till use.

Immune Reagents;

Specific immune priming of rabbits were donewith the PSP concentrations of 2.71 mg/l for PAPSP and1.8 md/l for KOPSP[12].The somatic antigens for both bacteria were prepared as heat killed as in [13].Complete Freund Ajuvant that from Difco,Co.Ltd. The test proposed immunogens were made as; one voliume of CFA mixed with one voliume the test proteins

Immunization Protocols

A mounthwise twice dosage of PSP-CFA in 2 mls amounts were dosaged in the first and second months to the rabbits .The specific immune priming was that of multisite injection protocol[14].

Rabbits:

A local breed rabbits brought from the local market were checked for ecto and endoparasites .As well as for pyrogens and found to be free off.They were acclimatized for housing conditions two weeks befor experimentation at an ad libitum conditions.Among which nine were elected and subdivided into two test and one control group each of three rabbits.

Samplings And Immune Function Tests

At the temination of the specific immune priming protocols,blood with and with out heparin were collected from the test and control rabbits by cardiac puncture for humoral and cellular immune tests.Sera were saved for serology and cytokine studies.Heparinized blood were used for leukocyte inhibitory factor[15] and for NBT phagocytosis[16].Appendix for test and control rabbits were collected and open up,washed from digesta and processed for separation of mucosal globulins[17].Mucosal leukocytes were separated by dextran 2% as in [18].IL 6 determinations were made as in the recommendation of the instruction of the manufacturer.Standared tube agglutination test were made as in [19].

Results

The NBT neutrophil phagocytosis percentages in PAPSPA(56.75 % for mucosal and36.5 for systemic) and KOPSPA(48.5% for mucosal , 41% for systemic)primed rabbits were higher than that of control rabbits(20% for mucosal,18% for the systemic).Leukocyte inhibitory cytoknine LIF % in PAPSP primed rabbits were56.6% for mucosal and 58.45 fo systemic as compared to control ,the mucosal was 90% and systemic was 86%.While for KOPSPA primed rabbits were; 88% for mucosal and 87.5% for systemic LIF as compared to normal control were 93% for mucosal and 89% for systemic responses.The IL6 concentration determinations were shownig that PAPSPA and KOPSPA primed rabbits have got higher IL6

concentration means than normal control rabbits. PAPS PA primed rabbits IL6 concentration means were 92.8 for mucosal and 72.7 for systemic responses. In KOPSPA primed rabbits the mucosal concentration means were 83.26 and for systemic were 76.79 as compared to normal control were 8.83 for mucosal and 9.25 for systemic responses. The humoral specific agglutinin titre levels for PAPS PA were 128 for mucosal and 320 for the systemic responses and KOPSP primed rabbits were 64 for mucosal and 640 for systemic responses as compared to control rabbits were 4 for mucosal and 20 for systemic responses. Mucosal agglutinins were resistant to treatment with 2ME, Tables 1 and 2.

Table – 1 : The immunogenicity of PAPS P in primed rabbits and controls.

Rabbits groups	NBT%	LIF%	IL6 pg/ml.	Agglutinin titres
PAPS PA				
M	46.75	56.6	2.8	128
S	36.6	58.45	72.7	320
Control				
M	20	93	92.8	4
S	18	89	72.7	20

Table – 2 : The immunogenicity of KOPSP in primed and control rabbits

Rabbit groups	NBT%	LIF %	IL6 pg/ml	Agglutinin titres	
KOPSPA					
M	48.5	88	48	64	
S	41	87	72	640	
Controls					
M	20	93	8.83	4	
S	18	89	9.25	20	

Discussion

The concept, application and continuity of the immunogenicity theme for bacterial protein antigens are still in the current mode of researchers all over the world [1-7]. Immunogenicity appeared to have two main facets. First that of theoretical immunologists which advocate that immunogenicity is denoted to self-nonself recognition theme [6]. While the second facet was that for most of the proper immunologists which can be summarized as the

ability of an antigen to initiate humoral and/or cellular conversion from the normal baseline immune functions to an optimized cellular immune reactions outcomes that are finalized by the optimum synthesis and production of cellular secretory proteins(antibodies,cytokines) concentrations and /or optimised depression of such secretory proteins[1-5,7]. Immunogenicity appeared to be essential for diagnosis and prophylaction of human infections as well as in cancer personalized prophylactic and therpeutic medicine.Hence the presnt study for Pseudomonas and Klebsiella immunogenicity may participate in developing of a porotype bacterial protein based vaccine and/ or in development for an autovaccine for multidrug resistant skin burn infection[20]

The antigenic make up of bacteria[1-5,7] like that of P.aeruginosa are formed from;flagella,pili,exotoxin A,exopolysaachride,LPS,OMP, hemolysin, elastases, proteasis, heat stable phospholipid,heat stable glycolipid,exoenzyme S and ribosome[21-23].While that of Klebsiella oxytoca are;capsule,somatic antigens and LPS[24].Hence, PAPSP and KOPSP proteins antigens are novel antigens prepared in this experimental settings.

PAPS was proved to be immunogenic in rabbits model,Table-1,promoting humoral and cellular immune responses both at mucosal and systemi ccompartments.These findings were in contradiction with that of[24].While,KOPSP immunogenicity,Table -2, were by promoting humoral immune responses both at mucosal and systemic level.[21].Different burn infecting bacterial protein preparations have shown different immune potential features,Table - 3,[25].The functional epitope mapping for PAPSPmay be T cell depndent and T cell independent,while that of KOPSP may be of T cell independent or Th2 cell depent B cell epitopes[26,27].**Since immunogenicity is an integral part for developing bacterins.Such findings holds the position of novelty,based upon the need for an autogenous bacterins for multidrug resistant skin burn bacterial infections [28,29].**

Table – 3 : The immune features of the study bacterial proteins.

Features	PAPSP[25]	KOPSP[25]
1-Chemical nature	Protein	Protein
2-Origin	Bacterial	Bacterial
3-Location	Intracellular	Intracellular
4-Processing method	Ultrasonication	Ultrasonication
5-Non-specific immune	Rise up of phagocytosis By	Rise up of phagocytosis by

function	NBT%	NBT %
6- Specific cellular immune function	Significant inhibition of LIF cytokine	Nonsignificant inhibition of LIF cytokines
7-Induction of cytokine network	Rise up of IL6 cytokine	Rise up of IL6 cytokines
8-Humoral Immune responses	Rise up of specific agglutinins	Rise up of specific agglutinins
9-Functional Epitope Mapping[26,27]	T cell dependent and T cell independent epitopes	T cell independent and /or, TH2 dependent B cell epitope
10-Immune system compartments	Mucosal and Systemic	Mucosal and systemic
11-Expected Immune potentials[20]	Prototype protein based vaccine and an autovaccine	Prototype protein based vaccine and an autovaccine.

Conclusions

P.aeruginosa PAPSP and *K.oxytoca* KOPSP protoplasmic sonicate proteins were found as lapin immunogens. PAPSP mediate humoral and cellular immune responses both at mucosal and systemic compartments. While, KOPSP mediate humoral immune responses both at mucosa and blood stream. These proteins may be of expected immune potentials as protein based vaccine candidate and as an autovaccine for multidrug resistant skin burn infections.

Ethical Issue

The housing, handling and interventions on rabbits were done following the international acts for regulation of housing, handling and interventions.

References

- 1- Tian G , Qin C , Hu J et al. 2023. Side chain functional groups in immunogenicity of bacterial surface glycans. *Molecules* 28:7112. doi.3390/molecules28207112.
- 2- Yang J , Zhang J , Dong J et al. 2023. De novo identification of bacterial antigens of clinical isolate by combining use of proteosurfaceomics, secretomics and Bascan techniques. *Front. Immunol.* 14:1274027. doi.10.3389/fimmu.2023.124027.
- 3- Alwis R de. Liang L , Taghavian J et al. 2021. The identification of novel immunogenic antigens as potential Shigella vaccine components. *Genomic Medicine.* 13:8. doi.101186/s13073-020-00824.
- 4- Maerz JK, Trastel C , lange A , et al. 2020. Bacterial immunogenicity is critical for the induction of regulatory B cells in suppressing inflammatory immune responses. *Front. Immunol.* 24:10:3093. doi.10.3389/fimmu.2019.03093.
- 5- Mahanty S , Pregel A , Garraud O. 2015. Immunogenicity of infectious pathogens and vaccine antigens. *BMC .Immunol.* 16(31):Article number, 31.
- 6- Pradeu T , Carosella ED. 2006. On the definition of a criterion of immunogenicity. *PANAS*, 103(47):17858-17861.

- 7- Dougan G ,Ghaem-Maghami M ,Ives SK, et al.2000.The immune responses to bacterial antigens encountered i-invivo-i at mucosal surfaces .Philo. Trans. Royl. Soc.Sci.London,355(1397): 705-712.
- 8- Bjor MJ , Iglewski HB , Ives SK et al.1978.Effect of iron on yeilds of exotoxin A in culture of *Pseudomonas aeruginosa*, PA-10,Inf.Immu.19(3)785-791.
- 9- Dolby MJ ,1958.The separation of histamine sensitizing factor from the protective antigen of *Brodetella pertussis*.Immunol.1:328.
- 10- Kwapnski JB .1972.Methodology of Immunochemical and Immunological researchs.Whiley-interscience,New York.267-316.
- 11- Shnawa IMS , AISadi MAK.2001.Gut mucosal immunoglobulin separation,patial charaterization and utility as infection probe Iraqi.J.Microbiol.13(3):
- 12- Bishop ML 1985.Clinical Biochemistry:Principles,Procedures,Correlations.2nd ed.Lippincott-Williams and Wilkins
- 13- Svanburg –Eden C , Kullhary R ,Martids S.1985.urinary immunoglobulin in healthy individuals and children with acute pyelonephritis.Scan.J.Immunol.21:305-313.
- 14- AL-Shahery MAN,Shnawa IMS,1989.The immunological adjuvanicity of sunflower ,Vet.Med.J. Giza.37(2):291-28.
- 15- Soberg M 1969.Interaction of human peripheral blood lymphocytes and granulocytes in the migration inhibition reactions,Acta Medica Scand.185:221.
- 16- Park BH , Fikring SM , Smithwick E J.1968.Infection and nitroblue tetrazolium reduction by neutrophils .A diagnostic aid .Lancet 2 :532-534.
- 17- Hassan A J.2002.Comparative Study on Some Aspects of the Immune System of Collard Dove and Rabbit,Ph.D.Thesis,College of Science,University of Babylon/IRAQ.
- 18- Skoog WA , Beck WS 1956.Studies on fibrinogen,dextran and phytohemagglutinin ,methods of isolating leukocytes from blood,Hematol.11:466-
- 19- Turgeon ML 2020.Immunology and Serology In Laboratory Medicine,Elsevier Health Sciences.
- 20- Shnawa IMS 2019.Vaccine Technology At A Glance.Boffin Access,UK.
- 21- Levinson W ,Chin-Hong P ,Joyce EA ,Nussbaum J, Schwartz B 2018.Rewiw of Medical Microbiology and Immunology:A Guide to Clinical Infectous Diseases 15th edLange MacGraw –Hill New York.
- 22-Brooks GF ,Morse S A , Mietzner TA,Miller S ,2016.Jawetz,Melnick and Adelberg's Medical Microbiology 27th ed ,Lange,MacGraw –Hill Education.
- 23-Hancock EW, Lam JS 1998.*Pseudomonas aeruginosa*: infection and immunity.Encyclopedia of Immunology Vo.4,2nd ed.Delves PJ,Roitt I eds .Academic Press.USA.2042-2045.
- 24-Toivaness P , Hansen DS, Mastre F ,1999.Somatic serogroups,capsular type and special *Klebsiella* in paients with ankylosing spondylitis.J.Clin.Microbiol.37(9)2808-2812.

25-Zheng D , Liwinski T , Elinary E .2020.Interaction between microbiota and immunity in health and Disease .Cell Research.30:492-506.

26-Gershoni JM , Roitburd-Berman A ,Siman-Tor DD, Freund NT , Weiss Y .2007.Epitope mapping.Drug Development 21:145-156.

27-Shnawa IMS 2016.Functional epitope mapping of a vaccine.Vaccines Vaccin.1(1):000101

28-Jin X et al. 2022.Recent adavances in vaccine development against multidrug resistant bacteria.Vaccines.10(9):1456.doi.10.3390/vaccines.10091456.

29-Smith R et al.2023.New insight into the immunogenicity of bacterial antigens: implications for vaccine development.J.Clin.Microbiol.61(5):e1425-23.doi.10.1128/JCM.01423-23.