

ETHNOBOTANICAL STUDY OF MEDICINAL PLANTS IN KEPILASGADHI RURAL MUNICIPALITY OF KHOTANG DISTRICT, NEPAL

ABSTRACT:

Background: Ethnobotany is a field that study traditional use of plants by local communities. In Nepal, approximately 80% of the population, particularly in rural areas, relies on traditional medicine due to their effectiveness and cultural significance.

Objectives: This study aims to investigate the ethnomedicinal knowledge in Kepilasgadhi Rural Municipality, Khotang, Nepal, documenting medicinal plants used by local communities, identify diseases treated, plant parts used and application method, evaluate the Use Value (UV), Fidelity Level (FL), Informant Consensus Factor (ICF) and Relative Frequency of Citation (RFC) as well as threats, and conservation practices to these medicinal plants.

Methods: Primary data were gathered through reconnaissance surveys, key informant interviews, structured questionnaires, focus group discussions, and field observations. Secondary data included previous research, journals, books, and reports.

Results: A total of 102 medicinal plants across 93 genera and 57 families were documented. Herbs were the most common life form (52.94%), with roots being the most used part (21.31%). Oral administration was predominant (72.54%), with inhalation being least common. Gastrointestinal disorders were most treated (54 plants), followed by skin issues. A total of 10 species showed highest FL of 100% for specific ailments. ICF values ranged from 0.11 to 0.75, with urologic disorders showing the highest consensus. *Paris polyphylla* had the highest RFC (0.98), while *Chilocostus speciosus* had the lowest (0.02). The findings highlight significant UV for plants like *Curcuma domestica* L (1.83), indicating their high importance in treating ailments and frequent use as a traditional medicine. Despite a rich ethnomedicinal heritage unsustainable harvesting, forest fires, deforestation, illegal collection have threatened these resources. Conservation efforts include cultivating high-demand species, reforestation, and selective harvesting.

Conclusion: Although Kepilasgadhi is rich in medicinal plants and traditional knowledge, declining youth interest threatens their preservation. Urgent initiatives are needed to educate younger generations about these practices.

Keywords: *Medicinal plants, Ethnobotany, Traditional medicine, Informant consensus factor, Use value.*

1. INTRODUCTION:

John W. Harshberger first introduced the term "ethnobotany" in 1896 to refer to the study of plants used by primitive and indigenous people. It explores the relationship between humans and plants, documenting traditional knowledge on how different ethnic groups or communities utilize local plant resources. Since the dawn of human civilization, people have relied on plants for various needs, including food, fodder, medicine, and fiber. Over time, through experience and experimentation, they discovered numerous properties and uses of plants, particularly their medicinal benefits (Ahmad et al., 2014; Bhattarai & Khadka, 2016). Approximately 80% of the global population, especially in the rural areas of developing and under developed countries, relies on traditional medicine as their main source of healthcare (Islam, 2006; Bibi et al., 2014) due to their effectiveness, the lack of modern medical options, and deeply rooted cultural preferences (Svarstad & Dhillon, 2000; Plotkin & Famolare, 1992; Taylor et al., 1995; Tabuti et al., 2003). Recent years have seen a decline in traditional medicinal practices due to decreased interest among the younger generation, urban migration, mass deforestation, changing life style, and traditional healers shifting to other jobs (Kadir et al., 2013). It is estimated that between 35,000 and 70,000 plant species are utilized in traditional medicine around the world (Farnsworth & Soejarto, 1991) and more than 50% of modern pharmaceutical drugs have origins rooted in traditional ethnomedicine (Jan et al., 2015; Sher et al., 2003). According to study conducted by Fabricant and Farnsworth (2001) 122 compounds derived from 94 plant species are used in medicine worldwide.

Nepal is a diverse nation with multiple ethnic groups, languages, and religions, reflecting varieties of culture and tradition (Mallik et al., 2020). With over 142 caste/ethnic groups and 129 spoken languages (CBS, 2021), the ethnic communities possess extensive traditional knowledge about the use of plants and plant parts. This indigenous knowledge has been passed down through generations, highlighting a long-standing tradition of utilizing plant resources (Acharya & Acharya, 2009). The traditional knowledge held by indigenous communities is crucial for the development of herbal remedies and the extraction of bioactive compounds, which could potentially result in the discovery of new drugs (Cox & Balick 1994, Fabricant & Farnsworth 2001). Although the use of medicinal plants in Nepal dates back to ancient times, the earliest documented evidence of their medicinal use is found in the Rig-Veda, written between 4500 and

1600 BC, which is considered one of the oldest repositories of human knowledge on plant-based medicine (Rokaya et al., 2010; Singh et al., 2012). Out of approximately 29 million inhabitants in Nepal, over 80% living in rural areas, who have limited access to government healthcare facilities and face high costs and unavailability of allopathic medicine, rely on traditional remedies as a primary or complementary treatment for illnesses (Sharma et al., 2004; Raut and Khanal, 2011). In Nepal a total of 6,653 species of angiospermic plants were documented, with 1,792 to 2,331 of them identified as potential medicinal and aromatic plants (Rokaya et al., 2010). With varied topography and microclimate within the small geographical area, Nepal harbors a rich diversity of medicinal and aromatic plants (Baral and Kurmi, 2006; Ghimire, 2008; Shrestha et al., 2022).

The traditional use of plants for medicinal purposes in Nepal is gaining popularity because these remedies are often free of side effects, readily accessible, affordable, and sometimes the only healthcare option available for those in need. (Acharya & Acharya, 2009). Numerous studies have demonstrated that medicinal plants have been traditionally used to treat a wide range of ailments, including fever, digestive issues, skeletal and muscular disorders, jaundice, blood purification, asthma, bone spurs, hemorrhoids, pneumonia, snake bites, dehydration, fatigue, liver disorders, tonsillitis, rheumatism, constipation, toothaches, high blood pressure, hemorrhage, skin diseases, and more (Bhattarai et al., 2010; Kutal et al., 2021; Shrestha & Pradhan, 1986; Sharma et al., 2004; Baral & Kurmi, 2006; Ghimire, 2008; Rana et al., 2015; Bhandari et al., 2021; Malla & Chhetri, 2009; Kunwar et al., 2013; Ambu et al., 2020; Mahato & Chaudhary, 2003). Medicinal plants are used not only for treating human ailments but also for addressing various diseases in animals (Rana et al., 2015). Some medicinal plants are also used for religious or ritual purposes (Ambu et al., 2020) due to their significant spiritual/religious value, particularly in Hinduism. For example, *Ocimum sanctum* (Tulsi) is regarded as one of the most sacred plants in Hindu culture. It is believed to attract positive energy, bring blessings, and strengthen one's connection to God.

In Nepal, ethnobotanical research began with Banerji's 1955 publication on medicinal and food plants. Until then, various studies have been conducted across various region of the country (Mahato & Chaudhary, 2003; Shrestha & Pradhan, 1986; Shrestha et al., 1998; Shrestha & Dillion, 2003; Mahato & Chaudhary, 2005; Malla & Chhetri, 2009; Uprety et al., 2010; Singh et

al., 2018) but, Kepilasgadhi Rural Municipality in Khotang District remains largely unexplored despite its significance for medicinal plant diversity. No scientific research has yet been conducted in this area. Given these considerations, the present study aims to document ethnomedicinal knowledge among people of Kepilasgadhi Rural Municipality. The main objective of the study are:


1. To investigate the medicinal plants utilized by the people of Kepilasgadhi Rural Municipality.
2. To identify the diseases treated by these medicinal plants.
3. To assess the plant parts/products used and their methods of application.
4. To evaluate the Use Value (UV), Informant Consensus Factor (ICF), Fidelity Level (FL) and Relative Frequency of Citation (RFC) of medicinal plants in Kepilasgadhi Rural Municipality.

2. MATERIALS AND METHODS:

2.1. STUDY AREA:

The study was conducted in Kepilasgadhi Rural Municipality, located in the Khotang district of Nepal (Figure 1), positioned at 27.33° N latitude and 86.83° E longitude. Kepilasgadhi Rural Municipality cover an area of 191.28 km² and is consists of 664 households with the population of 15,288 individuals, comprising 7,323 men and 7,965 women (CBS, 2021). The region experiences a maximum temperature of 21.75°C, a minimum temperature of 10.59°C, and an average annual rainfall of 263.29 mm.

The Rai community is the predominant ethnic group in Kepilasgadhi Rural Municipality, with a population of 9,654, speaking the Sampang language. Other ethnic groups residing in the rural municipality include Chhetri, Tamang, Kami, Damai/Dholi, Sherpa, Gurung, Newar, Koiri/Kushwaha, Hill Brahmin, Nachhiring, Gharti/Bhujel, Sunuwar, Sanyasi/Dashnami, Limbu,



Sarki.

Figure 1: Map of the study area

2.2. DATA COLLECTION

2.2.1. PRIMARY DATA COLLECTION

Primary data were systematically gathered using various methods, including a reconnaissance survey, key informant interviews, a structured questionnaire survey, Focus group discussions, and field observations.

2.2.1.1. Reconnaissance Survey

A reconnaissance survey was conducted in Kepilasgadhi Rural Municipality, through field visits and discussions with local leaders. This survey aimed to develop an initial understanding of the respondents' socio-economic status and healthcare needs. During this phase, permission to conduct the study was obtained by clearly communicating the study's objectives to the local authorities and community members.

2.2.1.2. Key Informant Interviews (KII)

A total of 20 key informants were purposively selected for in-depth interviews. This group included 10 traditional healers (Lama, Baidhyas, and local practitioners), 2 traders, 2 officials of Division forest office, Khotang, and 6 individuals representing the Community Forest User Groups (CFUG), model farmers, and teachers. The interviews focused on gathering detailed information about the local medicinal flora, including trees, shrubs, and herbs used in traditional medicine.

2.2.1.3. Questionnaire Survey

To ensure a representative and diverse sample, stratified random sampling approach was employed. The Household were first divided into distinct strata based on key characteristics, including ethnicity, caste, and economic class. From these strata, a total of 100 households were

randomly selected, representing 15% of the total households in the study area. One individual from each selected household was interviewed. This method ensured that the sample accurately reflected the population's diversity and provided robust data for analyzing the research objectives.

2.2.1.4. Field Observation

Field observations were conducted in various settings, including forests, fallow lands, and farmyards, with the assistance of local experts identify and authenticate the plant species. Specimens of accessible medicinal plants were collected, and photographed for future reference.

2.2.1.5. Focus Group Discussions (FGDs)

Three Focus group discussions were conducted, involving 9-12 participants each. Two discussions were held with the dominant Chhetri and Rai communities separately, and one with a mixed group comprising individuals from different ethnic backgrounds, traditional healers, and local experts. These discussions provided insights into the ethno-medicinal knowledge of bio-resources, including the effectiveness and trade of medicinal plants, the threats they face, and the conservation practices implemented.

2.2.2. SECONDARY DATA COLLECTION

Secondary data were sourced from previous research studies, academic journals, books, websites, and reports from relevant agencies, both published and unpublished. This comprehensive literature review provided a deeper understanding, interpretation, and analysis of the research context and findings.

2.3 DATA ANALYSIS

To systematically analyze the ethnomedicinal data collected, statistical software such as SPSS and Microsoft Excel were employed. The analysis focused on quantifying the significance of

various medicinal plant species and assessing the consensus among informants regarding their use. The following indices were calculated to evaluate the ethnobotanical knowledge: Use Value (UV), Fidelity Level (FL%), Informant Consensus Factor (ICF) and Relative Frequency of Citation (RFC).

2.3.1. Use Value (UV)

The Use Value (UV) index was calculated to determine the relative importance of each plant species based on the number of use reports provided by indigenous informants. This index reflects the significance of each species within the local medicinal practices. The UV was calculated using the formula (Tabuti et al., 2003).

$$UV = U/n$$

Where 'U' represents the total number of use of particular species reported by all respondents, and 'n' represents the total number of informants questioned for that particular plant species. The Use Value (UV) of a plant increases with the number of use provided by informants, indicating that the plant is of greater significance within the community. Conversely, a lower UV suggests less frequent use or recognition of the plant (Vitalini et al., 2013).

2.3.2. Fidelity Level (FL%)

The Fidelity Level (FL%) was employed to identify the most preferred plant species for treating specific ailments. Fidelity level quantifies the percentage of informants who endorse the use of a particular plant species for a specific health condition. The FL was calculated using the formula (Friedman et al., 1986).

$$FL (\%) = (N_p / N) * 100$$

Where 'N_p' represents the number of informants reporting the use of the plant for a specific illness, while 'N' denotes the total number of informants reporting the use of the plant for any health condition. A high Fidelity Level (FL) value indicates that the plant species is frequently

used by informants for treating a specific ailment category within the study area (Yaseen et al., 2015).

2.3.3. Informant Consensus Factor (ICF)

The Informant Consensus Factor (ICF) was calculated to assess the degree of agreement among informants regarding the use of plants for specific ailment categories. The ICF was calculated using the formula: (Trotter & Logan, 1986; Teklehaymanot, 2009)

$$ICF = \frac{Nur - Nt}{(Nur - 1)}$$

Where 'Nur' indicates number of use citations in each category, and 'Nt' indicates the number of species used. The ICF value ranges from 0 to 1, with higher values indicating greater agreement among informants, suggesting a higher potential for the identified species to be effective in treating the ailment (Trotter & Logan, 1986; Heinrich et al., 1998). High ICF values suggest that certain species are widely recognized within the community for their medicinal properties, making them prime species for further cultural and pharmacological research (Mbuni et al., 2020).

2.3.4. Relative Frequency of Citation (RFC)

The Relative Frequency of Citation (RFC) indicates the local significance of each species that was determined using the formula (Vitalini et al., 2013).

$$RFC = FC/N$$

where 'FC' represents the number of informants who reported the use of the species, and 'N' is the total number of informants who participated in the survey.

3. RESULTS:

3.1 Availability and diversity of used medicinal plants.

In the Kepilasangadhi Rural Municipality, a total of 102 medicinal plant species, representing 93 genera and 57 families, have been documented (Figure 2). The species composition includes 54

herbaceous plants, 22 tree species, 16 climbers, and 10 shrubs. Among these, the family Rosaceae is the most prevalent, with 7 species, followed by Poaceae with 6 species. Several families, including Ranunculaceae, Cucurbitaceae, Ericaceae, and Asteraceae, each contribute 5 species to the medicinal flora. Additionally, Lauraceae, Moraceae, and Zingiberaceae are each represented by 3 species. Families such as Saxifragaceae, Polygonaceae, Rutaceae, Orchidaceae, Liliaceae, Vitaceae, Urticaceae, Rubiaceae, Lamiaceae, Apiaceae, Piperaceae, Apocynaceae, and Fabaceae are each represented by 2 species. The remaining 34 families are represented by a single species each, highlighting the diverse botanical resources utilized within this ethnic community.




Figure 2: Plant families of medicinal plants

3.2 Ethnomedicinal uses of plants in Kepilasgadhi Rural Municipality and their Use Value

Table given below shows the list of ethnomedicinal plants used by the people Kepilasgadhi Rural Municipality and their their Use Value.

Table 1: Ethnomedicinal uses of plants and their Use Value (UV)

SN	Local name	Scientific name	Family	Parts used	Management	Life form	Distribution	Diseases treated	Method of preparation	Mode of application	Previous use reports	Use value (UV)
1	Pakhanbed	<i>Bergenia ciliata</i> (Haw.) Sternb.	Saxifragaceae	Root	W	H	A	Body pain/bone fracture/ pregnancy problems/ diarrhea/ dysentery	Powder/ decoction	O	wound, diabetes, piles, tumor, urinary problems, hearth, and respiratory diseases (Bhattarai & Khadka, 2016; Malla & chhetri, 2009; Thapa, 2012; Shah & lamichhane, 2017)	1.33
2	Padamchal	<i>Rheum acuminatum</i> Hook. f. & Thomson ex Hook	Polygonaceae	Root	W	H	A	Body pain/gastritis/ Indigestion/bone fracture	Decoction	O	Treatment of arthritis, rheumatism, blood purification, fever reduction, blood purification, constipation (Rokaya et al., 2010, Singh et al., 2012; Farnsworth, 1990)	1.75
3	Satuwa	<i>Paris polyphylla</i> Sm.	Melanthiaceae	Whole plants	W/C	H	R	Gastritis/vomit/burn /fever/headache/cut/ stomach infection/sore throat/snakebite	Paste	T/O	Gastritis, Stomach swelling, wounds, skin infections, reduce fever, snake bites (Singh & Singh, 2009; Hamilton, 2004; Hu et al., 2020)	1.23
4	Bethlauri	<i>Chilocostus speciosus</i> (J. König) C. Specht	Costaceae	Stem/ root	W	H	R	Jaundice/burning urination	Juice extraction	O		0.75
5	Sil timur	<i>Lindera neesiana</i>	Lauraceae	Fruit	W	T	A	Common cold/cough, snake bite/indigestion/ sore throat	Decoction	O		1.07
6	Boke timur	<i>Zanthoxylum acanthopodium</i> DC.	Rutaceae	Fruit	W	T	A	Gastritis/fever/tooth ache	Decoction/ 1-2 pieces eaten raw	O	Gastritis, Leech bite, toothaches, oral discomfort, indigestion, stomach pain (Chae et al., 2016; Singh & Singh, 2009; Brisdon & forman 2010)	0.92

7	Sarpagandha	<i>Rauwolfia serpentina</i> (L.) Benth. ex Kurz (RKM78-S17)	Apocynaceae	Root/ leaf	W /C	S	R	Body pain/uric acid/ herpes zoster	Powder	O	Reduce blood pressure, dysentery, labour pain, removal of opacities of the cornea (Acharya & Acharya, 2009; Singh & Singh, 2009)	1.09
8	Jatamansi	<i>Nardostachys grandiflora</i>	Caprifoliaceae	Root	W	H	R	Diarrhea/ dysentery/anti-poison	Juice extraction	O	insomnia, hysteria, convulsions (Kunwar et al., 2013)	0.81
9	Bikhuma	<i>Aconitum bisma</i>	Ranunculaceae	Root/ leaf	W	H	R	Headache/fever/ common cold/snakebite/ low pressure	Direct chewing/ Juice extraction	O	tonsillitis, sore throat, gastritis and debility, anti-inflammatory (Kunwar et al., 2010; Shyaula, 2011)	1.05
10	Saktigumba	<i>Pleione praecox</i> L.	Orchidaceae	Pseudo bulb	W	H	R	Body pain/ligament sprain/bone fracture/cut wound	Powder mixed with milk/paste	O/ T		0.8
11	Golkakri	<i>Coccinia grandis</i> (L.) Voigt	Cucurbitaceae	Root/ fruit	W	CL	R	Cut/wounds/ body pain/ dysentery	Paste/Direct chewing	O/ T	leprosy, jaundice, diabetes, tongue sores, Indigestion, (Ramachandran et al., 2014; Waisundara et al., 2015)	1.21
12	Indreni	<i>Trichosanthes ricuspidata</i> Lour	Cucurbitaceae	Seed	W	CL	R	Cut/wounds	Oil extraction	T	Abscess, tonsillitis (Bhandari et al., 2021)	1.33
13	Khiraula	<i>Polygonatum cirrhifolium</i> L.	Liliaceae	Root	W	H	A	Cut/wound/diarrhea / mensuration problem	Powder/ Paste/juice extraction	O		1.08
14	Ban karela	<i>Herpetospermum pedunculatum</i> (Ser.) C.B. Clarke	Cucurbitaceae	Seed	W	H	R	Cut/wounds	Oil extraction	T	Treatment of high blood pressure, joint pain, constipation (Ambu et al., 2020)	0.86
15	Ban lasun	<i>Allium carolinianum</i> L.	Amaryllidaceae	Bulb	W	H	R	Stone/stomach disorder/ headache	Powder/ paste	O	Altitudinal sickness, high blood pressure, indigestion, colds, coughs, and bronchitis (Bhatta & Kunwar, 2020; rana et al., 2015)	1.46
16	Ban mula	<i>Potentilla fulgens</i> Wall. ex Hook	Rosaceae	Root/ bulb	W	H	R	Stone/body pain/sore throat/stomach pain/high pressure	Decoction/ dried piece is chewed directly	O	Regulate blood sugar levels, manage diabetes, dysentery, and stomach ulcers (Kichu et al., 2015; Tene et al., 2007)	1.29
17	Bojho	<i>Acorus calamus</i> L.	Araceae	Rhizome	W	H	R	Common cold/allergy/sore throat	Dried/fresh roots chewed directly/ infusion	O	cut, common cold, vomiting, sore throat, fever, Scabies, reduce mental stress, high blood pressure, asthma (Rana et al., 2015; Bhattarai, 2017; Baral, 2005; Bennett & Prance, 2000; Chase et al., 2016)	1.43

18	Nagbeli	<i>Lycopodium clavatum L.</i>	Lycopodeaceae	Whole plants	W	CL	A	Allergy/fever/ headache	Paste/juice extraction	T/ O	burns, headache (Rana et al., 2015)	1.08
19	Nangre/nas jhar	<i>Hemiphragma heterophyllum L.</i>	Scrophulariaceae	Whole plants	W	H	A	Cut/wounds	Paste	O		1
20	Pinase jhar/ tinpate	<i>Clematis buchananiana DC.</i>	Ranunculaceae	Whole plants	W	CL	A	Sinusitis/allergy/nasal bleeding/vitiligo	Juice extraction/ paste	O/T		0.64
21	Haadjora	<i>Cissus quadrangularis L.</i>	Vitaceae	Whole plants	W	H	R	Bone fracture/bone disease/cuts/ wounds	Decoction/ paste	T	menstrual disorders, wound healing, bone disorders (Siddiqua & Mittapally, 2017)	0.9
22	Laligurans	<i>Rhododendron arboretum Sm.</i>	Ericaceae	Flower/ leaf	W	T	A	Allergy/throat obstruction/ gastritis/ fertility increase in women	Leaf/flowers mixed with Tea/honey	O	cuts, cure from having fish bone into neck, jaundice, tonsillitis, diarrhea (Rana et al., 2015; Anel & Fundiko, 2016; Badola & Pradhan, 2013; Bhandari et al., 2021)	0.77
23	Chimal	<i>Rhododendron campanulatum L.</i>	Ericaceae	Flower	W	T	A	Body ache/throat pain	Direct Chewing	O		0.77
24	Loth salla/ dhyangre salla	<i>Taxus wallichiana Zucc.</i>	Taxaceae	Whole	W/C	T	A	Diabetes/cancer/urine infection/womb problems/cold/ cough/fever	Paste/juice extraction/ powder	O	treat digestive, respiratory, nervous and skeletal systems, cough, fever, pain (Sharma & Garg, 2015; Juyal et al., 2014)	1.13
25	Pani amala	<i>Nephrolepis auriculata (L.) Trimen</i>	Nephrolepiaceae	Root/ bulb	W	H	A	Diabetes/ stomach problems and infestation worms	Direct chewing	O	diabetes, stomach problems, diuretic (Dhami, 2008; Bhatt & Kunwar, 2020)	1.73
26	Bot amala	<i>Phyllanthus emblica L.</i>	Euphorbiaceae	Bark/ seed	W/C	T	R	Sore throat/tonsils/ blood purify	Powder/ Direct Chewing	O		1.46
27	Sisnu	<i>Urtica hyperborean L.</i>	Urticaceae	Leaf/ stem	W	H	A	Blood pressure/chest problems/cuts/ wounds/ eye disorder	Powder/ paste	O	Diuretic and appetizer (Dhami, 2008; Bhatt & Shakya, 2015).	1.4
28	Allo/bhagresisnu	<i>Girardinia diversifolia</i>	Urticaceae	Root	W	H	A	Swollen body parts	Paste	T	Galactagogue, diabetes, high pressure, fever, asthma, toothache, paralysis, uterine bleeding, rheumatism (Shah & lamichhane, 2017; Shrestha & Dhillion, 2003; Thapa, 2012)	0.02

29	Charpate jhar	<i>Oldenlandia lineata</i> (Roxb.) Kuntze	Rubiaceae	Leaf	W	H	A	Cut/wounds	Paste	T		1.09
30	Dhade bikh	<i>Aconitum ferox.</i>	Ranunculaceae	Root	W	H	A	Cancer	Paste/ powder	T		0.02
31	Lajjawoti jhar	<i>Mimosa pudica L.</i>	Leguminaceae	Whole plants	W	H	R	Jaundice	Paste	O	dysentery, regulate menstrual cycles, inflammations, treat burning sensation, asthma, fatigue and blood diseases (Joseph et al., 2013; Ahmad et al., 2012)	0.02
32	Ukhu	<i>Saccharum officinarum L.</i>	Poaceae	Stem	C	H	A	Jaundice	Juice extraction	O	Jaundice, stomach disorders, seminal weakness, liver health, energy boost (Ambu et al., 2020; Malla & Chhetri 2009)	0.04
33	Titepati	<i>Artemisia indica Willd.</i>	Asteraceae	Leaf	W	S	A	Indigestion/fever/cut/ scabies/body pain	Juice extraction	O	cut and smallpox, high blood pressure, muscular swelling (Rana et al., 2015; Baniya et al., 2019; Bhattarai, 2020)	1.5
34	Dhasingre/ maxino	<i>Gaultheria fragrantissima Wall.</i>	Ericaceae	leaves/ seed	W	S	A	Bone fracture/body pain/ cold/cough	Oil extraction	T	back pain, headaches, fever, skin diseases, sore throats, tooth ache (Chevallier, 1996; Joshi & Subedi, 2016)	1.07
35	Kurillo	<i>Asparagus racemosus Willd</i>	Asparagaceae	Root/ shoot	W	H	A	Immunity power/skin disease/diarrhe/ lactation in women	Powder/ decoction	O	feed to cattle increase, alleviate menstrual cramps, reduce the fatigue, support fertility. (Rana et al., 2015; Mallik et al., 2020; Malla & Chhetri, 2009)	1.25
36	Bantarul	<i>Dioscorea pentaphylla L.</i>	Dioscoreaceae	Bulb/ tuber	W	CL	A	Backbone pain	Infusion	O	skin infections, stomach pain, constipation problems (Manjula et al., 2022)	0.04
37	Sugandhawal	<i>Valeriana jatamansi Jones.</i>	Valerianaceae	Root	W/C	H	A	Nerves problems	Powder	O	Cuts and wounds, Fever, Tonsillitis, indigestion, abdominal pain (Bhatta & Negi 2006; Acharya, 2012; Bhandari et al., 2021)	1
38	Dhupi salla	<i>Crytomerica japonica</i>	Cupressaceae	leaves	W	T	R	Unconsciousnes/vo miting	Paste	O	liver disease, anti-inflammatory (Shyur et al., 2008)	1.5

39	Sikre/lokta	<i>Daphne bholua</i>	Thymileoaceae	leaves	W/C	S	A	Anti-poison	Paste	T	Fever, Treatment of intestinal disorders, sinusitis, coughs and bronchitis (Malla & Chhetri, 2009; Shrestha & Dhillon, 2016)	0.36
40	Pudina	<i>Mentha spicata L.</i>	Lamiaceae	Whole plant	W	H	A	Indigestion, diarrhea/stomach pain	Paste	O	Stomach pain, relaxant, haemorrhage, influenza, Intestinal parasites (Kichu et al., 2015; Macia et al., 2005; Tene et al., 2006)	0.10
41	Jarelo	<i>Cissus adnata</i>	Vitaceae	Whole plants	W	H	A	Eye infection	Paste	T		0.02
42	JharMauro	<i>Oxalis corniculata</i>	Oxalidaceae	Whole plants	W	H	A	Snakebite/ eye infection	Paste	T		0.16
43	Dubo	<i>Cynodon dactylon (L.) Pers</i>	Poaceae	Whole plant	W	H	A	Cut/wounds/ blood pressure/typhoid/fever	Juice extraction/ paste	O	bleeding, piles, paste applied in fracture (Singh, 2014; Bhatt & Shakya, 2015).	1.5
44	Khaniyu	<i>Ficus semicordata Buch-Ham. ex Sm.</i>	Moraceae	Latex	W	T	R	Boils/ulcers	Infusion	T		0.16
45	Chiraito	<i>Swertia Chirayita (Roxb. ex Fleming) Karsten</i>	Gentianaceae	Whole plant	W/C	H	A	Headache/fever/ blood pressure/ common cold	Juice extraction/ plant mix with tea	O	Bronchitis (Bhandari et al., 2021)	1.5
46	Pahale	<i>Cuscuta reflexa Roxb.</i>	Convolvulaceae	Whole plants	W	CL	R	Bone fracture/ jaundice	Juice/oil extraction	O / T	Jaundice, regulate blood sugar levels, anti-inflammatory, Fever (Abbas et al., 2017; Badalo & Pradhan, 2013; Farnsworth, 1990)	1.43
47	Sikari lahara	<i>Periploca calophylla</i>	Apocynaceae	Whole plants	W	CL	R	Bone fracture/ligament sprain/body aches	Paste	O		0.76
48	Budo okhati/ thulo okhati	<i>Astilbe rivularis Buch-Ham. ex D. Don,</i>	Saxifragaceae	Root/ stem	W / C	H	A	Tonsils/body aches/ pneumonia/ pregnancy problems/ fracture	Decoction/ powder	O	Treatment of wound healing, anti-inflammatory (Chen et al., 2021)	1.09
49	Sungure kada	<i>Cirsium verutum (D. Don) Spreng.</i>	Asteraceae	Root/ latex	W	H	A	Eye pain/Measles/ appendix	Taken raw/ juice extraction	O	Body pain, Fever, indigestion, rheumatism, cough, asthma, bronchitis (Asase et al., 2010; Bhatta, 1999; Budha magar et al.,	1.04

											2020)	
50	Hade kaphal	<i>Myrica esculenta</i> Buch-Ham. ex D.Don.	Myricaceae	Bark/fruit	W	T	A	Dysentery/fracture	Decoction/paste	O/T	eye pain and measles, fever, abdominal pain (Manandhar, 1995a; Kunwar et al., 2010; Rai & Singh, 2015)	0.95
51	Bhainsi kada	<i>Clematis connata</i> DC.	Ranunculaceae	Bark	W	CL	A	Sprain/fracture	Powder	O	Muscular swelling and sprain, sinusitis (Bhandari et al., 2021)	0.76
52	Mel	<i>Pyrus pashia</i> Buch-Ham. ex Don.	Rosaceae	Fruit	C	T	A	Diarrhea/dysentery	Decoction/Juice extraction	O		0.92
53	Nase jhar	<i>Plantago lagopus</i> L.	Plantaginaceae	Leaf	W	H	A	Fever/cough/common cold/stomach problem	Juice extraction	O	eye infections, sore throat, anemia, diarrhea, gastrointestinal disorder (Prakash et al., 2021; Murad et al., 2011; Kumar et al., 2011)	1.02
54	Dabuke jhar	<i>Centella asiatica</i>	Apiaceae	Leaf	W	H	A	Jaundice/pressure	Direct chewing	O	Hepatic pain, insomnia (Tene et al., 2007)	0.24
55	Ainselu	<i>Rubus ellipticus</i> Sm.	Rosaceae	Root/bud	W	S	A	Tonsils/fever, diarrhea/sore throat/cough	Powder/Juice extraction	O	Dysuria, Pneumonia, Tonsillitis (Baral & Kurmi, 2006; Bantawa & Rai, 2009; Bhandari et al., 2021; Asase et al., 2014)	0.47
56	Bhuin ainselu	<i>Fragaria daltoniana</i> J. Gay	Rosaceae	Stem	W	H	A	Cut/wounds	Paste	T		0.16
57	Angeri	<i>Lyonia ovalifolia</i> L.	Ericaceae	Leaf	W	T	A	Allergy	paste	O	Skin diseases treatment (Rana et al., 2015)	0.02
58	Jhyau	<i>Permelia nepalensis</i>	Parmeliaceae	Whole plants	W	H	A	Skin burn/piles	paste	T		0.82
59	Chabo	<i>Piper mullesua</i> Buch-Ham. ex D.Don	Piperaceae	Leaf	W	CL	A	Painkiller/tonsils	Direct chewing	O	antidepressant, anti-inflammatory and anti-oxidant (Tisserand & Balacs, 1994).	0.84
60	Hande okhar	<i>Juglans Regia</i> L.	Juglandaceae	Bark/root	W/C	T	A	Toothache/gum problems	Taken raw	O	Diabetes, Kidnet ailment, liver ailment (Macia et al., 2005; Kichu et	0.65

											al., 2015)	
61	Pipla	<i>Piper longum</i> L.	Piperaceae	Fruit	W	Cl	A	Immunity power/fever/ sore throat	powder	O	chronic bronchitis, asthma, constipation, diarrhea, cholera, chronic malaria (Kumar et al., 2013; Khushbu et al., 2011)	1.5
62	Aduwa	<i>Zingiber officinale</i> Rosc.	Zingiberaceae	Rhizome	C	H	A	Common cold/fever Sore throat/tonsils/ asthma	Juice extraction/ mix with tea	O	Diarrhoea, sinusitis, cold and cough, goiter (Ambu et al., 2020; Bhattarai et al., 2009; Tamang et al., 2016)	0.36
63	Besar	<i>Curcuma domestica</i> L.	Zingiberaceae	Rhizome	C	H	A	Body pain/fever sore throat/common cold/ jaundice	Infusion/ powder	O	cough, Skin disease Treatment, Common cold, Fungal infection (Bhandari et al., 2021; Rana et al., 2015)	1.83
64	Lasun	<i>Allium sativum</i> L.	Alliaceae	Leaf/ bulb	C	H	A	Cancer/heart disease/ gastritis	Decoction/ Bulb are fried	O	Malaria, high blood pressure, spider & snake bite (Kichu et al., 2015; Mesfin et al., 2009; Tene et al., 2006; Shrestha & Dhillon, 2003)	0.65
65	Balu	<i>Pieris formosa</i> (Wall.)D.Don	Ericaceae	Leaf/ root	W	T	A	Allergy/headache	Paste	T		0.64
66	Tejpaat	<i>Cinnamomum tamala</i> L.	Lauraceae	Leaf/ bark	W/C	T	A	Corona/stomach ache	Decoction/ Infusion	O	diarrhea, nausea and vomiting, colic, diarrhoea, dysentery or cough. (Chakraborty & Das, 2010; Shah & Panchal, 2010)	1.09
67	Thado unneu	<i>Dryopteris cochleata</i> (Don) C.Chr.	Dryopteridaceae	Rhizome	W	H	A	Mental disorder	Infusion	I		0.06
68	Dautheniuro	<i>Diplazium maximum</i> (D.D on)C.Chr.	Athyriaceae	Bud	W	H	A	Osteopenia/joint pain	Direct chewing	O	stomach problem, smallpox, asthma, diarrhea, rheumatism, dysentery, fever, wounds, high blood pressure, (Rana et al., 2015; Abe & Ohtani, 2013; Kadir et al., 2014; Tag et al., 2012)	0.1
69	Kukur daine	<i>Smilax ovalifolia</i> L.	Smilacaceae	Root/ fruit	W	CL	A	Joint pain/ dysentery/ gastritis	Juice extraction/ paste	O		0.33
70	Naspati	<i>Pyrus communis</i>	Rosaceae	Fruits	C	T	A	Diarrhea/indigestion/nausea/ vomiting	Juice extraction	O	Weight loose, skin whitening, fight against diabetes, Cholesterol Lowering, Macular Degeneration	1.5

											(Abbas & Bano, 2020)	
71	Kagati	<i>Citrus aurantiifolia</i> (Cristm.) Swingle*	Rutaceae	Fruit	C	S	A	Headache/fever/ blood pressure/ common cold	Juice extraction	O	Diarrhea, stomach ache, headache treat nausea and resuscitate fainting individuals(Enejoh et al., 2015; Khan et al., 2010; Kunow, 2003)	0.06
72	Sugandhak okila	<i>Cinnamomum glaucescens</i> L.	Lauraceae	Seed	C	S	R	Anti-inflammatory/ toothache	Oil extraction	T	toothache, wound, joints pain treatment (Shrestha & Dhillion, 2003; Ambu et al., 2020)	0.01
73	Bamboo	<i>Bambusa vulgaris</i>	Poaceae	Shoots	W/C	S	A	Good digestion/ boosting immunity	Cooking	O	malaria, diabetes, reduce fats, dysentery, diarrhea, pile and epilepsy (Rajeshwari, 2012; Fitri et al., 2020; Ogunjinmi et al., 2009)	0.95
74	sunakhari	<i>Dendrobium amoenum</i>	Orchidaceae	Flower	W	H	A	Immunity power	Juice extraction	O	treat wounds, skin disease, reduce fever, nourish stomach (Wang., 2021; Ng et al., 2012; Xu et al., 2013) al. 2013)	0.95
75	Jai phul	<i>Jasminum</i> sp.	Oleaceae	Bud/ leaf	C	CL	R	Sore throat/stomach problem	Raw leaves/bud chewing	O		0.92
76	Tite karela	<i>Momordica charantia</i>	Cucurbitaceae	Fruit	C	CL	A	Blood pressure/diabetes	Cooking	O		0.92
77	Mewa	<i>Carica papaya</i> L.	Caricaceae	Latex/ root/ fruit	C	H	A	Indigestion/ kidney stone	Juice extraction	O	maintain blood pressure, used as vegetable or pickle, anemia, urinary discharges, asthma, ulcers, bronchitis (ambu et al., 2020; Kumar & Bhowmik, 2010; Anilakumar et al., 2015)	0.1
78	Majito	<i>Rubia manjith</i> Roxb. ex Fleming	Rubiaceae	Root	W	CL	A	Fever/chest pain	Powder/ paste	O	constipation, toothache, increase in urine output, intestinal parasites (Ayyanar & Ignacimuthu, 2011; Vij & Prashar, 2015)	0.16
79	Halhale	<i>Rumex nepalensis</i> Spreng.	Polygonaceae	Root	W/C	H	A	Cut/wounds/ring worm	paste	O		0.84

80	Dudhe lahara	<i>Ficus hederacea</i> Roxb.	Moraceae	Whole plants	W	CL	R	Body pain/bone fracture/diabetes	Juice extraction	O		0.86
81	Banmara	<i>Ageratina adenophora</i> (Spreng.) King & H. Rob	Asteraceae	Leaf	W	H	A	Cut/wounds	Paste/powder	T	Treatment of cuts and wounds, arthritis, rheumatism, indigestion, diarrhea, fever reduction (Badola & Pradhan, 2013; Bhattarai, 2017; Chaudhery et al., 2015;	1.45
82	Haadchur	<i>Viscum album</i> L.	Loranthaceae	Whole plants	W	T	R	Bone fracture/body pain/ irregular mensuration	Juice extraction	O		0.33
83	Makai	<i>Zea mays</i> L.* (RKM61-G1)	Poaceae	fruits	C	H	A	diabetes	decoction	O	given to buffaloes to stimulate milk production, stomach pain, dermatitis, diuretic, kidney ailment (Ambu et al., 2020; Tene et al., 2006; Manuel et al., 2004)	0.02
84	Bhutkesh	<i>Selinum wallichianum</i> (DC.) Raizada & Saxena	Apiaceae	Root	W	H	R	Stomach ache/common cold/blood pressure	powder	O	Cold, fatigue, nervousness (Bhandari et al., 2021; Ghimire et al., 2018)	0.65
85	Bishjara	<i>Aconitum ferox</i> L.	Ranunculaceae	Root	W	H	A	Snakebite/ diarrhea/ dysentery headache/cough/ cold	Powder/ paste	O	antibacterial, diuretic (Joshi, 2013)	1.09
86	Bukki phul	<i>Anaphalis contorta</i>	Asteraceae	Whole	W	H	R	Cold/cough/ gastritis	decoction	O		0.36
87	Ghiukumari	<i>Aloe vera</i> (L.) Burm.f.	Liliaceae	Leaf juice	C	H	R	Burn/uric acid/piles/constipation/ gastritis	Juice extraction	O/T	burn treatments, Treat constipation, diabetes, dermatitis (Manvitha & Bidya, 2014; Jones, 2007; Kathi & Victoria, 1999)	1.46
88	Jaringo	<i>Phytolacca americana</i> Roxb.	Phytolaccaae	Root/ stem/ leaf	W	H	R	Stomach problems	Cooking	O	Anorexia, Stomach pain, hepatic pain (Tene et al., 2006; Ambu et al., 2020)	1.13
89	Gogan	<i>Sauraria nepalensis</i> Roxb.	Saurariceae	Bark	W	T	A	Fever	Juice extraction	O		0.05

90	Gande jhar	<i>Ageratum conyzoides L.</i>	Asteraceae	Leaf	W	H	A	Cut/wound	Paste	O	pneumonia, asthma, gynecological diseases, skin diseases and wound healing, diarrhea, headaches (Okunade, 2002; Ming, 1999, Marques et al., 1988; Negrelle et al., 1988)	0.04
91	Alainchi	<i>Amomum subulatum Roxb.</i>	Zingiberaceae	Fruit	C	H	A	Nausea/indigestion	Extract in powder form/Mix with Tea	O	indigestion, reduce stomach cramps, reduce pain and swelling, dental health, stress relief (Bisht et al., 2011; Parveen et al., 2018)	0.33
92	Tulsi	<i>Ocimum tenuiflorum L.</i>	Lamiaceae	Whole Plants	C	H	R	Vomiting/joint pain/urinary diseases/toothache	Juice extraction/ powder	O	gastric problems, Blood pressure control, respiratory diseases, typhoid (Bhattarai et al., 2009; Ambu et al., 2020)	1.55
93	Rato akhle	<i>Achyranthes aspera L.</i>	Amaranthaceae	Root/ stem	W	H	A	Bleeding/dog bite/snake bite	Paste	T		0.77
94	Koiralo	<i>Bauhinia variegata (L.) Benth.</i>	Fabaceae	Bark/ flower	W	T	A	Dysentery/ diabetes/ constipation/diarrhea/ indigestion	Juice extraction	O	diarrhoea, dysentery, piles, gastritis, toothache, goiter (Ambu et al., 2020; Tamang, 2003; Thapa, 2012)	0.06
95	Kalo siris	<i>Albizia lebbek (L.) Benth.</i>	Fabaceae	Bark	W	T	A	snake bite/scorpion sting/ diarrhea	Juice extraction	O	Gingivitis, fish killing (Bhandari et al., 2021; Kichu et al., 2015)	1.5
96	Chutro	<i>Berberis aristata DC.</i>	Berberidaceae	Bark/ leaf	W	S	A	skin disease/ jaundice/ piles	paste/ juice extraction	O/T	kill intestinal worms, jaundice, treat fever, blood purification, toothache (Rana et al., 2015; Bhandari et al., 2021; Shrestha & Dhillion, 2003; Ambu et al., 2020)	1.4
97	Paiyu	<i>Prunus cerasoides D. Don.</i>	Rosaceae	latex	W	T	A	Jaundice	Juice extraction	O		0.06
98	Kodo	<i>Eleusine coracana (L.) Gaertn.</i>	Poaceae	seed	C	S	A	Diarrhea	fermentation	O	Gingivitis (Bhandari et al., 2021)	0.14
99	Siru	<i>Imperata cylindrica (L.) P. Beauv.</i>	Poaceae	root	W	H	A	Worms/ fever/ diarrhea	Juice extraction	O	anti-helminthic, indigestion, boils, diarrhea, dysentery, fever (Dhami, 2008; Bhatt & Shakya, 2015; Bhatt & Kunwar, 2020).	0.04

100	Farsi	<i>Cucurbita moschanta Duchesne ex Poir.</i>	Cucurbitaceae	Leaf	C	CL	A	Wounds cut	Paste	T	Antiparasite, diabetes, tumor, cholesterol, inflammation (Tene et al., 2006; Caili et al., 2006; Yang et al., 2002)	0.33
101	Kimbu	<i>Morus alba L.</i>	Moraceae	Leaf/fruit	W/C	T	R	Blood clotting/diabetes	Juice extraction	O	asthma, cough, bronchitis, constipation, wound healing, diabetes, burning sensation (Nadkarni, 1976; Anonymous, 1952; Arya, 1997)	0.1
102	Aaru	<i>Prunus persica Siebold & Zucc. 1845</i>	Rosaceae	Fruit/Barks	C	T	A	Wounds/stomach ache	Juice extraction	O	diabetes, obesity, hypertension, inflammation, cardiovascular diseases (Bento et al., 2022; Kant et al., 2018)	0.22

W= Wild
C= Cultivated
H= Herbs
S= Shrubs

O= Oral
T= Topical
I= Inhalation

T= Tree
CL= Climbers

A= Abundant
R= Rare

3.3. Life Form of Medicinal Plants:

Medicinal plants in the study area predominantly consist of herbs, which account for 52.94% (54 species) of the total documented flora. Trees represent 21.56% (22 species) of the medicinal plants, while climbers and shrubs constitute 15.68% (16 species) and 9.80% (10 species) respectively.





Figure 3: Life Form of Medicinal Plant

3.4. Parts Used:

Various parts of medicinal plants have been utilized to treat different ailments through a range of application methods (Figure 4). The root was the most frequently used part, accounting for 21.31% (26 species), followed by leaves at 18.86% (23 species). Whole plants were employed in 15.57% (19 species) of cases, while fruits were used in 12.30% (15 species). Stems and bark each contributed 7.4% (9 species), with rhizomes and seeds at 6.6% (8 species) and 4.1% (5 species)



respectively. Latex and flowers were the least utilized, at 3.3% (4 species) each.

Figure 4: Parts used in Medicinal Plants

3.5. Mode of Administration

The mode of administration of medicinal plants varied according to the species (Figure 5). Approximately 72.54% (74 species) are administered orally, either in raw form, as juice, or as a decoction. A smaller proportion, 7.84% (8 species), are used both orally and topically. Topical application alone accounts for 18.62% (19 species). Inhalation is the least common method, employed for only 0.98% (1 species). These varied methods highlight the versatility and depth of traditional medicinal application practices in the study area.




Figure 5: Mode of Administration

3.6 Number of Plants Treating a Disease:

The gastrointestinal disorders are the most treated category with 54 plants cited. Skin-related issues (cuts, wounds, burns) and headaches/fevers also have high citation counts which is 21 and 22 plants respectively. Other conditions like Common cold and Cough (15 plants) and respiratory issues (11 plants) have moderate representation, while categories like urologic disorders and general health with 2 plants each have minimal citations.




Figure 6: Number of Plants Treating a Disease

3.7. Fidelity Level Calculation

The study identifies several plant species with high Fidelity Levels (FL), indicating a strong preference by the local community for treating specific ailments (Table 2). Notably, *Aconitum bisma* and *Swertia chirayita* achieve an FL of 100% for treating fever and typhoid, while *Cissus*

quadrangularis also reaches 100% for bone fractures. In gastrointestinal disorders, Zanthoxylum acanthopodium shows a perfect FL for gastric issues, and Ageratina adenophora reaches 100% for treating cuts. In contrast, Cissus adnata has the lowest FL at 30.8% for treating eye cataracts, indicating less selection among informants regarding its use for this condition.

Table 2: Fidelity level (FL) of medicinal plants of the study area

Disease Category	Diseases Treated	Total Species	Diseases for Fidelity Level Calculation	Species Fidelity Level Calculation	Np	N	Fidelity Level (FL) (%)
Headache and Fever	Headache, Fever, Typhoid	20	Fever	<i>Paris polyphylla</i>	86	98	87.8
				<i>Aconitum bisma</i>	76	76	100
				<i>Swertia chirayita</i>	82	82	100
Skeletal Problems	Bone fracture, Ligament Sprain, Body pain	18	Bone fracture	<i>Bergenia ciliate</i>	56	68	82.4
				<i>Cissus quadrangularis</i>	42	42	100
			Ligament sprain	<i>Periploca calophylla</i>	24	46	52.2
				<i>Pleione praecox</i>	38	59	64.4
Gastrointestinal Disorders	Gastric, Constipation, Dysentery, Diarrhea, Jaundice	54	Gastric	<i>Paris polyphylla</i>	79	86	91.9
				<i>Zanthoxylum acanthopodium</i>	87	87	100
			Constipation	<i>Aloe vera</i>	28	53	52.8
				<i>Bauhinia variegata</i>	7	13	53.8
			Dysentery	<i>Pyrus pashia</i>	32	45	71.1
Skin Problems	Cuts, Wounds, and Burns	21	Cuts	<i>Ageratina adenophora</i>	46	46	100
				<i>Rumex nepalensis</i>	4	9	44.4
			Burns	<i>Parmelia nepalensis</i>	8	11	72.7
Common cold and Cough	Common cold, cough, and tonsillitis	15	Cough & cold	<i>Lindera neesiana</i>	19	19	100
				<i>Zingiber officinale</i>	64	64	100
Dermatological Disorders	Scabies, Skin Allergy, Vitiligo	9	Skin allergy	<i>Acorus calamus</i>	11	19	57.9
			Vitiligo	<i>Clematis buchananiana</i>	3	8	37.5
Snake Bite	Snake Bite	9	Snake bite	<i>Aconitum bisma</i>	67	67	100
				<i>Lindera neesiana</i>	36	42	85.7

Respiratory Diseases	Sore throat, Asthma	11	Asthma	<i>Zingiber officinale</i>	12	15	80
Odontological Disorders	Toothache, blood in gums	3	Toothache	<i>Cinnamomum glaucescens</i>	16	23	69.7
Cardio Vascular Diseases	Heart diseases, High/Low blood pressure	8	Blood pressure	<i>Momordica charantia</i>	18	27	66.6
				<i>Swertia chirayita</i>	88	92	95.7
Ophthalmic Diseases	Eye disease, eye cataracts, poor eyesight	4	Eye cataract	<i>Cissus adnata</i>	4	13	30.8
Mental Disorders (Neurological)	Epilepsy, nervous problem Unconsciousness	3	Nerves problem	<i>Valeriana jatamansi</i>	12	14	85.7
Metabolic Disorder	Diabetes	7	Diabetes	<i>Taxus wallichiana</i>	21	32	65.6
Gynecological Disorders	Pregnancy problems, mensuration problem	7	Pregnancy problems	<i>Rhododendron arboretum</i>	38	38	100
Immunity Boosters	Overall Immune system	4	Not specific	<i>Piper longum</i>	17	19	89.5
Urologic Disorders	Burning urination	2	Burning urination	<i>Chilocostus speciosus</i>	2	2	100
General	Kidney stone, uric acid	3	Uric acid	<i>Rauvolfia serpentina</i>	31	34	91.2

3.8 Informants Consensus Factor (ICF)

The Informant Consensus Factor (ICF) is found to range between 0.11 to 0.75. Urologic disorders have the highest ICF value of 0.75, based on 5 use reports for 2 species, followed by Gastrointestinal disorders with an ICF of 0.51, derived from 109 use reports for 54 species. In contrast, the category of snake bites exhibits the lowest ICF value of 0.11, with 10 use reports spread across nine plant species

Table 3: ICF values for medicinal plants used to cure a variety of ailments.

Disease Category	No. of species (Nt)	Use reports (Nur)	Informants Consensus Factor (ICF)
Headache and Fever	20	26	0.24
Skeletal Problems	18	21	0.15
Gastrointestinal Disorders	54	109	0.51
Skin Problems	21	30	0.31
Common cold and Cough	15	22	0.33

Dermatological Disorders	9	12	0.27
Snake Bite	9	10	0.11
Respiratory Diseases	11	17	0.38
Odontological Disorders	3	5	0.5
Cardio Vascular Diseases	8	10	0.22
Ophthalmic Diseases	4	5	0.25
Mental Disorders	3	4	0.33
Metabolic Disorder	7	9	0.25
Gynecological Disorders	7	11	0.40
Immunity Boosters	4	5	0.25
Urologic Disorders	2	5	0.75
General	3	5	0.5

3.9. Relative Frequency Citation (RFC)

Paris polyphylla has the greatest Relative Frequency of Citation (RFC) with a value of 0.98, indicating that the informants are widely aware of and utilize it. On the other hand, *Chilocostus speciosus* has the lowest RFC (0.02), indicating that it is not often cited and probably has little importance in the community. This contrast highlights the differences in the relative importance of the various plant species within the study area.

Table 4: Relative Frequency Citation (RFC) of the medicinal plants

Plant Species	FC	N	RFC
<i>Paris polyphylla</i>	98	100	0.98
<i>Aconitum bisma</i>	76	100	0.76
<i>Swertia chirayita</i>	82	100	0.82
<i>Bergenia ciliate</i>	68	100	0.68
<i>Cissus quadrangularis</i>	42	100	0.42
<i>Periploca calophylla</i>	46	100	0.46
<i>Pleione praecox</i>	59	100	0.59
<i>Paris polyphylla</i>	86	100	0.86
<i>Zanthoxylum acanthopodium</i>	87	100	0.87
<i>Aloe vera</i>	53	100	0.53
<i>Bauhinia variegata</i>	13	100	0.13

<i>Pyrus pashia</i>	45	100	0.45
<i>Ageratina adenophora</i>	46	100	0.46
<i>Rumex nepalensis</i>	9	100	0.09
<i>Permelia nepalensis</i>	11	100	0.11
<i>Lindera neesiana</i>	19	100	0.19
<i>Zingiber officinale</i>	64	100	0.64
<i>Acorus calamus</i>	19	100	0.19
<i>Clematis buchananiana</i>	8	100	0.08
<i>Aconitum bisma</i>	67	100	0.67
<i>Lindera neesiana</i>	42	100	0.42
<i>Zingiber officinale</i>	15	100	0.15
<i>Cinnamomum glaucescens</i>	23	100	0.23
<i>Momordica charantia</i>	27	100	0.27
<i>Swertia chirayita</i>	92	100	0.92
<i>Cissus adnata</i>	13	100	0.13
<i>Valeriana jatamansi</i>	14	100	0.14
<i>Taxus wallichiana</i>	32	100	0.32
<i>Rhododendron arboretum</i>	38	100	0.38
<i>Piper longum</i>	19	100	0.19
<i>Chilocostus speciosus</i>	2	100	0.02
<i>Rauwolfia serpentina</i>	34	100	0.34

4. DISCUSSION:

4.1. Diversity of Medicinal Plants and their Uses

This study identified a total of 102 plant species across 57 families, similar to the findings of Bhattarai & Khadka (2016) in Ilam, where 102 species from 94 genera and 64 families were reported to treat 56 human ailments. Bhattarai (2018) documented 30 species from 24 families and 29 genera within the Thami community in Ilam. In this study, the Rosaceae family has the highest representation by 7 species, followed by Poaceae with 6 species, aligning with the study

by Karki et al. (2023) in Dolakha District, where Rosaceae had 3 species. Additionally, 5 species each were found in the Ranunculaceae, Cucurbitaceae, Ericaceae, and Asteraceae families, while the Lauraceae, Moraceae, and Zingiberaceae families each had 3 species. Other families, including Saxifragaceae, Polygonaceae, Rutaceae, Orchidaceae, Liliaceae, Vitaceae, Urticaceae, Rubiaceae, Lamiaceae, Apiaceae, Piperaceae, Apocynaceae, and Fabaceae, were each represented by 2 species. These findings are inconsistent with a study in the Gulmi District (Acharya, 2012), where the Labiatae family had the highest representation with 8 species, followed by Leguminosae and Poaceae with 7 species each, and various other families with fewer species.

The current study revealed 17 different disease categories treated with medicinal plants, including a large number of species (54) used for gastrointestinal disorders, 21 for cuts, wounds, and burns, and approximately 20 for fever and headaches. The emphasis on gastrointestinal treatment is consistent with previous research, however the number of species varies. For instance, a study by Bhattarai & Khadka (2016) found 26 species for gastrointestinal difficulties, 17 for headaches and fever, and a more precise breakdown for other conditions like respiratory problems, dermatological issues, and snake stings. Another study by Acharya & Acharya (2007) diversifies the categories, identifying 36 species for gastrointestinal difficulties, 23 for Ear, Nose, and Throat (ENT) and ophthalmological concerns, and varied numbers for other conditions such as dermatological and genitourinary problems. The number of species employed for specific conditions varies throughout studies, which could be due to geographical variances in plant availability, traditional knowledge, and cultural practices. While gastrointestinal difficulties are always the most addressed, the distribution of plant use for other ailments shows a broader application of ethnomedicinal knowledge that is adapt to local health requirements and circumstances.

4.2. Life form, Plant parts used, and Mode of administration

The people in the study area primarily harvest herbaceous plants (52.94%) for remedy preparation. These could be due to ease of collecting, storing, transporting, and extracting active components from compared to other life forms (Shrestha & Dhillion, 2003). This finding is consistent with studies conducted in various parts of Nepal (Budha Magar et al., 2020; Luitel et

al., 2014; Ojha Khatri et al., 2021). The preference for herbaceous plants may also be due to their effectiveness and the relative ease of harvesting, transporting, and preparing remedies (Luitel et al., 2014; Mallik et al., 2020). Additionally, researches has shown that herbs often contain higher concentrations of secondary metabolites with therapeutic potential (Bhat & Karim, 2010; Mishra et al., 2014).

In the study area, the most frequently utilized plant parts are the root (21.31%), followed by leaves (18.86%). This pattern is consistent with other studies (Bhattarai & Khadka, 2016; Moore, 1994; Basualdo et al., 1995; Pradhan & Badola 2008; Chettri et al., 2014) in where underground parts like roots and rhizomes were heavily harvested. The preference for underground parts, such as roots, rhizomes, tubers, and bulbs, is often due to their higher concentration of bioactive compounds, as suggested by Srithi et al. (2009). However, the study conducted by Bhat et al. (2013), revealed that leaves were the most commonly used part (32%), followed by roots (24%) and the whole plant (13%). Study by Ahmad et al. (2014) in Pakistan also showed that leaves were mostly harvested plants in that study area. The consistent use of various plant parts across different areas demonstrates a common understanding of their therapeutic significance, while variances in utilization reflect each area's unique ecological and cultural environment.

Oral administration was the most common method of treatment, accounting for 72.54%, which suggests a higher prevalence of internal ailments in Kepilasgadhi Rural Municipality. However, careful attention to dosing is crucial for oral administration, as improper dosages can lead to severe internal complications. Similar findings have been reported in studies from Ethiopia (Kefalew et al., 2015; Tefera & Kim, 2019) and Nepal (Uprety et al., 2010; Rokaya et al., 2010), where oral application was the primary method of administration.

4.3. Fidelity Level (FL) and Informant Consensus Factor (ICF)

The study identified a total of 17 disease categories, with Informant Consensus Factor (ICF) values ranging from 0.11 to 0.75, and an average ICF of 0.33. This average is closely aligned with the findings of Yaseen et al. (2014) who reported an average ICF value of 0.38. The highest ICF value in our study was 0.75 for urologic disorders, which contrast to the significantly higher ICFs reported by Ragupathy et al. (2008) who found an ICF of 0.92 for jaundice, indicating strong consensus among informants for that ailment. In comparison, Uprety et al. (2010) reported

ICF values as high as 1.00 for ophthalmological problems, toothache, and kidney issues, reflecting very strong agreement among informants for these conditions. Rokaya et al. (2010), however, reported generally lower ICF values, such as an ICF of 0.40 for gastrointestinal ailments, which is notably lower than the 0.51 found in our study for the same category. The variations in ICF values across these studies may be attributed to the cultural and geographical diversity of the informant groups, and the extent of medicinal plant knowledge within the communities studied.

In this study, a high Fidelity Level (FL) of 100% was observed for ten plant species used to treat specific ailments. These include *Aconitum bisma* and *Swertia chirayita* for fever, *Cissus quadrangularis* for bone fractures, *Zanthoxylum acanthopodium* for gastric disorders, *Ageratina adenophora* for cuts, *Lindera neesiana* and *Zingiber officinale* for cold and cough, *Aconitum bisma* for snake bites, *Chilocostus speciosus* for burning urination, and *Rhododendron arboretum* for pregnancy problems. This high FL reflects a strong traditional reliance on these plants within our study area. Similarly, the study by Tumoro & Maryo (2016) also identified seven plant species with 100% FL. The high level of FL indicates that these plants have a long history of use in traditional medicine to treat certain diseases and are therefore expected to be trusted greatly by people in the community (Bibi et al., 2014). In contrast, *Cissus adnata* had the lowest FL of 30.8% for eye cataracts in our study, suggesting either less consensus or a wider variation in its use within the community, whereas Yaseen et al. (2014) reported lowest FL values of 50% for *Moringa oleifera* and *Sonchus asper* for specific ailments. *Moringa oleifera* was noted for a range of conditions, including eye disorders and gastric problems, while *Sonchus asper* was used for male sexual disorders and jaundice.

4.4. Use Value (UV) and Relative Frequency Citation (RFC)

The study reveals that RFC values for the plant species ranged from 0.02 to 0.98, with *Paris polyphylla* having the highest RFC value of 0.98, followed by *Swertia chirayita* (0.92) and *Zanthoxylum acanthopodium* (0.87). The high RFC values for these species can be attributed to their widespread distribution, easy availability, and traditional use in treating numerous diseases (Kayani et al., 2014). These high citation levels indicate that these species are not only well-known but also highly trusted and widely used within the community. Species of plants with high RFC values should be prioritized for pharmacological, phytochemical, and other biological

research in order to confirm and validate their medicinal properties (Mukherjee et al., 2012). Conversely, plants like *Chilocostus speciosus* (0.02), *Clematis buchananiana* (0.08), and *Rumex nepalensis* (0.09) exhibit much lower RFC values, suggesting that they are either less recognized or less frequently used for medicinal purposes. In a similar study by Faruque et al. (2018), RFC values ranged from 0.02 to 0.25, with the highest RFC recorded for *Rauvolfia serpentina* (0.25) and the lowest for *Scoparia dulcis* (0.20) indicating that the ailments treated by these plants may be less common or there is less consensus on their effectiveness in the area.

The range of Use Values (UV) from 0.01 to 1.83 reflects the diversity in the importance and frequency of use of various medicinal plants. Plants with higher Use Values, such as *Curcuma domestica* L with a UV of 1.83, are frequently utilized and highly valued for treating multiple ailments, including body pain, fever, sore throat, common cold, and jaundice. This indicates that they are integral to local medicinal practices and are often employed in traditional remedies. In contrast, plants with lower Use Values, like *Cinnamomum glaucescens* (UV of 0.01), are cited less frequently and may play a less significant role in local medicinal practices.

4.5. Medicinal plant trade, threats, and conservation practices

This area presents significant trade opportunities for medicinal and aromatic plants (MAPs), with several high-value species with good market value available, such as *Paris polyphylla*, *Swertia chiraita*, *Lindera neesiana*, *Berberis aristata*, *Astilbe rivularis*, *Piper longum*, *Bergenia ciliata*, and *Aconitum bisma*. Among these, *Paris polyphylla* and *Swertia chiraita* are in particularly high demand according to local traders. Many of the traders also cultivate and harvest these medicinal plants themselves. However, the sustainability of these resources is under threat due to unsustainable harvesting practices, animal grazing, forest fires, deforestation, illegal collection, and the impact of the COVID-19 pandemic. During the pandemic, widespread unemployment in the village led many to turn to the irregular collection and sale of MAPs, contributing to the decline of these resources in the forest. Similar threats have been documented in other studies, such as Chaudhary (1998), which identified deforestation, habitat encroachment, shifting cultivation, and forest fires as significant challenges, and Bibi et al. (2014), which highlighted grazing, urbanization, and the uprooting of medicinal plants as serious concerns.

To mitigate the threats to medicinal and aromatic plants (MAPs), villagers have adopted several conservation practices, including the domestic cultivation of high-demand species like *Paris polyphylla* and *Swertia chiraita* to reduce pressure on wild populations and maximize economic benefits. They also engage in reforestation efforts, sustainable harvesting techniques, such as selective and seasonal harvesting. Community-based forest management is also practiced, along with strict enforcement of rules against illegal collection.

5. CONCLUSION:

The study reveals that Kepilasgadhi Rural Municipality is exceptionally rich in medicinal plants, with some community members possessing extensive knowledge about their uses; however, these plants remain largely unexplored. This knowledge of medicinal plants, is mainly passed down through generations but, nowadays there is a noticeable decline in interest among younger generations regarding traditional medicinal plants. This shift interest poses a risk to the preservation of this valuable indigenous knowledge, urgent need for initiatives to engage and educate younger community members about the importance and benefits of traditional medicinal practices. Additionally, the medicinal plants in the study area are threatened by unsustainable harvesting practices, animal grazing, forest fires, deforestation, and illegal collection. Addressing these threats is crucial for maintaining the ecological balance, ensuring economic benefits, and conserving these vital resources for continued use.

ACKNOWLEDGEMENT

The authors would like to acknowledge the local people of the area for permitting us to conduct this study and for their invaluable assistance with providing essential information. We also extend our gratitude to the Division Forest Office, Khotang, Nepal, for their kind cooperation, support, and valuable suggestions throughout the study. Additionally, we are thankful to all the anonymous reviewers who contributed to the drafting of the manuscript.

CONFLICT OF INTEREST:

The Authors declare no conflict of interest.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES:

- Abbas SR, Bano S. Significance of pyrus communis as a medicinal plant [A Review].
- Abbas Z, Khan SM, Alam J, Khan SW, Abbasi AM. Medicinal plants used by inhabitants of the Shigar Valley, Baltistan region of Karakorum range-Pakistan. Journal of ethnobiology and ethnomedicine. 2017 Dec;13:1-5.
- Abe R, Ohtani K. An ethnobotanical study of medicinal plants and traditional therapies on Batan Island, the Philippines. Journal of ethnopharmacology. 2013 Jan 30;145(2):554-65.
- Acharya R, Acharya KP. Ethnobotanical study of medicinal plants used by Tharu community of Parroha VDC, Rupandehi district, Nepal. Scientific world. 2009;7(7):80-4.
- Acharya R. Ethnobotanical study of medicinal plants of Resunga Hill used by Magar community of Badagaun VDC, Gulmi district, Nepal. Scientific world. 2012 Sep 20;10(10):54-65.

- Acharya R. Ethnobotanical study of medicinal plants of Resunga Hill used by Magar community of Badagaun VDC, Gulmi district, Nepal. *Scientific world*. 2012 Sep 20;10(10):54-65.
- Adurosakin OE, Iweala EJ, Otike JO, Dike ED, Uche ME, Owanta JI, Ugbogu OC, Chinedu SN, Ugbogu EA. Ethnomedicinal uses, phytochemistry, pharmacological activities and toxicological effects of *Mimosa pudica*-A review. *Pharmacological Research-Modern Chinese Medicine*. 2023 Jun 1;7:100241.
- Ahmad H, Sehgal S, Mishra A, Gupta R. *Mimosa pudica* L.(Laajvanti): an overview. *Pharmacognosy reviews*. 2012 Jul;6(12):115.
- Ahmad M, Sultana S, Fazl-i-Hadi S, Ben Hadda T, Rashid S, Zafar M, Khan MA, Khan MP, Yaseen G. An ethnobotanical study of medicinal plants in high mountainous region of Chail valley (District Swat-Pakistan). *Journal of ethnobiology and ethnomedicine*. 2014 Dec;10:1-8.
- Ambu G, Chaudhary RP, Mariotti M, Cornara L. Traditional uses of medicinal plants by ethnic people in the Kavrepalanchok district, Central Nepal. *Plants*. 2020 Jun 17;9(6):759.
- Anilakumar KR, Kumar GP, Ilaiyaraja N. Nutritional, pharmacological and medicinal properties of *Momordica charantia*. *International Journal of Nutrition and Food Sciences*. 2015 Jan 1;4(1):75-83.
- Anonymous. *The Wealth of India, A Dictionary of Indian Raw materials*. Vol. 7. New Delhi: Council of Scientific and Industrial Research; 1952. p. 429-37
- Arya VS. *Indian Medicinal Plant*. Vol. 14. Chennai: Orient Longman Limited; 1997. p. 65-67.
- Asase A, Akwetey GA, Achel DG. Ethnopharmacological use of herbal remedies for the treatment of malaria in the Dangme West District of Ghana. *Journal of ethnopharmacology*. 2010 Jun 16;129(3):367-76.
- Ayyanar M, Ignacimuthu S. Ethnobotanical survey of medicinal plants commonly used by Kani tribals in Tirunelveli hills of Western Ghats, India. *Journal of ethnopharmacology*. 2011 Apr 12;134(3):851-64.

- Aziz MA, Khan AH, Adnan M, Ullah H. Traditional uses of medicinal plants used by Indigenous communities for veterinary practices at Bajaur Agency, Pakistan. *Journal of ethnobiology and ethnomedicine*. 2018 Dec;14:1-8.
- Badola HK, Pradhan BK. Plants used in healthcare practices by limboo tribe in south-west of Khangchendzonga biosphere reserve, Sikkim, India.
- Baniya AK, Karki S, Dhungana SM, Rizal G. Economic analysis on the production of large cardamom in Panchthar, Nepal.
- Baral HS, Inskipp C. Important Bird Areas in Nepal: key sites for conservation. *Bird Conservation Nepal*; 2005.
- Baral SR, Kurmi PP. A compendium of medicinal plants in Nepal. (No Title). 2006.
- Basualdo I, Zardini EM, Ortiz M. Medicinal plants of Paraguay: underground organs, II. *Economic Botany*. 1995 Oct 1:387-94.
- Bennett BC, Prance GT. Introduced plants in the indigenous pharmacopoeia of Northern South America. *Economic botany*. 2000 Jan 1:90-102.
- Bento C, Goncalves AC, Silva B, Silva LR. Peach (*Prunus persica*): Phytochemicals and health benefits. *Food Reviews International*. 2022 Nov 17;38(8):1703-34.
- Bhandari P, Gurung MB, Subedi CK, Chaudhary RP, Basnet K, Gurung J, Uprety Y, Neupane A, Shrestha KK. Traditional use of medicinal plants in the Chyangthapu-Phalaicha biological sub-corridor, Panchthar District, Kangchenjunga Landscape, Nepal. *Ethnobotany Research and Applications*. 2021 Oct 9;22:1-43.
- Bhat JA, Kumar M, Bussmann RW. Ecological status and traditional knowledge of medicinal plants in Kedarnath Wildlife Sanctuary of Garhwal Himalaya, India. *Journal of Ethnobiology and Ethnomedicine*. 2013 Dec;9:1-8.
- Bhat R, Karim AA. Tongkat Ali (*Eurycoma longifolia* Jack): a review on its ethnobotany and pharmacological importance. *Fitoterapia*. 2010 Oct 1;81(7):669-79.

- Bhatt MD, Kunwar RM. Distribution pattern and ethnomedicinal uses of plants in Kanchanpur district, Far-Western Nepal. *Ethnobotany Research and Applications*. 2020 Jul 24;20:1-21.
- Bhatt MD, Shakya BS. Medicinal plants in buffer zone area of Shuklaphanta Wildlife Reserve. *Himalayan Biodiversity*. 2015;3:20-30.
- Bhatt VP, Negi GC. Ethnomedicinal plant resources of Jaunsari tribe of Garhwal Himalaya, Uttaranchal.
- Bhatta LR. Ethnobotanical study in a village at Rukum district, Nepal. *Banko Janakari*. 1999;9(2):40-3.
- Bhattarai KR, Khadka MK. Ethnobotanical survey of medicinal plants from Ilam District, East Nepal. *Our Nature*. 2016 Jan 1;14(1).
- Bhattarai KR. Ethnobotanical study of plants used by Thami community in Ilam District, eastern Nepal. *Our Nature*. 2018 Dec 1;16(1).
- Bhattarai KR. Ethnobotanical survey on plants used in Mai Municipality of Ilam district, eastern Nepal. *Banko Janakari*. 2020 Dec 17;30(2):11-35.
- Bhattarai KR. Ethnomedicinal practices of the Lepcha community in Ilam, East Nepal. *Journal of Plant Resources*. 2017 Apr;15(1):31-44.
- Bhattarai S, Chaudhary RP, Quave CL, Taylor RS. The use of medicinal plants in the trans-himalayan arid zone of Mustang district, Nepal. *Journal of Ethnobiology and Ethnomedicine*. 2010 Dec;6:1-1.
- Bibi T, Ahmad M, Tareen RB, Tareen NM, Jabeen R, Rehman SU, Sultana S, Zafar M, Yaseen G. Ethnobotany of medicinal plants in district Mastung of Balochistan Province-Pakistan. *Journal of ethnopharmacology*. 2014 Nov 18;157:79-89.
- Bisht VK, Negi JS, Bhandari AK, Sundriyal RC. *Amomum subulatum* Roxb: Traditional, phytochemical and biological activities-An overview. *Afr. J. Agric. Res*. 2011 Oct 26;6(24):5386-90.
- Bridson D, Forman L. *The Herbarium Handbook*. Royal Botanic Gardens, Kew.

- Budha-Magar S, Bhandari P, Ghimire SK. Ethno-medicinal survey of plants used by Magar (Kham) community, Rolpa district, Western Nepal. *Ethnobotany Research and Applications*. 2020 Feb 26;19:1-29.
- Caili FU, Huan S, Quanhong LI. A review on pharmacological activities and utilization technologies of pumpkin. *Plant foods for human nutrition*. 2006 Jun;61:70-7.
- CBS. National population and housing census 2021, National report. Kathmandu: National Planning Commission. 2012.
- Chakraborty U, Das H. Antidiabetic and antioxidant activities of *Cinnamomum tamala* leaf extracts in STZ-treated diabetic rats. *Global Journal of Biotechnology & Biochemistry*. 2010;5(1):12-8.
- Chase MW, Christenhusz MJ, Fay MF, Byng JW, Judd WS, Soltis DE, Mabberley DJ, Sennikov AN, Soltis PS, Stevens PF. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical journal of the Linnean Society*. 2016 May 1;181(1):1-20.
- Chaudhary RP, Uprety Y, Joshi SP, Shrestha KK, Basnet KB, Basnet G, Shrestha KR, Bhatta KP, Acharya KP, Chettri N. Kangchenjunga Landscape Nepal: from conservation and development perspectives.
- Chaudhary RP. Biodiversity in Nepal: status and conservation. S. Devi, Saharanpur. 1998.
- Chettri D, Moktan S, Das AP. Ethnobotanical studies on the tea garden workers of Darjiling Hills. *Pleione* 8 (1): 124. 2014;132.
- Chevallier A. The encyclopedia of medicinal plants. 1996.
- Cox PA, Balick MJ. The ethnobotanical approach to drug discovery. *Scientific American*. 1994 Jun 1;270(6):82-7.
- Dhami N. Ethnomedicinal uses of plants in Western Terai of Nepal: a case study of Dekhatbhuli Vdc of Kanchanpur District. *Medicinal plants in Nepal: an anthology of contemporary research*. 2008:165-77.

- Enejoh OS, Ogunyemi IO, Bala MS, Oruene IS, Suleiman MM, Ambali SF. Ethnomedical importance of *Citrus aurantifolia* (christm) swingle. *The Pharma Innovation*. 2015 Oct 1;4(8, Part A):1.
- Fabricant DS, Farnsworth NR. The value of plants used in traditional medicine for drug discovery. *Environmental health perspectives*. 2001 Mar;109(suppl 1):69-75.
- Farnsworth NR, Soejarto DD. Global importance of medicinal plants. *The conservation of medicinal plants*. 1991 Jul 26;26(26):25-51.
- Farooq A, Amjad MS, Ahmad K, Altaf M, Umair M, Abbasi AM. Ethnomedicinal knowledge of the rural communities of Dhirkot, Azad Jammu and Kashmir, Pakistan. *Journal of ethnobiology and ethnomedicine*. 2019 Dec;15:1-30.
- Faruque MO, Uddin SB, Barlow JW, Hu S, Dong S, Cai Q, Li X, Hu X. Quantitative ethnobotany of medicinal plants used by indigenous communities in the Bandarban District of Bangladesh. *Frontiers in pharmacology*. 2018 Feb 6;9:40.
- Fitri A, Asra R, Rivai H. Overview of the traditional, phytochemical, and pharmacological uses of gold bamboo (*Bambusa Vulgaris*). *World Journal of Pharmacy and Pharmaceutical Sciences*. 2020 Jun 10;9(8):21.
- Friedman J, Yaniv Z, Dafni A, Palewitch D. A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev Desert, Israel. *Journal of ethnopharmacology*. 1986 Jun 1;16(2-3):275-87.
- Ghimire K, Adhikari M, Uprety Y, Chaudhary RP. Ethnomedicinal use of plants by the highland communities of Kailash Sacred Landscape, Far-west Nepal. *Acad J Med Plants*. 2018;6(11):365-8.
- Ghimire SK. Medicinal plants in the Nepal Himalaya: current issues, sustainable harvesting, knowledge gaps and research priorities. *Medicinal Plants in Nepal: an anthology of contemporary research*. 2008:25-42.
- Gürdal B, Kültür Ş. An ethnobotanical study of medicinal plants in Marmaris (Muğla, Turkey). *Journal of ethnopharmacology*. 2013 Mar 7;146(1):113-26.

- Hamilton AC. Medicinal plants, conservation and livelihoods. *Biodiversity & Conservation*. 2004 Jul;13:1477-517.
- Heinrich M, Ankli A, Frei B, Weimann C, Sticher O. Medicinal plants in Mexico: Healers' consensus and cultural importance. *Social science & medicine*. 1998 Dec 1;47(11):1859-71.
- Hu R, Lin C, Xu W, Liu Y, Long C. Ethnobotanical study on medicinal plants used by Mulam people in Guangxi, China. *Journal of ethnobiology and ethnomedicine*. 2020 Dec;16:1-50.
- Islam R. Role of plant medicine in health care and improving nutritional standard in rural areas of Bangladesh. In *National Seminar on Diversity of Medicinal Plants and Their Sustainable Utilization in Health Care and Improving Nutritional Standard in Rural Areas (Kolkata) 2006* (pp. 1-30).
- Jan SA, Shinwari ZK, Zeb A, Khalil AT, Shah SH. Ethnobotany and research trends in *Trachyspermum ammi* L.(Ajowan); A popular folklore remedy. *American-Eurasian Journal of Agricultural & Environmental Sciences*. 2015 Apr 22;15:68-73.
- Jones K. Dietary Aloe Vera supplementation and glycemic control in diabetes. *B5 srl Nutracos 2007*; 6-9.
- Joseph B, George J, Mohan J. Pharmacology and traditional uses of *Mimosa pudica*. *International journal of pharmaceutical sciences and drug research*. 2013 Apr 1;5(2):41-4.
- Joshi RK. Essential oil of flowers of *Anaphalis contorta*, an aromatic and medicinal plant from India. *Natural Product Communications*. 2013 Feb;8(2):1934578X1300800224.
- Joshi S, Subedi PC. Phytochemical and biological studies on essential oil and leaf extracts of *Gaultheria fragrantissima* Wall. *Nepal Journal of Science and Technology*. 2013;14(2):59-64.
- Juyal D, Thawani V, Thaledi S, Joshi M. Ethnomedical properties of *Taxus wallichiana* zucc.(Himalayan yew). *Journal of traditional and complementary medicine*. 2014 Jul 1;4(3):159-61.
- Kadir MF, Sayeed MS, Mia MM. Ethnopharmacological survey of medicinal plants used by traditional healers in Bangladesh for gastrointestinal disorders. *Journal of Ethnopharmacology*. 2013 May 2;147(1):148-56.

- Kadir MF, Sayeed MS, Setu NI, Mostafa A, Mia MM. Ethnopharmacological survey of medicinal plants used by traditional health practitioners in Thanchi, Bandarban Hill Tracts, Bangladesh. *Journal of Ethnopharmacology*. 2014 Aug 8;155(1):495-508.
- Kant R, Shukla RK, Shukla A. A review on peach (*Prunus persica*): an asset of medicinal phytochemicals. *Int J Res Appl Sci Eng Technol*. 2018 Jan;6(1):2186-00.
- Karki S, Dhital AP, Uprety Y, Ghimire SK. Medicinal plants and their use by an ethnic minority Jirel in Dolakha district, Central Nepal. *Ethnobotany Research and Applications*. 2023 Feb 16;25:1-29.
- Kathi JK, Victoria C. The Longwood Herbal Task Force and the Center for Holistic Pediatric Education and Research. *Aloe vera (Aloe vera)* 1999;1-24
- Kayani S, Ahmad M, Sultana S, Shinwari ZK, Zafar M, Yaseen G, Hussain M, Bibi T. Ethnobotany of medicinal plants among the communities of Alpine and Sub-alpine regions of Pakistan. *Journal of ethnopharmacology*. 2015 Apr 22;164:186-202.
- Kefalew A, Asfaw Z, Kelbessa E. Ethnobotany of medicinal plants in Ada'a District, East Shewa Zone of Oromia regional state, Ethiopia. *Journal of ethnobiology and ethnomedicine*. 2015 Dec 11:1-28.
- Khan IA, Abourashed EA. *Leung's Encyclopedia of Common Natural Ingredients*. John Wiley and Sons Publication, New Jersey, 2010, 422-423. 19.
- Khanal DP, Raut B, Kafle M. A comparative study on phytochemical and biological activities of two *Grewia* species. *Journal of Manmohan Memorial Institute of Health Sciences*. 2016 Sep 26;2:53-60.
- Khushbu C, Roshni S, Anar P, Carol M, Mayuree P. Phytochemical and therapeutic potential of *Piper longum* Linn a review. *International journal of research in Ayurveda and pharmacy*. 2011 Jan;2(1):157-61.
- Kichu M, Malewska T, Akter K, Imchen I, Harrington D, Kohen J, Vemulpad SR, Jamie JF. An ethnobotanical study of medicinal plants of Chungtia village, Nagaland, India. *Journal of Ethnopharmacology*. 2015 May 26;166:5-17.

- Kumar KS, Bhowmik D. Traditional medicinal uses and therapeutic benefits of *Momordica charantia* Linn. *International Journal of Pharmaceutical Sciences Review and Research*. 2010;4(3):23-8.
- Kumar M, Sheikh MA, Bussmann RW. Ethnomedicinal and ecological status of plants in Garhwal Himalaya, India. *Journal of Ethnobiology and Ethnomedicine*. 2011 Dec;7:1-3.
- Kumar S, Kamboj J, Sharma S. Overview for various aspects of the health benefits of *Piper longum* linn. fruit. *Journal of acupuncture and meridian studies*. 2011 Jun 1;4(2):134-40.
- Kunow MA. *Maya Medicine: Traditional Healing in Yucatan*. UNM Press, New Mexico, 2003, 117.
- Kunwar RM, Burlakoti C, Chowdhary CL, Bussmann RW. Medicinal plants in Far-west Nepal: indigenous uses and pharmacological validity. *Med Aromat Plant Sci Biotechnol*. 2010;4(1):28–42
- Kunwar RM, Mahat L, Acharya RP, Bussmann RW. Medicinal plants, traditional medicine, markets and management in far-west Nepal. *Journal of ethnobiology and ethnomedicine*. 2013 Dec;9:1-0.
- Kunwar RM, Shrestha KP, Bussmann RW. Traditional herbal medicine in Far-west Nepal: a pharmacological appraisal. *Journal of ethnobiology and ethnomedicine*. 2010 Dec;6:1-8.
- Kutal DH, Kunwar RM, Uprety Y, Adhikari YP, Bhattarai S, Adhikari B, Kunwar LM, Bhatt MD, Bussmann RW. Selection of medicinal plants for traditional medicines in Nepal. *Journal of Ethnobiology and Ethnomedicine*. 2021 Dec;17:1-1.
- Luitel DR, Rokaya MB, Timsina B, Münzbergová Z. Medicinal plants used by the Tamang community in the Makawanpur district of central Nepal. *Journal of Ethnobiology and ethnomedicine*. 2014 Dec;10:1-1.
- Macía MJ, García E, Vidaurre PJ. An ethnobotanical survey of medicinal plants commercialized in the markets of La Paz and El Alto, Bolivia. *Journal of ethnopharmacology*. 2005 Feb 28;97(2):337-50.
- Malla, B.; Chhetri, R.B. Indigenous knowledge on ethnobotanical plants of Kavrepalanchowk District. *Kathmandu Univ. J. Sci. Eng. Technol*. 2009, 5, 96–109.

- Mallik AR, Chaudhary S, Shrestha S. Useful valuable plants of Maithili community in Eastern Nepal: An ethnobotanical study. *Bangladesh Journal of Plant Taxonomy*. 2020 Jul 1;27(2):439.
- Manandhar NP. A survey of medicinal plants of Jajarkot district, Nepal. *J Ethnopharmacol*. 1995a;48(1):1–6.
- Manjula BL, Mishra S, Benny ME, Kumar S. *Dioscorea pentaphylla*: A pharmacological agent. *Yam*. 2022:9-17.
- Manvitha K, Bidya B. Aloe vera: a wonder plant its history, cultivation and medicinal uses. *Journal of Pharmacognosy and Phytochemistry*. 2014;2(5):85-8.
- Marques Neto JF, Costallat LT, Fernandes SR, Napoli MD, Samara AM. Efeitos do *Ageratum conyzoides*, Linèe no tratamento da artrose. *Rev. bras. reumatol*. 1988:109-4.
- Mbuni YM, Wang S, Mwangi BN, Mbari NJ, Musili PM, Walter NO, Hu G, Zhou Y, Wang Q. Medicinal plants and their traditional uses in local communities around Cherangani Hills, Western Kenya. *Plants*. 2020 Mar 5;9(3):331.
- Mesfin F, Demissew S, Teklehaymanot T. An ethnobotanical study of medicinal plants in Wonago Woreda, SNNPR, Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2009 Dec;5:1-8.
- Ming LC. *Ageratum conyzoides*: A tropical source of medicinal and agricultural products. Perspectives on new crops and new uses. 1999;1988:469-73.
- Mishra S, Aeri V, Gaur PK, Jachak SM. Phytochemical, therapeutic, and ethnopharmacological overview for a traditionally important herb: *Boerhavia diffusa* Linn. *BioMed research international*. 2014;2014(1):808302.
- Moore PD. Trials in bad taste. *Nature*. 1994 Dec 1;372(6505).
- Mukherjee PK, Nema NK, Venkatesh P, Debnath PK. Changing scenario for promotion and development of Ayurveda—way forward. *Journal of ethnopharmacology*. 2012 Sep 28;143(2):424-34.

- Murad W, Ahmad A, Gilani SA, Khan MA. Indigenous knowledge and folk use of medicinal plants by the tribal communities of Hazar Nao Forest, Malakand District, North Pakistan. *Journal of Medicinal Plants Research*. 2011 Apr 4;5(7):1072-86.
- Nadkarni AK. *Indian Materia Medica*. Vol. 1. Mumbai: Popular Prakashan; 1976. p.1292-94
- Negrelle RR, Sbalchiero D, Cervi AC. Espécies vegetais utilizadas na terapêutica popular no município de Curitiba, Paraná. Brasil. Univ. Federal do Paraná. 1988.
- Ng TB, Liu J, Wong JH, Ye X, Wing Sze SC, Tong Y, Zhang KY. Review of research on *Dendrobium*, a prized folk medicine. *Applied microbiology and biotechnology*. 2012 Mar;93:1795-803.
- Ogunjinmi AA, Ijeomah HM, Aiyelaja AA. Socio-economic importance of bamboo (*Bambusa vulgaris*) in Borgu local government area of Niger State, Nigeria. *Journal of Sustainable Development in Africa*. 2009;10(4):284-98.
- Ojha Khatri S, Chaudhary S, Shrestha N, Munankarmi NN. Ethnomedicinal study and phytochemical screening of selected plants in Jhule, Dolakha District, Nepal. *Vegetos*. 2021 Dec;34(4):834-46.
- Okunade AL. *Ageratum conyzoides* L.(asteraceae). *Fitoterapia*. 2002 Feb 1;73(1):1-6.
- Owoyele BV, Nafiu AB, Oyewole IA, Oyewole LA, Soladoye AO. Studies on the analgesic, anti-inflammatory and antipyretic effects of *Parquetina nigrescens* leaf extract. *Journal of ethnopharmacology*. 2009 Feb 25;122(1):86-90.
- Parveen U, Maaz M, Mujeeb M, Jahangir U. Biological and therapeutic uses of *Amomum subulatum* Roxb: A. *European Journal of Biomedical*. 2018;5(1):167-76.
- Plotkin MJ, Famolare L. Conclusions and recommendations. *Sustainable Harvest and Marketing of Rain Forest Products*. Island Press, Washington, DC. 1992:310-3.
- Pradhan BK, Badola HK. Ethnomedicinal plant use by Lepcha tribe of Dzongu valley, bordering Khangchendzonga Biosphere Reserve, in north Sikkim, India. *Journal of Ethnobiology and Ethnomedicine*. 2008 Dec;4:1-8.

- Prakash OM, Chauhan AS, Kudachikar VB. Traditional uses, nutrition, phytochemistry and various pharmacological properties of Indian wild pear. *International Journal of Functional Nutrition*. 2021 Sep 1;2(4):1-3.
- Pranay Bantawa PB, Ritu Rai RR. Studies on ethnomedicinal plants used by traditional practitioners, Jhankri, Bijuwa and Phedangma in Darjeeling Himalaya.
- Ragupathy S, Steven NG, Maruthakkutti M, Velusamy B, Ul-Huda MM. Consensus of the 'Malasars' traditional aboriginal knowledge of medicinal plants in the Velliangiri holy hills, India. *Journal of Ethnobiology and Ethnomedicine*. 2008 Dec;4:1-4.
- Rai R, Singh NB. Medico-ethnobiology in Rai community: a case study from Baikunthe Village development committee, Bhojpur, eastern Nepal. *J Inst Sci Technol*. 2015;1:127–32.
- Rajeshwari E. Evaluation of anti-microbial activity of *Bambusa vulgaris* leaves. *International Journal of Phytotherapy Research*. 2012;2(2):36-9.
- Ramachandran A, Prasath R, Anand A. The medical uses of *Coccinia grandis* L. Voigt: a review. *Int J Pharmacogn*. 2014;1:681-90.
- Rana SK, Oli PS, Rana HK. Traditional botanical knowledge (TBK) on the use of medicinal plants in Sikles area, Nepal. *Asian J Plant Sci Res*. 2015;5(11):8-15.
- Rokaya MB, Münzbergová Z, Timsina B. Ethnobotanical study of medicinal plants from the Humla district of western Nepal. *Journal of Ethnopharmacology*. 2010 Aug 9;130(3):485-504.
- Shah M, Panchal M. Ethnopharmacological properties of *Cinnamomum tamala*—a review. *International Journal of Pharmaceutical Sciences Review and Research*. 2010 Nov;5(3):141-4.
- Shah, S.; Lamichhane, D. Documentation of Indigenous Knowledge on plants used by Tamang Community
- Sharma H, Garg M. A review of traditional use, phytoconstituents and biological activities of Himalayan yew, *Taxus wallichiana*. *Journal of integrative medicine*. 2015 Mar 1;13(2):80-90.
- Sher H, Midrarullah G, Coll PG. Medicinal plants of Udigram, District Swat, Pakistan. *Pakistan Journal of Forestry (Pakistan)*. 2003;53(1).

- Shrestha PM, Dhillion SS. Medicinal plant diversity and use in the highlands of Dolakha district, Nepal. *Journal of ethnopharmacology*. 2003 May 1;86(1):81-96.
- Shrestha, KK, Ghimire, SK, Gurung, TN, Lama, YC, Aumeeruddy-Thomas, Y. Conservation of plant resources, community development, and training in applied ethnobotany at Shey-Phoksundo National Park and its buffer zone, Dolpa. WWF Nepal Program Kathmandu, Nepal. 1998.
- Shyaula SL. Phytochemicals, traditional uses and processing of *Aconitum* species in Nepal. *Nepal Journal of Science and Technology*. 2011;12:171-8.
- Shyur LF, Huang CC, Lo CP, Chiu CY, Chen YP, Wang SY, Chang ST. Hepatoprotective phytochemicals from *Cryptomeria japonica* are potent modulators of inflammatory mediators. *Phytochemistry*. 2008 Apr 1;69(6):1348-58.
- Siddiqua A, Mittapally S. A review on *Cissus quadrangularis*. *The Pharma Innovation, Journal*. 2017 Jul 1;6:329-34.
- Singh A, Singh PK. An ethnobotanical study of medicinal plants in Chandauli District of Uttar Pradesh, India. *Journal of Ethnopharmacology*. 2009 Jan 21;121(2):324-9.
- Singh AG, Kumar A, Tewari DD, Bharati KA. New ethnomedicinal claims from Magar community of Palpa district, Nepal. 2018.
- Singh AG, Kumar A, Tewari DD. An ethnobotanical survey of medicinal plants used in Terai forest of western Nepal. *Journal of ethnobiology and ethnomedicine*. 2012 Dec;8:1-5.
- Singh AG. Medicinally important plants of Palpa used as blood purifier. *Himalayan Biodiversity*. 2014;3:35-41.
- Srithi K, Balslev H, Wangpakapattanawong P, Srisanga P, Trisonthi C. Medicinal plant knowledge and its erosion among the Mien (Yao) in northern Thailand. *Journal of ethnopharmacology*. 2009 Jun 22;123(2):335-42.
- Svarstad H, Dhillion SS. Responding to bioprospecting: rejection or regulation. Responding to Bioprospecting, from Biodiversity in the South to Medicines in the North. Spartacus Forlag As, Oslo. 2000:9-15.

- Tabuti JR, Dhillion SS, Lye KA. Traditional medicine in Bulamogi county, Uganda: its practitioners, users and viability. *Journal of Ethnopharmacology*. 2003 Mar 1;85(1):119-29.
- Tag H, Kalita P, Dwivedi P, Das AK, Namsa ND. Herbal medicines used in the treatment of diabetes mellitus in Arunachal Himalaya, northeast, India. *Journal of Ethnopharmacology*. 2012 Jun 14;141(3):786-95.
- Tamang, G. An ethnobiological study of Tamang people. *Our Nat*. 2003, 1, 37–41.
- Tamang, R.; Sedai, D.R. Documentation of Ethnomedicinal Knowledge on Plant Resources Used by Baram Community in Arupokhari VDC, Gorkha District, Central Nepal. *Bull. Dept. Plant Res*. 2016, 38, 60–64.
- Taylor RS, Manandhar NP, Towers GH. Screening of selected medicinal plants of Nepal for antimicrobial activities. *Journal of ethnopharmacology*. 1995 Jun 5;46(3):153-9.
- Tefera BN, Kim YD. Ethnobotanical study of medicinal plants in the Hawassa Zuria District, Sidama zone, Southern Ethiopia. *Journal of ethnobiology and ethnomedicine*. 2019 Dec;15:1-21.
- Tene V, Malagon O, Finzi PV, Vidari G, Armijos C, Zaragoza T. An ethnobotanical survey of medicinal plants used in Loja and Zamora-Chinchipe, Ecuador. *Journal of ethnopharmacology*. 2007 Apr 20;111(1):63-81.
- Thapa, S. Medico-ethnobotany of Magar Community in Salija VDC Parbat district, Central Nepal. *Our Nat*. 2012, 10, 176–190.
- Tisserand R, Balacs T. Nutmeg: A review of the toxicology and pharmacology of nutmeg and its essential oil. *Int. J. Aromather Winter*. 1994;6:28-38.
- Trotter RT, Logan MH. Informant consensus: a new approach for identifying potentially effective medicinal plants. In *Plants and indigenous medicine and diet* 2019 Dec 16 (pp. 91-112). Routledge.
- Tumoro G, Maryo M. Determination of informant consensus factor and fidelity level of ethnomedicinal plants used in Misha Woreda, Hadiya Zone, Southern Ethiopia. *International Journal of Biodiversity and Conservation*. 2016 Dec 31;8(12):351-64.

- Uprety Y, Asselin H, Boon EK, Yadav S, Shrestha KK. Indigenous use and bio-efficacy of medicinal plants in the Rasuwa District, Central Nepal. *Journal of ethnobiology and ethnomedicine*. 2010 Dec;6:1-0.
- van Andel T, Fundiko MC. The Trade in African Medicinal Plants in Matonge-Ixelles, Brussels (Belgium). *Economic Botany*. 2016 Dec;70:405-15.
- Vij T, Prashar Y. A review on medicinal properties of *Carica papaya* Linn. *Asian Pacific Journal of Tropical Disease*. 2015 Jan 1;5(1):1-6.
- Vitalini S, Iriti M, Puricelli C, Ciuchi D, Segale A, Fico G. Traditional knowledge on medicinal and food plants used in Val San Giacomo (Sondrio, Italy)—An alpine ethnobotanical study. *Journal of Ethnopharmacology*. 2013 Jan 30;145(2):517-29.
- Waisundara VY, Watawana MI, Jayawardena N. *Costus speciosus* and *Coccinia grandis*: Traditional medicinal remedies for diabetes. *South African Journal of Botany*. 2015 May 1;98:1-5.
- Wang YH. Traditional uses and pharmacologically active constituents of *Dendrobium* plants for dermatological disorders: a review. *Natural Products and Bioprospecting*. 2021 Oct;11:465-87.
- Xu J, Han QB, Li SL, Chen XJ, Wang XN, Zhao ZZ, Chen HB. Chemistry, bioactivity and quality control of *Dendrobium*, a commonly used tonic herb in traditional Chinese medicine. *Phytochemistry reviews*. 2013 Jun;12:341-67.
- Yang JO, Oh SR, Lee HK, Kim CJ, Song KB. Isolation of anticomplementary substances from *Cucurbita moschata* Duch. *Journal of food science*. 2002 May;67(4):1348-51.
- Yaseen G, Ahmad M, Sultana S, Alharrasi AS, Hussain J, Zafar M. Ethnobotany of medicinal plants in the Thar Desert (Sindh) of Pakistan. *Journal of ethnopharmacology*. 2015 Apr 2;163:43-59.