

# LC-MS BASED TARGETED METABOLITE PROFILING, AAS BASED MINERAL PROFILING IN METHANOL LEAF EXTRACT OF *Terminalia bellerica* (Gaertn.)

## ABSTRACT

**AIM:** Natural products in medicinal plants are one of the important sources for modern drug discovery. Plant-based formulations play an important role in management of numerous diseases. To discover the medical potential of natural products, it is important to understand the bioactive compounds present in medicinal plants. Hence in this study, it is planned to assay the volatile compounds and minerals in the leaves of *Terminalia bellerica*

**STUDY DESIGN:** In the present study, metabolites in leaves of *Terminalia bellerica* have been studied by AAS and LC-MS techniques.

**RESULTS:** AAS assay recorded the leaves of *T.bellerica* are rich in iron content (260mg/kg) when compared with Zn, Mn and copper content. LC-MS assay revealed chebulinic acid, chebulagic acid, punicafolin, myricetin rutine, galloyl punicalagin are in highest amount.

**CONCLUSION:** This plant could provide reservoir of iron that have greater therapeutic potential with fewer side effects than to synthetic drugs. Due to rich phyto nutrients in leaves of *T.bellerica*, it may be used in the development of health supplements. To develop new drugs in pharmaceutical industry, compounds in leaves of *T.bellerica* lead a pathway. Further isolation of novel compounds leads to the development of pharmacologically important drugs of specific function.

**Key words:** Medicinal herbs, *Terminalia bellerica* leaves, Phytoceutics, AAS assay, GC-MS assay.

## 1. Introduction:

Plants have served as an essential source of drugs and remedies against diseases and health disorders since ancient times. Health benefits may be credited to the presence of the various phytochemicals like polyphenols, terpenes, anthocyanins, flavonoids, alkaloids and glycosides. Plants of antimicrobial activity can be related to their essential oils (1) or other isolated compounds such as alkaloids (2), flavonoids, tannins, phenolic acids (3), among

other chemical classes including naturally occurring peptides (4,5). These compounds possess their antimicrobial activity through various mechanisms such as depressing the nuclear or ribosomal enzyme(s) synthesis, altering the membrane structure and the electron flow, or affecting the metabolic activity of the microbial cell, as well as inhibiting the secretion of their toxins (6,7,8).

Natural products from medicinal plants are one of the important sources for modern drug discovery. Triphala is an ayurvedic drug prepared by mixing dried fruit powder of *Terminalia chebula*, *Terminalia bellerica* and *Phyllanthus emblica* in the ratio of 1:1:1 and also its individual constituent's fruit powder exhibit immuno-modulatory activity. The important constituents of the Triphala are chebulagic acid (CA), chebulinic acid (CI), gallic acid. These compounds inhibit TNF $\alpha$  induced pro-angiogenic and pro-inflammatory activities in retinal capillary endothelial cells by inhibiting p38, ERK and NF $\kappa$ B phosphorylation. Polyphenol containing plants, of *Terminalia* species, have reportedly shown various health benefits and applications in pharma industry. *Terminalia* have a wide spectrum of pharmacological activities including antioxidant, anti-inflammatory, anti-cancer, hepatoprotective, and antimicrobial activities. Polyphenols have been also proven to lower the risk of cardiovascular diseases, enhance liver regeneration, and increase life expectancy.

Traditional medicinal plants practices are used for the treatment of infectious diseases and they have very low level risk compared to synthetic drugs. Natural compounds in medicinal plants are essential sources for drug discovery (9) and are reported that natural products take up to 35% of the global medicine market (10). To discover the medical values of natural products, it is important to understand the ethno pharmacological uses of various medicinal plants (11). Although the modern medicines is quickly growing, large population still preferring herbal medicines due to the effectiveness, medical effectiveness, enhancing cost of modern medicines and cultural preferences (12,13,14). Based on WHO reports, 80% of the global population depends on traditional medicine, and 60% of the Indian rural population use herbal medicines (14). These natural medicines are generally safe, cost-effective, and efficient (14). Apart from the medical values, plants are also widely used as food, health care products (15), veterinary medicine (16), have extensive impacts on daily life. Herbal medicine is still the mainstay of health care in several developing countries. The World Health Organization has estimated that more than 80 % of the world's population in developing countries depends primarily on herbal medicine for basic healthcare needs.

*Terminalia bellirica* Roxb. is a deciduous tree that is widely distributed in the tropical regions. The fruit extract of *T.bellirica* has shown hepatoprotective(17) and the main phyto constituents reported are tannins such as chebulic acid, chebulinic acid, chebulagic acid, gallic acid, corilagin, ellagic acid and flavonoids, sterols, amino acids, fructose, resin, fixed oils etc. *T.bellirica* contains different chemical constituents in different parts such as stem bark contains arjungenin and its glycosides, belleric acid, belleriosides. Ripe fruits are used as astringent in combination

with chebulic myrobalan (*Terminalia chebula*) and *Phyllanthus emblica* as the famous Triphala drug of Ayurveda are also useful in eye problems like cataract, glaucoma, progressive myopia, and conjunctivitis.

Secondary plants metabolites possess various compounds differ widely in terms of structure, biological properties and wide mechanisms of actions (10). Polyphenols have been also proven to lower the risk of cardiovascular diseases, enhance liver regeneration, and increase life expectancy. These secondary metabolites include flavonoids, phenols, phenolic glycosides, saponins, and glycosides (18). *Terminalia* species exhibit nutraceutical value with numerous health benefits in the treatment of some diseases (19). For example, fruits of *T.bellirica* (Gaertn.) Roxb usually form Triphala, the well-known polyherbal formulation in Ayurvedic has pharmacological applications as a laxative, detoxifying, and rejuvenating effects (20). Thus, our present study aims to analyze the phytochemical composition of *T.bellirica* leaves and their antimicrobial activity. In this study, methanol extract of leaves of *T.bellirica* is analyzed by LC-MS techniques for the phytochemicals profile and antibacterial assay is also carried.

## **2.Methods and materials:**

### **2.1.Preparation of extract:**

Fresh mature plant leaves of *T.bellirica* were collected from trees growing in Orathanadu taluk, Thanjavur District, Tamilnadu. The leaves of *T.bellirica* (250 g) were air-dried, ground, and extracted with 95% methanol at room temperature for 8 hours (6 × 500 ml). The methanol extracts were filtered, and reduced under vacuum at 40 °C.

### **2.2. LC-MS assay:**

Leave extract was analyzed by shimadzu LCMS2010 / LCMS-QP8000α and the solvents used were 0.5% (v/v) acetic acid (A) and 100% methanol (B). The isocratic elution was as follows: (i) 55% of solvent A, from 0 to 10 min, (ii) 65%, from 11 to 20 min (iii) 35%, at 21-30 min of total run time. The PDA detector (UPLC LG 500 nm) was monitored at 340 nm and the column temperature was maintained at 30 °C. The mass spectrometer (MS) was operated in the positive ionization mode with the mass range of 150 m/z to 1000 m/z, the capillary voltage of 3.50 kV, cone voltage of 30 V, extractor voltage of 3V, the gas flow of 30 L/Hr and collision gas flow of 0.18 mL/Min. The mass spectrometry (MS) was determined for phytochemicals using a shimadzu LCMS-2010 / LCMS- QP8000α plus instrument by direct injection. The experiment was carried out in NIFTEM, Thanjavur, Tamilnadu. Detection of the ions was performed in electrospray ionization (ESI) and quadrupole ion trap mass analyzer (21).

### **2.3. AAS Analysis:**

Leaves extract was taken in a precleaned and constantly weighed silica crucible and heated in a muffle furnace at 400C till there was no evolution of smoke. The crucible was cooled at room temperature in a desiccator and carbon-free ash was moistened with concentrated sulphuric acid and heated on a heating mantle till fumes of

sulphuric acid ceased to evolve. The crucible with sulphated ash was then heated in a muffle furnace at 60°C till the weight of the content was constant (~2–3 h). One gram of sulphated ash obtained above was dissolved in 100 mL of 5% HCl for Atomic Absorption Spectroscopy (AAS). Standard solution of each element was prepared and calibration curves were drawn for each element using AAS (22).

### 3. Results and discussion

*T.bellirica* has a diverse pharmacological spectrum and has been used in Ayurveda, Siddha, Chinese medicine etc, because of having important phytoconstituents (23,24,25). LC-MS analysis of *T.bellirica* leaves recorded 21 compounds and out of these chebulinic acid, chebulagic acid, quercetin and bellericoside were found to be higher in the leaves (Fig 1,2,3). Chebulinic acid is known to possess numerous biological activities including anti tumor activity, anti antherogenic, anti fibrotic, anti inflammatory, antiulcer, antioxidant, hepatoprotective, antidiabetic and antiviral etc. Chebulagic acid, ellagitannins and proantho cyanidins are the major chemical constituents present in the leaves extract of *T.bellirica*.

Chebulagic acid is benzopyran tannin and an antioxidant that has many potential uses in medicine. It has been found to be immunosuppressive (26,27), hepatoprotective (28) and a potent alpha-glucosidase inhibitor (29,30), a human gut enzyme useful in diabetic studies has been shown to be active against *Staphylococcus aureus* and *Candida albicans* (31). It is formed from geraniin through a glutathione-mediated conversion(32). Gallo-tannic acid, bellericanin, ellagic acid, gallic acid, termilignan, thanni lignan, flavone and anolignan B, tannins, ellagic acid, ethyl gallate, galloyl glucose and chebulaginic acid, phenyllembin, sitosterol, mannitol, glucose, fructose and rhamnose were also found in it. Because of these compounds, it showed many of the pharmacological activities such as antisecretory, analgesic, antihypertensive, antidiarrhoeal activity, antimicrobial activity, antidiabetic, antioxidant(33,34) antiulcer, antipyretic, hepatoprotective, anticancer, angiogenesis, antidepressant-like activity (35). It is useful in the treatment of gastric ulcer, constipation, general debility, piles etc. still, and it has actually not been explored comprehensively. These compounds are traditionally used in relief in a cough, asthma, indigestion, dental problems, sore throat and wounds (36).

Glucosides, tannins, gallic acid, ellagic acid, ethyl gallate, galloyl glucose, chebulanic acid are mainly responsible for its wide therapeutic actions. It has anti-HIV1, antimalarial and antifungal activity. It is used as antioxidant, antimicrobial, antidiarrheal, anticancer, antidiabetic, antihypertensive and hepatoprotective agent (37,38). It also possesses analgesic, antipyretic and anti-ulcerogenic effect and antimicrobial activity (39,40). Many phyto constituents have been isolated from the leaves, chebulagic acid, gallic acid have antimicrobial activity of *T.bellerica* leaves. Proteins, three hydrolysable tannins, gallic acid, ellagic acid and methyl gallate, one flavone,

luteolin, two flavonol aglycones, quercetin and kaempferol and four flavonol glycosides, quercetin 3-O-[6"-a-L-rhamnopyranosyl]-(1g6)-B-D-glucopyranoside (rutin), quercetin-3-O-a-L-rhamnopyranoside, quercetin 3-O-B-D-glucopyranoside and kaempferol 3-O-B-D-glucopyranoside saponins, tannins, amino acids, proteins, alkaloids, carbohydrates and flavonoids. 4-hydroxy-(2-methylbutanol) benzoic acid were present in *T.bellerica* leaves. Apart from vitamins, non-nutritional components of wild green vegetables, notably phenolic compounds are also known to possess powerful radical scavenging properties against reactive oxygen species (14).

Chemically complex compounds from herbs have great therapeutic potential as they have fewer side effects compared to synthetic drugs and also low chances of developing resistance. Bacteria may develop resistance to medicinal plants treatment if only one active ingredient with a specific target is involved, a condition similar to an antibiotic. Glucosides, tannins, gallic acid, ellagic acid, ethyl gallate, gallic acid, gallic acid, gallic acid, gallic acid are having wide therapeutic actions. They are used as antioxidant, antimicrobial, antidiarrheal, anticancer, antidiabetic, antihypertensive and hepatoprotective agent. *T.bellerica* also possess analgesic, antipyretic and anti ulcerogenic effect and antimicrobial activity(40). The micro minerals included copper (Cu), zinc (Zn), iron (Fe) and manganese (Mn) are shown in chart 1 and is expressed as mg/g dry plant material, except for Cu which is expressed as µg/g dry plant material. *T.bellerica* contained the highest amount of iron. The search for alternative nutritious foods may hold the key to nourishing many people by rich sources of vitamins, namely β-carotene, ascorbic acid, riboflavin, folate as well as minerals such as iron, calcium and phosphorous. Dietary diversification is critical in order to mitigate the widespread nutrient deficiencies of vitamin A, iron, and zinc. The dietary requirement for a micronutrient refers to the intake level, which meets specified criteria for adequacy; however, threshold levels remain undefined for most dietary nutrients. Nutritional qualities of *T.bellerica* leaves provide critical insights about its potential in adding diversity and may provide the ultimate weapon against dietary deficiency. Based on the analysis, it has been validated that the *T.bellerica* leaves recorded antibacterial activity due to potent bioactive components.

#### **4. Conclusion:**

In the leaves of *T.bellerica*, chebulinic acid, chebulagic acid, punicalagin, myricetin rutine, galloyl punicalagin are present in higher amount. Based AAS study, this plant could provide reservoir of iron which have great therapeutic potential with fewer side effects than to synthetic drugs. Due to rich phyto nutrients leaves of *T.bellerica*, may be used in development of health supplements. *T.bellerica* leaves are yet to be actively explored on the molecular docking studies for drug development in the pharma industry.

#### **8. REFERENCES:**

1. Varghese SA, Siengchin S, Parameswaranpillai J. Essential oils as antimicrobial agents in biopolymer-based food packaging - a comprehensive review. *Food Biosci.* 2020; 38
2. Casciaro B, Mangiardi L, Cappiello F, Romeo I, Loffredo MR, Iazzetti A, Calcaterra A, Goggiamani A, Ghirga F, Mangoni ML, et al. Naturally-occurring alkaloids of plant origin as potential antimicrobials against antibiotic-resistant infections. *Molecules* 2020; 25(16):3619
3. Takó M, Kerekes EB, Zambrano C, Kotogán A, Papp T, Krisch J, Vágvolgyi C. Plant phenolics and phenolic-enriched extracts as antimicrobial agents against food-contaminating microorganisms. *Antioxidants* 2020; 9(2)
4. Datta S, Roy A. Antimicrobial peptides as potential therapeutic agents: a review. *Int. J. Pept. Res. Therapeut.* 2021;27(1):555–577.
5. Seal T, Chaudhuri K. Nutritional analysis of some selected wild edible plants consumed by the tribal people of Meghalaya state in India. *Int. J. Food Sci. Nutr.* 2016; 1(6):39–43.
6. Redo M, Rios J, Villar A. A review of some antimicrobial compounds isolated from medicinal plants reported in the literature 1978–1988. *Phytother Res.* 1989;3(4):117–125.
7. Ultee RA, Slump G, Steging EJ, Smid. Antimicrobial activity of carvacrol toward *Bacillus cereus* on rice, *J Food Prot*, 2000 May; 63(5):620-4.
8. Sasidharan I, Sundaresan A, Nisha VM, Kirishna MS, Raghu KG, Jayamurthy P. "Inhibitory effect of *Terminalia chebula* Retz. fruit extracts on digestive enzyme related to diabetes and oxidative stress". *J Enzyme Inhib Med Chem.* 2012; 27 (4):578–86.
9. Harvey AL, Edrada-Ebel R, Quinn RJ. The re-emergence of natural products for drug discovery in the genomics era, *Nat. Rev. Drug Discovery* 2015;14:111–129.
10. Calixto JB, The role of natural products in modern drug discovery. *Acad. Bras Cienc* 2019;91:3: e20190105.
11. Buenz EJ, Verpoorte R, and Bauer BA . The ethnopharmacologic contribution to bioprospecting natural products. *Annu. Rev. Pharmacol. Toxicol* 2018; 58: 509–530.
12. Heinrich M. Ethnobotany and its role in drug development. *Phytother. Res* 2000; 14: 479–488.

13. Tabuti JR, Lye KA, and Dhillion SS. Traditional herbal drugs of Bulamogi, Uganda: plants, use and administration. *J. Ethnopharmacol.* 2003;88: 19– 44.
14. Amalraj A, and Gopi S. Medicinal properties of *Terminalia arjuna* (Roxb.) Wight & Arn.: A review. *J. Tradit. Complement Med.* 2017; 7: 65–78.
15. Kim H, and Song MJ. Ethnomedicinal practices for treating liver disorders of local communities in the southern regions of Korea. *Evid. Based Complement Alternat. Med.* 2013; 8:691-76.
16. Upadhyay B, Singh KP and Kumar A. Ethno-veterinary uses and informants consensus factor of medicinal plants of Sariska region, Rajasthan, India. *J. Ethnopharmacol.* 2011;133:14–25.
17. Seal T, Pillai B, Chaudhuri K. Evaluation of nutritional potential of five unexplored wild edible plants consumed by the tribal people of Arunachal Pradesh State in India. *J. Food Nutr. Res.* 2017; 5(1):1–5.
18. Fan C, Dong Y, Xie Y, Su Y, Zhang X, Leavesley D, et al. Shikonin reduces TGF- $\beta$ 1-induced collagen production and contraction in hypertrophic scar-derived human skin fibroblasts. *Int. J. Mol. Med.* 2015; 36: 985–991.
19. Cock IE, and Rayan P. Ascorbic acid potentiates the *Giardia duodenalis* growth inhibitory activity of pure *Terminalia ferdinandiana* Exell compounds. *Parasitol. Res.* 2020; 119:1125–1137.
20. Intharuksa A, Ando H, Miyake K, Sirisa-Ard P, Mikage M, and Sasaki Y. Molecular analysis of *Terminalia* spp. distributed in Thailand and authentication of crude drugs from *Terminalia* plants. *Biol. Pharm. Bull.* 2016; 39:492–501.
21. Devi PN, Kaleeswari S & Poonkothai M. Antimicrobial activity and phytochemical analysis of fruit extracts of *Terminalia bellerica*. *Int J Pharm Pharm Sci*, 2014; 6(5): 639-642.
22. Khan AU, Hassan A and Gilani. Pharmacodynamic evaluation of *Terminalia bellirica* for its anti hypertensive effect, *J Food Drug Anal*, 2008;16:6-14.

23. Kumar N, & Khurana SM. Phytochemistry and medicinal potential of the Terminalia bellirica Roxb. (Bahera). *Indian Journal of Natural Products and Resources (IJNPR)* [Formerly *Natural Product Radiance (NPR)*]. 2018; 9(2): 97-107.
24. Badoni H, Sharma P, Waheed SM & Singh Saumya. Phytochemical analyses and evaluation of antioxidant, antibacterial and toxic properties of Emblica officinalis and Terminalia bellirica fruit extracts. *Asian J Pharm Clin Res*. 2016; 9(6): 96-102.
25. Ruddaraju LK, Pammi SVN, Guntuku GS, Padavala VS, Kolapalli VRM. A Review on Anti-Bacterials to Combat Resistance: From Ancient Era of Plants and Metals to Present and Future Perspectives of Green Nano Technological Combinations. *Asian J. Pharm. Sci*. 2020; 15:42–59.
26. Abraham A, Mathew L & Samuel S. Pharmacognostic studies of the fruits of Terminalia bellirica (Gaertn.) Roxb. *Journal of Pharmacognosy and Phytochemistry*, 2014; 3(2).
27. Hamada, Shin-ichi, Kataoka, Takao, Woo, Je-Tae, Yamada, Atsushi, Yoshida, Takashi, Nishimura, Toshio, Otake, Noboru; Nagai, Kazuo. "Immunosuppressive Effects of Gallic Acid and Chebulagic Acid on CTL-Mediated Cytotoxicity" *Biological & Pharmaceutical Bulletin*. 1997; 20 (9):1017–1019.
28. Kinoshita S, Inoue Y, Nakama S, Ichiba T, Aniya Y. "Antioxidant and hepatoprotective actions of medicinal herb, Terminalia catappa L. from Okinawa Island and its tannin corilagin". *Phytomedicine* 2007; 14 (11):755–762.
29. Gina Porras, François Chassagne, James T Lyles, Lewis Marquez et al. Natural Products in Antibiotic Drug Discovery, *Chem Rev*. 2021; 121(6):3495-3560.
30. Abdus Satter MM, Khan MML, Jabin SA, Abedin N, Islam MF, Shaha B. Nutritional quality and safety aspects of wild vegetables consume in Bangladesh. *Asian Pac. J. Trop. Biomed* 2016; 6(2):125–131.
31. Pin-Shern Chen and Jih-Heng Li. Chemopreventive effect of punicalagin, a novel tannin component isolated from Terminalia catappa, on H-ras-transformed NIH3T3 cells. *Toxicology Letters* 2006; 163(1):44-53

32. Tanaka T, Kouno I and Nonaka GI. Glutathione-mediated conversion of the ellagitannin geraniin into chebulagic acid, *Chemical and pharmaceutical bulletin*. 1996;44(1):34-40.

33. Basu T, Panja, S, Ghate NB, Chaudhuri D, and Mandal N. Antioxidant and antiproliferative effects of different solvent fractions from *Terminalia belerica* Roxb. fruit on various cancer cells. *Cytotechnology*. 2017; 69: 201– 216.

34. Rastogi S, Pandey MM & Rawat AKS. Phytochemical analysis, phenolic content and antioxidant parts of *Terminalia belerica* (Gaertn.) Roxb.-A comparative study 2018;23-26.

**Table 1: Compounds identified in LC-MS assay**

35. Saraswathi MN, Karthikeyan M, Kannan M and Rajasekar S, *Terminalia bellirica* Roxb.-A phytopharmacological review, *Int J Pharm Biomed Res*, 2012; 3(1): 96-99.

36. Dawe A, Talom B, Kapche G, Siddiqui K, Yakai F, Talla E, et al. Termiglaucescin, a new polyhydroxy triterpene glucoside from *Terminalia glaucescens* with antioxidant and antiinflammatory potential. *Z. Fur Naturforschung Section C-a J. Biosci.* 2017;72: 203–208.

37. Kokate CK, Purohit AP and Gokhale SB, *Pharmacognosy*, 39th edn, (Nirali Prakashan), 2008: 260-261.


38. Anand KK, Singh B, Saxena AK, Chandan BK, Gupta VN and Bhardwaj V. The Hepatoprotective principle in the fruits of *Terminalia bellirica*-bioassay guided activity, *Pharmacol Res.* 1997; 36: 315-321.

39. Biradar YS, Jagatap S, Khandelwal KR and Singhania SS. Exploring of antimicrobial activity of triphala mashi—an ayurvedic formulation, *Complement Alternat Med.* 2008;5(1):107-113.

40. Ody P, *The Complete Medicinal Herbal: A Practical Guide to the Healing Properties of Herbs*. Skyhorse Publishing Inc.; New York, NY, USA 2017: 1–271.

S.No	Tentatively identified compounds	m/z
1.	7- Hydroxy3,4(methylene dioxyflavan)	270
2.	Luteolin	286
3.	Beta sitosterol	414
4.	Termilignan	296
5.	Anolignan	266
6.	Gallic acid	170
7.	Ellagic acid	302
8.	Methyl Gallate	184
9.	Phyllembin	198
10.	Chebulinic acid	956
11.	Chebulagic acid	955
12.	3-0- alpha glucopyranosylL-aminopyranose	326
13.	Hexahydroxydiphenic acid	338
14.	Arjungenin	505
15.	Bellericoside	666
16.	Quercetine	302
17.	Kempherol	286
18.	Rutin	610
19.	Quercetin – 3-07-0-alpha rhamnopyranoside	756
20.	Quercetin-3-beta-glucoside	464
21.	Dihydroclerodin	436
22.	Quinic acid	191
23.	Mailc acid	133
24.	Galloyl hexoside	933
25.	puigluconin	801
26.	Grahatin	951
27.	corilagin	633
28.	Galloyl punicalagin	1235
29.	Trigaloxyl hexoside	635
30.	Elgatannin	967
31.	Punicafolin	937
32.	Quercetin coumaroyl glucoside	609
33.	Vitexin	431
34.	Chebulic acid	355
35.	Cuarictin	935
36.	Myricetin rutine	935
37.	Myricetin glucoside	479


**Chart 1: Minerals Assay by AAS in leaves of *T.bellerica***




**Fig 1. Chebulagicacid, C<sub>41</sub>H<sub>30</sub>O<sub>27</sub>, Exact Mass:954.66**  
MSES+

2:TOF


1.05e6



**Fig 2: Bellericoside, C<sub>36</sub>H<sub>58</sub>O<sub>11</sub>, Exact Mass: 666.8**



**Fig 3: quercetin 3-O-7-O- $\alpha$ -L-rhamnopyranoside, C<sub>33</sub>H<sub>40</sub>O<sub>20</sub> Exact Mass: 756.7**



UNDER PEER REVIEW

UNDER PEER REVIEW