

EFFECT OF TILLAGE PRACTICES AND HYDROGEL APPLICATIONS ON YIELD AND YIELD ATTRIBUTING TRAITS OF MAIZE (*ZEA MAYS L.*) OVER THE YEARS

Abstract:

This study investigated the effects of different tillage practices and hydrogel applications on maize yield in sandy loam soils at the Agricultural Research Station, Karimnagar, PJTSAU, Telanganastate over a period of four years (2015-2018) during the Kharif season. The experiment was designed as a split plot with four main treatments *i.e.*, Conventional tillage (CT), Conventional tillage with residue mulching (4 t/ha), Zero tillage (ZT) and Zero tillage with residue mulching (4 t/ha). These were combined with three sub-treatments involving hydrogel application *i.e.*, Control (no hydrogel), Hydrogel at 2.5 kg/ha and Hydrogel at 5.0 kg/ha. The study was replicated three times to ensure reliability of the results. The combination of zero tillage with residue mulching yielded the highest maize grain yield (4435 kg/ha). This suggests that this practice is more effective in retaining soil moisture, which is critical under rainfed conditions. The addition of hydrogel, regardless of the rate (2.5 kg/ha or 5.0 kg/ha), did not lead to any significant improvement in maize yield. The use of zero tillage combined with residue mulching is beneficial for maize production in sandy loam soils under rainfed conditions, primarily due to enhanced moisture retention. However, hydrogel application did not provide any additional yield benefits in this context.

Keywords: Maize, tillage, hydrogel, mulching, conventional tillage and zero tillage.

Introduction:

“Maize (*Zea mays* L.) is one of the most versatile emerging crop showing wider adaptability under varied agro-climatic conditions. Globally, maize is known as queen of cereals because it has the highest genetic yield potential among the cereals. In India, maize is the third most important food crops after rice and wheat. Maize in India, contributes nearly 9 % in the national food basket and its production was 24.26 M.T. from 9.06 M.ha with a productivity of 2.68 t/ha” (Parihar et al., 2011). “Projected demand for maize production by 2050 in India is around 121 M.T.. Maize is predominantly grown as a rainfed crop in India. Currently rainfed yields (1.9 tonnes/ha) are much lower than irrigated yields (3.5 tonne/ha) in India. This indicates huge untapped yield potential in rainfed maize production system. Maize is cultivated around 4.2- 4.5 lakh hectares in rainfed situations in Telangana state” (Agricultural Statistics at a Glance 2022).“Conservation agriculture is on priority to sustain the productivity of cropping system in India. The interest in conservation tillage systems has increased in response to the need to limit erosion, promote water conservation and reduce weed seed bank. The yields on the zero tillage plots approached those of the conventional tilled plots when N was added. Had a higher level of N fertilizer been applied it is possible that the yields on zero tillage plots would have equaled or exceeded those of the conventional tilled plots. The net mineralization rate was greater with conventional tillage than the zero tillage. Soil temperature at 5 and 10 cm depths was less on zero tillage than on conventional tillage plots. Water content in soil was greater on zero tillage

compared to conventional tillage plots. The drought had a considerable effect on socio-economic, agricultural and environmental aspects” (Chaithra and Sridhara, 2018). “Mulching has been proved to be useful material in conserving moisture and enhancing the productivity of maize. Leaf mulch also provides benefit regarding increasing infiltration rate, lowers the soil temperature and improves fertilizer availability and increase crop yield” (Dushouyuet *al.*, 1995). “The hydrogel used in agriculture is mainly polyacrylamides with acrylic acid as basic units” (Bai et al., 2010). Recently, hydrogels have included cellulose (Mi et al., 2018), protein (Kong et al., 2019), and starch (Mahmoodi-Babolan et al., 2019) in their structures. “The hydrogel gradually releases up to 95% of its stored water when its surroundings begin to dry out. But, when comes in contact with water again, it gets replenished. This process can last up to 2-5 years, by which time biodegradable hydrogel decomposes to CO₂, water and ammonia and potassium ions, without any residue, thus, environment friendly” (Trenkel, 1997). “And also improve the crop growth by increasing water holding capacity in soil and delay the wilting point in drought stress” (Boatright *et al.*, 1997). In this context, the present study was undertaken to study the effect of conservation tillage and hydrogel in maize yield performance under rainfed conditions.

Materials and Methods:

A field experiment was conducted during two *Kharif* seasons of 2015 to 2018 at Agricultural Research station, Karimnagar PJTSAU, Telanganastate under rainfed conditions. The soils are red sandy loam with medium in organic carbon (0.65 %), available N (213.1 kg ha⁻¹), available P (42.4 kg ha⁻¹) and available K (415 kg ha⁻¹). This experiment was laid out in split plot design with 4 main plots *i.e.*, M1-Conventional till, M2- Conventional till + residue mulching (4 t/ha), M3- Zero tillage, M4- Zero tillage + residue mulching (4 t/ha) and three sub plot *i.e.*, S1- Control (no hydrogel), S2- Hydrogel 2.5 kg/ha, S3-Hydrogel 5.0 kg/ha in three replications. Proper management practices like nutrient management (200:60:50 NPK Kg/ha) weed management by herbicides and pest management by pesticides were. Biometric observations like length of cobs, grains row per cob, number of grains per row, test weight, grain and Stover yield were recorded after harvesting of crop. Economics of each treatment was calculated on the basis of the nearest market prices of inputs and outputs.

Results and discussion:

First year: The grain yield resulted significantly higher under zero tillage + mulching (2195 kg/ha) on par with conventional tillage (2148 kg/ha). Due to low yields, the returns are negative. The yield attributes, cob length & girth, kernel rows & no. & 100-grain weight were also higher with zero tillage + mulching. The hydrogel application did not show significant difference on grain yield and yield attributes.

Second year: The grain yield and BC ratio resulted significantly higher under zero tillage (7663 kg/ha) on par with zero tillage + mulching (7616 kg/ha). The yield attributes cob length, kernel no. were also higher with zero tillage & zero tillage + mulching. The hydrogel application did not show significant difference on grain yield and yield attributes

Third year: Grain yield and BC ratio resulted significantly higher under conventional tillage + mulching (5046 kg/ha) on par with zero tillage + mulching (5298 kg/ha). The hydrogel application did not show significant difference on grain yield and yield attributes.

Forth year: Significantly higher grain yield and BC ratio resulted under zero tillage + mulching (2632 kg/ha) on par with conventional tillage + mulching (2506 kg/ha). The hydrogel application did not show significant difference on grain yield and yield attributes.

Pooled data: Higher maize grain yield (4435 kg/ha) and BC ratio resulted under zero tillage + mulching when compared to without mulching in conventional tillage and Zero tillage tillage (3977 kg/ha) under rainfed conditions. Significantly lowest grain yield was observed with Conventional tillage (3154 kg/ha). The hydrogel application (@ 2.5 or 5 kg/ha) did not show significant difference on grain yield and yield attributes. Similar results were obtained with findings of Khadem *et al.*, 2010, Gunes *et al.*, 2016, Kumaret *al.*, 2017, Kumaret *al.*, 2020 and Shubham *et al.*, 2023 in terms of yield and yield attributes of maize and other crops.

Conclusion:

The zero tillage + mulching recorded higher maize grain yield when compared to conventional or zero tillage without mulch and conventional with mulch under rainfed conditions due to long term availability of moisture in the soil. To overcome from prolonged dry spells in rainfed situation mulching is useful.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

I am hereby declaring that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.,) and text-to-image generators have been used during writing or editing of manuscripts.

References:

Agricultural Statistics at a Glance 2022. <https://desagri.gov.in>.

Bai W, Zhang H, Liu B, Wu Y, Song J Q. 2010. Effects of super-absorbent polymers on the physical and chemical properties of soil following different wetting and drying cycles, Soil Use Management. 26: 253-260.

Boatright JL, Balint DE, Mackay WA, Zajicek JM. 1997. Incorporation of a hydrophilic polymer into annual landscape beds. Journal of Environmental Horticulture.15:37-40.

Chaithra G M and Sridhara S. 2018. Growth and yield of rainfed maize as influenced by application of super absorbent polymer and Pongamia leaf mulching International Journal of Chemical Studies. 6(5): 426-430.

Dushouyu T, Enping WM, Wqingshan A. 1995. Study of the Overall Effect of Straw Mulching and Ploughing in and Techniques for Its Applications. Ningxia Journal of Agriculture and Forestry Science and technology. 5(2):10-14.

Gunes A, NurulKitir, MetinTuran, ErdalElkoca, ErtanYildirim, Nazmiye et al. 2016. Evaluation of effects of watersaving superabsorbent polymer on corn (*Zea mays* L.) yield and phosphorus fertilizer efficiency. Turkish J Agri. Forestry. 40:365-378.

- Khadem SA, Galavi M, Ramrodi M. 2010. Effect of animal manure and super absorbent polymer on corn leaf relative water content, cell membrane stability and leaf chlorophyll content under dry condition. *Australian Journal of Crop Science*. 4(8):642-647.
- Kong W, Li Q, Li X, Su Y, Yue Q, Gao B. 2019. A biodegradable biomass-based polymeric composite for slow release and water retention. *Journal of Environmental Management*. 230 : 190-198.
- Kumari S, Solanki NS, Dashora LN, Upadhyay B. 2017. Effect of superabsorbent polymer and plant geometry on growth and productivity of maize (*Zea mays* L.). *Journal of Pharmacognosy Phytochemistry*. 6(4):179-181.
- Kumar Naik A H, Chaithra G M, Kiran Kumar N, Madhu G, Nataraja M, Umesha S and Madhu BM. 2020. Effect of hydrogel on growth, yield and Economics of rainfed castor. *The Pharma Innovation Journal*. SP-9(7): 36-39.
- Mahmoodi Babolan N, Nematollahzadeh A, Heydari A, Merikhy A. 2019. Bioinspired catecholamine/starch composites as superadsorbent for the environmental remediation. *International Journal of Biological Macromolecules* 125 : 690-699,
- Mi Hao Yong, Jing Xin, Politowicz Alexander L, Chen Edward, . Huang Han-Xiong, Turng Lih Sheng . 2018. Highly compressible ultra-light anisotropic cellulose/graphene aerogel fabricated by bidirectional freeze drying for selective oil absorption. *Carbon* 132 :199-209.
- Parihar C M, Jat S L, Singh A K, Sai Kumar R, Hooda K S, Chikkappa G K, Singh D K. 2011. *Maize Production Technologies in India*. 1-34.

Shubham Jamwal, Joy Dawson and Cheenu Kashyap.2023. Effect of Phosphorus and Hydrogel on Growth and Yield of Maize (*Zea mays* L.). International Journal of Environment and Climate Change. 13(9): 2122-2128.

Trenkel ME. 1997. Controlled release and stabilized fertilizers in Agriculture. International Fertilizer Industry Association, Paris, France. 1-156.

UNDER PEER REVIEW

UNDER PEER REVIEW