

Recruitment index as a new approach to assessing the resilience of natural mangrove ecosystems: A case study in the coastal of West Muna Regency

Abstract

This study aims to determine the resilience of mangrove ecosystems with a recruitment index approach on the coast of West Muna Regency. The research method used quadrant transects (10 x 10 m²). Analysis of the resilience index is based on the recruitment index (RI) with the equation $RI = \ln ((0.7 \times DSd/DT.) + (Dsp/DT))$. The results showed that the mangrove ecosystem in all stations had a high resilience with an RI value of 1.43 for station I, 1.38 for station II, 1.07 for station III, and 1.05 for station IV. Meanwhile, mangrove species with very high resilience categories are *B. gymnorrhiza* (RI = 1.70) and *B. cylindrica* (RI = 1.63). In comparison, the resilience of *S. alba* (RI = 0.90) and *R. stylosa* (RI = 1.22) is classified as medium and high resilience, respectively. Based on the RI value, the mangrove ecosystem on the coast of West Muna Regency is in the high resilience category (RI = 1.25 ± 0.20). It indicates that the mangrove ecosystem on the coast of West Muna Regency has good regeneration and growth abilities, so it can recover even when under ecological pressure.

Keywords: Mangrove ecosystem, mangrove management, recruitment index, resilience.

Introduction

Mangrove ecosystems play an essential role in marine life (Nordhaus et al., 2006) and coastal communities by provisioning various ecosystem services (Owuor et al., 2019; Handayani et al., 2020). Some essential services of mangrove ecosystems include the providing of biodiversity (Sievers et al., 2023; Rahman et al., 2024a), carbon sequestration (Choudhary et al., 2024; Rahman et al., 2024b), coastal protection (Damastuti et al., 2023; Morris et al., 2023; van Hespén et al., 2023; Anu et al., 2024), wind and wave storm breakers (Sunkur et al.,

2023; Rahman et al., 2024c), as well as providers of firewood, honey, fruit, fodder, and traditional medicine (Nyangoko et al., 2022).

The availability of mangrove ecosystem services triggers socio-ecological interactions that cause the degradation of mangrove ecosystems, such as declining density, species diversity, and mangrove area (Rahman et al., 2020a). However, mangrove ecosystems can maintain ecological conditions or renew themselves after being pressured by anthropogenic activities formed in these ecological social interactions. The ability to recover is known as resilience (Holling, 1973). Resilience is the ability of environmental systems to cope with disturbances and changes and return to stable conditions or adapt to new situations (Holling et al., 2002). Studies on the resilience of mangrove ecosystems that have been reported analyze the level of resilience based on ecological pressures caused by several factors, such as sea level rise (Yulianti et al., 2013) and storm waves (Lagomasino et al., 2021). The determination of the level of resilience in the study refers to several ecological aspects, especially the density, thickness, and diversity of species (Irwansyah et al., 2023).

One aspect that needs to be understood in the mangrove ecosystem resilience study is the ability of ecosystems to renew and maintain existing mangrove plant populations. It can be measured through the recruitment index, which is one of the indicators of the health of the mangrove ecosystem. The recruitment index in mangrove ecosystems can compare the number of newly grown mangrove plants (seedlings) and mangrove saplings with the number of adult mangroves (Rahman et al., 2019). The availability of seedlings and their ratio to mangrove density represents the ability of mangroves to be resistant and resilient to ecological disturbances such as marine debris inputs, changes in water quality, timber, land conversion, and other environmental stresses. In addition, it can also indicate the resilience of mangrove ecosystems, especially in ecological aspects.

Research on the recruitment index concerning the resilience of mangrove ecosystems has rarely been conducted. This research can help evaluate mangrove ecosystems' health and ability to maintain existing mangrove plant populations. In addition, the results of this study can also be the basis for formulating appropriate management strategies to reduce damage to mangrove ecosystems and increase the resilience capacity of mangrove ecosystems in the future. The research could also provide deeper insights into ways to improve the health of mangrove ecosystems and prevent further damage. For example, by improving environmental quality, reducing human pressure on ecosystems, or protecting critical mangrove areas, as Rahman et al. (2020a) reported.

Materials and Methods

Description of Study Sites

This research was conducted in the mangrove ecosystem area of West Muna Regency. The study area was divided into four observation stations, namely Maginti District (Station I), Tiworo Tengah District (Station II), Tiworo Kepulauan District (Station III), and Sawerigadi District (Station IV) (Figure 1). The location selection is a representation of the mangrove ecosystem on the coast of West Muna Regency with a varying area of 372.73 ha at station I, 534.31 ha at station II, 1033.04 ha at station III, and 286.65 ha at station IV.

The dominant mangrove species in the mangrove ecosystem along the coast of West Muna Regency are *B. cylindrica*, *B. gymnorhiza*, *R. apiculata*, *R. mucronata*, *R. stylosa*, and *S. alba* (Rahman et al., 2020b). The species has been degraded due to its use as timber, residential areas, and aquaculture from 2014 to 2017 (Rahman et al., 2020a).

Commented [MA1]: Study Area

Commented [MA2]: Mention the district and are. Try to give a location map.

Commented [MA3]: On what basis 4 stations was selected?




Figure 1. Map of the study site

Commented [MA4]: Rewrite the title
Location map of West Muna Regency with four studied stations

Sampling Methods

Mangrove ecosystem data was collected using a quadrant plot measuring 10 x 10 m², placed by random sampling. The number of quadrants at each location adjusts to the area and condition of the mangrove ecosystem so that 300 quadrants are obtained with a distribution of 60 at station I, 90 at station II, 110 at station III, and 40 at station IV.

Data was collected by counting the number of mangrove stands in all categories. Species identification was carried out based on the morphological characteristics of mangroves, including roots, leaves, flowers, and fruits, as referred to in the manual according to Noor et al. (2006) and Bengen et al. (2022).

The mangrove vegetation structure was observed across three categories: trees, saplings, and seedlings. Each category is classified based on diameter size: D = 1-3 cm for seedlings, D = 3-5 cm for saplings, and D > 5 cm for the tree category. The classification of D > 5 cm as trees is based on observations showing that mangroves can reproduce at this diameter, particularly for the *Rhizophora* types (**Figure 2**).



Figure 2. *Rhizophora* mangroves ($D \geq 5$ cm) produce propagules as a source of new individuals

Data Analysis

Mangrove Density

Mangrove density in each type of stand was analyzed regarding the equation according to Bengen et al. (2022), namely as follows:

$$D = \frac{\sum Ni}{A} \quad (1)$$

Where D is mangrove density (stands/ha), N_i is the number of mangroves of each type found in the plot (stands), and A is the area of the plot (ha).

The mangrove density status is classified into five categories according to the criteria by Rahman et al. (2019) as in Table 1 below:

Table 1. The classification of mangrove density status (Rahman et al. 2019)

Density (stands/ha)	Criteria
<500	Very rare
>500 – 1000	Rare
>1000 – 1500	Medium
>1500 – 2000	Dense
>2000	Very dense

Mangrove Resilience and Recruitment Index

The recruitment index in mangrove ecosystems can refer to the comparison between the number of newly growing mangrove plants (seedlings) and mangrove seedlings (saplings) to the number of mature mangroves (Rahman et al. 2019).

Ecologically, each mangrove seedling has a 60 – 80% survival potential or an average of 70% to grow into saplings until they reach a tree type (Bengen et al. 2022). Based on this, the mangrove ecosystem recruitment index can be formulated as follows:

$$RI = \text{Ln} \left(\left(0.7 \times \frac{DSd}{DT} \right) + \left(\frac{DSp}{DT} \right) \right) \quad (2)$$

Where:

RI = value is the recruitment index

0.7 = the average value of the probability that mangrove seedlings can live to reach tree size (Bengen et al., 2022).

DT = the density of mangroves in the tree category with $DT \neq 0$ (stands/ha)

DSd = the density of mangroves in the seedling category (stands/ha)

DSp = the density of mangroves in the sapling category (stands/ha).

Either of DSd and DSp can be equal to zero but not for both.

The criteria for resilience and management actions are based on the value of the Mangrove Recruitment Index (RI.), namely (1) Very low resilience ($RI \leq 0$): in this condition, management of the mangrove ecosystem is with complete protection. Efforts to improve mangrove conditions are rehabilitation based on a modification of ecological conditions according to the ability of mangrove species to live; (2) Low resilience ($0 < RI \leq 0.5$): In this condition, the management of the mangrove ecosystem is conservation and rehabilitation. Rehabilitation efforts can be carried out in the original habitat according to its natural conditions in the ecosystem; (3) Moderate resilience ($0.5 < RI \leq 1.0$): in this condition, mangrove management is conservation with particular uses such as ecotourism or fruit utilization so that mangroves can remain sustainable; (4) High resilience ($1.0 < RI \leq 1.5$): in this condition, mangrove management is conservation with limited use. Limited use means that under certain conditions, mangroves can be cut down in small quantities to fulfill the need for wood; (5) Very high resilience ($RI > 1.5$): in this condition, mangrove management is conservation with medium utilization. Medium utilization means that under certain conditions, the mangrove ecosystem can be converted into aquaculture areas or other areas with a maximum area of 20% of the total ecosystem area (Table 2).

Table 2. Resilience criteria and management actions based on Recruitment Index (RI) values.

<i>R.I.</i> Value	Criteria of Resilience	Management Strategy
$RI \leq 0$	Very low resilience	Protection and rehabilitation
$0 < RI \leq 0.5$	Low resilience	Conservation and rehabilitation
$0.5 < RI \leq 1.0$	Medium resilience	Conservation with a special activity
$1.0 < RI \leq 1.5$	High resilience	Conservation with low utilization
$RI > 1.5$	Very high resilience	Conservation with medium utilization

Result and Discussions

Mangrove Density

Six dominant species are found in the mangrove ecosystem on the coast of West Muna Regency, including *B. cylindrica*, *B. gymnorhiza*, *R. apiculata*, *R. mucronata*, *R. stylosa*, and

S. alba. These findings align with the report of Rahman et al. (2020). This species dominates in all mangrove categories (trees, saplings, and seedlings). In the tree category, mangrove density ranges from 617 to 829 stands/ha. According to Rahman et al. (2019), this value is in the low category. The highest density was found in Maginti District (station I) and Sawerigadi District (station IV), namely 829 stands/ha and 808 stands/ha, respectively. In contrast, the smallest densities were found at stations II and III, namely 617 stands/ha and 688 stands/ha, respectively (Table 3).

In the sapling category, the highest density of mangroves was found at Tiworo Tengah District (station II) namely 1393 stands/ha. In comparison, the lowest was found at station III of 1011 stands/ha. Meanwhile, the highest and smallest density of mangroves in the seedling category was found at Maginti District (station I), namely 3155 stands/ha, and Tiworo Kepulauan District (station III), namely 1421 stands/ha (Table 3).

Table 3 also shows that the highest tree species density is *S. alba* (299 stands/ha), and the lowest is *B. gymnorhiza* (31 stands/ha) found in Sawerigadi District (station IV). In the sapling category, the mangrove species with the highest density was *R. stylosa* (433 stands/ha), which was found in Tiworo Tengah District (station II), while the lowest was *B. cylindrica* (97 stands/ha), which was found in Maginti District (station I). In the seedling category, the highest density was *R. stylosa*, namely 736 stands/ha, and the lowest was *B. cylindrica*, namely 37 stands/ha, each of which was found in Maginti District (station I) and Sawerigadi District (station IV).

According to Ilman et al. (2016), the low density of mangrove ecosystems in an area is caused by the conversion of land functions into several uses, such as ponds, wharves, and settlements. Arifanti et al. (2021) also reported a similar thing and found that mangrove degradation in Indonesia occurs due to land conversion to cultivation, agriculture, settlements, and mining activities.

In addition, Rahman et al. (2020) state that the degradation of mangrove forests on the coast of West Muna Regency, specifically at stations I to IV, is caused by land conversion into ponds and settlements. Some infrastructure was also built, such as docks and roads, which were carried out by converting mangrove forest areas.

Table 3. Mangrove density on the coast of West Muna Regency

Species	Station I			Station II			Station III			Station IV		
	D.T.	DSp	DSd	DT	DSp	DSd	DT	DSp	DSd	DT	DSp	DSd
<i>B. cylindrica</i>	43	97	330	38	108	166	34	113	67	57	138	37
<i>B. gymnorrhiza</i>	105	178	467	59	201	133	34	167	154	31	123	87
<i>R. apiculata</i>	180	250	667	106	221	267	144	198	224	75	223	179
<i>R. mucronata</i>	120	143	543	74	163	189	121	133	216	65	227	155
<i>R. stylosa</i>	225	367	736	178	433	427	137	167	312	281	381	584
<i>S. alba</i>	156	233	412	162	267	315	218	233	448	299	133	512
Total	829	1268	3155	617	1393	1497	688	1011	1421	808	1225	1554

Recruitment Index

The highest mangrove species recruitment index (RI) at each station was dominated by *Bruguiera cylindrica* and *Bruguiera gymnorrhiza* species, while the lowest was dominated by *Sonneratia alba*. This disparity is due to the different fertilization processes between these mangrove types. *Bruguiera* species reproduce primarily through propagules, which have a better growth ability than the round-shaped fruit of *Sonneratia* species (Hogarth, 2007). According to Bengen et al. (2022), the propagules of *Bruguiera* mangroves, when they fall into the ecosystem areas, are more likely to grow naturally because they easily embed in the mud sediments, which are the typical habitat for these mangroves. Conversely, the fruit of *Sonneratia* species disperses more quickly due to the current and wave patterns found in the ecosystem area. This rapid dispersal is also related to the presence of *Sonneratia* species in open zones that directly face ocean currents and waves (Bengen et al., 2022).

The highest RI values were at stations I and II, namely *B. cylindrica*, with respective values of 2.03 and 1.77). The highest RI at stations III and IV was *B. gymnorrhiza*, with R.I. values of 2.09 and 1.78, respectively (Figure 3). The lowest R.I. values at stations I, II, and IV

were *S. alba*, namely RI = 1.21 at station I, RI = 1.10 at station II, and RI = 0.50 at station IV. The lowest R.I. at station III were *R. mucronata* and *R. apiculata*, with R.I.s of 0.85 and 0.90, respectively (Figure 3).




Figure 3. Mangrove species recruitment index at each station along the coast of West Muna Regency

Figure 3 also shows that the highest RI. values for total mangrove forests were found at Maginti district (station I) and Tiworo Tengah district (station II), with values of 1.43 and 1.38. At the same time, stations III and IV were the lowest, with R.I. values of 1.07 and 1.05, respectively. Meanwhile, the highest R.I. values for mangrove species on the coast of West Muna Regency were *B. gymnorhiza* and *B. cylindrica*, with RI = 1.70 and RI = 1.63, respectively. In comparison, the lowest were *S. alba* (RI = 0.90) and *R. stylosa* (RI = 1.22) (Figure 4).




Figure 4. Recruitment Index of mangrove species on the coast of West Muna Regency

Mangrove Resilience

Mangrove ecosystem resilience is a form of adaptation to various disturbances outside the ecosystem. Disruption to mangrove ecosystems can occur due to human intervention, such as land conversion, and natural disturbances, such as climate change and sea level rise (Alongi 2008; Yulianti et al. 2013).

Based on the RI. value, the mangrove ecosystem on the coast of West Muna Regency is in the high resilience category ($RI = 1.25 \pm 0.20$) (Table 3). The resilience value is based on each stand category's total density (Table 2). The findings indicate that the mangrove ecosystem on the coast of West Muna Regency has good regeneration and growth abilities, so it can recover even when under ecological pressure.

The resilience criteria also indicate that mangroves must be managed wisely, namely conservation-based management with limited use. According to Rahman et al. (2020), the coastal communities of West Muna Regency still often use mangroves as firewood and building materials in residential areas in the mangrove ecosystem area. Conservation-based management with limited use is still included in the criteria for a sustainable management mechanism because it maintains ecological sustainability, supports economic life, and harmonizes the social life of coastal communities in mangrove ecosystem areas or between

communities and government stakeholders. Yulianda (2007) states that conservation does not mean complete protection; instead, it uses ecosystems wisely to form a balance between social and ecological systems.

Ali et al. (2020) reported that good social-ecological interactions in managing mangrove ecosystems could achieve sustainable management. Mangroves provide ecosystem services to support people's lives. At the same time, humans as a social system will protect mangrove ecosystems for the sustainability of mangrove ecosystem services by providing direct and indirect economic benefits (Handayani et al. 2020; Nazar et al. 2021).

Table 4. Resilience criteria and management strategies for mangrove ecosystems in the coastal areas of West Muna Regency

Locations	RI value	Criteria	Management Strategy
Station I	1.43	High resilience	Conservation with low utilization
Station II	1.38	High resilience	Conservation with low utilization
Station III	1.07	High resilience	Conservation with low utilization
Station IV	1.05	High resilience	Conservation with low utilization
Total	1.25±0.20	High resilience	Conservation with low utilization

Conclusion

The findings revealed that the mangrove ecosystems at all stations exhibited high resilience, with RI values of 1.43 for station I, 1.38 for station II, 1.07 for station III, and 1.05 for station IV. *Bruguiera gymnorrhiza* (RI = 1.70) and *Bruguiera cylindrica* (RI = 1.63) demonstrated very high resilience among the mangrove species. In contrast, *Sonneratia alba* (RI = 0.90) were categorized as having medium resilience. Overall, the mangrove ecosystem along the coast of West Muna Regency falls into the high resilience category, with an average RI value of 1.25 ± 0.20 . This indicates that the mangrove ecosystem in this region has strong regeneration and growth capabilities, enabling it to recover effectively even under socio-ecological pressure.

References

- Ali, M., Muliani., Adrianto, L., Hariyadi, S., Rahman. 2020. Resiliensi sistem sosial ekologi kawasan desa pesisir Kabupaten Subang. *Grouper*. 11(2): 33 - 44
- Alongi, D.M. 2008. Mangrove forests: resilienci, protection from tsunamis, and responses to global climate change. *Estuarine, Coastal, and Shelf Science*. 76: 1-13.
- Anu, K., Henna, P.K., Sneha, V.K., Busheera, P., Jumana, M., Anu, A. 2024. Mangroves in environmental engineering: Harnessing the multifunctional potential of nature's coastal architects for sustainable ecosystem management. *Results in Engineering*. 21, 101765. <https://doi.org/10.1016/j.rineng.2024.101765>
- Arifanti, V.B., Novita, N., Subarno, Tosiani, A. 2021. Mangrove deforestation and CO₂ emissions in Indonesia. *IOP Conference Series: Earth and Environmental Science*, 874(1), 012006. <https://doi.org/10.1088/1755-1315/874/1/012006>.
- Bengen, D.G., Yonvitner., Rahman. 2022. *Pedoman Teknis Pengenalan dan Pengelolaan Ekosistem Mangrove*. Bogor (I.D.). IPB Press. 76p.
- Choudhary, B., Dhar, V., Pawase, A.S. 2024. Blue carbon and the role of mangroves in carbon sequestration: Its mechanisms, estimation, human impacts and conservation strategies for economic incentives. *Journal of Sea Research*. 199, 102504. <https://doi.org/10.1016/j.seares.2024.102504>
- Damastuti, E., van Wesenbeeck, B.K., Leemans, R., de Groot, R.S., Silvius, M.J. 2023. Effectiveness of community-based mangrove management for coastal protection: A case study from Central Jawa, Indonesia. *Ocean and Coastal Management*. 283, 106498. <https://doi.org/10.1016/j.ocecoaman.2023.106498>

- Handayani, S., Bengen, D.G., Nurjaya, I.W., Adrianto, L., Wardiatno, Y. 2020. Pemetaan jasa ekosistem mangrove pada wilayah rehabilitasi di Pesisir Sayung, Kabupaten Demak. *JUPI* 25(4): 574 – 583.
- Hogarth, J. P. 2007. *The Biology of Mangroves and Seagrasses*. Oxford University Press. America.
- Holling, C.S. 1973. Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*. 40 (50): 1-23.
- Holling, C.S., Gunderson, L.H., Ludwig, D. 2002. In Quest of Theory of Adaptive Change. In: Gunderson LH, Holling CS (editor). *Panarchy: understanding transformations in human and natural systems*. Washington (DC): Island Press.
- Ilman, M., Dargusch, P., Dart, P., Onrizal. 2016. A historical analysis of the drivers of loss and degradation Indonesia's mangroves. *Land Use Policy*. 54: 448-459.
- Irwansyah, I., Asgar, M.A., Daris, L., Massiseng, A.N.A., Alpiani, A., Masriah, A. 2023. Level of mangrove ecosystem resilience in Pallime waters, Cenrana District, Bone Regency. *Jurnal Akuatiklestari*. 7(1): 52-59.
- Lagomasino, D., Fatoyinbo, T., Castaneda-Moya, E., Cook, B.D., Montesano, P.M., et al. 2021. Storm surge and ponding explain mangrove dieback in southwest Florida following Hurricane Irma. *Nature communications*. 12, 4003. <https://doi.org/10.1038/s41467-021-24253-y>
- Morris, R.L., Fest, B., Stokes, D., Jenkins, C., Swearer, S.E. 2023. The coastal protection and blue carbon benefits of hybrid mangrove living shorelines. *Journal of Environmental Management*. 331, 117310, <https://doi.org/10.1016/j.jenvman.2023.117310>
- Nazar, F., Rahman., Nopiana, M., Rifqi, M. 2021. Analisis akar masalah kemiskinan masyarakat pesisir dengan pendekatan sistem sosial ekologi. *Eqien*. 8(1): 86 – 93.

- Nordhaus, I., Wolff, M., Diele, K. 2006. Litter processing and food intake of the mangrove crab *Ucides cordatus* in a high intertidal forests in Northern Brazil. *Estuarine, Coastal, and Shelf Science*. 67(1-2): 239-250.
- Noor, Y.R., Khazali, M., Suryadiputra, I.N.N. 2006. *Panduan Pengenalan Mangrove di Indonesia*. Bogor (ID): PHKA/WI – PI.
- Nyangoko, B.P., Berg, H., Mangrora, M.M., Shalli, M.S., Gullstrom, M. 2022. Local perceptions of changes in mangrove ecosystem services and their implications for livelihoods and management in the Rufiji Delta, Tanzania. *Ocean and Coastal Management*. 219, 106065, <https://doi.org/10.1016/j.ocecoaman.2022.106065>
- Owuor, M.A., Mulwa, R., Otieno, P., Icely, J., Newton, A. 2019. Valuing mangrove biodiversity and ecosystem services: A deliberative choice experiment in Mida Creek, Kenya. *Ecosystem Services*. 40, 101040, <https://doi.org/10.1016/j.ecoser.2019.101040>
- Rahman., Wardiatno, Y., Yulianda, F., Rusmana, I. 2019. Production ratio of seedlings and density status of mangrove ecosystem in coastal areas of Indonesia. *Advances in Environmental Biology*. 13(6): 13 – 21.
- Rahman., Wardiatno, Y., Yulianda, F., Rusmana, I. 2020a. Socio-ecological system of carbon-based mangrove ecosystem on the coast of West Muna Regency, Southeast Sulawesi, Indonesia. *AAEL Bioflux*. 13(2): 518-528.
- Rahman., Wardiatno, Y., Yulianda, F., Rusmana, I. 2020b. Sebaran spesies dan status kerapatan ekosistem mangrove di pesisir Kabupaten Muna Barat. *JPSL*. 10(3): 461-478. <http://dx.doi.org/10.29244/jpsl.10.3.461-478>
- Rahman., Lokollo, F.F., Manuputty, G.D., Hukubun, R.D., Krisye., Wawo, M., Wardiatno, Y. 2024a. A review on the biodiversity and conservation of mangrove ecosystems in Indonesia. *Biodiversity and Conservation* **33**, 875-903. <https://doi.org/10.1007/s10531-023-02767-9>

- Rahman., Ceantury, A., Tuahatu, J.W., Lokollo, F.F., Supusepa, J., Hulopi, M., Permatahati, Y.I., Lewerissa, A., Wardiatno Y. 2024b. Mangrove ecosystem in Southeast Asiaregion: mangrove extent, blue carbon potential and CO₂ emission in 1996-2020. *Science of the Total Environment* 915(3): 1-12. <https://doi.org/10.1016/j.scitotenv.2024.170052>
- Rahman., Maryono., Saiful., Wardiatno, Y. 2024c. *Penilaian Jasa Ekosistem Mangrove: Pendekatan Ekologi, Sosial, dan Ekonomi*. Bogor (I.D.). IPB Press. 98p.
- Sievers, M., Brown, C.J., McGowan, J., Turschwell, M.P., Buelow, C.A., Holgate, B. et al. 2023. Co-occurrence of biodiversity, carbon storage, coastal protection, and fish and invertebrate production to inform global mangrove conservation planning. *Science of the Total Environment*. 904, 166357. <https://doi.org/10.1016/j.scitotenv.2023.166357>
- Sunkur, R., Kantamaneni, M., Bokhoree, C., Ravan, S. 2023. Mangroves' role in supporting ecosystem-based techniques to reduce disaster and adapt to climate change: A review. *Journal of Sea Research*. 196, 102449. <https://doi.org/10.1016/j.seares.2023.102449>
- van Hespen, R., Hu, Z., Borsje, B., De Dominicis, M., Friess, D.A., Jevrejeva, S., Kleinhans, M.G., Maza, M., van Bijsterveldt, C.E.J., der Stocken, T.V., van Wesenbeeck, B., Xie, D., Bouma, T.J. 2023. Mangrove forests as a nature-based solution for coastal flood protection: Biophysical and ecological considerations. *Water Science and Engineering*. 16(1): 1-13. <https://doi.org/10.1016/j.wse.2022.10.004>
- Yulianda, F. 2007. Ekowisata bahari sebagai alternatif pemanfaatan sumber daya pesisir berbasis konservasi. *Seminar Sains Departemen Manajemen Sumber daya Perairan*. Bogor: FPIK-IPB.
- Yulianti, P., Wardiatno, Y., Samosir, A. M. 2013. Mangrove ecosystem resilience to sea level rise: a case study of Blanakan Bay, Subang Regency, West Java, Indonesia. *Aquatic Science and Management*. 1(1): 63 – 71.