

# Genetic Variation for Tolerance to High Temperatures in Tomato Using Critical Sterility Temperature.

## ABSTRACT

Tomato (*Solanum lycopersicum* L.) reproduction is influenced by temperature, prominently impacting yield. The ideal daytime range is between 25°C and 30°C and the nighttime temperature is around 20°C for tomato and deviations of a few degrees from these temperatures can have negative effects. The study assessed the effect of temperature on tomato pollen germination and aimed to ascertain the critical sterility temperature for pollen germination. In an open field, pollen from 10 genotypes (Anagha, Manuprabha, Vellayani Vijay, Akshaya, Nandi, IHR26372, Pusa Rohini, Arka Sourabh, Arka Rakshak, and Arka Vikas) underwent in-vitro temperature exposure ranging from 32 to 40°C at 2°C intervals. Variations in in-vitro pollen germination percentage were observed across the genotypes, notably revealing a sharp decline after 36°C. This decline indicated a pivotal shift in pollen germination and viability. Genotypes, Anagha and Arka Sourabh, were identified as temperature tolerant and susceptible, respectively. The findings present an opportunity to incorporate temperature-dependent pollen response functions and identified genotypes into tomato breeding programs, offering significant implications for enhancing yield under varying temperature conditions.

Commented [71]: IN CURSIVES

*Keywords: Temperature, Pollen germination, Tomato, Critical temperature, Pollen selection*

## 1. INTRODUCTION

The tomato (*Solanum lycopersicum* L.) is a globally significant horticultural crop with substantial economic value, but it faces increasing challenges from rising temperatures due to global climate change. When temperatures exceed 25 °C, there is a notable decrease in fruit quantity, weight, and seed count per fruit (1). Exposure to extreme heat (45 °C) for short periods can induce programmed cell death (PCD), release of cytochrome c, and production of caspase-like enzymes (2). The plant's reproductive phase is typically more susceptible to high temperatures than its vegetative phase (3,4). Elevated temperatures during flowering significantly affect pollen viability and germination, leading to reduced fruit set and yield (1). Various crops, including bell pepper, corn, tomato, cowpea, and soybean, exhibit sensitivity to above-optimal temperatures during fruit set (5, 6,7, 8,9). Numerous studies have focused on heat stress effects on pollen development (10,11,12,13). Research by (11) revealed that heat stress causes male sterility in tomatoes, but male sterile plants grown at 29 °C could produce fruit when pollinated by pollen developed at 25 °C. In tomatoes, pollen germination and tube development are impaired at temperatures above 30 °C (14). Research indicates that the ideal temperature for pollen germination in vitro ranges from 15 to 22 °C, while 25 °C is optimal for in vivo germination (15, 16). Pollen viability and germination success can be negatively affected by temperatures exceeding 30°C. The male reproductive organs are particularly susceptible to heat stress, which can lead to a decrease in the quantity of pollen grains produced and released by the anther, as well as reduced pollen viability and germination rates

Commented [72]: IN CURSIVES

Commented [73]: HERE INSERT (STAMENS)

(17,18). Pollen germination and viability are especially vulnerable to high temperatures, making them key indicators of heat tolerance.

To assess high-temperature tolerance in tomato, researchers employ various physiological and biochemical tests, with the pollen germination percentage assay playing a crucial role. These tests provide direct insights into how well tomato plants reproduce under heat stress conditions (11). By examining pollen responses to elevated temperatures, plant breeders can identify and choose varieties with enhanced tolerance characteristics, thus improving breeding programs aimed at developing heat-resistant tomato cultivars (19). This approach offers a rapid and efficient screening method to aid in the selection of robust tomato varieties capable of thriving in warmer environments. This study seeks to examine the methodology for evaluating tomato varieties' high-temperature tolerance using pollen germination assays.

**Commented [74]:** ELIMINATE

## 2. MATERIAL AND METHODS

### 2.1 Plant growth

The study was carried out at the Plant Physiology Department of the Agriculture College in Vellayani, situated at 8°5'N latitude, 76°9'E longitude, and 29 meters above sea level. The research involved ten tomato varieties: Anagha, Manuprabha, Vellayani Vijay, Akshaya, Nandi, IIHR26372, Arka Sourabh, Arka Rakshak, and Arka Vikas. These varieties were grown in pots containing an equal mixture of farmyard manure (FYM), sand, and soil. The pots were positioned in an open field, and the plants received nutrients and pest control treatments as recommended by Kerala Agricultural University.

**Commented [75]:** Pusa Rohini,? It is missing

**Commented [76]:** HOW MANY POTS OR TOMATO PLANTS FOR EACH VARIETY (GENOTYPE) WERE USED?

### 2.2 Temperature treatments

Mature pollen grains were harvested from the chosen genotypes between 8:00 and 10:00 AM. The collected pollen was then exposed to various temperatures (ranging from 32°C to 40°C, in 2°C increments) for two hours. This incubation took place in a BOD incubator (Rotary shaker cum BOD, Rotek, ROSI-1) using a standardized pollen germination medium in petri plates.

**Commented [77]:** IT IS MISSING TO POINT OUT HOW MANY REPEATS OF EACH VARIETIES (GENOTYPES)

### 2.3 Pollen germination measurements

After incubation, pollen germination was assessed using a compound microscope (Leica DC 7.5 V, 10X). The germination percentage was calculated by determining the ratio of germinated pollen to the total pollen count per field of view, expressed as a percentage. The replicated data on maximum pollen germination underwent analysis using two-way ANOVA through GRAPES statistics (20). The critical temperature, where only 20-30% of the pollen grains germinated, was also determined for each variety.


**Commented [78]:** BEFORE INCUBATION IT COULD BE BETTER TO THE APPLY TRETRAZOLIUM TEST, ALSO KNOWN AS TZ TEST.

## 3. RESULTS AND DISCUSSION

### 3.1 Pollen germination

Pollen grains started germinating in about 30 minutes on contact with the in vitro pollen germination medium. The effect of five constant temperature regimes from 32 to 40 °C at 2 °C intervals on the in vitro pollen germination of ten tomato genotypes after 2hrs incubation was observed and expressed as the percentage of germinated pollen grains (Table 1). The pollen germination percentage decreased significantly with increase in temperature. Figure 1 shows the variation for pollen germination in response to temperature of two genotypes for clarity. Maximum germination percentage ranged from 24.57 % (Manuprabha) to 46.63% (Anagha), with a mean of 33.01 % were observed at 32°C. After 36°C, the mean pollen germination

percentage decreased below 20 %. The pollen germination observed for the tomato genotypes at a temperature of 40°C was very low; and the germination rates were below 1% in three genotypes (Arka sourabhv, Arka Vikas and Pusa Rohini). Differences among pollen germination means were found statistically significant and the highest germination percentages were observed in Anagha (24.63 %) followed by Vellayani Vijay (22.79 %) and IIHR 26372 (21.5 %) and the three genotypes which showed least mean pollen germination were Pusa rohini (15.13%), Manuprabha (15.37 %) and Arka Sourabhv (16.04%). Fig. 1 shows the variation in pollen germination in response to temperature of two tomato genotypes for clarity.



**Fig. 1. In vitro pollen germination responses to temperature of two tomato genotypes (Anagha and Arka Saurabhv).**

### 3.2 Critical sterility temperature

Critical temperature was assessed as the temperature where 20-30% of the pollen only germinates. Critical temperatures for pollen germination differed among genotypes. There was considerable variability in the critical temperatures for pollen germination among the genotypes. The genotypes Arka Sourabhv, Arka Vikas, Pusa Rohini, Manuprabha and Akshay showed less than 20 % germination percentage after 34°C while Vellayani vijay, Anagha, Arka Rakshak, IIHR 26372 and Nandi has showed pollen germination percentage less than 20 only after 36°C. The mean pollen germination % at each temperature were taken and was observed that germination percentage was 21.04 % at 36°C and decreased drastically to 9.29 % at 38°C. So 36°C was identified as the critical temperature for pollen selection.

### 3.3 Discussion

Temperature plays a crucial role in influencing plant growth, pollen production, and reproductive success. Most plant species thrive within a specific temperature range for pollen germination, with tomato preferring 20-30°C (11). Studies have demonstrated that temperatures outside this range significantly diminish germination rates and pollen tube length (21). This research corroborates previous findings, observing a marked decrease in pollen germination under elevated temperature conditions. For instance, (12) reported a considerable decline in tomato pollen viability and germination when daytime temperatures

**Commented [79]:** IT IS IMPORTANT TO INCLUDE A SECOND GRAPHIC, THAT IS, A GRAPHIC INCLUDING ALL THE RESULTS WITH ALL THE GENOTYPES



#### 4. CONCLUSION

Significant genetic variations were noted in tomato genotypes regarding pollen germination and critical sterility temperature. 36°C was identified as critical temperature for pollen selection where 20% pollen germination was observed. Beyond critical temperature, pollen germination percentage decreased below 20 % except in Manuprabha, Arka Sourabhv, and Arka vikas in which the percentage germination reduced below 20 after 34°C itself. Anagha, Vellayani vijay and IIHR 26372 which showed higher germination at the critical sterility temperature was selected as tolerant and Arka Saurabhv, Manuprabha and Pusa Rohini with least germination percentage as susceptible. The sensitivity of pollen germination rate and tube growth makes them valuable indicators of thermal tolerance, reflecting the ability of pollen to function in elevated temperatures. Assessing the performance of different tomato varieties in controlled high-temperature environments can provide valuable insights into their thermal tolerance mechanisms. However, further studies are required to confirm heat tolerance with respect to various attributes contributing to tolerance mechanisms.

#### REFERENCES

1. Peet MM, Sato S, Gardner RG. Comparing heat stress effects on male-fertile and male-sterile tomatoes. *Plant cell environ.* 1998; 21: 225-231.
2. Qu G, Liu X, Zhang Ya-Li, Yao D, Ma Q, Yang M, et al. Evidence for programmed cell death and activation of specific caspase-like enzymes in the tomato fruit heat stress response. *Planta.* 2009; 229. 1269-79.
3. Ruan YL, Jin Y, Yang YJ, Li GJ, Boyer JS. Sugar input, metabolism, and signaling mediated by invertase: roles in development, yield potential, and response to drought and heat. *Molecular Plant.* 2010;3(6):942–955.
4. Zinn KE, Tunc-Ozdemir M, Harper JF. Temperature stress and plant sexual reproduction: uncovering the weakest links. *J. Exp. Bot.* 2010; 61: 1959-1968.
5. Erickson AN, Markhart AH. Flower developmental stage and organ sensitivity of bell pepper (*Capsicum annuum* L.) to elevated temperature. *Plant Cell Environ.* 2002; 25:123–130.
6. Herrero MP, Johnson RR. High temperature stress and pollen viability of maize. *Crop Sci.* 1980; 20:796–800.
7. Sato S, Peet MM, Thomas JF. Determining critical pre and postanthesis periods and physiological processes in *Lycopersicon esculentum* Mill. exposed to moderately elevated temperatures. *J. Exp. Bot.* 2002; 53:1187–1195.
8. Hall AE. Comparative ecophysiology of cowpea, commonbean and peanut. 2004;p. 271–325. In H.T. Nguyen and A. Blum (ed.) *Physiology and biotechnology integration for plant breeding*, Marcel Dekker, New York.
9. Ferris R, Wheeler TR, Hadley P, Ellis RH. Recovery of photosynthesis after environmental stress in soybean grown under elevated CO<sub>2</sub>. *Crop Sci.* 1998; 38:948–955.
10. Raja MM, Vijayalakshmi G, Naik ML, Basha PO, Sergeant K, Hausman JF, et al. Pollen development and function under heat stress: from effects to responses. *Acta Physiologiae Plantarum.* 2019;41(4):47.
11. Pressman E, Peet MM, Pharr DM. The effect of heat stress on tomato pollen characteristics is associated with changes in carbohydrate concentration in the developing anthers. *Ann. Bot.* 2002;90(5):631–636.
12. Firon N, Shaked R, Peet MM, Pharr DM, Zamski E, Rosenfeld K. Pollen grains of heat tolerant tomato cultivars retain higher carbohydrate concentration under heat stress conditions. *Sci. Hortic.* 2006;109(3):212–217.

13. Frank G, Pressman E, Ophir R, Althan L, Shaked R, Freedman M, et al. Transcriptional profiling of maturing tomato (*Solanum lycopersicum* L.) microspores reveals the involvement of heat shock proteins, ROS scavengers, hormones, and sugars in the heat stress response. *J. Exp. Bot.* 2009;60(13):3891–3908.
14. Vasil IK. Physiology and culture of pollen. *Int. Rev. Cytol.* 1987;107:127–174.
15. Kakani VG, Reddy KR, Koti S, Wallace TP, Prasad PVV, Reddy VR, et al. Differences in *in vitro* pollen germination and pollen tube growth of cotton cultivars in response to high temperature. *Ann. Bot.* 2005;96(1):59–67.
16. Dempsey W. Effects of temperature on pollen germination and tube growth. *Rep. Tomato Genet. Coop.* 1970; 20:15–16.
17. Alsamir M, Ahmad K, Mahmood NM, Trethowan R. Identification of High-Temperature Tolerant and Agronomically Viable Tomato (*S. lycopersicum*) Genotypes from a Diverse Germplasm Collection. *Advances in crop science and Technology.* 2017; 5:299.
18. Rieu I, Twell D, Firon N. Pollen development at high temperature: From acclimation to collapse. *Plant Physiology*: 2017; pp. 01644.2016.
19. Sato S, Kamiyama M, Iwata T. Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. *Ann. Bot.* 2006; 97: 731-738.
20. Gopinath et al. grapesAgri1: Collection of Shiny Apps for Data Analysis in Agriculture. *Journal of Open Source Software.* 2021; 6(63), 3437, <https://doi.org/10.21105/joss.03437>
21. Paupière MJ, van Heusden AW, Bovy AG. The metabolic basis of pollen thermo-tolerance: perspectives for breeding. *Metabolites* 2014; 4:889–920.
22. Dane F, Hunter AG, Chambliss OL. Fruit set, pollen fertility and combining ability of selected tomato genotypes under high temperature field conditions. *J. Am. Hort. Sci.* 1991; 116(5): 906-910.
23. Marine JP, Haperen PV, Rieu I, Richard GFV, Yury MT, Arnaud GB. Screening for pollen tolerance to high temperatures in tomato. *Euphytica* 2017; 213: 130-138.
24. Salem AM, Kakani VG, Koti S, Reddy KR. Pollen-based screening of soybean genotypes for high temperatures. *Crop Sci.* 2007; 47: 219.
25. Song G, Chen Q, Tang, Canming. The effects of high-temperature stress on the germination of pollen grains of upland cotton during square development. *Euphytica.* 2014; 200.
26. Reddy KR, Kakani VG. Screening Capsicum species of different origins for high temperature tolerance by *in vitro* pollen germination and pollen tube length. *Sci. Hortic.* 2007;112(2):130–135.
27. Ranasinghe CS, Waidyarathna KP, Pradeep APC, Meneripitiya MSK. Approach to screen coconut varieties for high temperature tolerance by *in-vitro* pollen germination *Cocos*, 19 (2010), pp. 01-1.