

EFFECT OF POTASSIUM AND ZINC ON SOIL PROPERTIES AND FRACTION OF POTASSIUM AND ZINC IN SOIL

Abstract

A field experiment entitled “Effect of potassium and zinc on yield and nutrient content of chickpea and fractions of potassium and zinc in soil” was carried out on Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand during *rabi* season of the year 2020-21. The experiment comprising nine treatment combinations consisted of three levels of potassium (0, 20 and 40 kg/ha) and three levels of zinc (0, 2.5 and 5.0 kg/ha). These treatments were evaluated in Randomized Block Design (factorial) with four replications. The soil of the experimental plot was loamy sand in texture, alkaline in reaction, low in organic carbon and available nitrogen, medium in available phosphorus, potassium and zinc. Results of the experiment revealed that the available K_2O in soil was significantly increased due to application of 40 kg K_2O/ha (K_2) and was found to be at par with application of 20 kg K_2O/ha (K_1). All other chemical properties of soil (pH, EC, organic carbon, available N, P_2O_5 , S, Fe, Mn, Zn and Cu) were not differ significantly due to application of different levels of potassium. Levels of zinc also failed to produce its significant effect on chemical properties of soil except available Zn in soil as it was significantly increased with the treatment of Z_2 (5.0 kg Zn/ha) than control but was found to be at par with treatment Z_1 (2.5 kg Zn/ha). Moreover, no any chemical property of soil differ due to the interaction of different levels of potassium and zinc. The water soluble K and exchangeable K fractions in soil significantly increased with the treatment of K_2 (40 kg K_2O/ha) otherwise non-exchangeable K, total K and different zinc fractions in soil did not differ significantly by the different levels of potassium. Different potassium fractions and total Zn fraction in soil were found to be non-significant under the different zinc levels but all the zinc fractions in soil (water soluble plus exchangeable Zn, carbonate bound Zn, Fe/Mn oxide bound Zn, organically bound Zn and residual Zn) significantly increased with the treatment of Z_2 (5.0 kg Zn/ha) and was remained at par with treatment Z_1 (2.5 kg Zn/ha). The interaction between potassium and zinc failed to influence significantly on potassium and zinc factions in soil.

Key Point: Potassium, Zinc

Introduction

Potassium (K) is one of the important nutrient among 17 vital nutrients required for the growth and reproduction of crops. Potassium has a great buffering action and stabilizes various enzymes system. It acts as a major cation for the maintenance of cation-anion balance. Its beneficial action on crop quality shows better utilization of nitrogen and increased protein formation.

Potassium application has been neglected in many countries, including India, which has resulted in soil K depletion in agricultural ecosystems and a decline in crop yields (Regmi *et al.*, 2002). Among essential nutrient elements, Potassium is the third major element taken up

by the plants. Plants absorb it in larger amounts as compared to other mineral elements except nitrogen. It has utmost importance for imparting drought and disease resistance and has synergistic effect on crops with nitrogen and phosphorus (Das, 1999). Higher yields and crop quality can be obtained at optimal N:K nutritional ratios. It is an essential macronutrient required for proper development of plants. Potassium has been considered as the “quality element” for crop production. Pulse growing Agroecological region of India vary widely in their K supplying capacity. Light textured alluvial soils, red and lateritic soils and shallow black soils with low levels of available K and even black cotton soil needs K supplementation to enhance the productivity. Based on the number of field studies, it can be suggested that the application of 20-40 kg K₂O/ha and foliar application of 1-2% of KNO₃ is beneficial for higher pulse production (Ali and Srinivasa Rao, 2001).

Most of the Potassium in soil exists in various insoluble rocks, minerals and sedimentary materials. Based on its availability to plants, soil potassium can be classified into three main groups *i.e.*, unavailable, readily available and slowly available K. The bulk of soil potassium (about 98% of total K) usually exists in unavailable form and is found in primary (micas and feldspars) and secondary (illite group) clay minerals (Attoe and Truog, 1945). The readily available K constitutes only 1-2% of total K and exists in soil in two forms, *viz.*, solution and exchangeable K adsorbed on soil colloidal surface (Brady and Well, 2002). It is not a constituent of organic structures, but regulates enzymatic activities (over 60 enzymes require K for activation), translocation of photosynthesis and considerably improves seed yield of chickpea if applied as a fertilizer (Samiullah and Khan, 2003, Singh *et al.*, 1994 and Verma, 1994).

Among the micronutrients, zinc plays a vital role in plant growth and development. Zinc has been the micronutrient needed by crops especially pulses in sufficiently large quantity. Zinc also catalyses the biosynthesis of indole acetic acid, acting as metal activator of the enzyme, there by ultimately increasing the crop yield. Moreover, it controls the equilibrium between CO₂, water and carbonic acid in plant metabolism and helps in synthesis of nucleic acids, proteins and stimulates seed formation. Its deficiency retards photosynthesis and nitrogen metabolism.

The plants exhibited lower rate of protein synthesis and protein accumulation under zinc deficiency. Zinc also plays important role in physiological process of plants through synthesis of hormones essential for growth and reproduction. Zinc plays an important role in metabolism both in plants as well as in animals by acting as essential component of enzyme,

RNA, electron carrier *etc.*, and acts as a functional, structural and regulatory cofactor of a large number of enzymes. The proteinases, peptidases, carbonic, dehydrogenase, anhydrase *etc.*, are the examples of the metallo-enzymes in which zinc is an integral part. Zinc is an essential component of RNA polymerase and provides structural integrity to ribosomes. Soil is the principal source of zinc for plants. The accumulation of zinc in edible parts of plant serves as zinc source for primary consumers. Unfortunately, about 50% of Indian soils are deficient in zinc and expected to further increase up to 63% by 2025 which imparting zinc malnutrition in population especially in children (Shukla *et al.*, 2014), Singh (2010) reported that one third of the world population is at the risk of zinc malnutrition due to inadequate dietary intake of zinc resulting from wide spread hidden hunger of zinc in seeds and feeds. Also, the intensive cropping systems of high yielding varieties have led to depletion of micronutrients, especially zinc.

Materials and methods


A field experiment was carried out on Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand (Gujarat) during *rabi* season of the year 2020-21. The soil of the experimental plot was loamy sand in texture, alkaline in reaction, low in organic carbon and available nitrogen, medium in available phosphorus, potassium and zinc. The experiment comprising nine treatment combinations consisted of three levels of potassium (0, 20 and 40 kg/ha) and three levels of zinc (0, 2.5 and 5.0 kg/ha). These treatments were evaluated in Randomized Block Design (factorial) with four replications. The fertilizer application was done with fixed dose of nitrogen at 20 kg/ha and phosphorus at 40 kg/ha. Potassium and Zinc application was done according to the treatments. The nutrient of N, P, K and Zn were applied by using source of Urea, DAP, MOP and Zinc Sulphate respectively. The experimental data recorded for Physico-chemical Properties and Fractions of Potassium and Zinc were statistically analyzed for level of significance.

Table 1 Methods adopted for analysis of potassium fractions in soil

Sr. No.	K fractions	Method	References
1.	Water soluble K	Flame emission spectroscopy	Richards (1954)

2.	Exchangeable K	Neutral N NH ₄ OAC	Hanway and Heidel (1952)
3.	Non-exchangeable K	HNO ₃ extractable	Wood and De Turk (1941)
4.	Total K	HF-HClO ₄ digestion	Page <i>et al.</i> (1982)

Flow chart 1 : Flow chart of sequential fraction scheme by Miller *et al.* (1986)



Result and Discussion

EFFECT OF POTASSIUM AND ZINC ON CHEMICAL PROPERTIES OF SOIL

The soil properties (pH, EC and OC) analysis after harvest of chickpea were not differ significantly could be ascribed to the buffering capacity of soil. Similar results were found by Keram *et al.* (2012) and Sutariya *et al.* (2013).

Table 2 Effect of potassium and zinc on available N, P₂O₅, K₂O and S in soil after harvest of chickpea crop

Treatments	Soil chemical properties		
	pH (1:25)	EC (dS/m) (1:25)	OC (%)
K levels (kg/ha)			
K₀	7.75	0.140	0.39
K₁	7.79	0.142	0.39
K₂	7.81	0.145	0.40
S. Em±	0.05	0.002	0.01
CD (P = 0.05)	NS	NS	NS
Zn levels (kg/ha)			
Z₀	7.80	0.141	0.38
Z₁	7.78	0.142	0.40
Z₂	7.77	0.144	0.40
S. Em±	0.05	0.002	0.01
CD (P = 0.05)	NS	NS	NS
K × Zn	NS	NS	NS
CV%	2.38	3.88	4.40

A perusal of the data presented in Table 3 revealed that available N, P and S in soil after harvest of chickpea did not differ significantly by the different levels of potassium as well as zinc. But the treatment K₂ (40 kg K₂O/ha) and treatment Z₂ (5.0 kg Zn/ha) recorded maximum value of available N in soil (140 and 140 kg/ha, respectively). Numerically maximum available phosphorus (50.1 and 50.0 kg/ha, respectively) was found in treatment K₂ (40 kg K₂O/ha) and Z₂ (5.0 kg Zn/ha) than other treatments.

The significant increase in available potassium in soil by the application of 20 and 40 kg K₂O/ha might be attributed to solubilization of native status of potassium and such increase in available potassium status of the soil at harvest of the crop may also be due to direct addition of potassium to available pool of the soil. Similar results were also reported by Tandon (1987), Gajghane *et al.* (2015), Kadam *et al.* (2016) and Jat *et al.* (2017).

Table 3: Effect of potassium and zinc on available N, P₂O₅, K₂O and S in soil after harvest of chickpea crop

Treatments	Available nutrients
------------	---------------------

	N	P ₂ O ₅	K ₂ O	S
	(kg/ha)			(mg/kg)
K levels (kg/ha)				
K ₀	135	48.8	219	15.7
K ₁	138	49.5	228	16.2
K ₂	140	50.1	232	16.5
S. Em±	2	0.6	3	0.2
C.D. (P = 0.05)	NS	NS	9	NS
Zn levels (kg/ha)				
Z ₀	136	49.1	225	15.7
Z ₁	137	49.4	226	16.2
Z ₂	140	50.0	228	16.5
S. Em±	2	0.6	3	0.2
C.D. (P = 0.05)	NS	NS	NS	NS
K × Zn	NS	NS	NS	NS
CV%	4.33	3.98	4.54	5.03

A perusal of the data presented in Table 3 revealed that available Fe, Mn and Cu in soil after harvest of chickpea did not differ significantly by the different levels of potassium as well as zinc. But the treatment Z₂ (5 kg K₂O/ha) and recorded maximum value of available Zn in soil (0.78 mg/ha, respectively). But numerically maximum available (Fe 1.35 and 1.34 mg/kg, 4.04 and 4.03 mg/kg, Mn 2.12 and 2.11 mg/kg and Cu 1.35 and 1.34 mg/kg respectively) was obtained under treatment of application of 40 kg K₂O/ha (K₂) and 5.0 kg Zn/ha (Z₂) than other treatments.

Table. 4: Effect of potassium and zinc on available Fe, Mn, Zn and Cu in soil after harvest of chickpea

Treatments	DTPA extractable micronutrient (mg/kg)			
	Fe	Mn	Zn	Cu
K levels (kg/ha)				
K ₀	3.77	2.03	0.73	1.29
K ₁	3.92	2.07	0.75	1.32
K ₂	4.04	2.12	0.76	1.35
S. Em±	0.08	0.03	0.01	0.02
CD (P = 0.05)	NS	NS	NS	NS
Zn levels (kg/ha)				
Z ₀	3.78	2.03	0.71	1.31
Z ₁	3.92	2.08	0.75	1.32
Z ₂	4.03	2.11	0.78	1.34
S. Em±	0.08	0.03	0.01	0.02
CD (P = 0.05)	NS	NS	0.03	NS
K × Zn	NS	NS	NS	NS
CV%	6.65	4.66	5.33	3.95

1.2 EFFECT OF POTASSIUM AND ZINC ON FRACTIONS OF POTASSIUM AND ZINC IN SOIL

1.2.1 Potassium fractions in soil

The data pertaining to the different potassium fractions *viz.*, water soluble, exchangeable, non-exchangeable and total fractions in 0-15 cm depth of soil as influenced by application of potassium and zinc is presented in Table 5.

1.2.1.1 Water soluble K

1.2.1.1.1 Effect of potassium

The data presented in Table 5 indicated the significant effect of different levels of potassium on water soluble K in soil. The range of water soluble K was varied 25.35-27.59 mg/kg due to application of potassium. The treatment of application of 40 kg K₂O/ha recorded significantly highest water soluble K (27.59 mg/kg) than rest of the treatments.

1.2.1.1.2 Effect of zinc

The water soluble K did not differ significantly due to the treatment containing application of zinc (Table 5). The water soluble K was varied from 25.95 -26.66 mg/kg by the application of different levels of zinc. But maximum water soluble K (26.66 mg/kg) was found in treatment of application of 5.0 kg Zn/ha than other treatments.

1.2.1.1.3 Interaction effect

The interaction effect between different levels of potassium and zinc was found to be non-significant with respect to water soluble K fractions in soil after the harvest of the chickpea Table 5.

1.2.1.2.1 Effect of potassium

The data presented in Table 5 indicated that the exchangeable K in soil was significantly influenced with the treatment containing application of potassium. The exchangeable K was varied from 72.03-75.61 mg/kg by the application of potassium. The treatment of application of 40 kg K₂O/ha (K₂) recorded significantly highest exchangeable K (75.61 mg/kg) than rest of the treatments.

1.2.1.2.2 Effect of zinc

The treatment containing application of zinc did not show any significant effect on exchangeable K in soil. The exchangeable K was varied from (73.98-74.89 mg/kg) by the different levels of zinc. But maximum exchangeable K was (74.89 mg/kg) observed by treatment of application of 5.0 kg Zn/ha (Z₂) Table 6.

1.2.1.2.3 Interaction effect

The interaction between different levels of potassium and zinc did not show any significant effect on exchangeable K in soil (Table 6).

1.2.1.3 Non-exchangeable K (mg/kg)

1.2.1.3.1 Effect of potassium and zinc

The non-exchangeable K was not differ significantly by the application of potassium and zinc. But it was obtained higher non-exchangeable K (602 and 593 mg/kg, respectively) with the treatment K₂ (40 kg K₂O/ha) and treatment Z₂ (5.0 kg Zn/ha) as compared to other treatments.

1.2.1.3.2 Interaction effect

The interaction effect of different levels of potassium and zinc was found to be non-significant on non-exchangeable K in soil (Table 6).

1.2.1.4 Total K

1.2.1.4.1 Effect of potassium and zinc

Similarly, different levels of potassium and zinc fail to exert significant effect on total K fraction in soil. The higher value of total K *i.e.*, 35446 and 35011 mg/kg, respectively were recorded under treatment K₂ (40 kg K₂O/ha) and treatment Z₂ (5.0 kg Zn/ha) than other levels of potassium and zinc.

1.2.1.4.2 Interaction effect

The interaction effect of treatment combination of different levels of potassium and zinc was not found to be significant in relation to total K fraction in soil (Table 6).

Table 5: Effect of potassium and zinc on potassium fraction in 0-15 cm depth of soil

Treatments	Water soluble K	Exchangeable K	Non-exchangeable K	Total K
	(mg/kg)			
K levels (kg/ha)				
K ₀	25.35	72.03	578	32861
K ₁	26.15	75.25	587	34530
K ₂	27.59	75.61	602	35446
S. Em±	0.26	0.94	7.2	756
CD (P = 0.05)	0.77	2.75	NS	NS
Zn levels (kg/ha)				
Z ₀	25.95	73.98	585	33442
Z ₁	26.48	74.03	590	34384
Z ₂	26.66	74.89	593	35011
S. Em±	0.26	0.9	7.2	705
CD (P = 0.05)	NS	NS	NS	NS
K × Zn	NS	NS	NS	NS
CV%	3.47	4.39	4.25	7.13

1.2.2 Fractions of zinc in soil

The data pertaining to the different fractions of zinc viz., water soluble plus exchangeable Zn, carbonate bound Zn, Fe/Mn oxide bound Zn, organically bound Zn residual Zn, and total Zn in 0-15 cm depth of soil as influenced by different levels of potassium and zinc is presented in Table 6.

1.2.2.1 Water soluble plus exchangeable zinc

1.2.2.1.1 Effect of potassium

The result revealed that there was no significant effect on water soluble plus exchangeable Zn in soil observed with the different potassium levels. The water soluble plus exchangeable Zn ranged from 0.147 to 0.153 mg/kg observed under the application of different levels of potassium. Numerically maximum value of water soluble plus exchangeable Zn was found in treatment K₂ (40 kg K₂O/ha) than other treatments.

1.2.2.1.2 Effect of zinc

The perusal data presented in Table 6 indicated that different levels of zinc resulted significant increased in form of water soluble plus exchangeable Zn (0.156 mg/kg) in 0-15 cm depth of soil by the treatment of application of 5.0 kg Zn/ha (Z₂) which was found statistically at par with the treatment Z₁ (2.5 kg Zn/ha).

1.2.2.1.3 Interaction effect

The interaction between different levels of potassium and zinc did not show any significant effect on water soluble plus exchangeable Zn in soil Table 6.

1.2.2.2 Carbonate bound Zinc

1.2.2.2.1 Effect of potassium

There was no significant difference observed on carbonate bound zinc due to the treatment containing potassium application (Table 6). The carbonate bound zinc was varied from (1.23 to 1.28 mg/kg) by the potassium. But numerically maximum carbonate bound zinc was observed with treatment of application of 40 kg K₂O/ha (K₂) as compared to other potassium levels.

1.2.2.2.2 Effect of zinc

Results given in Table 6 revealed that treatment of application of 5.0 kg Zn/ha (Z₂) significantly increased the carbonate bound zinc in soil (1.29 mg/kg) than control (Z₀) but was remained at par with treatment Z₁ (2.5 kg Zn/ha).

1.2.2.2.3 Interaction effect

The non-significant interaction effect was found on carbonate bound zinc fractions in soil at harvest of chickpea with the treatment combination of different levels of potassium and zinc (Table 6).

1.2.2.3 Fe/Mn oxides bound Zinc

1.2.2.3.1 Effect of potassium

The treatment containing application of different levels of potassium did not show any significant effect on Fe/Mn oxides bound zinc fraction in soil. But it was observed Fe/Mn oxides bound zinc fraction range from (7.46 to 7.90 mg/kg) with the levels of potassium up to 40 kg K₂O/ha (K₂) Table 6. Moreover, maximum value of Fe/Mn oxides bound zinc (mg/kg) was found under treatment K₂ (40 kg K₂O/ha).

1.2.2.3.2 Effect of zinc

The data presented in Table 6 indicated the Fe/Mn oxides bound zinc fractions were influenced significantly with the different levels of zinc. Significantly higher in soil (7.93 mg/kg) was recorded under the treatment of application of 5.0 kg Zn/ha which was found to be at par with treatment Z₁ (2.5 kg Zn/ha).

1.2.2.3.2 Interaction effect

The interaction effect of different levels of potassium and zinc was found to be non-significant on Fe/Mn oxides bound zinc fraction in soil Table 6.

1.2.2.4 Organically bound zinc

1.2.2.4.1 Effect of potassium

Organically bound zinc fraction in soil was not affect significantly due to the different levels of potassium (1.5). But organically bound zinc was varied from (2.91 to 3.05 mg/kg) with the treatment of application of potassium up to 40 kg K₂O/ha. Numerically maximum organically bound zinc (3.05 mg/kg) was found under the treatment K₂ (40 kg K₂O/ha) as compared to other treatments.

1.2.2.4.2 Effect of zinc

Table 6 showed that the significantly increased due to application of different levels of zinc. Significantly increased organically bound zinc fraction in soil was (3.05 mg/kg) found with the treatment containing application of 5.0 kg Zn/ha (Z₂) than control but it was remained at par with treatment (Z₁) 2.5 kg Zn/ha.

1.2.2.4.3 Interaction effect

The non-significant interaction effect was found on organically bound zinc fraction in soil after harvest of chickpea with the treatment combination of different levels of potassium and zinc.

1.2.2.5 Residual zinc

1.2.2.5.1 Effect of potassium

A perusal of the data presented in Table 6 revealed that residual zinc fraction in soil did not differ significantly by different potassium levels. However, the residual zinc fraction ranged from (60.30 to 64.04 mg/kg) due to different levels of potassium up to 40 kg K₂O/ha. The higher potassium level *i.e.*, 40 kg K₂O/ha (K₂) recorded numerically maximum residual zinc in soil (64.29 mg/kg) than rest of the treatments.

1.2.2.5.2 Effect of zinc

Different levels of zinc had significant effect on residual zinc fraction in soil (Table 4). The treatment containing application of 5.0 kg Zn/ha recorded significant increased in in soil (64.64 mg/kg) than control but was found to be at par with treatment Z₁ (2.5 kg Zn/ha).

1.2.2.5.3 Interaction effect

The interaction between different levels of potassium and zinc did not show any significant effect on residual zinc fraction in soil Table 6.

1.2.2.6 Total zinc

1.2.2.6.1 Effect of potassium and zinc

The data indicated in Table 6 revealed that different levels of potassium and zinc fail to produce any significant effect on total zinc fraction in soil. But, total zinc fraction in soil is varied from 75.38 to 77.73 mg/kg due to different potassium levels and its maximum value was found in treatment K₂ (40 kg K₂O/ha). Whereas total zinc fraction was range from 74.66 to 78.23 mg/kg with the different levels of zinc and its maximum value was observed in treatment Z₂ (5.0 kg Zn/ha).

1.2.2.6.2 Interaction effect

The interaction between different levels of potassium and zinc did not show any significant difference with respect to total zinc fraction in soil after harvest of chickpea (Table 6).

Table. 6 Effect of potassium and zinc on fraction of zinc 0-15 cm depth of soil

Treatments	Water soluble plus exchangeable	Carbonate bound fraction	Fe/Mn oxides bound fraction	Organically bound fraction	Residual fraction	Total Zn
	(mg/kg)					
K levels (kg/ha)						
K₀	0.148	1.23	7.46	2.91	60.30	75.38

K₁	0.150	1.25	7.68	2.95	62.63	76.53
K₂	0.153	1.28	7.90	3.05	64.29	77.73
S. Em±	0.002	0.02	0.14	0.04	1.20	1.11
CD (P = 0.05)	NS	NS	NS	NS	NS	NS
Zn levels (kg/ha)						
Z₀	0.146	1.22	7.44	2.88	59.89	74.66
Z₁	0.150	1.26	7.70	2.98	62.44	76.75
Z₂	0.156	1.29	7.93	3.05	64.39	78.23
S. Em±	0.002	0.02	0.14	0.04	1.20	1.11
CD (P = 0.05)	0.006	0.06	0.40	0.12	3.51	NS
K × Zn	NS	NS	NS	NS	NS	NS
CV%	5.09	5.53	6.24	4.89	6.68	5.05

References

- Ali, M., and Srinivasa Rao, Ch. (2001). Role of potassium fertilization in improving productivity of pulse crops. In *Proceeding of International Symposium held in New Delhi, India* (pp. 261-267).
- Attoe, O. J. and Truog, E. (1945). Exchangeable and acid soluble K as regards availability and reciprocal relationships. *Soil Science Society of Soil America Proceeding*, 10, 81-86.
- Brady, N. C. and Well, R. R. (2002). *The Nature and Properties of Soils (13th Ed.)*. Personal Education Pvt. Ltd, New Delhi.
- Das, P.C. (1999). Plant Nutrients. In: *Manures and Fertilizers (2nd Ed.)*. Kalyani Publishers, New Delhi, India, pp: 5-10.
- Gajghane, P. G., Toncher, S. S. and Rutz, M. M. (2015). Effect of Potassium and Sulphur Levels on Soils Fertility Status after Harvest of Mustard! *Plant Archives*, 15 (1), 347-351.
- Jat, Ramswaroop., Naga, Sita Ram., Choudhary, Rajsingh. and Mohammad, Irfan. (2017). Effect of Potassium and Sulphur on Quality of Sesame (*Sesamum indicum* L.)! *International Journal of Current Microbiology and Applied Sciences*, 6 (4), 1876-1878
- Kadam, Y. B., Kharche, V. K., Borkar, V. S., Konde, N. M. and Gabhne, V. V. (2016). Effect of sources and levels of potassium in cotton as influenced fractions of soil potassium

under vertisols in Vidarbha region of Maharashtra! *An Asian Journal of Soil Science*, 11 (2), 246-257.

- Keram, K. S., Sharma, B. L. and Sawarkar, S. D. (2012). Impact of Zn application on yield, quality, nutrients uptake and soil fertility in a medium deep black soil (vertisol). *International Journal of Science, Environment and Technology*, 1(5), 563-571.
- Regmi, A., Ladha, J., Pasuquin, E., Pathak, H., Hobbs, P., Shrestha, L. and Duveiller, E. (2002). The role of potassium in sustaining yields in a long-term rice-wheat experiment in the Indo-Gangetic Plains of Nepal. *Biology and Fertility of Soils*, 36(3), 240-247.
- Samiullah and Khan, N. A. (2003). Physiological investigation on interactive effect of P and K on growth and yield of chickpea. *Indian Journal of Plant Physiology*, 8(2), 165-170.
- Shukla, A. K. and Tiwari, P. K. (2014). Progress Report of AICRP on Micro- and Secondary Nutrients and Pollutant Elements in Soils and Plants. IISS, Bhopal.
- Singh, M. V. (2010). Detrimental effect of zinc deficiency on crops productivity and human health. First Global Conference on Biofortification, Harvest Plus, Washington, USA.
- Singh, O. N., Singh, R. S. and Singh, J. P. (1994). Supplementing fertilizer potassium to chickpea. *Journal of Potash Research*, 10, 83-85.
- Sutariya, G. S., Ramdevputra, M. V., Ansodaria, V. V. and Akbari, K. N. (2013). Effects of potassium and zinc nutrition on yield and quality of forage sorghum. *Indian Journal of Agricultural Research*, 47(6), 540-544.
- Tandon H. L. S. (1987). Phosphorus research and agricultural production in India! FDCO, New Delhi.
- Verma, V.K. (1994). Response of irrigated chickpea (*Cicer arietinum* L.) to NPK in light soils and its economics. *Advances in Plant Science*, 7(1), 138-142.