

# **Pedological study under cocoa trees to improve production in the Soubré region (southwest of Ivory Coast).**

## **ABSTRACT**

Maintaining the fertility of cultivated soils in tropical countries such as Côte d'Ivoire remains a major concern for producers because of crop failures. The aim of this study is to determine the soil characteristics of a cocoa plantation in the Soubré region (south-west of Côte d'Ivoire: 6° - 7° N latitude and 5° - 7° W longitude), in order to improve production. In the centre of the plantation, a section of land (50 m × 50 m) was identified. Five soil pits were opened on this section of land, one at each corner and one in the centre. Soil samples were taken from each of the pits, in the 0-20 cm and 20-40 cm layers. Composite samples were then taken at these two depths for the analytical data. The pH, sum of exchangeable bases (S), cation exchange capacity (CEC), saturation rate of exchangeable bases (V), particle size fraction and organic matter content were measured and compared with threshold values known from the literature. The results show that pH (5.6 - 6.8) and assimilable phosphorus [P<sub>ass</sub> (130 - 350 ppm)] increase with depth and are in good proportion. Concentrations of potassium (K<sup>+</sup> : 2.9 - 2.3 cmol/kg), calcium (Ca<sup>2+</sup> : 29.1 - 47.2 cmol/kg) and magnesium (Mg<sup>2+</sup> : 2.7 - 22 cmol/kg), as well as chemical properties (S : 35 - 73 cmol/kg, CEC : 48 - 80 cmol/kg and V : 73 - 90 %) are relatively good for cocoa production. However, organic matter (1.38%) and total nitrogen (N = 0.1 - 0.05%) are low. And high levels of clay (48%) and silt (23%) were noted from the first 20 cm depth. Spreading organic matter in the form of compost could therefore be recommended in order to correct the defective characteristics of the soils caused by the abundance of these two mineral particles: clay and silt, on the one hand, and deficiencies in nitrogen and organic matter, on the other.

**Key words** : cocoa tree - physical and chemical properties - soils - Soubré - Ivory Coast

## **1. INTRODUCTION**

In Ivory Coast, cocoa plays an essential role in the economic and social prosperity of the population [1]. Since the 1970s, the country has remained the world's leading producer, with production estimated in 2022 at 2.4 tonnes [2]. Cocoa provides nearly 40% of Côte d'Ivoire's export earnings and contributes over 20% of the country's GDP [3]. Some 500,000 producers and their families benefit directly from the socio-economic advantages of this sector [4]. Despite this, cocoa production in Côte d'Ivoire faces certain difficulties [5]. Moreover, the Ivorian cocoa farmer has found himself in a situation where he can no longer reproduce his traditional system, due to the lack of forest and fallow land. At the same time, aging plantations continue to produce more or less satisfactory harvests under degraded, nutrient-depleted soils. This situation can compromise the sustainability of cocoa farming and is detrimental to the Ivorian economy [6]. During the 1990s, attempts were made to find solutions, including the adoption of improved seeds and, more specifically, the use of chemical fertilizers to make plantations more profitable. However, this approach has encountered obstacles in its application [7], due to the inaccessibility of fertilizers to farmers because of their high cost [8]. In addition to their unavailability, chemical fertilizers are water and soil pollutants. Faced with this situation, it appeared necessary to find other alternatives, in particular the use of organic matter [9]. Indeed, studies have shown the importance of organic matter such as compost in agriculture, which justifies its use [10]. As part of this study, the Centre de Production Biologique Durable (CPBD), an agricultural company

located in Soubré, wishes to encourage the use of compost in cocoa plantations where yields appear to be falling, with a view to improving production. To do this, it is important to assess the initial soil characteristics of these plantations, in order to ensure sustainable management of this input, which is compost. In concrete terms, this involves : - determining the chemical and physical properties of soils under cocoa trees; - comparing chemical and physical property values with threshold values or reference standards for cocoa farming.

## 2. MATERIALS AND METHODS

### 2.1. Description of study site

The study was carried out at the beginning of the wet season, in 2021, in the cocoa plantation of the Centre de Production Biologique Durable (CPBD: agricultural inputs research center) in the Soubré locality (6° - 7° N latitude and 5° - 7° W longitude), 412 km from Abidjan (Côte d'Ivoire). The equatorial climate is characterized by high rainfall totalling nearly 1,600 mm annually, and good atmospheric humidity [11]. The mean annual temperature ranges from 25°C to 27°C, with little variation of less than 3.5°C [12]. This is an area of dense forest in steep decline under increasing anthropic pressure [13]. The experimental site is a ten-year-old cocoa field located at the top of a gentle slope (2%). The soils encountered are essentially strongly to moderately desaturated ferralsols [14].

### 2.2. Soil characterization of the cocoa farm

The cocoa farm's soils were morphologically and physico-chemically characterized. A square-shaped portion of 2500 m<sup>2</sup> (50 m × 50 m) was delimited in the center of the experimental plot. Five points on this portion were chosen for the opening of five soil pits, one at each corner and one in the center. The profiles were described. The 0 - 20 cm and 20 - 40 cm layers of each pit were sampled. A composite sample was then taken for each depth level. The composite soil samples were sent to the laboratory for analysis. The following parameters were determined: pH, organic matter (OM), assimilable phosphorus, exchangeable bases (Ca, Mg and K), total nitrogen (Nt), cation exchange capacity (CEC), sum of exchangeable bases (S), base saturation rate (V).

**Data analysis:** chemical soil analysis results were compared with threshold values, reference standards or results from previous work.

**Table 1.** Soil parameters and normal threshold values

soil parameters	Normal threshold values
A (%)	10 - 20
L (%)	< 30
A+L (%)	< 30
S (%)	20 - 30
M.O (%)	2 - 3
Nt (%)	> 1
C/Nt	8 - 10
Water pH	6 - 7.5
Pass (ppm)	> 20
Ca <sup>2+</sup> (cmolkg <sup>-1</sup> )	2 - 3
Mg <sup>2+</sup> (cmolkg <sup>-1</sup> )	0.25 - 0.35
K <sup>+</sup> (cmolkg <sup>-1</sup> )	0.15 - 0.35
Na <sup>+</sup> (cmolkg <sup>-1</sup> )	> 1
S (cmolkg <sup>-1</sup> )	> 10

CEC (cmolk <sup>-1</sup> )	5 - 10
V (%)	50

A: clay; L: silt; S: sand. Sources: ([16]; [17]; [18])

### 3. RESULTS

#### Physical parameters: average proportions of sand, silt and clay in soils from the site studied

The results of the particle size analysis of the soils studied are presented in Table 2. Almost 45% of sand is concentrated in the superficial layers (0 - 20 cm) of the soil. Its content decreases in the underlying horizon (20 - 40 cm), reaching around 30%. This is an acceptable level for most soils. From the surface horizon (0 - 20 cm) to that below (20 - 40), silt contents are almost identical, at 26% versus 23% respectively. In contrast to silt and sand contents, clay proportions increase with depth, with higher contents estimated at 48%.

**Table 2.** Average proportions of sand, silt and clay in soils at the study site as a function of depth.

Physical particles	Depths (cm)	
	0 – 20	20 – 40
Sand (%)	45	29
Silt (%)	26	23
Clay (%)	29	48

#### 3.1. Organic matter, pH and available phosphorus (Pass) in cocoa farm soils.

Table 3 summarizes the average contents of organic matter, nitrogen and assimilable phosphorus, as well as the pH values and C/N ratios according to depth. Soil pH values show the soil to be acidic (pH = 5.6) at the surface (0 - 20 cm) and weakly acidic (pH = 6.8) at depth (20 - 40 cm). Overall, organic matter and nitrogen are poorly represented in the soil layer (0 - 40 cm). Organic matter levels are moderate (1.38%) on the surface (0 - 20 cm), but drop to 0.69% in the first 20 cm of the underlying layers. Superficial horizons are depleted of nitrogen (0.1%), which tends to disappear below (0.05%). Compared with nitrogen, assimilable phosphorus, sufficiently supplied at the surface, accumulates in the deep layer at levels 3 times higher (133 ppm vs. 352.7 ppm). On the other hand, the C/N ratio remained stable at all depths (-20 cm or -40 cm), with values close to 11 or even 12.

**Table 3.** Average values for organic matter, nitrogen, pH<sub>water</sub> and assimilable phosphorus (Pass) in soils at the study site, by depth.

Chemical elements	Depths (cm)	
	0 – 20	20 – 40
MO (%)	1.38	0.69
N (%)	0.1	0.05
C/Nt	12	11
Water pH	5.6	6.8
Pass (ppm)	133.1	352.7

### 3.2. Characteristics of the adsorbent complex

Table 4 shows the characteristics of the adsorbent complex in the soil at the study site: alkalis ( $K^+$  and  $Na^+$ ), alkaline earths ( $Ca^{2+}$  and  $Mg^{2+}$ ), sum of exchangeable bases (S), saturation rate (V) and cation exchange capacity (CEC). In terms of alkalinity, potassium ( $K^+$ ) is evenly distributed along the soil profile, from top to bottom, with average levels of around  $2.6 \text{ cmol.kg}^{-1}$ . These levels are considered to be well above the norm, which is  $0.35 \text{ cmol.kg}^{-1}$ . On the contrary, the deep layers are enriched in sodium with a high concentration ( $1.4 \text{ cmol.kg}^{-1}$ ). In terms of alkaline earth elements ( $Ca^{2+}$  and  $Mg^{2+}$ ), the soils at the study site are highly enriched in calcium ( $Ca^{2+}$ ). Moreover, levels rise from  $29 \text{ cmol.kg}^{-1}$  to  $47 \text{ cmol.kg}^{-1}$  in the 0 - 20 cm and 20 - 40 cm layers respectively. Magnesium ( $Mg^{2+}$ ) is well represented at the surface ( $2.7 \text{ cmol.kg}^{-1}$  vs.  $0.5 \text{ cmol.kg}^{-1}$  = threshold value). In addition, it accumulates in the underlying layers with high levels, above the norm ( $21.8 \text{ cmol.kg}^{-1}$ ).

### 3.3. Sum of exchangeable bases (S) and saturation rate (V)

S varies almost twofold from  $35 \text{ cmol.kg}^{-1}$  to  $73 \text{ cmol.kg}^{-1}$  from one layer (0 - 20 cm) to the other (20 - 40 cm). Overall, both horizons, 0 - 20 cm and 20 - 40 cm, are highly saturated with exchangeable bases, with average contents of 73% and 91% respectively, compared with the reference threshold values (50 - 90%).

### 3.4. Cation exchange capacity (CEC)

There was a significant difference between the various CEC values at depths of 0 - 20 cm and 20 - 40 cm. The value  $48 \text{ cmol.kg}^{-1}$  was observed on the surface (0 - 20 cm). Below, double this value was recorded (20 - 40 cm). However, both layers showed capacities well above the normal threshold required for cocoa farming ( $10 - 20 \text{ cmol.kg}^{-1}$ ).

**Table 4.** Adsorbent complex characteristics of study plot soils by depth.

Chemical elements	Depths (cm)	
	0 – 20	20 – 40
CEC ( $\text{cmol.kg}^{-1}$ )	48	80
S ( $\text{cmol.kg}^{-1}$ )	34.8	72.7
V (%)	72.5	90.8
$Ca^{2+}$ ( $\text{cmol.kg}^{-1}$ )	29.1	47.2
$Mg^{2+}$ ( $\text{cmol.kg}^{-1}$ )	2.7	21.8
$K^+$ ( $\text{cmol.kg}^{-1}$ )	2.9	2.3
$Na^+$ ( $\text{cmol.kg}^{-1}$ )	0.1	1.4

### 3.5. Relationships between chemical elements in the soils of the study site

The proportion of ions bound to the clay-humus complex (CAH) provides information on the possibility of antagonism. Poor calcium nutrition can occur if the ratio of  $Mg^{2+}/Ca^{2+}$  and  $K^+/Ca^{2+}$  is greater than 1 each. In  $Mg^{2+}$ -poor soils, this element may be deficient if the  $K^+/Mg^{2+}$  ratio is greater than 1. In the case of the present study, the ratios obtained are all less than 1, with the exception of

**K<sup>+</sup>/Mg<sup>2+</sup>.** The concentration ratios between the various chemical elements in the adsorbent complex are shown in Table 5 below.

**Mg<sup>2+</sup>/Ca<sup>2+</sup> ratio :** The ratio of Mg<sup>2+</sup> to calcium ions in the surface horizons (0 - 20 cm) and at depth (20 - 40 cm) hovers around 0.09 and 0.46 respectively, all of which are less than 1.

**K<sup>+</sup>/ Ca<sup>2+</sup> ratio :** Concentrations of potassium ions relative to calcium (K<sup>+</sup>/ Ca<sup>2+</sup>) give a value relatively similar to that of the Mg<sup>2+</sup>/Ca<sup>2+</sup> ratio in humus-bearing surface horizons. At depth, however, the Mg<sup>2+</sup>/Ca<sup>2+</sup> ratio is almost 10 times greater than that of K<sup>+</sup>/ Ca<sup>2+</sup>. This shows that K<sup>+</sup> ions are likely to disappear in the presence of Mg<sup>2+</sup> ions (or else express a deficiency of potassium ions in the deeper layers).

**K<sup>+</sup>/Mg<sup>2+</sup> ratio :** At the surface, the K<sup>+</sup>/Mg<sup>2+</sup> ratio is 1.07 times greater than 1. This figure also decreases almost 10-fold in the underlying horizons. On the surface, K<sup>+</sup> ions are more prevalent than Mg<sup>2+</sup> ions. This observation highlights the risk of massive loss of Mg<sup>2+</sup> ions to the K<sup>+</sup> ion profile. At all depth levels, calcium ions are strongly represented, testifying to the good chemical fertility of the soils at the site studied.

Table 5. Average chemical element content ratios of study site soils by depth.

chemical element ratios	Depth (cm)	
	0 – 20	20 – 40
K <sup>+</sup> / Ca <sup>2+</sup>	0.01	0.05
Mg <sup>2+</sup> /Ca <sup>2+</sup>	0.09	0.46
K <sup>+</sup> /Mg <sup>2+</sup>	1.07	0.11

## 4. DISCUSSION

### 4.1. Organic matter characteristics of the soils studied

The C/N ratio of soils is an indicator of the ability of organic matter to decompose more or less rapidly in soils. Those obtained from the soils studied, at around 11 and 12, confirm good mineralization of organic matter and show soils in good conditions to ensure cocoa tree development [19]. Such a situation leads to the over-rapid release of nitrogen into the environment [20], compromising its availability, in the soil solution due to the intense leaching process undergone by tropical soils, particularly those in south-west Côte d'Ivoire, which are under conditions of heavy rainfall. Furthermore, the nitrogen concentrations obtained ( $0.1 - 0.05\% \leq 1\%$ ) indicate soils depleted in this element ([16] ;[21]). Like nitrogen, organic matter is poorly represented ( $< 1\%$ ). This low level of organic matter could be linked to the ageing of the plants, which return little litter to the soil [22], in addition to the rapid mineralization of organic matter. These inadequate soil organic matter and nitrogen levels indicate a decline in soil fertility, with organic matter and nitrogen as factors limiting crop yields. This situation can compromise cocoa production, since to produce one tonne of cocoa, at least 45 kgN/ha/year and soils with a minimum organic matter content of 3% are required ([23] ; [24]). The pH values recorded on the surface and at depth show the acidic ( $5 < \text{pH} = 5.6 < 6.8$ ) and weakly acidic ( $5.6 < \text{pH}_{\text{water}} = 6.8 < 7$ ) character of these soils respectively. These pH conditions do not represent a constraint for the development of cocoa trees, which can reproduce on acidic (pH 4.5- 6) or slightly basic (pH 6.7-7.5) soils [25]. What's more, these pH conditions make phosphorus available. Moreover, phosphorus bound to clays by calcium, and that complexed by organic matter, exists in appreciable quantities only in slightly acidic soils with pH values between 5.5 and 7 [26]. Many authors consider this range of pH values favorable to the development of crop plants and cereal crops,

particularly rainfed rice [27]. Assimilable phosphorus levels in the soils studied are above 100 ppm, indicating good nutrition of cocoa trees in this element ([22];[28]).

#### **4.2. Characteristics of the adsorbent complex in the soils studied**

CEC is a good indicator of potential soil fertility. The values obtained in the present study are considered to be good, and well above the threshold described by Sawadogo [29], placing these soils in the category of fertile soils capable of ensuring better cocoa production. The good cation exchange capacity recorded partly explains the high saturation of exchangeable bases noted in these soils, reflecting their high nutrient content. Contrary to Van [30], in ferralsols in humid tropical zones, due to heavy rainfall, leaching processes are very pronounced. By eliminating exchangeable cations from surface horizons, these pedological processes deplete the specific surface area of clays in exchangeable bases in soils under cocoa trees in south-western Côte d'Ivoire [23].

#### **4.3. Physical characteristics of the soils studied**

The upper layers of the region's soils have a high sand content (45%), well above the threshold recommended by Quittet [16]. At this proportion, the defective character of the sands is present but, at the same time, attenuated by the presence of silt and clay. These horizons showed good drainage, air circulation and easy root penetration during in situ morphological characterization. On the other hand, the underlying layers (20 - 40 cm) are enriched in clay (48%) due to partial desaturation of the colloidal complex of the surface layers in lime ( $\text{Ca}^{2+}$ ), which, dispersing in the soil solution, migrates downwards, making this layer an accumulation layer [16]. The large amount of clay observed in the underlying layers (20 - 40 cm) is also probably due to the weathering of the parent material. The connection between the origin of the clays in the soil and the material had already been pointed out in the work of Millot [31]. However, according to Boyer [32], soils with more than 40% clay are unfavorable for cocoa cultivation, as they promote waterlogging that can asphyxiate roots and lead to rootlet necrosis. Defective silt-related properties appear in soils with a silt content of 30% or more. The average silt content of the soils studied tends towards the critical threshold, regardless of soil depth. The average clay content evolves in the opposite direction to that of the sand. On the other hand, in the 20-40 cm horizon, the clay content increases, with higher contents around 48%. Above 20% clay content, clay-related defects increase and worsen. The risk of compaction and asphyxiation is high in these soils under cocoa trees, from the first 20 cm down, due to the high clay and silt content. The use of compost is therefore recommended.

### **5. CONCLUSION**

The present work was carried out in the south-west region of Côte d'Ivoire (Soubré, 6° - 7° north latitude and 5° - 7° west longitude). The aim was to develop an appropriate and sustainable fertilization strategy for cocoa. The aim is to characterize some essential chemical and physical properties of soils under cocoa with a view to improving and increasing cocoa production. On the surface, soils are depleted in organic matter and nitrogen, which can limit yields. On the other hand, soil pH conditions favor the availability of assimilable phosphorus and cocoa production. The characteristics of the adsorbent complex are good. A high clay content is recorded from the first 20 cm depth. As things stand, there is no need to apply mineral fertilizer to the soil. However, the use of organic fertilizer (compost) is recommended to improve organic and nitrogen stocks, essential for better yields.

### **REFERENCES**

1. Hervé K. S., Coulibaly N. G. A. and Alloueboraud W. A. (2018). Characterisation of cocoa orchards and diseases in Côte d'Ivoire: the case of Abengourou, Divo and Soubré departments. *Journal of Animal & Plant Sciences*, vol. 35, no. 3, pp. 5706-5714.
2. Marine J. (2023). Newspaper, *Le Monde Afrique*. African economy. In Ivory Coast, rising cocoa prices are not benefiting farmers; Published on 12 September 2023 at 18:00 ; [https://www.lemonde.fr/afrique/article/2023/09/12/en-cote-d-ivoire-la-hausse-des-cours-du-cacao-ne-profite-pas-aux-planteurs\\_6189063\\_3212.html#:~:text=La%20C3%B4te%20d'Ivoire%2C%20qui,resserrer%20son%20offre%20cette%20ann%C3%A9e](https://www.lemonde.fr/afrique/article/2023/09/12/en-cote-d-ivoire-la-hausse-des-cours-du-cacao-ne-profite-pas-aux-planteurs_6189063_3212.html#:~:text=La%20C3%B4te%20d'Ivoire%2C%20qui,resserrer%20son%20offre%20cette%20ann%C3%A9e).
3. International Cocoa Organization (2019). Quarterly bulletin of cocoa statistics, vol. XLV, n°1, cocoa year, 2018-2019.
4. Dehevels O. (2003). Cocoa planting / replanting dynamics in Côte d'Ivoire: comparison of technical choices with Olympe. Olympe" seminar Montpellier, France, September 2003. 13 p.
5. Tano, Assi M. (2012). Cocoa Crisis and Producer Strategies in the Sub-Prefecture of Méadji in South-West Côte d'Ivoire. PhD thesis, Université Toulouse le Mirail, Toulouse, France, 261 p.
6. Konan A., Yaméogo I., Assiri A. and Ehoughban V. (2015). Sustainable cocoa farming technical manual: for the attention of the technician. February edition. 165 p.
7. Léonard and Vimard, (2005). Crisis and reshaping of pioneer agriculture in Côte d'Ivoire, Editions IRD 213, Fayette street 75010 Paris, Karthala 22-24, boulevard Arago 75013 Paris, 361 p.
8. Honfoga, Gbenoukpo B. (2007). Towards efficient private fertiliser supply and distribution systems for sustainable agricultural intensification in Benin. 2007. PhD thesis. Centre for Development Studies, University of Groningen. 412 p.
9. Adjagodo A., Tchibozo M. A. D., Kelome N. C. and Lawani, R. (2016). Flows of pollutants linked to human activities, risks to surface water resources and the trophic chain worldwide: bibliographical summary. *International journal of biological and chemical sciences*, Vol 10, N°3, pp. 1459-1472.
10. Jama B., Palm C. A., Buresh R. J., Niang A., Gachengo C., Nziguheba G., et Amadalo B. (2000). *Tithoniadiversifolia* as a green manure for soilfertility improvement in western Kenya: areview. *Agroforestrysystems*, 2000, vol. 49, no 2, pp. 201-221.
11. Kitabala M. A., Tshala U. J., Kalenda M. A., Tshijika I. M., and Mufind K. M. (2016). Effects of different doses of compost on tomato (*Lycopersiconesculentum* Mill) production and profitability in the town of Kolwezi, Lualaba Province (DR Congo). *Journal of Applied Biosciences*, vol. 102, pp. 9669–9679.
12. Eldin M. (1971). Côte d'Ivoire's natural environment: Climate. ORSTOM dissertation n° 50 ; Paris, France, pp. 106 – 107.
13. Sorokoby V. M., Saley M. B., Kouame K. F., Djagoua M. E., Bernier M., Affian K. and Biemi J. (2010). Use of Landsat ETM+ and SIRS images for lineament and the matic mapping of Soubré-Meagui (south-west Côte d'Ivoire): contribution to the management of groundwater resources. *Remote sensing*, Vol. 9 (3-4), pp. 209-223.
14. Brou T. Y. (2005). Climate, socio-economic changes and landscapes in Côte d'Ivoire. HDR, University of Lille, 212 p. XI, pp 221- 237.
15. Perraud A. (1971). The natural environment of the Ivory Coast: Soils. ORSTOM dissertation n° 50 ; Paris, France, pp. 271 – 27.
16. Quittet E., 1967. *Agriculture*, Tome 1, 15ième Edition. Dunod, Paris, pp. 5 – 6.
17. Horneck D.A., Sullivan D.M., Owen J.S., and Hart J.M.. 2011. *Soil Test Interpretation Guide*. 12p. Archival copy. For current version, see: <https://catalog.extension.oregonstate.edu/ec1478>.

18. Quemada M., and Cabrera M. L., 1995. CERES-N model predictions of nitrogen mineralized from cover crop residues. *Soil Science Society of American Journal* 59: 1059–1065.
19. Snoeck D., Koko L., Joffre J., Bastide P. et Jagoret P. (2016). Cacao Nutrition and Fertilization : Sustainable Agriculture Reviews, Vol. 19, Chapter 4. E. Lichtfouse (Ed.). Cham, Springer International Publishing: pp.155-202.
20. Jadin, P. and Lotode R. (1981). The use of 'soil diagnosis' to estimate the fertiliser requirements of cocoa farms in Côte d'Ivoire.  
Website: [https://horizon.documentation.ird.fr/exl-doc/pleins\\_textes/pleins\\_textes\\_5/b\\_fdi\\_02-03/00858.pdf](https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_5/b_fdi_02-03/00858.pdf)
21. Bassolé Z., Yanogo P.I. and Idani F.T. (2023). Characterisation of leached tropical ferruginous soils and eutrophic tropical brown soils for agricultural use in the Goundi-Djoro lowland (Burkina Faso). *Int. J. Biol. Chem. Sci.* 17(1): 247-266.
22. Koko K.L. ; R. G. Yoro , K. Ngoran and A. ASSA. (2008). Evaluation of soil fertility under cocoa trees in south-west Côte d'Ivoire. *Agronomie africaine*, 20 (1) : 81 – 95.
23. Smith A. J. (1980). Soils classification and the cocoa grower. *Cocoa Growers' Bull.*, 30 (5), pp. 5 - 10.
24. **Jadin Pierre** and **Snoeck Jacques**. 1985. La méthode du diagnostic sol pour calculer les besoins en engrais des cacaoyers. *Café, Cacao, Thé*, **29** (4) : 255-272. Website : <https://agritrop.cirad.fr/462882/>
25. Appiah M. R., K. Ofori-Frimpong, A. A. Afrifa, M. K. Abekoe and D. Snoeck. (2006). Improvement of soil fertility management in cocoa plantations in Ghana. FSP Regional Cacao scientific and technical final report. CRIG (Cocoa Research Institute of Ghana), Ghana, 22 p.
26. Bélanger N. (2023). Agroforestry and sustainable development. Module 3.4. Availability of phosphorus and aluminium in tropical soils. <https://env3114.telug.ca/module-3/3-4-la-biodisponibilite/#:~:text=Dans%20ce%20sens%2C%20la%20disponibilit%C3%A9,est%20retenu%20par%20l'aluminium>. Accessed on 13 December 2023.
27. Landon J. R. (1991). Booker Tropical Soil Manual, A handbook for soilsurvey and agricultural land evaluation in the tropics and subtropics, paperback, longman, Booker Tate limited, Oxon, Royaume Uni, 474 p.
28. Jadin P. (1992). Cocoa agronomy at IRCC. Montpellier, France, IRCC/CIRAD, IRCC studies and work, 44 p.
29. Sawadogo H. (2006). Organic and phosphate fertilisation in the Zaï cropping system in the Sudano-Sahelian environment of Burkina Faso. PhD thesis, Gembloux University Faculty of Agronomic Sciences, Belgium, 219 p.
30. Van Wambeke A. (1974). Management properties of ferralsols. FAO Soils Bulletin, no 23, 129 p.
31. Millot G. (1964). Clay geology: weathering, sedimentology, geochemistry. Masson, Paris, 449 p.
32. Boyer J. (1982). Les sols Ferralitiques. "Facteurs de fertilité et utilisation des sols," Tome X. Initiations-documentations techniques n° 52. ORSTOM Paris. 384 p.