

## Original Research Article

### Adoption of Cold reeling, as a tool of selection in breeding of “Nistari” a multivoltine breed of mulberry silkworm (*Bombyx. mori*) in West Bengal, India.

#### Abstract:

Mulberry silkworm (*B.mori*) is an economic insect reared for silk. Economic traits of silkworm are highly influenced and deteriorated by fluctuating environmental conditions arising due to global warming. Artificial selection has always played a crucial role in silkworm domestication to improve economic traits. Filament length is important from the perspective of mulberry sericulture. Nistari lines (Marked and Plain) which is a ruling multivoltine breed in West Bengal, India, has short filament length and economic traits are observed to be deteriorated. Selection of parents with high filament length is hard as the pupa corresponding with higher filament length is sacrificed during the process of reeling in hot temperature in conventional reeling. “Cold reeling is an alternative reeling, which is devoid of hot temperature, thereby enabling the rescue of pupae for further metamorphosis. Cold reeling can be adopted to select parents with higher filament length in mulberry silkworm for successive generations. The present study made an attempt to increase the filament length of Marked and Plain lines of Nistari, by selection of parents with high filament length for consecutive generation by implementing cold reeling.

#### Keywords

Filament length, *Bombyx mori*, Selection, Cold reeling, Nistari lines.

#### 1.Introduction:

The domestic silkworm is an economically important insect [1]. Mulberry silkworms are reared for silk. More than 95% of natural silk, all over the world is obtained from this insect [2]. Fluctuating climatic conditions affects the success of cocoon production [3] and decline in economic traits. Economic traits are also affected by inbreeding depression and rearing conditions. Artificial selection plays an important role in moulding economic traits of choice for desired characters. During the process of domestication, silkworm have evolved rapidly under human-preferred selection [4]. In silkworm domestication, different modes of artificial selections come to play. Conventional selection plays a pivotal role where economic traits are selected visually on the basis of shape, size, colour, weight, larval marking etc. Marker assisted selection also plays an important role in silkworm breeding. Some of the widely employed markers are SSR markers and ISSR markers [5]; [6]; [7]; [8]. Filament length is an important economic character. It is the fibrous length reeled out from a single cocoon [9]. Filament length varies in different breeds of silkworm. Environmental factors affect filament length [10]. Application of selection pressure, in economic traits like cocoon weight of parental lines was shown to improve the performance of cocoon weight of hybrids [11]. The scenario of mulberry sericulture in West Bengal, India is dominated by Cross breeds of

Nistari lines Marked and Plain. Nistari is broadly classified into Nistari Marked and Nistari Plain lines based on presence of larval marking. Nistari Marked lines possess black larval marking which is devoid in the Plain lines.

Nistari lines are characterized for its high survival in drastic climatic conditions. One of the downsides of this preferred breed among many is low filament length. In this study an attempt was made to improve the filament length of Nistari lines, by subjecting selection pressure in different generations. Cold reeling was adopted where single cocoons are reeled manually without subjection of high temperature to rescue the pupa for further metamorphosis unlike the conventional reeling. Gradual improvement in filament length was observed. Without selection pressure, gradual increase in filament length was absent, but the filament length was higher from the initial generation, when selection was initiated.

## **2. Materials and Methodology:**

Cocoons (5-6 day aged after spinning) of Nistari Marked and Plain lines with its racial characteristics were selected based on visual observation. The cocoons were labelled individually and placed in small chambers separately. The cocoons were soaked for 10-15 minutes in 0.25% NaOH solution. The cocoons were reeled individually in eprouvette in 0.25% NaOH solution. Filament length of each cocoon was calculated. The individual live pupa after reeling was dipped in 0.1% glacial acetic acid for 1 min and later washed thoroughly in tap water and dried with paper towel. The sex of the pupae was recorded and kept for further metamorphosis. 70-75 cocoons of both Marked and Plain lines were reeled as described above. 20-25 each of male and female pupae with highest filament length were selected as parent for next generation. The average filament length of the corresponding males and females' pupae were taken as filament length of the particular generation. Selection for six generations was conducted in a similar protocol as described.

Selection was retracted after six generations. Cold reeling was conducted as described above from generation 7 to generation 13 without selection. Average filament length of unselected cocoons was recorded from equal numbers of male and female cocoons in each generation.

## **3. Results:**

The Cocoons of Nistari lines Plain and Marked were cold reeled. The average filament length of equal males and females were taken (Table 1 & 2). In Nistari Marked lines it could be observed that there is a gradual improvement of Filament length from generation 1 (G1) to generation 6 (G6). There was an overall increase in filament length from 378m to 454m (Table 1). In Nistari Plain line the average filament length increases from 348m to 361m though a decline could be seen in generation 4 (Table 2) during the period.

<b>Generation</b>	<b>Average Filament Length (m)</b>
G1	378
G2	384
G3	409
G4	412
G5	430

G6	454
----	-----

Table 1: Average of selected longest filament length of males and females of Nistari Marked lines from generation 1 to generation 6, by conducting cold reeling which shows a gradual increase in filament length over the generations (G represents generation).

Generation	Average Filament Length (m)
G1	348
G2	379
G3	395
G4	379
G5	452
G6	461

Table 2: Average of selected longest filament length of males and females of Nistari Plain lines from generation 1 to generation 6, by conducting cold reeling which shows a gradual increase in filament length over the generations (G represents generation).

Generation	Average Filament Length (m)
G7	452
G8	427
G9	436
G10	426
G11	467
G12	448
G13	431

Table 3: Average filament length of males and females of Nistari Marked line from generation 7 to generation 13, by conducting cold reeling without subjecting to selection (G represents generation).

Generation	Average Filament Length (m)
G7	420
G8	417
G9	447
G10	422
G11	467
G12	442
G13	437

Table 4: Average filament length of males and females of Nistari Plain line from generation 7 to generation 13, by conducting cold reeling without subjecting to selection (G represents generation).

Filament length was also estimated without selection, by cold reeling in both the lines of Nistari (Fig 3 & Fig 4). It can be observed that in absence of selection there is no gradual increase in the length of the filament length, but changes over generations. But it is also prominent from the study that even in absence of selection, an improvement of filament length can be observed from the initial generation of G1 in both the lines of Nistari.

#### 4. Discussion:

Artificial selection has always played a key role in shaping traits of choice. In domesticated animal and plant breeding, a breeder can change genetic characteristics of a population in two ways. The first way is by selection of individuals as parents and the second way is controlling of mating of parents which includes inbreeding and outbreeding [12], which illustrates the importance of selection in breeding. Studies have shown the adoption of directional selection to produce insect strains with higher tolerance to extreme environmental conditions, such as low humidity, for extended periods [13], has adopted Directional selection to improve desiccation resistant *Anastrephaludens* strains. Economic insects are domesticated for economic traits and selection plays a crucial role in improving the economic traits. Evidence on increase in cocoon weight by selection was observed in silkworm [14]. Gradual improvement of economic traits is pivotal for silkworm rearing to increase profit of sericulture industry [15].

Filament length is an important economic trait in silkworm. High filament length is an important characteristics of an elite silkworm breed. Among the mulberry silkworm breeds bivoltine breeds have higher filament length than multivoltine breeds [16]. Filament length like other traits can be improved by selection. In a silkworm breed, parent with higher filament length from a population can be selected over generations to yield progeny with higher filament length.

Conventional selection for high filament length is hard, unlike other silkworm economic traits. Estimation of filament length requires reeling of the cocoons. The process of reeling in mulberry silkworm (*B. mori*) involves cooking up of the cocoons in hot water [17]. During the process of reeling the metamorphosing pupae is killed. This makes it hard to select better parents with longer filament length. "Cold reeling" is an alternative reeling where the subjection of heat during the process of reeling is avoided. Here the cocoons are reeled individually, in alkaline solution, thereby rescuing the pupae and allowing it for further metamorphosis to silkworm moths and oviposition. Cold reeling is labour intensive and time-consuming comparative to the conventional reeling. The process of Cold reeling facilitates conventional artificial selection for filament length. It enables the selection of the pupa corresponding to high filament length as parent for next generation.

It was observed from the present study that filament length could be improved by selection. A gradual increase in filament length could be observed in both Nistari Marked and Plain lines with continuous subjection of selection pressure from generation 1 to generation 6 (Fig 1 & 2). Without the subjection of selection pressure gradual improvement was not observed (G7 to G13), but the filament length in both the Nistari lines, Marked and Plain were higher than the filament length of initial generation (G1) (Fig 3 & 4). Hence, the experiment on selection implies that cold reeling could be employed as a selection tool to improve filament length in mulberry silkworm "Nistari".

Nistari, plays a leading role in mulberry sericulture of West Bengal, India. The Farmers' field are dominated by Nistari and its cross breeds. Multivoltine x Bivoltine hybrids (Nistari x SK6.7) and Multivoltine x Multivoltine (Nistari x M12W), crosses are widely preferred

depending upon the seasons. Improving the filament length of Nistari can be one of the factors to improve production of raw silk in West Bengal.

The present study was conducted on “Nistari” a multivoltine breed, but similar attempts can be conducted to other parental silkworm breeds. The commercial sericulture sector depends on hybrids, viz., double hybrids, bivoltine hybrids, multivoltine x bivoltine hybrids, multivoltine x multivoltine hybrids depending upon the climatic conditions. Enhancing the filament length of the parental breeds is expected to improve the filament length of the hybrids too.

Economic insects like mulberry silkworm due to continuous inbreeding and domestication are usually associated with deterioration of economic traits and fluctuating environmental conditions. Economic traits suffer from inbreeding depression [18]. Filament length also suffers the same fate of inbreeding depression and decline due to harsh and unpredictable environmental factors like temperature and humidity. Directional selection can be used to restore those characters lost during the mass rearing process in insects [13]. The process of cold reeling can also be employed to rejuvenate filament length suffering the above factors. In the present study an attempt has been made to improve filament length by selection, adopting cold reeling in the multivoltine breed “Nistari”. The observations from the study, presents an increase of filament length which implies the application of cold reeling as a “selection tool” in Nistari and the possibility to adopt it in other breeds of mulberry silkworm rearing.

## **5. Conclusion:**

Selection plays a significant role in silkworm breeding. In the present study filament length of “Nistari” lines was improved by adopting cold reeling. Nistari being the backbone of mulberry sericulture in West Bengal, improving the filament length is expected to increase the raw silk production. Similar approach can also be adopted for other silkworm breeds.

## **References:**

[1] Fang SM, Zhou QZ, Yu QY, Zhang Z. Genetic and genomic analysis for cocoon yield traits in silkworm. *Sci Rep.* 2020;10(1):5682. <https://doi.org/10.1038/s41598-020-62507-9>

[2] Collard BC, Jahufer Y, Brouwer JB, Pang ECK. An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: The basic concepts. *Euphytica* 2005;142:169–196. <https://doi.org/10.1007/s10681-005-1681-5>

[3] Ram RL, Maji C, Bindroo BB. Impact of climate change on sustainable sericultural development in India. *Int. J. Agric. Innov. Res.* 2016;4(6):1110-1118.

[4] Zhu YN, Wang LZ, Li CC, Cui Y, Wang M, Lin YJ, Zhao RP, Wang W, Xiang H. Artificial selection on storage protein 1 possibly contributes to increase of hatchability

during silkworm domestication. PLoS Genet. 2019;15(1):1-19.  
<https://doi.org/10.1371/journal.pgen.1007616>

[5] Sreekumar S, Ashwath SK, Salathia M, Kumar SN, Qadri SMH. Detection of Single Nucleotide Polymorphism (SNP) DNA markers linked to cocoon traits in the mulberry silkworm, *Bombyx mori* (Lepidoptera: Bombycidae). European Journal of Entomology. 2011; 108:347-354. DOI: 10.14411/eje.2011.043

[6] Reddy KD, Abraham EG, Nagaraju J. Microsatellites in the silkworm, *Bombyx mori*: Abundance, Polymorphism strain characterization. Genome. 1999a; 42:1057-1065. <https://doi.org/10.1139/g99-027>

[7] Reddy KD, Nagaraju J, Abraham EG. Genetic characterization of the silkworm *Bombyx mori* by Simple Sequence Repeat (SSR)-Anchored PCR. Heredity. 1999b; 83:681-687. <https://doi.org/10.1046/j.1365-2540.1999.00607.x>

[8] Nagaraju J, Reddy KD, Nagaraju GM, Sethuraman BN. Comparison of multilocus RFLPs and PCR-based marker systems for genetic analysis of the Silkworm, *Bombyx mori*. Heridity. 2001; 86: 588-97. <https://doi.org/10.1046/j.1365-2540.2001.00861.x>

[9] Kweon HY, Lee KG, Park KY, Kang SW, Kang PD, Kim MJ, Kim KY. Cocoon Filament Quality of a Special Silkworm Variety, Golden Silk. Int. J. Indust. Entomol. 2012; 24 (1): 17-21. <http://dx.doi.org/10.7852/ijie.2012.24.1.017>

[10] Rahmathulla VK. Management of Climatic Factors for Successful Silkworm (*Bombyx mori* L.) Crop and Higher Silk Production: A Review: 2012: Psyche: doi:10.1155/2012/121234

[11] Khaligh M, Seidavi A, Gharahveysi S, Das T. Measurement of response to selection in silkworm commercial hybrids against individual selection of parental lines. Int J Trop Insect Sci. 2024; 44: 31–39. Measurement of response to selection in silkworm commercial hybrids against individual selection of parental lines.

[12] Falconer DS. 1990. Introduction to quantitative genetics. 3<sup>rd</sup> ed. New York: Wiley; p. 438.

[13] Tejada MT, Gordillo JA, Fong LQ, Dávila DO, Fleischer DF. Directional selection to improve the sterile insect technique: Survival and sexual performance of desiccation resistant *Anastrepha ludens* strains. Evolutionary Applications. 2017;10:1020–1030. <https://doi.org/10.1111/eva.12506>

[14] Mirhosseini SZ, Nematollahian S, Ghanipoor M, Seidavi A. Comparison of phenotypic and genetic performance of local silkworm groups and two commercial lines. Biol. Res. 2010;43:411–416. <http://dx.doi.org/10.4067/S0716-97602010000400005>

[15] Hemmatabadi ZN, Seidavi A, Gharahveysi S. A review on correlation, heritability and selection in silkworm breeding. *Journal of Applied Animal Research*.2016; 44:1, 9-23. <https://doi.org/10.1080/09712119.2014.987289>

[16] Sharma K, Bali K. Analysis of heterosis in some bivoltine silkworm hybrids of *Bombyx mori* L. *Journal of Entomology and Zoology Studies*.2019; 7(5): 01-08

[17] Chattopadhyay D, Munshi R, Chakravorty D. Studies on distribution of filament length and non-broken filament length for tropical tasar and muga silk cocoons vis-à-vis mulberry silk cocoons. *The Journal of The Textile Institute*.2018; 109 (9):1202-1207. <https://doi.org/10.1080/00405000.2017.1422307>

[18] Doreswamy J, Gopal S. Inbreeding effects on quantitative traits in random mating and selected populations of the mulberry silkworm, *Bombyx mori*. *Journal of Insect Science*.2012; 12(140): 1-14. <https://doi.org/10.1673/031.012.14001>

UNDER PEER REVIEW