

# Original Research Article

## Simple Geographic Distribution of Patient with Stroke According to their Medical Care Financing System in a Private University Hospital

---

### ABSTRACT

**Aims:** To reveal the geographical distribution of stroke patients in a private university hospital in Jakarta, Indonesia according to their address and further divide them into three sub group of using universal health coverage (UHC) vs other type of insurance ((OTI) vs private non insurance patient (private)

**Study design:** simple retrospective cross-sectional study

**Place and Duration of Study:** Sample: Medical record from patients of the Department of Neurology, *Rumah Sakit* Universitas Kristen Indonesia, Jakarta Indonesia; a private university hospital in Jakarta Indonesia, between January 2021 and December 2023.

**Methodology:** We analyzed medical record of 162 patients (90 men, 72 women; age range 15-87 years (mean age 61.12 SE± .82) with history of stroke and at the present routinely underwent routine medical control (outpatient) or current diagnosis of stroke (inpatient in hospital). We further divided them based on their medical care financing system (UHC vs OTI vs private) and using the GPS service made available by Google Maps™ we simply try to make a geographical distribution of those patients based on their address/residence location

**Results:** Geographical distribution of patients using UHC were closer to the hospital compared to the other 2 non UHC medical care financing users.

**Conclusion:** Our analysis regarding stroke patients in our hospital (2021-2023), we found that 55.6% male patients, 42% were in the range of age between 55-64 years old, most of the patient (94.4%) using BPJS as their medical care financing system, with 27.7% came from sub district Makkasar which is near from our hospital. Further analysis using the help of Google maps™ we can show that the UHC or BPJS health referral system has facilitated patients receiving health services at the health facility closest to their residence.

*Keywords: referral, social security agency for health, chronic, rehabilitation*

### 1. INTRODUCTION

*Badan Penyelenggara Jaminan Sosial Kesehatan* (BPJS Kesehatan, or 'Social Security Agency on Health') is a social security agency of Indonesia, Social Security Fund as a Form of Constitutional Responsibility for the Right to Health [1], and was aimed at providing universal health care to all of its citizens [2] which launched in January 2014. The Indonesian government launched a universal health care system called the *Jaminan Kesehatan Nasional* (JKN, "National Health Insurance"). Covering around 95.77% or 268 million of people by February, 2024 [3], it is the world's second most extensive insurance system or to

be more precisely is called publicly funded health insurance (PFHI) program after the programme in India with population coverage of 500 million. [4].

As part of the implementation of JKN, on the first day of 2014, PT Askes (Persero) was shifted from a state-owned company into BPJS Kesehatan, a state appointed public agency, which becomes the sole provider of JKN [5]. It is hoped that the entire citizen of Indonesia will be covered in the year 2019 [6]. In 2016, the BPJS program suffered from a horrible deficit of more than six trillion IDR. However, the deficit expanded up to 32 trillion in only three years [7]. In response, the government issued a policy that increases the monthly premium for access using system dynamics method [8]. Expert opinion regarding the government policy saw that the move as adding more burden on low to middle-income citizens [9].

BPJS Kesehatan, or Badan Penyelenggara Jaminan Sosial Kesehatan, is a state/government-managed health insurance program established in Indonesia that offers protection for its members, as long as the pay [8,9], and it provides equal access to health facilities for everyone [1,2], with no exception to those who live in rural, remote and even poor areas [10]. BPJS covers hundreds of diseases [11], including chronic conditions like stroke [12] and Indonesia is the leading country in the number of stroke patients throughout Asia with a stroke rate of 3,382.2/100,000 people and stroke-related mortality in the south east Asian Region was the highest in Indonesia [13]. Since the launching of BPJS, the early detection of stroke has become more effective and efficient [14].

In 2022, it estimated that healthcare costs for stroke patients in Indonesia, covered by BPJS Kesehatan, ranging from the smallest Rp 1,937,400 to the highest Rp 207,675,700, averaging Rp 8,487,531 per patient [15]. BPJS Kesehatan's actual total spending on stroke treatment and in combination with patient's rehabilitation in 2022 estimated as much as Rp 2,883,269,825,972. The biggest group benefiting from this were those receiving Beneficiary Contribution Assistance (PBI) from the government, which in 2019 estimated reached >96 million or roughly 46.5% of all National Health Insurance participants.

The aim of this simple pre-eliminary study is to reveal the geographical distribution of stroke patients in a private university hospital in Jakarta, Indonesia according to their address and further divide them based on their medical care financing system

## **2. METHODOLOGY**

During June to July 2024 at the dept. of Community Medicine, we cross sectionally analyzed medical record of 162 patients (90 men, 72 women) with history of stroke and at the present routinely underwent routine medical control (outpatient) or having current diagnosis of stroke (inpatient in hospital) in a private university hospital located in Jakarta, Indonesia; all 162 patients were patients from January 2021 to December 2023. We further divided them based on their medical care financing system (BPJS vs OTI vs private) and using the GPS service made available by Google Maps™ we simply try to make a geographical distribution of those patients based on their address/residence location.

Because we only dealing with data of medical record, consent obtained from the director of the hospital (no. 378/DIRUT/RSU.UKI/06.2024) with cover letter from the dean to the faculty of Medicine (no, 0623/UKI.F5.D/PPM.00.01/2024) prior to data collection and after all the necessary data collected, then it was further processed and presented in tables and figures in the result section.

### 3. RESULTS AND DISCUSSION

This simple retrospective cross sectional study of 162 patients of stroke revealed an interesting phenomenon. First lets started with demographic condition of these patient.

Table 1. Demographic characteristics

Characteristics		N (162)	%
Year	2021	71	43.8
	2022	55	34.0
	2023	36	22.2
Gender	Men	90	55.6
	Women	72	44.4
Age	15 – 24	1	0.6
	25 – 34	0	0
	35 – 44	4	2.5
	45 – 54	29	17.9
	55 – 64	68	42
	65 – 74	49	30.2
	>75	11	6.8
Medical care financing system	BPJS	153	94.4
	Private	8	4.9
	OTI	1	0.7
The patient comes from the sub-district ( <i>kecamatan</i> )			
	Cakung	2	1.2
	Cipayung	7	4.3
	Ciracas	4	2.5
	Duren sawit	10	6.2
	Jatinegara	17	10.5
	Kramat Jati (where the hospital exists)	36	22.2
	Makkasar	44	27.2
	Matraman	0	0
	Pasar Rebo	1	0.6
	Pulogadung	1	0.6
	Area outside east Jakarta	40	24.7

Within three years (2021-2023) there were gradual reduction on the number of patients with stroke, and this perhaps due to the tiered hospital/health facility referral system. within those three years, there are more and more primary health care facilities, whose service levels are actually below hospitals, but are actually become more capable to handle stroke patients themselves, so they no longer need referrals to higher hospitals.

Based on the sizes and medical services coverage, the health facilities in Indonesia are divided into two tiers: primary care providers and secondary care providers [17,18]. Primary care providers consist of private physicians, community clinics, primary health care facilities/centers (Puskesmas), and class D-hospitals [19,20], whilst secondary care providers categorized as specialized or even sub-specialized service (which cannot have conducted in primary care providers) and consists of classes A to C-hospitals with class A as

the highest tier [21]. The higher the hospital “tier”, the more coverage of health services are provided.

Real efforts from all stakeholder are needed to organize better health promotion programs down to the community level with a more rational and a better distribution of neurologists and neurointerventionalists which is needed in order to tackle this preventable disease [22]. All community based hospitals should have a ready to use CT scan machine and stroke units to minimize in-hospital mortality rate [23]. Intensifying telemedicine for stroke patients [24] is a very promising effort as an addition to the improvement of integrated acute stroke management system [25].

Regarding the gender of the patient with stroke in our hospital, there are more male patient than female patient (55.6 % Vs 44.4%). Our findings also in line with the findings of Zamzam et al [26] where those researchers found out the incidence of stroke higher in male gender (OR=1.48; 95% CI= 1.07 to 2.03  $p=0.017$ ).

Many study justified that age-adjusted burden of stroke is greater in men compared to women (4.4 vs 3.9 per 1,000 person-years) [27]. Nonetheless, the lifetime risk of stroke is higher in women than in men [28] and furthermore, women are at increased risk of stroke in the very old age group [29]. Since overall, women always have a higher life expectancy than men, the overall stroke burden is therefore higher in women [30]. This give us insight to analyze further in our future study regarding the impact of stroke on individual burden post hospitalization, psychologically, economically and also clinically.

Age is an essential non-modifiable risk factor for stroke [31,32]. The age distribution in our patient are as follow. The youngest patient is 15 years old and the oldest 87 years old with mean age 61.12 years (SE±.825, SD± 10.59). previous report from Kusuma in 2009 stated that the mean age of stroke patients found in Indonesia is 58.8 years [33]. According to the INTERSTROKE study, the mean age of stroke was 62.2 years overall compared to 59.6 and 56.6 in South and Southeast Asia, respectively. [34] In other part of the world, the risk of stroke among European populations in 6 countries study revealed that median age was 73 years and the distribution based on gender was varied more than 2-fold in men and women. On average, higher rates of stroke were observed in eastern and lower rates in southern European countries [35]. while in the United States, according to a 2012 study, the mean age of stroke in the United States is 74.3 years, with a standard deviation of 13.6. Men tend to have strokes at a younger age than women, with a mean age of 71.4 years for men and 76.9 years for women [36]. The mean age of stroke patients in Africa varies by study and location: In Sub Saharan Africa region, including Somalia, stroke represents a significant portion of the chronic disease burden in previous studies veiling the whole region (not including data from Somalia), the overall pooled prevalence of ischemic stroke among all strokes was 61.4%, with a mean age range of 55 – 62.8 [37] while a study from South Africa revealed the mean age of the patients was 57.7 years [38].

The relative risk definitely increases with age, the incidence doubling with each decade after the age of 45 years and over 70% of all strokes occur above the age of 65. Of the estimated 795, 000 new or recurring strokes that occur in the United States each year, approximately 145, 000 will result in death [39]. Pathobiologically, aging is associated with shifts in the cerebral arteries’s anatomy and physiology that actually can be used to predict the relative risk of stroke [40]. These patho-bio-anatomical shifts include:

- (1) endothelial dysfunction which release inflammatory mediators [41] that facilitates chronic neuro-inflammation [42],

- (2) impaired cerebral blood flow autoregulation [43] that lead to microvascular injury [44] which further lead to brain dysfunction and permanent cognitive or other higher brain function impairment [45] and
- (3) Impaired neurovascular coupling, even reaching the zone outside the affected area [46], which definitely lead to a decline in cortical function and connectivity to other part of the brain [47].

The three factors above showed us how keeping a stable blood supply to the brain is essential for survival because it supplies the brain with sufficient oxygen, glucose and nutrients, which are necessary for the brain's homeostasis and to keep its function properly.

Based on their financing system, most of stroke patients in our hospital using BPJS as their primary funding source for their medical expenses (94.4%) in comparison with 4.9% that uses their own pocket money/private patient and only 0.7% considered as patient using other type of insurances. Other literature regarding the percentage of stroke patients in Indonesia who use BPJS compared to other insurance or private costs is difficult to find on the internet. Most of the available literature is regarding studies on the use of BPJS in various types of chronic diseases, such as cancer [48], diabetes [49] and including stroke [14,15]. This shows that BPJS has been used widely in Indonesia and the coverage includes many types of diseases, various medical procedures/treatments as well as various types of health services from the first level (primary care) to the most advanced level (secondary or tertiary care in tier hospital) [16-20].

In 2022, healthcare costs for stroke patients in Indonesia, covered by BPJS Kesehatan, varied widely from Rp 1,937,400 to Rp 207,675,700, averaging Rp 8,487,531 per person. BPJS Kesehatan's total spending on stroke treatment and rehab in 2022 was reaching Rp 2,883,269,825,972, [15] one of the disadvantage is, as revealed by Rachmawati et al that the hospital costs exceeded the BPJS health insurance tariff leading to financial losses [50]

Most of our patient comes from the sub-district (*kecamatan*) Makkasar (27.2%) followed by Kramat Jati (22.2%), Jatinegara (10.5%), Duren sawit (6.2%), Cipayung (4.3%), Ciracas (2.5%), Cakung (1.2%), with Pasar Rebo and Pulogadung each 0.6%. These are sub district located in east Jakarta. There are 40 patients (24.7%) came from outside east Jakarta. This shows us that patient's referral system is organized with the aim of providing good quality healthcare, so that the purpose of service is achieved without having to use expensive costs [51], including in the context of distance.

Implementing this type of referral system makes health services in primary health facilities must be strengthened because First Level Health Facilities (FKTP) act as the first contact and filter for referrals to health services. FKTP is required to handle 144 disease diagnoses entirely according to the Indonesian Doctor Competency Standard (SKDI), which follows the Clinical Practice Guidelines.

We further evaluate the distance from the patient's address to out hospital. In order to estimate the distance (km) from the home address to the hospital, we used the directions feature from Google Maps and then entered the home address as the place of origin and the hospital address as the destination. The total distance shown by Google Maps is then recorded for further analysis to determine the distribution of distance from house to hospital based on the financing scheme. Mean distance of patient's address which use BPJS was 6.41km (Median 4.0km). On the other hand, for private patients, the mean distance from their home to our hospital was 12.87km (median 8.85km) and for OTI, there was only one

patient and the distance from his/her house to our hospital was 11.4km. Data showed in the following box plot diagram presented in fig. 1.

Distance to a health facility can affect how common or to which extent people use health services. [52] Delays in care-seeking for life threatening illness such as stroke may lead to more severe or even fatal outcomes [53]. a relationship between travelling further and having worse health outcomes cannot be ruled out and should be considered within the healthcare services location debate. [54]

The result of our evaluation showed that the distance from a primary healthcare facility, based on the patient care financing system, was significantly different between group of BPJS vs private vs OTI (using Kruskal Wallis non parametric test,  $p=0.022$ ). in general, BPJS patient's home were closer to our hospital compared to private patients and OTI patient.




Fig. 1. (A) To estimate the distance between patient's house to the hospital, we use Route feature available in Google Maps™.(B) We put the patient's address as complete as possible in the place of origin and put our hospital (RSU UKI) in the destination. we chose to simulate using a vehicle/driving and the results showed in kilometres. All data regarding distance were put in Microsoft Excel™ and then further exported to SPSS for further processing. (C) Box plot diagram of distance distribution of stroke patients, from their home to the hospital (in Km), based on their medical care financial system (BPJS, OTI or private)

This simple study actually is part of our student's thesis. By incorporating Google Maps™ with the patient's address and the hospital, we can develop simple health related geographic information system (GIS) by using data that is attached to a unique location, and it can be used to help people understand spatial patterns and relationships; in this context is the distribution of the patient in relation our hospital location. Post-covid pandemic, Medical education is become rapidly changing. Everyone in the business realize that COVID pandemic, globally and in Indonesia, had changed and forced the *civitas academica*, including lecturers, staffs and students, to make so many adjustments in order to make sure that learning is held in a proper manner, as best as possible [55]. During their educational process, medical students usually developed their own style of reading and learning in order

to address those issue (pile of textbooks / papers / lecture notes they must read). Those style of reading and learning also supported by advancement of information and communication technologies [56]. The educational goals of using technology in medical education include facilitating basic knowledge acquisition, improving decision making, enhancement of perceptual variation, improving skill coordination, practicing for rare or critical events, learning team training, and improving psychomotor skills. Different technologies can address these goals. [57]

This simple study showed us that the incorporation of information technology into existing data and combine it with other tools, e.g., maps, can make the data more informative to all stakeholder. Geographic Information System (GIS) is actually a trusted computer system that collects, stores, analyzes, and displays real time any kind of data (including health related data) in relation to the Earth's surface [58]. GIS is used in many industries and sciences, including health sector, to help stakeholder acknowledge spatial patterns and relationships [59]. GIS can also be used to inspect disease distribution (mapping, routing and even clustering) [60] and pinpoint areas with improving priority resource requirements [61], identifying communities, e.g., poor and underserved prone sub-population [62], or gaps in healthcare access [63], helping in prioritizing in strategic planning and effective-efficient response of local authorities [64]. GIS take part in an important role in governing the exact point where and when stakeholder start to surveillance [65], spatially mapping the risk pattern [66], the accessibility of health service and arbitrating health equity [67], increasing the quality of care [68], and even finding more cost-effective approach of health intervention in accordance to distance [69]. Our findings justify the previous last statement.

Geographical information systems (GIS) can be very helpful in the health related domain, but unfortunately there are some unsolved problems which limits its use, including:

- 1) its dependencies on the amount of data quality and availability, especially when doing field survey [70]. e.g., when preparing accurate data on disease transmission, then data regarding certain environmental condition and specific demographic information become a pre-requisition;
- 2) its vulnerability regarding data security and confidentiality because it usually deals with large and open source data [71], sometimes security devices become too prone for the potency of skimming, hacking and penetration. There is always concerns about data security and confidentiality;
- 3) its weaknesses in data interoperability and integration especially in emergency setting [72]; which include different type of data within the same file or handling the metadata, especially when the data needed in a short time;
- 4) its inconsistencies and even unwanted errors in the spatial scales of the data [73]; this condition can be produced by a number of unexpected factors, such as different nomenclature conventions, various data formats/programme, or even non-identical scales used by different sources. These discrepancies and even inconsistencies can make it hard to integrate or overlay datasets, and can lead to errors in analysis;
- 5) during its early year of application, its limitation in operating statistical functions needed was obvious; e.g., in order to emphasize analysis or even to withdraw any conclusion based on numerous data available, so in order to overcome the limitations to understanding spatial change between two time periods, spatial data

may be collected at multiple time and then the data tested using newer version of statistical test (raster GIS) [74],

- 6) its expensive cost and complexity which prevent single end user to buy the software product, e.g., Software costs: GIS software can cost between \$600 and \$17,000 per year, depending on the type and quality of the software, and the size of the project. For the hardware costs: Implementing GIS can involve additional hardware costs [75];
- 7) its lacking in quality and consistency of input addresses which in turn can be a source of error in the geocoding process [76];
- 8) its time consuming: The process of collecting, storing, and analyzing information using GIS can be time consuming. It may need a long time to obtain complete information regarding a certain specific set of data due to the vastness of the data availability and its related infrastructure [77];
- 9) as lesson learned from COVID-10 pandemic, GIS have limitation regarding disease outbreak detection [78]: GIS may not be able to detect rapidly evolving disease outbreaks due to challenges still continue in data aggregation, knowledge discovery, and dynamic expression.
- 10) its limitation in health disparities: GIS may not be able to directly identify localized health disparities; further careful assessment of the data and applications were required to properly interpret the findings in understanding the relationship between vulnerable populations and environmental burden [79].




Fig. 2. Geographic distribution of patient's residence based on their medical care financing system

#### 4. CONCLUSION

From three years of stroke patient's analysis, we found that more male patients (55.6%), with 42% were in the range of age between 55-64 years old, most of the patient (94.4%) using BPJS as their medical care financing system, with 27.7% came from sub district Makkasar which is near from our hospital. Further analysis using the help of Google maps™ we can show that the UHC or BPJS health referral system has facilitated patients receiving health services at the health facility closest to their residence.

#### ETHICAL APPROVAL (WHERE EVER APPLICABLE)

None to declare

#### REFERENCES

1. Usman SF. Social Security Fund as a Form of Constitutional Responsibility for the Right to Health. *Jurnal Jaminan Kesehatan Nasional (JJKN)*, 2021; 1(2): 95-106. <https://doi.org/10.53756/jjkn.v1i2.43>
2. Perdana NR, Adhasari G, G. Mahadewi EP. Challenges and Implementation of Universal Health Coverage Program in Indonesia . *International Journal of Health and Pharmaceutical (IJHP)*, 2022;2(3), 589–596. <https://doi.org/10.51601/ijhp.v2i3.97>
3. Anonymous. JKN coverage reached 95.77 percent of Indonesia's population: Ministry. *Antara*. 2024 March 28<sup>th</sup>. <https://en.antaranews.com/news/309495/jkn-coverage-reached-9577-percent-of-indonesias-population-ministry>
4. Kamath R, Brand H. A Critical Analysis of the World's Largest Publicly Funded Health Insurance Program: India's Ayushman Bharat. *Int J Prev Med*. 2023 Feb 18;14:20. [https://doi.org/10.4103/ijpvm.ijpvm\\_39\\_22](https://doi.org/10.4103/ijpvm.ijpvm_39_22).
5. Kunarti S, Sudrajat T, Handayani SW. Transformation of Social Security Administrative Body (BPJS) within Social Security Reform in Indonesia. *SHS Web of Conferences*, 2018 54, 03017. <https://doi.org/10.1051/shsconf/20185403017>
6. NATIONAL TEAM for the acceleration of poverty reduction: Secretariat of the Vice President of the Republic of Indonesia. *The Road to National Health Insurance (JKN)*. 2015. [https://www.tnp2k.go.id/images/uploads/downloads/FINAL\\_JKN\\_road%20to%20national%20health%20insurance.pdf](https://www.tnp2k.go.id/images/uploads/downloads/FINAL_JKN_road%20to%20national%20health%20insurance.pdf)
7. Asyrofi D, Ariutama IG. Deficit of Health Social Security Fund in National Health Insurance Program: A Case Study of BPJS Kesehatan. *Jurnal Ekonomi dan Studi Pembangunan*.2019; 11. 2019. <https://doi.org/10.17977/um002v11i22019p116>.
8. Kurnianingtyas D, Santosa B, Siswanto N, Pham DT. Reforming premium amount in the Indonesian National Health Insurance System program using system dynamics. *Cogent Engineering*, 2021;8(1). <https://doi.org/10.1080/23311916.2021.1938368>

9. Wibawa F, Madya F. "The Public Policy Dynamics of Increasing Health Insurance Rates in a Developing Country" in The 3rd International Conference on Governance, Public Administration, and Social Science (ICoGPASS), KnE Social Sciences, 2022; 49–69. <https://doi.org/10.18502/kss.v7i9.10927>
10. Wenang S, Schaeffers J, Afdal A, Gufron A, Geyer S, Dewanto I, Haier J. Availability and Accessibility of Primary Care for the Remote, Rural, and Poor Population of Indonesia. *Front Public Health*. 2021 Sep 21;9:721886. <https://doi.org/10.3389/fpubh.2021.721886>.
11. Sambodo NP, Bonfrer I, Sparrow R, Pradhan M, van Doorslaer E. Effects of performance-based capitation payment on the use of public primary health care services in Indonesia. *Social Science & Medicine*, 2023; 327: 115921. <https://doi.org/10.1016/j.socscimed.2023.115921>.
12. Prayitno L, Novita N, Esther S, Mahadewi EP. The Usage Of Government Formulary For BPJS Inpatients With Ischemic Stroke In Indonesia. *International Journal of Health and Pharmaceutical (IJHP)*, 2024;4(3):491–8. <https://doi.org/10.51601/ijhp.v4i3.347>
13. Pandian JD, Padma Srivastava MV, Aaron S, Ranawaka UK, Venketasubramanian N, Sebastian IA, Injety RJ, Gandhi DBC, Chawla NS, Vijayanand PJ, Rangamani S, Kalkonde YV. The burden, risk factors and unique etiologies of stroke in South-East Asia Region (SEAR). *Lancet Reg Health Southeast Asia*. 2023 Oct 9;17:100290. <https://doi.org/10.1016/j.lanseas.2023.100290>.
14. Perkasa AN. A Comparison of the Clinical Picture of Stroke Before and After Implementation of National Health Insurance (BPJS) in RSUP Dr Sardjito Yogyakarta. Thesis. 2020. Universitas Gadjah Mada. <https://etd.repository.ugm.ac.id/penelitian/detail/186579>
15. Santoso M, Adisasmito WB, Hasibuan SR. Cost of Illness: Stroke Patients with Medical Rehabilitation . *Jurnal Keterampilan Fisik*, 2024;9(1), 1–9. <https://doi.org/10.37341/jkf.v9i1.422>
16. Noerjoedianto D, Solida A, Mekarisce AA. Analysis of Contribution Assistance Recipients' Utilization of National Health Insurance Participants. *Journal of Applied Nursing and Health*, 2022; 4(2):215-23
17. Handayani PW, Dartanto T, Moeis FR, Pinem AA, Azzahro F, Hidayanto AN, et al. The regional and referral compliance of online healthcare systems by Indonesia National Health Insurance agency and health-seeking behavior in Indonesia. *Heliyon*, 2021; 7(9): e08068. <https://doi.org/10.1016/j.heliyon.2021.e08068>
18. Sukartini T, Arifin H, Kurniawati Y, Pradipta RO, Nursalam N, Acob JRJ. Factors Associated with National Health Insurance Coverage in Indonesia. *F1000Res*. 2022 Sep 14;10:563. <https://doi.org/10.12688/f1000research.53672.2>.
19. Wulandari RD, Laksono AD, Rohmah N, Ashar H. Regional differences in primary healthcare utilization in Java Region-Indonesia. *PLoS One*. 2023 Mar 27;18(3):e0283709. <https://doi.org/10.1371/journal.pone.0283709>.

20. Fanda RB, Probandari A, Yuniar Y, Hendarwan H, Trisnantoro L. The availability of essential medicines in primary health centers in Indonesia: achievements and challenges across the archipelago. *The Lancet Regional Health-Southeast Asia*, 2024; 22:100345. <https://doi.org/10.1016/j.lansea.2023.100345>
21. Darmasurya D, Anindita R, Ikhda MC. Did The Horizontal Referral Policy in Indonesia Create Cost Savings? *Jurnal Jaminan Kesehatan Nasional*, 2021; 1(1): 15-32. <https://doi.org/10.53756/jjkn.v1i1.14>
22. Venketasubramanian N, Yudiarto FL, Tugasworo D. Stroke Burden and Stroke Services in Indonesia. *Cerebrovasc Dis Extra*. 2022;12(1):53-57. <https://doi.org/10.1159/000524161>.
23. Jarman B, Aylin P, Bottle A. Acute stroke units and early CT scans are linked to lower in-hospital mortality rates. *BMJ*. 2004 Feb 14;328(7436):369. <https://doi.org/10.1136/bmj.328.7436.369>.
24. Gao L, Tan E, Kim J, Bladin CF, Dewey HM, Bagot KL, Cadilhac DA, Moodie M. Telemedicine for Stroke: Quantifying the Long-Term National Costs and Health Benefits. *Front Neurol*. 2022 Jun 20;12:804355. <https://doi.org/10.3389/fneur.2021.804355>.
25. Lazarus G, Permana AP, Nugroho SW, Audrey J, Wijaya DN, Widyahening IS. Telestroke strategies to enhance acute stroke management in rural settings: A systematic review and meta-analysis. *Brain Behav*. 2020 Oct;10(10):e01787. <https://doi.org/10.1002/brb3.1787>.
26. Zamzam M, Qomaruddin MB, Kurniavie LE, Herdiani N. Factors influencing stroke in Indonesia based on the Indonesia Family Life Survey 5. *Journal of Public Health in Africa*. 2023 May 25;14(S2):2568. <https://doi.org/10.4081/jphia.2023.2568>
27. Rissanen I, Basten M, Exalto LG, Peters SAE, Visseren FLJ, Geerlings MI; UCC-SMART-Study Group. Sex differences in modifiable risk factors for stroke incidence and recurrence: the UCC-SMART study. *J Neurol*. 2024 Jun;271(6):3347-3358. <https://doi.org/10.1007/s00415-024-12268-6>.
28. Yoon CW, Bushnell CD. Stroke in Women: A Review Focused on Epidemiology, Risk Factors, and Outcomes. *J Stroke*. 2023 Jan;25(1):2-15. <https://doi.org/10.5853/jos.2022.03468>.
29. Manwani B, Finger C, Lisabeth L. Strategies for Maintaining Brain Health: The Role of Stroke Risk Factors Unique to Elderly Women. *Stroke*, 2022; 53. <https://doi.org/10.1161/STROKEAHA.121.036894>.
30. Ospel J, Singh N, Ganesh A, Goyal M. Sex and Gender Differences in Stroke and Their Practical Implications in Acute Care. *J Stroke*. 2023 Jan;25(1):16-25. <https://doi.org/10.5853/jos.2022.04077>.
31. Soto-Cámara R, González-Bernal JJ, González-Santos J, Aguilar-Parra JM, Trigueros R, López-Liria R. Age-Related Risk Factors at the First Stroke Event. *Journal of Clinical Medicine*. 2020; 9(7):2233. <https://doi.org/10.3390/jcm9072233>

32. Meaghan Roy-O'Reilly, Louise D McCullough, Age and Sex Are Critical Factors in Ischemic Stroke Pathology, *Endocrinology*, 2018; 159 (8): 3120–31, <https://doi.org/10.1210/en.2018-00465>
33. Kusuma Y, Venketasubramanian N, Kiemas L, Misbach J. Burden of Stroke in Indonesia. *International journal of stroke: official journal of the International Stroke Society*. 2009;4: 379-80. <https://doi.org/10.1111/j.1747-4949.2009.00326.x>.
34. Pandian JD, Padma Srivastava MV, Aaron S, Ranawaka UK, Venketasubramanian N, Sebastian IA, et al. The burden, risk factors and unique etiologies of stroke in South-East Asia Region (SEAR). *Lancet Reg Health Southeast Asia*. 2023 Oct 9;17:100290. doi: <https://doi.org/10.1016/j.lansea.2023.100290>.
35. European Registers of Stroke (EROS) Investigators; Heuschmann PU, Di Carlo A, Bejot Y, Rastenyte D, Ryglewicz D, Sarti C, Torrent M, Wolfe CD. Incidence of stroke in Europe at the beginning of the 21st century. *Stroke*. 2009 May;40(5):1557-63. <https://doi.org/10.1161/STROKEAHA.108.535088>.
36. Kissela BM, Khoury JC, Alwell K, Moomaw CJ, Woo D, Adeoye O, Flaherty ML, Khatri P, Ferioli S, De Los Rios La Rosa F, Broderick JP, Kleindorfer DO. Age at stroke: temporal trends in stroke incidence in a large, biracial population. *Neurology*. 2012 Oct 23;79(17):1781-7. <https://doi.org/10.1212/WNL.0b013e318270401d>
37. Arslan E, Cetinkaya O. Analysis of the demographic characteristics and clinical profile of acute ischemic strokes admitted to the emergency centre in the Somalia population. *Afr J Emerg Med*. 2022 Sep;12(3):216-224. <https://doi.org/10.1016/j.afjem.2022.04.010>.
38. Tribelhorn S, Motara F, Lewis C M. Stroke: A retrospective review of the incidence and epidemiology in a South African academic hospital emergency department. *SAMJ, S. Afr. med. J*, 2021 Dec; 111( 12 ): 1205-1210. <http://dx.doi.org/10.7196/samj.2021.v111i12.15682>.
39. Kelly-Hayes M. Influence of age and health behaviors on stroke risk: lessons from longitudinal studies. *J Am Geriatr Soc*. 2010 Oct;58 Suppl 2(Suppl 2):S325-8. <https://doi.org/10.1111/j.1532-5415.2010.02915.x>.
40. Kizhisseri M, Gharai S, Boopathy SR, Lim RP, Mohammadzadeh M, Schluter J. Differential sensitivities to blood pressure variations in internal carotid and intracranial arteries: a numerical approach to stroke prediction. *Sci Rep*, 2023; 13: 22319. <https://doi.org/10.1038/s41598-023-49591-3>
41. Dri E, Lampas E, Lazaros G, Lazarou E, Theofilis P, Tsioufis C, Tousoulis D. Inflammatory Mediators of Endothelial Dysfunction. *Life*. 2023; 13(6):1420. <https://doi.org/10.3390/life13061420>
42. Pacinella G, Ciaccio AM, Tuttolomondo A. Endothelial Dysfunction and Chronic Inflammation: The Cornerstones of Vascular Alterations in Age-Related Diseases. *International Journal of Molecular Sciences*. 2022; 23(24):15722. <https://doi.org/10.3390/ijms232415722>
43. Carlson A, Mayer A, Cole C, van der Horn H, Marquez J, Stevenson T, Shuttleworth C. Cerebral autoregulation, spreading depolarization, and implications for targeted

therapy in brain injury and ischemia. *Reviews in the Neurosciences*. 2024;35(6): 651-678. <https://doi.org/10.1515/revneuro-2024-0028>

44. Toth P, Szarka N, Farkas E, Ezer E, Czeiter E, Amrein K, et al. Traumatic brain injury-induced autoregulatory dysfunction and spreading depression-related neurovascular uncoupling: Pathomechanisms, perspectives, and therapeutic implications. *Am J Physiol Heart Circ Physiol*. 2016 Nov 1;311(5):H1118-H1131. <https://doi.org/10.1152/ajpheart.00267.2016>.
45. Wang S, Roman RJ, Fan F. Duration and magnitude of bidirectional fluctuation in blood pressure: the link between cerebrovascular dysfunction and cognitive impairment following spinal cord injury. *J Neurobiol Physiol*. 2020;2(1):15-18. <https://doi.org/10.46439/neurobiology.2.008>.
46. Staehr C, Giblin JT, Gutiérrez-Jiménez E, Guldbrandsen HØ, Tang J, Sandow SL, Boas DA, Matchkov VV. Neurovascular Uncoupling Is Linked to Microcirculatory Dysfunction in Regions Outside the Ischemic Core Following Ischemic Stroke. *J Am Heart Assoc*. 2023 Jun 6;12(11):e029527. <https://doi.org/10.1161/JAHA.123.029527>.
47. Mukli P, Pinto CB, Owens CD, Csipo T, Lipecz A, Szarvas Z, et al. Impaired Neurovascular Coupling and Increased Functional Connectivity in the Frontal Cortex Predict Age-Related Cognitive Dysfunction. *Adv. Sci*. 2024, 11, 2303516. <https://doi.org/10.1002/adv.202303516>
48. Nancy, Maysaroh T, Unggulian HG. Adult Cancer Patients' Quality of Life After Outpatient Chemotherapy with JKN Coverage. *Jurnal Jaminan Kesehatan Nasional*, 2023; 3(1):130-42. <https://doi.org/10.53756/jjkn.v3i1.144>
49. Putri LP, Mawarni D, Trisnantoro L. Challenges of Shifting Diabetes Mellitus Care From Secondary- to Primary-Level Care in Urban and Rural Districts: A Qualitative Inquiry Among Health Providers. *J Prim Care Community Health*. 2020 Jan-Dec;11:2150132720924214. <https://doi.org/10.1177/2150132720924214>.
50. Rachmawaty R, Sinrang AW, Wahyudin E, Bukhari A. Evaluation of health care quality among insured patients in Indonesian mother & child hospital: A secondary data analysis. *Gac Sanit*. 2021;35 Suppl 2:S613-S618. <https://doi.org/10.1016/j.gaceta.2021.10.100>.
51. Riati L, Heryana A, Muda CAK, Mahadewi EP. The Analysis of JKN-KIS Hospital Outpatient Referral System Implementation Subsequent to Online Referral Application. In *Proceedings of the 1st International Conference on Health (ICOH 2019)*, 2020;pp 29-33 SCITEPRESS – Science and Technology Publications, Lda. ISBN: 978-989-758-454-1 <https://doi.org/10.5220/0009562500290033>
52. Bai R, Gao J, Dong W. Is Travel Time Associated with Health Service Utilization in Northwest China? Evidence from Shaanxi Province. *Int J Gen Med*. 2022 May 13;15:4949-4957. <https://doi.org/10.2147/IJGM.S360582>.
53. Oldenburg CE, Sié A, Ouattara M, Bountogo M, Boudo V, Kouanda I, Lebas E, Brogdon JM, Lin Y, Nyatigo F, Arnold BF, Lietman TM; Étude CHAT Study Group. Distance to primary care facilities and healthcare utilization for preschool children in

- rural northwestern Burkina Faso: results from a surveillance cohort. *BMC Health Serv Res.* 2021 Mar 9;21(1):212. <https://doi.org/10.1186/s12913-021-06226-5>.
54. Kelly C, Hulme C, Farragher T, et al. Are differences in travel time or distance to healthcare for adults in global north countries associated with an impact on health outcomes? A systematic review. *BMJ Open* 2016;6:e013059. <https://doi.org/10.1136/bmjopen-2016-013059>
  55. Daroedono E, Siagian FE, Alfarabi M, Cing JM, Arodes ES, Sirait RH, et al. The impact of COVID-19 on medical education: our students perception on the practice of long distance learning. *International Journal of Community Medicine and Public Health*, 2020; 7 (7): 2790-2796
  56. Siagian FE, Maryanti E. E-Reading Vs Traditional Reading: Can Internet, Social media and Gadgets Bridge the Gap between Reading and Learning among Medical Students? *Jurnal Ilmu Kedokteran*, 2018; 11 (2): 52-9. <http://dx.doi.org/10.26891/JIK.v11i2.2017.52-59>
  57. Guze PA. Using Technology to Meet the Challenges of Medical Education. *Trans Am Clin Climatol Assoc.* 2015;126:260-70.
  58. Davenhall WF, Kinabrew C. GIS in Health and Human Services. *Springer Handbook of Geographic Information.* 2012:557–78. [https://doi.org/10.1007/978-3-540-72680-7\\_29](https://doi.org/10.1007/978-3-540-72680-7_29).
  59. Scott LM. Spatial Pattern, Analysis of. *International Encyclopedia of the Social & Behavioral Sciences.* 2015. <https://doi.org/10.1016/B978-0-08-097086-8.72064-2>.
  60. Murad A, Khashoggi BF. Using GIS for Disease Mapping and Clustering in Jeddah, Saudi Arabia. *ISPRS International Journal of Geo-Information.* 2020;9: 328. <https://doi.org/10.3390/ijgi9050328>.
  61. Robin TA, Khan MA, Kabir N. Using spatial analysis and GIS to improve planning and resource allocation in a rural district of Bangladesh. *BMJ Global Health* 2019;4:e000832. <https://doi.org/10.1136/bmjgh-2018-000832>
  62. Mohammed RN, Khawari A, Shaguy JA, Abouzied A. A GIS-based approach to identifying communities underserved by primary health care services-An Afghanistan case study. *Front Public Health.* 2023 Sep 21;11:1209986. <https://doi.org/10.3389/fpubh.2023.1209986>.
  63. Geofana A. GIS Application for Health Facility Coverage Mapping in Temanggung Regency, Central Java Province. *JP2WD.* 2021;5(3):160-73. <https://doi.org/10.29244/jp2wd.2021.5.3.160-173>
  64. Holdstock D. *Strategic GIS Planning and Management in Local Government.* 1st ed. CRC Press, 2016. Web. 15 Oct. 2022.
  65. Osheyor J, Biu P, Nwasike G, Tula O, Ezeigweneme A, Gidiagba J. A review of GIS applications in public health surveillance. *World Journal of Advanced Research and Reviews.* 2024;21: 30-039. <https://doi.org/10.30574/wjarr.2024.21.1.2684>.
  66. Joseph PV, Balan B, Rajendran V, Prashanthi DM, Somnathan B. Probability Mapping to Determine the Spatial Risk Pattern of Acute Gastroenteritis in

- Coimbatore District, India, Using Geographic Information Systems (GIS). *Indian J Community Med.* 2015 Jul-Sep;40(3):188-92. <https://doi.org/10.4103/0970-0218.158865>.
67. Kim Y, Byon YJ, Yeo H. Enhancing healthcare accessibility measurements using GIS: A case study in Seoul, Korea. *PLoS ONE*, 2018;13(2): e0193013. <https://doi.org/10.1371/journal.pone.0193013>
  68. Biu P, Nwasike C, Nwaobia N, Ezeigweneme C, Gidiagba J. GIS in healthcare facility planning and management: A review. *World Journal of Advanced Research and Reviews.* 2024; 21: 012-019. <https://doi.org/10.30574/wjarr.2024.21.1.2682>.
  69. Hawthorne T, Timothy & Kwan MP., Mei-Po. (2012). Using GIS and Perceived Distance to Understand the Unequal Geographies of Healthcare in Lower-Income Urban Neighborhoods. *The Geographical journal.* 2012;178: 18-30. <https://doi.org/10.2307/41475787>.
  70. Wang F, Reinhardt W. Spatial data quality concerns for field data collection in Mobile GIS - art. no. 64201C. *Proceedings of SPIE - The International Society for Optical Engineering.*6420. 2016. <https://doi.org/10.1117/12.712733>.
  71. Solymosi R, Buil-Gil D, Ceccato V, Kim E, Jansson U. Privacy challenges in geodata and open data. *Area.* 2023;55: 1-29. <https://doi.org/10.1111/area.12888>.
  72. Abdalla R, Tao C, Li J. Challenges for the Application of GIS Interoperability in Emergency Management. 2007; 10. [https://doi.org/1007/978-3-540-72108-6\\_25](https://doi.org/1007/978-3-540-72108-6_25).
  73. Heuvelink GBM, Burrough PA, Stein, A. Propagation of errors in spatial modelling with GIS, *International Journal of Geographical Information Systems*,1989; 3(4): 303-22. [https://www.geos.ed.ac.uk/~gisteac/gis\\_book\\_abridged/files/ch14.pdf](https://www.geos.ed.ac.uk/~gisteac/gis_book_abridged/files/ch14.pdf)
  74. Schlagel JD, Newton CM. A GIS-Based Statistical Method to Analyze Spatial Change. *Photogrammetric Engineering & Remote Sensing*, 1996; 62(7): 839-44
  75. Anonymous. How Much is GIS Software? Comparing Costs and Features. Downloaded from <https://sitemap.com/articles/how-much-is-gis-software-comparing-costs-and-features/>
  76. Wong DW, So BKL, Zhang P. Addressing quality issues of historical GIS data: an example of Republican Beijing. *Annals of GIS*, 2012; 18(1), 17–29. <https://doi.org/10.1080/19475683.2011.647074>
  77. Peuquet D. Time in GIS and geographical databases. 2005. [https://www.geos.ed.ac.uk/~gisteac/gis\\_book\\_abridged/files/ch08.pdf](https://www.geos.ed.ac.uk/~gisteac/gis_book_abridged/files/ch08.pdf)
  78. Osiecki KM, Kim S, Chukwudozie IB, Calhoun EA. Utilizing Exploratory Spatial Data Analysis to Examine Health and Environmental Disparities in Disadvantaged Neighborhoods. *Environ Justice.* 2013 Jun;6(3):81-87. <https://doi.org/10.1089/env.2013>.

UNDER PEER REVIEW