

STUDIES ON GENETIC VARIABILITY, CORRELATION AND PATH ANALYSIS IN FORAGE SORGHUM

ABSTRACT

Ten genotypes *i.e.*, CSV-15, Pusa Chari-6, HC-308, Pant Chari-4, G-48, HJ-513, UP Chari-4, HC-171, SSG-59-3 and ICSV-700 studied for genetic variability, correlation and path analysis for days to 50 % flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant, leaf area, leaf stem ratio, total soluble solids, protein content, fiber content, HCN content and green fodder yield. The analysis of variance for the experiment with fifty five treatments for all the thirteen attributes *viz.*, days to 50% flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant, leaf area, leaf stem ratio, total soluble solids, protein content, fiber content, HCN content and green fodder yield revealed significant differences among the material used in the present investigation, which indicated that wide spectrum of variation among the genotypes. Genotypic and phenotypic coefficient of variation was high (more than 25%) observed for leaf stem ratio and green fodder yield, which indicating that more variability and scope for selection in improving these characters. As a percentage of mean for leaf area, leaf stem ratio, leaf breadth, stem girth, number of leaves per plant, total soluble solids, and green fodder production, there is a strong heritability combined with a high genetic progress. This suggested that these qualities were highly heritable and that they could be enhanced by selecting high-performing genotypes. At the genotypic and phenotypic levels, green fodder output showed a strong, consistent, and positive connection with plant height, days to 50% blooming, leaf length, leaf breadth, stem girth, number of leaves per plant, and leaf area. These traits could be regarded as crucial yield components in sorghum forage. Leaf breadth displayed high order of direct effect on green fodder yield per plant followed by leaves per plant, leaf length and stem girth at phenotypic and genotypic level, which indicated that the contribution of individual attributes to fodder yield is of importance in planning a sound breeding programme for developing for high yielding varieties.

Key words: *sorghum bicolor*, variability, correlation, path coefficient analysis

INTRODUCTION

Sorghum, scientifically known as *Sorghum bicolor*, is a versatile and resilient cereal grain that has been cultivated for millennia. Originating from Africa, it has spread across the globe due to its adaptability to diverse climates and its numerous uses ranging from food and feed to industrial

applications. It belongs to the grass family, Poaceae, and is classified under the genus *Sorghum*. Sorghum is an annual plant that typically grows between 2 to 5 meters in height, although dwarf varieties exist. The plant has a robust root system, which aids in its ability to withstand drought and grow in regions with limited water availability. Sorghum produces large terminal panicles that contain seeds varying in size and color, depending on the variety. The cultivation of sorghum dates back thousands of years, primarily in Africa where it was domesticated. From there, it spread to Asia and eventually to the Americas during the Columbian Exchange. Sorghum played a crucial role in food security in regions with harsh climates and poor soil conditions, where other crops struggled to thrive. Its ability to grow in semiarid and arid regions made it a staple crop for many communities around the world. There are numerous varieties of sorghum cultivated globally, each adapted to different environmental conditions and intended uses. These varieties can be broadly categorized into grain sorghum, sweet sorghum and forage sorghum. Grain sorghum is primarily grown for its seeds, which are used for human consumption, animal feed and industrial purposes. Sweet sorghum is grown for its sweet juice, which can be processed into syrup or fermented into biofuels. Forage sorghum, on the other hand, is used as fodder for livestock. Sorghum is cultivated in a variety of agroecological zones, from tropical to temperate regions. The crop is adaptable to different soil types but thrives in well drained soils with good organic matter content. Its tolerance to heat and drought makes it suitable for regions where other cereal crops might fail. Sorghum is typically sown directly into the soil, either broadcast or in rows, depending on the farming system and equipment available. The economic importance of sorghum extends beyond its role as a staple food. It is used in various industrial applications such as the production of biofuels, biodegradable plastics and alcoholic beverages. In recent years, sorghum has gained attention as a potential crop for sustainable bioenergy production due to its high biomass yield and efficient use of water. Sorghum is highly nutritious, containing essential nutrients such as carbohydrates, proteins, fiber, vitamins, and minerals. It is gluten free, making it suitable for individuals with gluten intolerance or celiac disease. The grain's nutritional profile varies slightly among different varieties, with some varieties being richer in certain nutrients than others. Its deep root system improves soil structure and water retention, reducing erosion and enhancing soil fertility. Additionally, sorghum's efficient use of water makes it a resilient crop in the face of climate change and water scarcity, offering food security to millions of people worldwide. India is one of the largest producers of jowar globally and production in India has fluctuated over the years based on various factors such as weather conditions, government policies, and market demand. Productivity of sorghum in India varies across different regions and is influenced by agricultural practices, irrigation availability and use of modern farming techniques. The area under sorghum cultivation in India has seen some fluctuations, but it remains an important crop in several states, particularly Maharashtra, Karnataka,

Telangana, and Madhya Pradesh. Uttar Pradesh also cultivates sorghum, although it is not the dominant crop compared to states in central and southern India. Sorghum production in U.P. contributes to the overall national production figures, but specific data for recent years would be needed to provide precise numbers and area under sorghum cultivation in UP is comparatively smaller than in states where sorghum is a major staple crop. It has good nutritional value, especially when harvested at the right stage of growth, offering protein, fiber, and other essential nutrients. The need for genetic improvement of sorghum for increased yields and nutritive value of fodder is having high priority. It can be used as part of a crop rotation strategy to improve soil health and reduce pests and diseases in subsequent crops. Some varieties of sorghum have been studied for their potential in biofuel production, contributing to its versatility and potential economic benefits beyond traditional forage use. It can contribute to soil conservation and reduce erosion due to its extensive root system, helping to maintain soil structure and fertility over time. “The fodder sorghum is grown in 8.3 million ha mainly in Western U.P., Haryana, Punjab and Rajasthan and fulfils more than 65 per cent of the fodder demand during *kharif* season. The area under fodder cultivation is estimated to be about four per cent of the gross cropped area, which remained static for the last four decades. The traditional grazing lands are gradually diminishing because of urbanization, expansion of cultivable area, grazing pressure and industrialization *etc.* In India, only 4.4% area is under fodder crops, out of which fodder sorghum is grown on 2.3 million ha. India faces a net deficit of 36% and 11% of green fodder and dry fodder, respectively. To reduce the demand and supply gap, the production and productivity of fodder crops needs to be enhanced. The horizontal expansion of cultivable area under fodder crops is difficult due to severe competition from food crops. Apart from vertical expansion, utilization of non cultivable areas for pastures is one of the most viable options to balance the demand. Seed are the best propagating material for wide spread regeneration of marginal and uncultivable lands. One of the reasons reported to stumble the green fodder production is non availability of quality seed in sufficient quantities. As per estimation only 25-30 per cent of required quantity of quality seed is available in cultivated fodders and produce abundant. The great quality seed is pre-essential to improve the production and fodder yield of sorghum” (Kumar *et al.*, 2022).

MATERIAL AND METHODS

The experimental material for the present investigation was comprised of ten promising diverse parents *viz.*, CSV-15, Pusa Chari-6, HC-308, Pant Chari-4, G-48, HJ-513, UP Chari-4, HC-171, SSG-59-3 and ICSV-700 and their all possible 45 F₁'s, developed through crossing ten parental lines in diallel mating design (excluding reciprocals). All genotypes were evaluated in a complete randomized block design with three replications at Crop Research Center of Sardar

Vallabhbhai Patel University of Agriculture and Technology, Meerut, during *kharif* season 2022 and 2023. The observations were recorded on ten traits *viz.*, days to 50 % flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant, leaf area, leaf stem ratio, total soluble solids, protein content, fiber content, HCN content and green fodder yield. The data recorded on all these traits were subjected to various statistical and biometrical analyses like to work out analysis of genetic variability, correlation and path analysis. Seeds of ten parents and F₁'s sown by hand dibbling method and the length of each row was kept 4m with inter and intra row distances of 30cm and 10cm, respectively. The coefficients of variation, correlation and path coefficient analysis calculated by the formula given by **Burton (1952)** and **Johnson *et al.* (1955)**, **Crumpacker and Allard (1962)**, **Robinson *et al.* (1949)**, **Fisher (1918)** and **Dewey and Lu (1959)**

RESULTS AND DISCUSSION

The analysis of variance for the experiment with fifty five treatments for all the thirteen attributes *viz.*, days to 50% flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant, leaf area, leaf stem ratio, total soluble solids, protein content, fiber content, HCN content and green fodder yield revealed significant differences among the material used in the present investigation, which indicated that wide spectrum of variation among the genotypes. The variance due to treatments was further partitioned in to their orthogonal components *i.e.*, parents, crosses and parents vs crosses. Parents and crosses recorded highly significant differences for all the thirteen traits whereas parent's vs crosses were found highly significant differences for all the characters except protein content and fiber content (Table-1). Estimates of coefficient of variation *viz.*, genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) along with heritability and genetic advance as percent of mean for different characters are presented in table-2. High percentage of genotypic coefficient of variation (more than 25%) observed for leaf stem ratio (28.31) and green fodder yield (37.69) whereas moderate (10-25%) noted for leaf breadth (16.36), stem girth (16.80), leaves per plant (13.38), leaf area (21.34) and total soluble solids (16.51). Low genotypic coefficient of variation (<10%) recorded for days to 50% flowering (5.94), plant height (8.45), leaf length (7.54), protein content (7.31), fiber content (2.00) and HCN content (5.11) which indicating that more variability and scope for selection in improving these characters. Similar results were found by **Singh *et al.* (2023)** and **Shafiqurrahman *et al.* (2024)**. High percentage of phenotypic coefficient of variation (more than 25%) estimated for leaf stem ratio (30.39) and green fodder yield (38.31) while moderate (10-25%) showed for leaf breadth (16.79), stem girth (17.65), leaves per plant (14.95), leaf area (21.77) and total soluble solids (17.77). Low genotypic coefficient of variation (<10%) revealed for days to 50% flowering (6.39), plant height (8.97), leaf length (8.16), protein content (7.79), fiber content (3.26) and HCN content (7.11), indicated that

these traits are more influenced by environmental factors. Earlier researchers **Wadikar et al. (2018)** and **Dev et al. (2019)**. High heritability (> 60%) exhibited for days to 50% flowering (86.52), plant height (88.69), leaf length (85.29), leaf breadth (94.94), stem girth (90.67), leaves per plant (80.09), leaf area (96.08), leaf stem ratio (86.75), total soluble solids (86.34), protein content (88.04) and green fodder yield (96.82) whereas recorded medium heritability (30-60%) for fiber content (37.66) and HCN content (51.72) suggested that these characters are under genotypic control. Similar observations were also reported by **Diwakar et al. (2016)**, **Singh et al. (2016)** and **Jain et al. (2017)**. Expected genetic advance expressed as percentage of mean revealed high (> 20%) for leaf breadth (32.84), stem girth (32.96), leaves per plant (24.67), leaf area (43.09), leaf stem ratio (54.31), total soluble solids (31.36) and green fodder yield (76.40) while moderate genetic advance as percentage of mean (10-20%) observed for days to 50% flowering (11.39), plant height (16.39), leaf length (14.34) and protein content (14.12). Low genetic advance expressed as percentage of mean (<10%) was found for fiber content (2.53) and HCN content (7.57). This indicated that these traits were highly heritable and selection of high performing genotypes is possible to improve these attributes. High heritability coupled with high genetic advance for these characters have also been reported earlier by **Srivastava et al. (2019)**, **Singh et al. (2019)** and **Sirohi et al. (2019)**. Results on correlation coefficients, computed at phenotypic and genotypic both levels for all possible paired combinations among ten attributes and the values are given in Table-3. The genotypic correlation, in general was similar in sign and slightly higher in magnitude than their phenotypic correlation. At the genotypic level, the estimates of correlation coefficients were generally similar to that observed at the phenotypic level, though, their magnitude was slightly lower than the corresponding phenotypic correlations. Here, also green fodder yield per plant showed significant and positive correlation with stem girth (0.76) followed by leaf area (0.67), leaf breadth (0.63), leaves per plant (0.61), leaf length (0.53), plant height (0.39) and days to 50% flowering (0.15) whereas showed significant negative correlation with leaf stem ratio (-0.17) and protein content (-0.19). Green fodder yield exhibited significant stable and positive correlation with days to 50% flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant and leaf area at genotypic and phenotypic level. These characters may be considered as important yield component in forage sorghum. These results are similar to earlier reports of **Dubey et al. (2016)**, **Singh et al. (2017)**, **Sirohi et al. (2019)** and **Dev et al. (2019)**. The results obtained at both genotypic and phenotypic levels are presented in table-4. At genotypic level, leaf breadth (0.67) displayed high order of direct effect on green fodder yield followed by leaves per plant (0.65), leaf length (0.60) and stem girth (0.43). However the character which contributed indirect effects toward green fodder yield per plant observed stem girth through leaf area; leaves per plant via leaf length, leaf breadth, stem girth and leaf area; leaf area through days to 50% flowering, plant height, leaf length, leaf breadth, stem

girth, leaves per plant, leaf stem ratio and HCN content. At phenotypic level, leaf length (0.59) displayed high order of direct effect on green fodder yield followed by stem girth (0.47), leaf breadth (0.43) and leaves per plant (0.42) whereas, the attribute which contributed indirect effects toward green fodder yield per plant observed leaf breadth through stem girth, leaves per plant and leaf area; stem girth via plant height and leaf area; leaf area through days to 50% flowering, plant height, leaf length, leaf breadth, stem girth, leaves per plant and HCN content. Magnitudes of residual effects at both phenotypic and genotypic level were found to be low. Leaf breadth displayed high order of direct effect on green fodder yield per plant followed by leaves per plant, leaf length and stem girth at phenotypic and genotypic level, which indicated that the contribution of individual attributes to fodder yield is of importance in planning a sound breeding programme for developing for high yielding varieties. These findings are in accordance with the results obtained in sorghum by **Malik *et al.* (2013)**, **Girish *et al.* (2016)**, **Malaghan and Kajjidoni (2019)**, **Sirohi *et al.* (2019)** and **Srivastava *et al.* (2019)**.

Conclusion

The leaf stem ratio and green fodder output showed a significant (more than 25%) genotypic and phenotypic coefficient of variation, suggesting greater variability and room for selection to improve these traits. As a percentage of mean for leaf area, leaf stem ratio, leaf breadth, stem girth, number of leaves per plant, total soluble solids, and green fodder production, there is a strong heritability combined with a high genetic progress. This suggested that these qualities were highly heritable and that they could be enhanced by selecting high-performing genotypes. At the genotypic and phenotypic levels, green fodder output showed a strong, consistent, and positive connection with plant height, days to 50% blooming, leaf length, leaf breadth, stem girth, number of leaves per plant, and leaf area. These characters could be regarded as significant. Leaf breadth displayed high order of direct effect on green fodder yield per plant followed by leaves per plant, leaf length and stem girth at phenotypic and genotypic level, which indicated that the contribution of individual attributes to fodder yield is of importance in planning a sound breeding programme for developing for high yielding varieties.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

Acknowledgement. I would like to thanks my all endeavours.

Conflict of interest. None

REFERENCES

- Burton, G.W. (1952).** Quantitative inheritance in grasses. Proc. 6th Int. Grassland Cong.1:227-283.
- Crumpacker, D.W. and Allard, R.W. (1962).** A diallel cross analysis of heading date in wheat. *Helgardia* 32: 275-318.
- Robinson, H.F., Comstock, R.E. and Harvey, P.H. (1949).** Genotypic and phenotypic correlation in corn, their importance in selection, *Agron, J.*, 43: 282-287.
- Johnson, H.W., Robinson, H.F. and Camstock, F. (1955).** Genotypic and phenotypic correlation in soybean and their implications in selection. *Agron. J.*, 47: 477-483.
- Allard, R.W. (1956).** Estimation of prepotency from lilma bean diallel cross data. *Agron. J.* 48: 537-543.
- Dewey, D.R. and Lu, K.H. (1959).** A correlation and path coefficient analysis components of crested wheat grass. *Journal of Agronomy*, 51: 515-518.
- Sirohi, M., Singh, S.K., Chand, P., Kumar, M., Gupta, S. and Kumar, M. (2019).** Character association and path analysis in forage Sorghum. *Journal of Experimental Zoology India*, 22(1): 1469-1473.
- Srivastava, M., Singh, S.K., Chand, P., Kumar, M., Kumar, M., Gangwar, H.K. and Dev, A. 2019.** Studies on path analysis in forage sorghum [*Sorghum bicolor* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*, 8(5): 993-996.
- Girish, G., Kiran, S. B., Lokesh, R., Vikas, Kulkarni, V., Rachappa, V., Yogesh, L. N. and Talwar, A. M. 2016.** Character association and path analysis in advanced breeding lines of *rabi* sorghum [*Sorghum bicolor* (L.) Moench]. *Journal of Applied and Natural Science*, 8(1): 35–39.

Malaghan, S. and Kajjidoni, S.T. 2019. Character association and path analysis of grain yield in rabi sorghum [*Sorghum bicolor* (L.) Moench]. *International Journal of Chemical Studies*, 7(1): 2309-2313.

Malik, A., Singh, S.K., Chand, P., Singh, B. and Singh, D.K. 2013. Correlation and path analysis in forage sorghum. *Progressive Agriculture*, 13(1): 234-238.

Singh, A. and Singh, S.K. (2017). Studies on combining ability in forage sorghum for yield and quality parameters. *Journal of plant development sciences*,9(8): 785-782.

Dubey, M.K., Pandey, P.K., Shrotria, P.K., Chauhan, P. and Pandey, G. 2016. Assessment of variability, correlation and path analysis for yield and yield related traits in sorghum genotypes. *Green Farming*, 7(6): 1291-1295.

Dev, A., Singh, S. K., Chand, P., Kumar, M., Kumar, M., Poonia, M. and Srivastava, M. 2019. Genetic variability, character association and path analysis in forage sorghum. *Journal of Pharmacognosy and Phytochemistry*, 8(5): 1135-1139.

Singh, S.K., Gangwar, L.K. and Chaudhary, M. (2019). Studies on character association and path analysis in forage sorghum. *International Journal of Chemical Studies*, 7(5): 2939-2942.

Srivastava, M., Singh, S.K., Chand, P., Kumar, M., Kumar, M., Gangwar, H.K. and Dev, A. 2019. Studies on path analysis in forage sorghum [*Sorghum bicolor* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*, 8(5): 993-996.

Diwakar, A., Ranwah, B.R., Sharma, D.B. and Sinha, S.K. (2016). Study of variability parameters in forage sorghum [*Sorghum bicolor* (L.) Moench]. *Research of Environment Life Science*,9(5): 528-530.

Jain, S.K., Elangovan, M. and Patel, N.V. (2017). Correlation and path coefficient analysis for agronomical traits in forage sorghum [*Sorghum bicolor* (L.) Moench] *Indian Journal. Plant Genet. Resou*,23(1): 15-18.

Singh, S.K., Singh, A. and Kumar, R. 2016. Genetic variability, character association and path analysis in forage sorghum. *Progressive Agriculture*, 16(2): 214-218.

Dev, A., Singh, S. K., Chand, P., Kumar, M., Kumar, M., Poonia, M. and Srivastava, M. 2019. Genetic variability, character association and path analysis in forage sorghum. *Journal of Pharmacognosy and Phytochemistry*, 8(5): 1135-1139.

Wadikar, P. B., Ubale, D. L., Magar, M. R. and Thorat, G. S. 2018. Genetic variability studies in sweet sorghum [*Sorghum bicolor* (L.) Moench]. *Int. J. Curr. Microbiol. App. Sci. Special*, 6: 920-923.

Singh, S.K., Singh, M., Gangwar, L.K. and Singh, B. (2023). Study on path analysis in forage sorghum [*Sorghum bicolor* (L.) moench]. *International Journal of Farm Sciences* 13(1): 81

Table-1. Analysis of variance for fodder yield and yield components in forage sorghum (*Sorghum bicolor* L. Moench)

Source of variation	df	Days to 50% flowering	Plant height (cm)	Leaf length (cm)	Leaf breadth (cm)	Stem girth (mm)	Leaves per plant	Leaf area (cm) ²	Leaf stem ratio	Total soluble solid (%)	Protein content (%)	Fiber content (%)	HCN content (ppm)	Green fodder yield (g/plant)
Replication	2	0.55	403.46	2.73	0.26	0.29	0.11	663.47	0.03	2.54	1.04	0.57	694.66	1718.96
Treatment	54	86.70**	2544.44*	115.95**	2.76*	18.78*	9.47*	15063.03*	0.08*	12.49*	1.21*	1.39**	406.01*	42492.95*
Parents	9	106.95*	2163.56*	89.90*	1.45*	11.25*	9.37*	10106.12*	0.05*	13.10*	1.21*	1.77**	1093.41**	8641.11**
Crosses	44	74.69**	1940.55*	106.80**	2.71*	17.86*	8.83*	13615.09*	0.07*	12.51*	1.23*	1.34**	252.15*	40343.43*
Parents vs Crosses	1	432.99*	32533.03**	753.40**	16.74**	127.09**	38.56**	123384.44**	0.02*	5.85**	0.22	0.04	989.16*	441735.78**
Error	108	4.28	103.77	6.31	0.05	0.62	0.72	202.20	0.04	0.63	0.05	0.49	96.34	460.62

*, ** significant at 5% and 1% level, respectively

Table- 2. Phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), heritability and genetic advance as % of mean in forage sorghum (*Sorghum bicolor* L. Moench)

Sl. No.	Genotypes	GCV (%)	PCV (%)	Heritability (%)	Genetic Advance	Genetic Advance as % of mean
1.	Days to 50% flowering	5.94	6.39	86.52	10.04	11.39
2.	Plant height (cm)	8.45	8.97	88.69	55.33	16.39
3.	Leaf length (cm)	7.54	8.16	85.29	11.50	14.34
4.	Leaf breadth (cm)	16.36	16.79	94.94	1.91	32.84
5.	Stem girth (mm)	16.80	17.65	90.67	4.83	32.96
6.	Leaves per plant	13.38	14.95	80.09	3.15	24.67
7.	Leaf area (cm) ²	21.34	21.77	96.08	142.12	43.09
8.	Leaf stem ratio	28.31	30.39	86.75	0.10	54.31
9.	Total soluble solid (%)	16.51	17.77	86.34	3.81	31.60
10.	Protein content (%)	7.31	7.79	88.04	1.20	14.12
11.	Fiber content (%)	2.00	3.26	37.66	0.69	2.53
12.	HCN content (ppm)	5.11	7.11	51.72	15.05	7.57
13.	Green fodder yield (g/plant)	37.69	38.31	96.82	239.92	76.40

Table-3. Estimates of Genotypic (G) and Phenotypic (P) correlation coefficients for different traits in forage sorghum (*Sorghum bicolor* L. Moench)

Parameters		Days to 50% flowering	Plant height (cm)	Leaf length (cm)	Leaf breadth (cm)	Stem girth (mm)	Leaves per plant	Leaf area (cm) ²	Leaf stem ratio	Total soluble solid (%)	Protein content (%)	Fiber content (%)	HCN content (ppm)	Green fodder yield (g/plant)
Days to 50% flowering	G	1.00	0.07	0.20**	0.19*	0.15*	0.09	0.20**	0.22**	-0.06	-0.19*	-0.05	-0.03	0.15*
	P	1.00	0.07	0.17*	0.17*	0.15*	0.09	0.18*	0.21**	-0.09	-0.18*	-0.02	-0.02	0.15*
Plant height (cm)	G		1.00	0.24**	0.33**	0.44**	0.13	0.34**	-	-0.07	-0.08	-0.03	-0.10	0.39**
	P		1.00	0.20**	0.31**	0.40**	0.13	0.32**	-	-0.04	-0.06	-0.05	-0.13	0.36**
Leaf length (cm)	G			1.00	0.47**	0.60**	0.43**	0.74**	-0.04	-0.02	-0.10	-0.04	-0.01	0.53**
	P			1.00	0.42**	0.54**	0.34**	0.69**	-0.04	-0.09	-0.09	-0.08	0.02	0.48**
Leaf breadth (cm)	G				1.00	0.73**	0.55**	0.94**	0.22**	-0.16*	-0.14	0.09	0.09	0.63**
	P				1.00	0.69**	0.48**	0.91**	0.20**	-0.14*	-0.12	0.04	0.08	0.61**
Stem girth (mm)	G					1.00	0.63**	0.80**	-0.01	-0.07	-0.04	-0.14	0.14	0.76**
	P					1.00	0.52**	0.76**	-0.02	-0.06	-0.02	-0.05	0.11	0.73**

Leaves per plant	G						1.00	0.55**	0.10	-0.01	0.05	0.03	0.13	0.61**	
	P						1.00	0.50**	0.12	-0.05	0.05	0.03	0.07	0.54**	
Leaf area (cm) ²	G							1.00	0.12	-0.11	-0.15	0.03	0.11	0.67**	
	P							1.00	0.12	-0.10	-0.13	-0.03	0.07	0.64**	
Leaf stem ratio	G								1.00	-0.18*	-0.05	-0.09	0.10	-0.17*	
	P								1.00	-0.15*	-0.04	-0.04	0.08	-0.18*	
Total soluble solid (%)	G									1.00	0.01	-0.07	0.13	-0.07	
	P									1.00	0.09	-0.06	0.09	-0.06	
Protein content (%)	G										1.00	-0.13	-0.04	-0.19*	
	P										1.00	-0.08	-0.08	-0.16*	
Fiber content (%)	G											1.00	-	0.58**	0.01
	P											1.00	-	0.20**	0.01
HCN content (ppm)	G												1.00	-0.08	
	P												1.00	-0.06	
Green fodder yield (g/plant)	G														1.00
	P														1.00

*, ** significant at 5% and 1% level, respectively

Table-4. Estimates of direct and indirect effect of different characters on green fodder yield per plant in forage sorghum (*Sorghum bicolor* L. Moench)

Parameters	Days to 50% flowering	Plant height (cm)	Leaf length (cm)	Leaf breadth (cm)	Stem girth (mm)	Leaves per plant	Leaf area (cm) ²	Leaf stem ratio	Total soluble solid (%)	Protein content (%)	Fiber content (%)	HCN content (ppm)
------------	-----------------------	-------------------	------------------	-------------------	-----------------	------------------	-----------------------------	-----------------	-------------------------	---------------------	-------------------	-------------------

Days to 50% Flowering	G	0.08	-0.05	0.03	-0.05	0.02	0.06	0.83	- 0.04	0.04	0.04	0.02	0.02
	P	0.03	-0.01	0.02	0.07	0.07	0.02	0.60	- 0.05	0.03	0.03	0.01	0.03
Plant height (cm)	G	0.07	-0.06	0.09	-0.06	0.06	0.09	0.44	0.04	0.04	0.01	0.01	0.11
	P	0.02	-0.01	0.03	0.13	0.19	0.03	0.16	0.05	0.02	0.01	0.02	0.02
Leaf length (cm)	G	0.01	-0.05	0.60	-0.17	0.08	0.38	0.99	0.09	0.03	0.02	0.01	0.01
	P	0.05	-0.02	0.59	0.18	0.09	0.08	0.26	0.01	0.04	0.01	0.03	-0.04
Leaf breadth (cm)	G	0.01	-0.02	-0.06	0.67	0.01	0.36	0.92	- 0.04	0.08	0.03	-0.03	-0.05
	P	0.05	-0.03	-0.06	0.43	0.02	0.11	0.29	- 0.05	0.05	0.02	-0.02	-0.01
Stem girth (mm)	G	0.01	-0.02	-0.27	-0.12	0.43	0.41	0.36	0.03	0.04	-0.01	0.05	-0.09
	P	0.05	-0.04	0.08	0.30	0.47	0.12	0.24	0.07	0.02	0.02	0.02	-0.01
Leaves Per Plant	G	0.08	-0.08	-0.20	-0.09	0.08	0.65	0.30	- 0.02	0.01	-0.01	-0.01	-0.08
	P	0.03	-0.01	0.05	0.21	0.09	0.42	0.16	- 0.03	0.01	-0.01	-0.01	-0.01
Leaf area (cm) ²	G	0.01	-0.02	-0.19	-0.10	0.51	0.36	-0.17	- 0.02	0.06	0.03	-0.01	-0.07
	P	0.06	-0.03	0.11	0.39	0.35	0.12	-0.32	- 0.03	0.03	0.02	0.02	-0.01
Leaf stem ratio	G	0.01	0.01	0.06	-0.06	-0.02	0.06	0.53	- 0.21	0.01	0.01	0.03	-0.06
	P	0.07	0.02	-0.07	0.09	-0.01	0.03	0.04	- 0.25	0.05	0.08	0.02	-0.01
Total soluble solid (%)	G	-0.06	0.05	0.03	0.05	-0.01	0.04	-0.47	0.03	-0.05	0.03	0.06	-0.10
	P	-0.03	0.01	-0.01	-0.06	-0.02	-0.01	-0.03	0.03	-0.03	-0.02	0.03	-0.01
Protein	G	-0.01	0.05	0.16	0.04	0.01	0.03	-0.05	0.01	0.01	-0.22	0.05	0.02

content (%)	P	-0.05	0.01	-0.01	-0.05	-0.01	0.01	-0.04	0.01	0.01	-0.18	0.04	0.01
Fiber content (%)	G	-0.05	0.02	0.06	-0.26	-0.02	0.02	0.16	0.02	0.09	0.03	-0.08	0.37
	P	-0.01	0.01	-0.01	0.02	-0.02	0.07	0.01	0.01	0.02	0.01	-0.03	0.03
HCN Content (ppm)	G	-0.03	0.01	0.01	-0.26	0.01	0.08	0.49	- 0.02	-0.09	0.01	0.22	-0.40
	P	0.05	0.01	0.04	0.03	0.05	0.01	0.52	- 0.02	-0.03	0.02	0.08	-0.56

Residual values (G) = 0.02, Residual values (P) = 0.02 Bold values indicate direct effects *, ** significant at 5% and 1% level, respectively