

## Effect of planting dates and interval of reproductive pruning on yield and yield attributing characters of Yam bean (*Pachyrrhizus erosus* L.)

### Abstract

An experiment on “Effect of planting dates and interval of reproductive pruning on yield and yield attributing characters of Yam bean (*Pachyrrhizus erosus* L.)” was conducted at Department of vegetable Science, College of Horticulture, Dapoli, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli during year 2023-24. Two factors were studied during the investigation using factorial randomized block design viz. a) Planting dates (D) D<sub>1</sub>-1<sup>st</sup> week of June, D<sub>2</sub>-3<sup>rd</sup> week of June, D<sub>3</sub> -1<sup>st</sup> week of July, D<sub>4</sub> -3<sup>rd</sup> week of July and b) Interval of reproductive pruning (P) P<sub>1</sub> – Weekly, P<sub>2</sub> – Fortnightly, P<sub>3</sub> – No pruning. The results showed that weekly reproductive pruning (P<sub>1</sub>) with planting date 3<sup>rd</sup> week of June (D<sub>2</sub>) gave best effect on yield per plant, average weight of tuber, yield per plot, yield per hectare and length and diameter of tuber respectively. Those values were 642.03 g, 407.70 g, 12.90 t ha<sup>-1</sup>, 21.50 cm and 17.53 cm.

**Keywords:** Yam bean, planting dates, reproductive pruning, yield

### 1.Introduction

Yam Bean (*Pachyrrhizus spp.*) is one of the multifunctional, underutilized minor tuber crops belongs to the family Leguminosae, under the subfamily Papilionaceae. There are five species under the genera *Pachyrrhizus* viz. *P. erosus*, *P. tuberosus* and *P. ahipa* are cultivated species and *P. ferrugineus*, *P. panamensis* are wild species. Among this *P. erosus* is mostly cultivated all over the world. Yam beans have refreshing, crispy ice-white, fruit-flavored tuber which can eaten raw or as cooked in various sweet dishes and in numerous savory dishes worldwide. Tubers have very low-calories as well as high-quality dietary fiber (Oligofructose insulin) and antioxidants and some amount of minerals and vitamins. Yam bean tubers are also rich in vitamin C which is a powerful water-soluble antioxidant that helps the body scavenge

harmful free radicals, thereby offering protection from cancers, inflammation, viral cough and cold. It contains a healthy amount of potassium which is responsible for keeping our heart in good shape [2].

Yam bean contributes to ensuring food security by providing food and feeds products, enrichment of other foods that are deficient in major nutrients with beneficial phytochemicals and bioactive compounds that reduce the risk of diseases. Yam bean cultivation is now becoming more popular among the farmers of the Konkan region due its nutritional properties. Yam bean can withstand in high rainy conditions due to their hardy nature. The lateritic soils of the Konkan region are generally sandy clay loam in texture with pH 5.0-6.0, highly base leached and sesquioxide soils favour the production of yam bean. Successful yam bean production in many regions depends upon selecting suitable times for sowing by the specific environment. Further, it is highly nutritious therefore increasing demand from consumers. Thus, there is good scope for increasing the production and productivity of yam bean in the Konkan region [5].

## 2. Materials and methods

The field trial was conducted at field of College of Horticulture, Dapoli, Dr. Balasaheb Sawant Konkan Krishi Vidyapeeth, Dapoli, Dist- Ratnagiri (M.S) during *Kharif* season of the year 2023- 2024. Experiment was carried out in Factorial Randomized Block Design with three replications. Two factors were studied during the investigation viz. a) Planting dates (D) D<sub>1</sub>-1<sup>st</sup> week of June, D<sub>2</sub>3<sup>rd</sup> week of June, D<sub>3</sub> -1<sup>st</sup> week of July, D<sub>4</sub> -3<sup>rd</sup> week of July and b) Interval of reproductive pruning (P) P<sub>1</sub> – Weekly, P<sub>2</sub> – Fortnightly, P<sub>3</sub> – No pruning Individual plot size was 3 m x 3 m. The spacing between the ridges was 60cm and seed to seed distance was 20 cm respectively. In between two replications spacing 1m was added to maintain plants and avoid the nutrients uptake from the other block. Single seed was sown on the ridge at the spot of fertilizer application which were made as per the desired spacing at the depth of 2-3 cm. on four different planting dates i.e. 1<sup>st</sup> week of June, 3<sup>rd</sup> week of June, 1<sup>st</sup> week of July and 3<sup>rd</sup> week of July respectively. 6 to 7 weeks after sowing yam bean starts flowering. Removal of flower buds is done at intervals of weekly, fortnightly and no pruning. Removal of flower bud is a practice for better production of yam bean tubers. The flower buds were removed at the purple colour and particularly opened stage.

The harvesting of the tubers started 120 days after planting. Yield per plant, average tuber weight, yield per hectare, length and diameter of tuber were observed as parameters. The data obtained in the present investigation were statistically analysed by the method suggested by

Panase and Sukhatme [16]. The standard error (S.E.) of means was worked and a critical difference (CD) at 5% was also worked out whenever the result was significant.

### **3. Result and discussion**

#### **3.1 Yield per plant (g)**

In present investigation, the combined effect of planting date and reproductive pruning was significant on tuber yield per plant (g). The highest yield was recorded in D<sub>2</sub>P<sub>2</sub> (642.03 g), which was at par with the treatment D<sub>2</sub>P<sub>1</sub> (607.83 g), the lowest yield per plant was recorded in D<sub>4</sub>P<sub>3</sub> (250.13 g). The interaction effect of planting dates and interval of reproductive pruning may give more yield per plant due to the June often coincides with favourable weather conditions, including adequate rainfall and sunlight. The similar trend observed by Karhale *et al.* [12] in *kharif* sorghum, Bobade *et al.* [6] in *kharif* green gram, Lawrence *et al.* [13] in pigeon pea and Nisha Kumari *et al.* [15] in sweet potato. Pruning practice could increase the tuber yield of plant due to plants shifts energy from reproductive parts to vegetative parts (leaf and tuber). The similar finding was reported by the Mardhiana *et al.* [14] in cucumber. Similar findings were reported by the Gao, *et al.* [9] in *Helianthus tuberosus* L

#### **3.2 Average tuber weight (g)**

The combined effect of planting date and reproductive pruning was significant on average tuber weight (g). The highest average tuber weight was found to be in treatment D<sub>2</sub>P<sub>1</sub> (407.70 g) which was at par with D<sub>1</sub>P<sub>2</sub> (387.17), D<sub>3</sub>P<sub>1</sub> (386.67 g), D<sub>2</sub>P<sub>2</sub> (382.83 g) and D<sub>1</sub>P<sub>1</sub> (331.36 g). The lowest were recorded in D<sub>4</sub>P<sub>3</sub> (183.13 g). This may due to the Pruning practice could increase the tuber yield of plant due to plants shifts energy from reproductive parts to vegetative parts and the June often coincides with favourable weather conditions, including adequate rainfall and sunlight, which are essential for healthy plant growth and tuber development as compared to July planting.

#### **3.3 Yield per hectare (tonnes)**

The interaction effect of planting dates and interval of reproductive pruning on tuber yield per hectare (t ha<sup>-1</sup>) was found to be significant. The highest yield was recorded in D<sub>2</sub>P<sub>1</sub> (12.90 t ha<sup>-1</sup>) which was at par with D<sub>2</sub>P<sub>2</sub> (10.41 t ha<sup>-1</sup>). However, the lowest yield was obtained in D<sub>4</sub>P<sub>3</sub> (2.74 t ha<sup>-1</sup>). The planting dates 1<sup>st</sup> week of June and 3<sup>rd</sup> week of June give maximum yield due

to better growth and thrive well in heavy rain as compared to 1<sup>st</sup> week of July and 3<sup>rd</sup> week of July where more mortality of plants was observed. Similar findings were reported by the Kang *et al.* [11] in soybean, Canavar and Kaynak [7] in peanut, Bashir *et al.* [3] in rice, Karhale *et al.* [12] in *kharif* sorghum and Bobade *et al.* [6] in *kharif* green gram. While the Pruning practices increase the yield per hectare. Similar findings were reported by Belford *et al.* [4] in yam bean, Adjahossou, [1] in yam bean, Chakraborty *et al.* [8] in mungbean and Gibregwergis *et al.* [10] in potato.

### **3.4 Tuber length (cm)**

The combined effect of planting dates and interval of reproductive pruning on tuber length was found to be significant. The highest tuber length was found in D<sub>2</sub>P<sub>1</sub> (21.50 cm) which at par with D<sub>2</sub>P<sub>2</sub> (20.77 cm). However, the lowest tuber length was recorded in D<sub>4</sub>P<sub>1</sub> (16.17 cm). This may due to the planting at the right time can help to avoid stress from extreme weather conditions and allowing plants to allocate more energy to tuber growth and regular pruning can encourage bushier growth and increase leaf area, enhancing photosynthesis and ultimately leading to larger tubers.

### **3.5 Tuber diameter (cm)**

The interaction effect of planting dates and interval of reproductive pruning on tuber diameter was found significant. The highest tuber diameter was reported in D<sub>2</sub>P<sub>2</sub> (17.53 cm) which at par with D<sub>2</sub>P<sub>1</sub> (16.38 cm). However, the lowest tuber diameter was reported in D<sub>4</sub>P<sub>3</sub> (4.57 cm). This may due to the optimal planting dates and strategic reproductive pruning help to maximize resource allocation and growth conditions for yam beans. This approach allows the plants to thrive and produce larger and healthier tubers.

## **4. Conclusion**

Among the different treatments it may concluded that for better yield and yield attributing characters of yam bean, should be planted in 3<sup>rd</sup> week of June with weekly reproductive pruning i.e. **D<sub>2</sub>P<sub>1</sub>**. under Konkan agro-climatic conditions.

**Table 1. Effect of planting dates and interval of reproductive pruning and their interaction on yield per plant (g) in yam bean.**

Treatments	Yield per plant (g)			
	Interval of reproductive pruning (P)			
Planting dates (D)	P1	P2	P3	Mean
<b>D1</b>	490.73	486.10	453.00	<b>476.61</b>
<b>D2</b>	607.83	642.03	393.82	<b>547.89</b>
<b>D3</b>	475.67	406.67	286.87	<b>389.73</b>
<b>D4</b>	403.72	296.17	250.13	<b>316.67</b>
<b>Mean</b>	<b>494.48</b>	<b>457.74</b>	<b>345.95</b>	<b>432.72</b>
	F test	S.E m $\pm$	CD at 5%	
<b>D</b>	<b>SIG</b>	20.65	60.57	
<b>P</b>	<b>SIG</b>	17.89	52.46	
<b>DxP</b>	<b>SIG</b>	35.77	104.90	

**Table 2. Effect of planting dates and interval of reproductive pruning and their interaction on average tuber weight (g) of yam bean.**

Treatments	Average tuber wt. (g)			
	Interval of reproductive pruning (P)			
Planting dates (D)	P1	P2	P3	Mean
<b>D1</b>	331.16	387.17	194.50	<b>304.20</b>
<b>D2</b>	407.70	382.83	215.37	<b>335.30</b>
<b>D3</b>	386.67	293.33	189.33	<b>289.77</b>
<b>D4</b>	210.00	140.00	133.33	<b>161.11</b>
<b>Mean</b>	<b>333.88</b>	<b>300.83</b>	<b>183.13</b>	<b>272.61</b>
	F test	S.E m $\pm$	CD at 5%	
<b>D</b>	<b>SIG</b>	16.59	48.67	
<b>P</b>	<b>SIG</b>	14.37	42.15	
<b>DxP</b>	<b>SIG</b>	28.74	84.29	

**Table 3 Effect of planting dates and interval of reproductive pruning and their interaction on yield per hectare (ton) of yam bean.**

Treatments	Yield per ha. (ton)			
	Interval of reproductive pruning (P)			
Planting dates (D)	P1	P2	P3	Mean
<b>D1</b>	8.45	9.31	8.40	<b>8.72</b>
<b>D2</b>	12.90	10.41	7.18	<b>10.16</b>
<b>D3</b>	10.27	7.49	4.12	<b>7.29</b>
<b>D4</b>	4.77	6.14	2.74	<b>4.55</b>
<b>Mean</b>	<b>9.09</b>	<b>8.33</b>	<b>5.61</b>	<b>7.67</b>
	F test	S.E m $\pm$	CD at 5%	
<b>D</b>	<b>SIG</b>	0.51	1.49	
<b>P</b>	<b>SIG</b>	0.44	1.29	
<b>DxP</b>	<b>NS</b>	0.88	2.58	

**Table 4 Effect of planting dates and interval of reproductive pruning and their interaction on length (cm) of yam bean.**

Treatments	Length (cm)			
	Interval of reproductive pruning (P)			
Planting dates (D)	P1	P2	P3	Mean
<b>D1</b>	19.17	17.90	17.40	<b>18.15</b>
<b>D2</b>	21.50	20.77	19.73	<b>20.66</b>
<b>D3</b>	18.43	16.30	17.07	<b>17.26</b>
<b>D4</b>	16.13	18.40	16.17	<b>16.9</b>
<b>Mean</b>	<b>18.80</b>	<b>18.34</b>	<b>17.59</b>	<b>18.24</b>
	F test	S.E m $\pm$	CD at 5%	
<b>D</b>	<b>SIG</b>	0.29	0.83	
<b>P</b>	<b>SIG</b>	0.25	0.72	
<b>DxP</b>	<b>SIG</b>	0.49	1.44	

**Table 5 Effect of planting dates and interval of reproductive pruning and their interaction on diameter (cm) of yam bean tuber.**

Treatments	Diameter (cm)			
	Interval of reproductive pruning (P)			
Planting dates (D)	P1	P2	P3	Mean
<b>D1</b>	13.23	15.38	10.53	<b>13.04</b>
<b>D2</b>	16.38	17.53	8.53	<b>14.14</b>
<b>D3</b>	13.67	9.10	6.33	<b>9.70</b>
<b>D4</b>	9.36	5.83	4.57	<b>6.58</b>
<b>Mean</b>	<b>13.16</b>	<b>11.96</b>	<b>7.49</b>	<b>10.87</b>
	F test	S.E m $\pm$	CD at 5%	
<b>D</b>	<b>SIG</b>	0.32	0.92	
<b>P</b>	<b>SIG</b>	0.28	0.79	
<b>DxP</b>	<b>SIG</b>	0.55	1.59	

## 5. Reference

1. Adjahossou, D. F. (2006). Effect of flower pruning on the tuber yield of four yam bean accessions (*Pachyrhizus erosus* L.). *Cahiers Agricultures*, **15**(2): 213-219.
2. Anonymous (2014c). *Annual report of FAO-2013-14*. <https://www.fao.in>
3. Bashir, M. U., Nadeem, A., Iqbal, A. and Zaman, H. (2010). Effect of different sowing dates on yield and yield components of direct seeded coarse rice (*Oryza sativa* L.). *Pakistan J. Agricultural Science*, **47**(4): 361-365.
4. Belford, E. J., Karim, A. B. and Schroder, P. (2001). Exploration of the tuber production potential of yam bean (*Pachyrhizus erosus* L.) under field conditions in Sierra Leone. *J. Applied botany*, **75**(18): 31-38.
5. Bhattacharyya, T., Haldankar, P. M., Haldavanekar, P.C. and Burondkar, M. M. (2019). Impact of climate change on horticulture in Konkan, Maharashtra: activities and strategies. *Indian J. Fertilizers*, **17**(3): 258-273.
6. Bobade, B. R., Asewarand, B. V. and Vikhe, R. R. (2018). Influence of dates of sowing on *kharif* green gram (*Vigna radiate* L. Wilezek) varieties under varied weather conditions. *Trends in Biosciences*, **11**(6): 794-796.

7. Canavar, O. and Kaynak, M. A. (2008). Effect of different planting dates on yield and yield components of peanut (*Arachis hypogaea* L.). *Turkish J. Agriculture and Forestry*, **32**(6): 521-528.
8. Chakraborty, R., Chowdhury, T. E., Sumon, J. I., Mostofa, M. and Rahman, M. L. (2015). Contribution of flower removal on the performance of mungbean. *European J. Experimental Biology*, **5**(3): 66-71.
9. Gao, K., Zhang, Z., Zhu, T. and Coulter, J.A. (2020). The influence of flower removal on tuber yield and biomass characteristics of *Helianthus tuberosus* L.in a semi- arid area. *Industrial Crops and Products*. **150**(2): 15-20.
10. Geregwegis, F. H., Gebremicheal, M., Gebremendhin, H. and Assefa, A. (2021). The Effect of flower removal and earthing up on tuber yield and quality of potato (*Solanum Tuberosum* L.). *J.Agricultural Sciences*, **66**(2): 121-137.
11. Kang, y. k., Kim, H. T., Cho, n. c. and kim, Y. C. (2001). Effect of planting date and plant density on yield and quality of soybean forage in Jeru. *Korean J. crop Science*, **46**(2): 95-99.
12. Karhale, M. B., Jaybhaye, P. R., asewar, B. V. and Shinde, P. B. (2014). Effect of different sowing dates on growth and yield of kharif sorghum hybrids. *J. Agriculture and Veterinary Science*, **7**(12): 05-08.
13. Lowrence, K., Sharma, M. B. and Longchar, A. (2020). Effect of planting dates on the performance of promising pigeon genotypes under NEHZ. *International J. Economic Plants*, **7**(1): 6-8.
14. Mardhiana Pradhana, Adiwena, M., Dwisantoso, K., Rizza, W. and Anas, A. (2017). Effect of yield and growth of cucumber (*Cucumis sativus*) mercy variety in the acid soil of North Kalimantan, Indonesia. *Cell Biology and Development*, **1**(1): 13-17.
15. Nisha Kumari Meena, R. S. Rathore and Mithlesh Kumari Meena. (2020). Effect of planting dates and plant spacing on growth and yield attributes of sweet potato (*Ipomea batatas* Lam.) cv. Co-3-4. *International J. Current. Microbiology. Applied. Science*, **9**(4): 2602-2628.
16. Panse, V. G and P.V. Sukhatme (1995). Statistical methods for Agriculture workers. ICAR, New Delhi.