
Length – Weight Relationship of Rohu (*Labeo rohita*) advanced fingerlings under sodic soil condition

ABSTRACT

The present study was conducted at the Instructional Fish Farm, College of Fisheries, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya, Uttar Pradesh, in earthen ponds (8m×8m×1m). The experiment lasted for 90 days, using *Labeo rohita* advanced fingerlings (avg. wt. 30.0 ± 0.5g) under sodic soil conditions. Five treatments were taken in triplicate, with 50 fingerlings per pond. T1 served as the control with traditional feed only (MOC+RB in 1:1 ratio, 5% of total fish body weight). T2 and T3 were fertilized with cattle dung (20 and 15 tonnes/ha/yr) respectively. T4 and T5 received poultry droppings (10 and 7.5 tonnes/ha/yr) respectively. Results showed negative allometric growth ($b < 3$), indicating fish becoming slimmer with length increase. Final weight gain was highest in T4 (110.9±0.9 gm) and T2 (108.7±0.88 gm), and lowest in T1 (70.7±0.49 gm).

Keyword: Cattle dung, poultry dropping, *Labeo rohita*

INTRODUCTION

Aquaculture is the fastest-growing food production industry in the world. It is extending, intensifying, and expanding in almost every region of the earth. The increasing worldwide population is driving the demand for aquatic food items. The bulk of the principal fishing fields have achieved their maximum fishing capacity, and the production of the capture fisheries has stagnant, indicating that the worldwide demand for aquatic food will be satisfied through aquaculture. Aquaculture is considered as a potential solution to the global supply-demand imbalance for aquatic food (Subasinghe *et al.*, 2008). Fish is the principal source of protein for over 950 million people globally, and they are an important element of the diets of many more. Fish offer approximately 16% of the animal protein consumed by people worldwide and are a good source of minerals (Pradeepkiran *et al.*, 2019).

Carp dominate freshwater aquaculture, with three basic species. Rohu (*Labeo rohita* Hamilton 1822: Cypriniformes, Cyprinidae) is well-known for its flavor and popularity among the general population. Rohu is also regarded as a viable aquaculture species in India,

Bangladesh, Myanmar (Burma), and Pakistan. Farmed Rohu are omnivores that require a high protein diet (30-35%). Aquaculture's key constraints include feed prices, disease outbreaks, and seed quality, all of which assure the long-term viability of fish production. This species' high fertility (2million eggs/kg), external fertilization, and domestication make it excellent for intensive breeding (Rasal and Sundaray, 2020).

Labeo rohita is the eleventh most popular species in aquaculture. This species, which is categorized as Indo-Gangetic, is native to the riverine regions of northern and central India. For mono and composite fish farming systems (polyculture) in India, *L. rohita* is considered a suitable species due to its fast development, distinct eating niche and behavior, fertility, and market demand. The body is bilaterally symmetrical and slightly extended. It has a more arched dorsal shape than ventral, and it is coated in cycloid scales. The back is blue, while the flanks and belly are silver. Rohu's favorite food as an infant was zooplankton. Rohu feeds largely on phytoplankton and periphyton during its juvenile and adult stages of life (Jhingran and Pullin, 1985).

The length-weight relationship (LWR) is important method in fish biology research to determine the fish's growth form, maturity, reproductive state, and overall well-being. LWR is a valuable method for estimating biomass and assessing stock. LWR can also be used to examine intra-species variation based on the body shape of fish specimens obtained from various environments or geographical regions (Moutopoulos and Stergiou, 2002). Fish species growth rates vary depending on their genetic makeup, food availability, and current environmental conditions.

Labeo is a highly significant fish species, widely consumed across the country for its excellent nutritional value and easy availability. However, its population has recently been affected. Notably, the ray-finned fish, *Schizothorax richardsonii*, is also found in the river, preferring the cold, mountainous waters and feeding mainly on aquatic plants and detritus (Krishan & Tarana, 2010). This species commands high market value due to its taste and nutritional quality. The overall fish production in the river has decreased, raising serious concerns about the well-being of the fish species inhabiting these waters.

Experimental site

The present study was conducted at Instructional Fish Farm College of Fisheries, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya Uttar Pradesh. The experimental set up earthen pond (8m×8m×1m). Experimental fish *Labeo rohita* advanced fingerlings (avg.wt. 30.2± 0.5 g) were also brought from Instructional Fish Farm of College of Fisheries.

Experimental details

The experiment was conducted for 90 days period in the earthen pond. Each group were having three (triplicate) earthen pond, each tank stocked with 50 *Labeo rohita* advanced fingerlings (avg. wt. 30.0 ± 0.5g) under sodic soil condition. Total six treatments were taken in triplicate the details of treatments and control T1 was taken as control where no cattle dung and poultry dropping was added in control (T1) only traditional feed MOC+RB in 1:1 was given @ 5% of total fish body weight present in the pond.T2 was fertilized with (20 Tonnes/ha/yr) cattle dung and traditional feed was given. T3 was fertilized with cattle dung (15 Tonnes/ha/yr), T4 was added with poultry dropping (10 Tonnes /ha/yr), T5 was added with poultry dropping (7.5 Tonnes /ha/yr). Single Super Phosphate (SSP) was applied 25kg/ha and urea was added @ 20kg/ha per month. After 15 days of pond preparation fishes were stocked in ponds water quality parameters were measured and fish were fed with control feed for one week. For acclimatization fish were stocked into the pond for 30 minutes

Condition factor: The Condition factor (K) generally, used for determining the physiological state of a fish, including reproductive capacity. The heavier the fish for a length, the higher its condition factor (K). It is calculated by using the formula by (Fulton 1904):

where K is the condition factor, W is the body weight of fish in grams, and L is the total length in centimetres (Bagenal *et al* 1978).

$$K = W \times 100/L^2$$

Where, W = weight (g), L = length (cm) and 100 is a factor to bring the value of K near unity (Froese, 2006).

RESULTS AND DISCUSSION

Length-weight relationship

The total length (TL) was estimated as a distance from tip of snout up to the end of caudal fin with the help of measuring scale, while fish weight was recorded with the help of a digital balance (corrected up to 0 mg) after eliminating water and mucus from the fish body. For any

biological organism the Length-Weight relationship is generally non-linear and expressed in the form of parabolic equation $W = aL^b$ where weight (W) is proportional to a certain power (b) of the length (L) and 'a' is the intercept. Values of the exponent 'b' provide information on fish growth. When 'b'=3, increase in weight is isometric. When the value of 'b' is other than 3, weight increase is allometric (positive if 'b' >3, negative if 'b' <3) (Berhan *et al.*, 2019). Relationship between these two variables were adjusted by transforming them into linear regression (Ricker 1975).


Table 1. Length-Weight relationship (LWR) different treatments

Treatment	a	b	Rsqr
T1	0.079824	2.301808	0.890889
T2	0.020759	2.816166	0.941575
T3	0.032692	2.66518	0.923328
T4	0.021111	2.823435	0.944928
T5	0.033908	2.657976	0.909125


T1= Traditional Feed +soil base ,T2 =Traditional feed+ cattle dung @20T /ha ,T3= Traditional feed+ cattle dung @15T /ha, T4= Traditional feed+ poultry dropping @ 10 T/ha,T5 Traditional feed+ poultry dropping @ 7.5 T /ha

During the present study, 'b' values ranged from 2.30-2.82 with corresponding 'r' values as 0.89-0.94 at the selected sites. Individuals showed negative *L. rohita* negative as it less than 3 allometric growth at the selected sites. During the present study the values of $b < 3$ clearly indicate that fishes are becoming slimmer with increase in length i.e the weight of fish is lower than cube of its length. The final weight gain observed was significantly higher and was maximum T4 (110.9±0.9 gm) followed by T2 (108.7±0.88 gm).The weight gain was minimum in T1 (70.7±0.49 gm). The introduction of poultry dropping boosted the protein content of the diet, which has been shown to improve fish growth performance (Steffens, 1989). Many other workers have reported similar findings in cold water cyprinids where fish is becoming slimmer


with increase in weight (Dar *et al.*, 2012). The values of correlation coefficient depicted a strong positive correlation between length and weight indicating an increase in length with corresponding increase in weight. The similar findings have been reported by Singh *et al.*, (2024) reported that the relationship between length and weight in *L. rohita*, with regression coefficients ranging from 0.242 to 3.234 across treatments. The findings indicated negative allometric growth. Gupta *et al.*, (2005) reported that the value of 'b' was noted less than '3' in the case of *S. richardsonii* thriving in the same habitat, thus showing a negative allometric growth. Similar results were also revealed by Pandey *et al.*, (1998) for *L. rohita* (0.917-1.04), *C. catla* (0.903-1.07) and *C. mrigala* (0.976-1.031) from a sodic soil pond in Uttar Pradesh. Mir *et al.* (2012) reported the condition factor of male rohu to vary between 1.20 to 1.51 while it was 1.26 to 1.40 in case of female rohu from six different rivers of Ganga basin.



L-W relationship
T3



L-W relationship
T4



UNDER PEER REVIEW




Fig 1. Graph showing L-W relationship

All treatments in the study exhibited negative allometric growth, with 'b' values ranging from 2.30 to 2.82 and 'r' values between 0.89 and 0.94, indicating that fish became slimmer as they increased in length. The highest weight gain was observed in T4 (110.9±0.9 gm) due to the use of poultry droppings, which enhanced dietary protein content. In contrast, the control group (T1) showed the lowest weight gain (70.7±0.49 gm). Similar findings were reported by Ujjania *et al.* (2012) and others, showing negative allometric growth in *Labeo rohita* and cold water cyprinids. The study concluded that poultry droppings significantly improve fish growth performance, despite the negative allometric growth pattern.

CONCLUSION

The study concluded that the use of poultry droppings significantly enhanced fish growth due to increased dietary protein content. All the treatment have negative allometric growth pattern suggests that fish became slimmer with length increase, emphasizing the importance of optimized feeding strategies for better growth performance in *Labeo rohita* under sodic soil conditions.

Acknowledgment

The authors are thankful to UPCAR, Lucknow, for financial support to carry out this work through project. The authors are also grateful to the Hon'ble Vice Chancellor of ANDUAT, Kumarganj, Ayodhya, for their support in carrying out this experimental work.

REFERENCES

- Bagenal, T., 1978. Methods for assessment of fish production in fresh waters. No. 597.052632 M4 1978.
- Berhan, A., Birhanu, B., Misikire, T. and Abraham, A., 2019. Length-weight relationships and condition factor of Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) (Cichlidae) in Koka Reservoir, Ethiopia. *International Journal of Fisheries and Aquatic Research*, 4(1), pp.47-51.
- Dar, S.A., Najar, A.M., Balkhi, M.H., Rather, M.A. and Sharma, R., 2012. Length weight relationship and relative condition factor of *Schizopyge esocinus* (Heckel, 1838) from Jhelum River, Kashmir. *Journal of Research and Development*, 1, pp.29-36.
- Froese, F., 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22, pp.241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805>.
- Fulton, T.W., 1904. The rate of growth of fishes. 22nd Annual Report of the Fishery Board of Scotland, 3, pp.41-241.
- Gupta, S. and Sharma, A.K., 2018. Effect of organic manures on survival rate and growth performance of *Labeo rohita* (Hamilton) fingerlings. *Journal of Aquaculture Research & Development*, 9(2), p.1000461.
- Jangampalli Adi Pradeepkiran, 2019. Aquaculture role in global food security with nutritional value: a review. *Translational Animal Science*, 3(2), pp.903-910.
- Jhingran, V.G. and Pullin, R.S.V., 1985. A hatchery manual for the common, Chinese and Indian major carps. *ICLARM Studies and Reviews*, 11, p.191.

- Jhingran, V.G., 1985. Fish and Fisheries of India. 3rd Edition. Hindustan Publishing Corporation, New Delhi, 727 pp.
- Krishan, N.R. and Tarana, N. (2010). Analysis of morphometric character of *Schizothorax richardsonii* (Gray, 1832) from the Uttarkashi District of Uttarakhand State, India. Saudi Journal of Biological Sciences, 10, pp. 536–540.
- Mir, J.I., Mir, F.A., Chandra, S. and Singh, A.K., 2012. Condition factor and length-weight relationship of *Schizopyge curvifrons* (Heckel, 1838) from the Jhelum River in Kashmir, India. Journal of Applied Ichthyology, 28(2), pp.283-285.
- Singh, M.B., Prasad, L., Kumar, D., Kanaujia, S., Pathak, A., Soni, M., Dubey, S. & Kalaria, K.K., 2024. Assessing the effectiveness of cow manure and gypsum soil amendments in mitigating sodic soil conditions and their influence on the length-weight relationship of *Labeo rohita*. Uttar Pradesh Journal of Zoology, 45(7), pp.118-125.
- Gautam, A., Saroj, J., Singh, C.P., Patel, P.K., Singh, V., Kumar, R. & Lavkush, 2024. Length-weight relationship of male and female *Sperata seenghala* from River Gomti and River Saryu in Eastern Uttar Pradesh, India. Uttar Pradesh Journal of Zoology, 45(15), pp.254-260.
- Moutopoulos, D.K. and Stergiou, K.I. (2002). Length-weight relationships of fish species from the Aegean Sea (Greece). *Journal of Applied Ichthyology*, 18(3), pp. 200-203.
- Pandey, B.N., Prakash, S. and Mishra, A.K., 1998. Growth and survival of Indian major carps in sodic soil ponds. *Aquaculture*, 167(3-4), pp.275-284.
- Rasal, K.D. and Sundaray, J.K., 2020. Status of genetic and genomic approaches for delineating biological information and improving aquaculture production of farmed rohu, *Labeo rohita* (Ham, 1822). *Reviews in Aquaculture*, 12(4), pp.2466-2480.
- Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish populations. *Bulletin Fisheries Research Board of Canada*, pp.1-382.
- Subasinghe, R., Soto, D. and Jia, J., 2009. Global aquaculture and its role in sustainable development. *Reviews in Aquaculture*, 1, pp.2-9.
- Ujjania, N.C., 2012. Comparative age and growth of Indian major carp in selected water bodies of Southern Rajasthan. *Catla catla Research Journal of Recent Sciences*, 1, pp.17-22.

UNDER PEER REVIEW