

MORPHOLOGICAL FACTORS AND FORCES ACTING IN NORMAL AND SICKLED ERYTHROCYTES IN SICKLE CELL ANAEMIA: A MATHEMATICAL MODEL

Abstract

We carried out a theoretical study that considers the morphological factors and forces acting on a normal and sickled blood cell. Sickle cell disease is associated with vaso-occlusion which creates blood flow crises as a result of loss in shape memory of the normal red blood cell. Certain forces and chemical compositions acting on the cell could retain the shape memory. The problem of a second-order partial differential equation modeled was solved to get the forces and chemical composition required to restore the sickled red blood cells to their original shape. The results showed that an increase in chemical reaction increased the RBC growth by creating a stretching force that causes the sickle cell to regain elasticity with improved oxygen.

Keywords: Blood Viscosity, Stretching Force, Chemical Reaction, Sickle cell, Sickle cell disease, Shear Force,

Nomenclature:

x, y	Coordinates position of the RBC	λ	Depth constant
r	Radius of the RBC	d	RBC Diameter
A	RBC Surface area	a	RBC Thickness
μ	Blood Viscosity	θ_i	Bending angle of the RBC
μ_p	Internal Blood Viscosity	d_o	Initial length/diameter of the RBC
ρ	RBC Density	d_i	RBC increased length/diameter.
I	Energy stored in the Blood Cells	A_i	Increased RBC Membrane cross sectional area
E	Chemical reaction constant	L	Well-defined Erythrocyte Boundaries

1 Introduction.

Sickle cell disease (SCD) was discovered in the past 50 years by [1] as the first molecular disease. In the United States, about 50,000 persons are having SCD hospitalized regularly for sickle cell attacks. Sickle cell disease (SCD) was the reason an average of 75,000 persons were

hospitalized between 1989 and 1993, with an annual cost of 475 million dollars while 600 African-Americans have been diagnosed with SCD in the United States [1]. Sickle cell anemia disease (SCD) also known as a molecular disease is a genetic hematological disorder that is genetically characterized by inhomogeneous or heterogeneous cell morphology, anomalous rheology of the red blood cell, and crisis caused by vaso-occlusion or closure/blockage of the blood vessels.

The circulatory system of the human body supplies a sufficient amount of nutrients and oxygen to the body cells and carries away waste from the cells through blood transportation. The biological fluid called Blood is made up of proteins, plasma, platelets, and deformable cells. The hemoglobin in the RBC carries oxygen into the cell while in its normal state contains separate hemoglobin beads that maintain its protein quaternary structure. SCD is a genetic disorder that is autosomal recessive and created when glutamic acid is substituted with valine in the subunit of the hemoglobin gene. The substitution creates the production of abnormal hemoglobin (HbS) [2]. A single substance change (β -globin mutation) causes the hemoglobin to form long rods which changes the red blood cell into a sickle shape [3, 4]. SCD is a disorder of a group of inherited blood that produces abnormally shaped RBCs in the body that is sickled in shape. The human RBCs when passing through a narrow blood vessel, undergo repeated large elastic deformations. This flexibility of the RBCs which is large is attributed primarily to the cell membrane since there are no organelles and filaments in the cell. The RBC membrane has a two – dimensional (2D) structure, which comprises a cytoskeleton and lipid bilayer together. The lipid bilayer contains different types of cholesterol, phospholipids, sphingolipids, and essential membrane proteins, such as band-3 and glycophorin. Sickle cell has a shorter life span when compared to healthy red blood cells (RBCs). The natural life span of the RBC is 100 – 120 days; while the

sickle cell is 10 – 20 days. The shortage of red blood cells known as anemia causes fatigue, reduced breath, and delay in the growth of children. Healthy RBCs have a biconcave shape while sickle cells are rigid and sticky. These sickle cells come together and stick to the wall of the blood vessels creating an obstruction called vaso-occlusion events in the small vessels, subsequently reducing the oxygen in the blood supplied to various organs. This creates and manifests periodic seasons of pains called crises, which could last for hours or days and could result in the damage of the organs in the body such as the eyes (blindness), kidneys (renal failure), bones (avascular necrosis), lungs (pulmonary hypertension) and brain (strokes) [5]. Since the sickle cell has a short life span, the spleen then handles large numbers of RBCs and becomes enlarged and fibered. The immune function will decline, causing the body to become vulnerable to infections. In an attempt to compensate for the lost RBC, more RBCs are produced by the bone marrow which grows larger causing weakened bones. Other signs include: jaundice (yellow skin or eyes) caused as a result of increased heme destruction. Healthy RBC has a biconcave shape, with a mean diameter of 7.8 μ m. The lipid bilayer membrane has an associated cytoskeleton created by spectrin proteins interconnected by short actin filaments. The lipid bilayer is considered to be almost viscous with area-preserving membrane [6], while the elasticity of RBC is attributed to the spectrin network attached, just as the integrity of the entire RBC when subjected to severe deformations in the capillaries with a diameter of 3 μ m. [7] discussed how the blood flow in a sickle cell patient could be improved with morphological effect with other techniques for improved flow of blood [8-13] and also discussed using various mathematical models.

The major component of RBCs is the hemoglobin which is responsible for the transportation of oxygen. Adult hemoglobin (HbA) consists of four protein hemoglobin chains; 2 – α chains, 2 –

β chains and β chains. The subunit contains the HBB gene. The HBB gene undergoes several mutations, responsible for sickle cell disease with individuals having two copies of the HBB gene. The SCD is developed when both mutated-producing abnormal β – globin is copied. The two copies could be mutated differently, creating two different kinds of abnormal β - subunits. The combinations in these mutations produce different kinds of sickle cell disease (HbSs and HbSc disease; HbS/b-0 and HbS/b+ thalassemia). SCD is more severe and common and is caused by two copies of the same mutation producing mutated HbS with each copy coming from a parent. The two parents carry each copy of this mutated gene but don't show any symptoms. This pattern is called autosomal recessive inheritance [14]. At the early stage of sickle cell anemia (SCA), there is intracellular polymerization of sickle cell hemoglobin (HbS) under de-oxygenation conditions resulting in an increase in intracellular viscosity and stiffness causing great damage to the RBC membrane [15, 16, 17]. The HbS polymerization process at the molecular scale has been characterized by a double nucleation mechanism while at the cellular scale; sickle RBCs have a remarkable heterogeneity in density, rigidity, and morphology [18]. At the micro-vascular scale, the HbS polymerization creates entrapment of the RBCs in capillaries [19, 20]. The HbS-creating polymers under conditions of low oxygen are referred to as sickling (gelation). As these polymers fill the membrane, they destroy the cells into sickle shape. Apart from oxygen tension, other hemoglobin present will affect the sickling process. Normal adult hemoglobin decreases sickling whereby heterozygote parents which produce both mutated HbS and NHA do not develop the disease [21, 22]. Sickle red blood cells become depleted with potassium and then dehydrated and abnormally dense. Dense sickle cells have more severe membrane abnormalities than cells with normal hydration. A high percentage of cells in dense fractions have lost their normal phospholipid asymmetry and display on the outer membrane

leaflet, Phosphatidylserine (PS) [23]. Hemoglobin polymerization, which increases erythrocyte rigidity, is important but not the cause of an onset episode of an induced sickle cell (SS) vaso-occlusion. The lack of a relationship between the percentage of dense cells and the incidence of sickle cell crisis [21] suggests that factors other than intracellular polymerization might be involved [24]. An Increase in the adhesion of sickle cells could be an added factor first demonstrated by [25] and confirmed later in both static and dynamic systems with cultured endothelial cells obtained from human and other mammalian sources [26]. It is not been estimated in a living micro-vascular interconnection that the increase in adherence of sickle cells to the endothelium wall will contribute to vaso-occlusion. The erythrocytes in patients with SCD have heterogeneous density, morphological characteristics, and function but no demonstration for a congener donation of separate sickle cell class to adhesion and occlusive events in a perfused microvasculature. In a microcirculatory event, sickle cell vaso-occlusion, contributions from micro-vascular factors like topography, vessel wall features, and predominant rate of the wall shear, will be critical to the micro-vascular occlusion of individual sickle cell class adhesion. No direct microcirculatory study has been performed to show specific sites of sickle cell adhesion and their topographical characteristics [27]. Nitric oxide (NO) reacts at rates with oxy-hemoglobin and deoxy-hemoglobin to create nitrate plus met-hemoglobin [28, 29] and iron-nitrosyl hemoglobin respectively [30] However, the rate of NO scavenging is reduced to 1,000-fold by sequestering hemoglobin in the red cell membrane [31, 32, 33]. The amount of NO consumed by cell-free and intra-erythrocyte hemoglobin shows that when there is physical compartmentalization of hemoglobin within erythrocytes, Nitrogen Oxide (NO) produced by endothelial cells will reach concentrations within the smooth muscle that will activate Guanylate cyclase and cause vasodilation [34, 35]. This effect causes pulmonary and systemic hypertension

[36], decreases organ perfusion [37, 38], esophageal (smooth muscle), and increases mortality, which happens after infusions of stroma-free hemoglobin in animals or humans as an oxygen-carrying, artificial blood substitute. Mutations of the Heme-pocket which reduces the hemoglobin–NO affinity, increase the effect of hypertension of cell-free hemoglobin. Furthermore, hemoglobin breaks up into dimers when released into the plasma. These smaller species extravagated from the vascular lumen to positions between endothelial cells and smooth muscle, and may magnify NO scavenging [39, 40].

This sickle cell anemia/molecular disease is associated with certain hematological disorders inherited genetically leading to crises such as splenic sequestration, vaso-occlusion, hemolysis, and others. In cell morphology, the shape structure, form, and size of the cell change as a result of the de-oxygenation of the RBC while in normal rheology, high viscosity of oxygenated blood. [1] did a proposal attributing the disease to abnormal molecules of the hemoglobin within the erythrocytes. A deliquescent amino acid called glutamic is replaced by a hydrophobic amino acid called valine at the B-6 chain in the molecule of sickled hemoglobin [41]. These intracellular sickled hemoglobin molecules become polymerized in the hypoxic condition which then alters the functioning of the membrane cell and micro-spread [42]. SCD is characterized by heterogeneous and irregular cell morphologies, abnormal rheology decreased cell deformability, and finally, vaso-occlusion crises which cause morbidity and mortality in patients with SCD. The double nucleation model characterizes the polymerization process of the sickled hemoglobin molecule that is caused by homogeneous nucleation of bulk solutions of the sickled hemoglobin molecule [43, 44]. As discussed by [45], groups of heterogeneous cell density in the suspension of a sickled red blood cell are broken into four parts as a result of the concentrated intercellular mean corpuscular hemoglobin. These include sickled red blood cells 1 and 2 with average

concentrated intercellular mean corpuscular hemoglobin made up of reticulocytes and discocytes. The reticulocytes are slightly immature red blood cells with their count measuring the number of cells in the blood while the discocyte is a discoid-shaped form of the RBC. Furthermore, sickled RBCs 3 and 4 with highly concentrated intercellular mean corpuscular hemoglobin are made of irreversible sickle cells and rigid discocytes in association with rigid heterogeneous cells [46, 47, 48] and abnormal blood rheology [49, 50, 51]. According to the study done by [52, 53, 54], hydroxyurea is a treatment drug applied to patients with SCD which is targeted at the intracellular sickle hemoglobin molecule polymerization process responsible for the vaso-occlusion crisis such that the fetal hemoglobin reduces the timing of the sickling process. Several researchers have used mathematical models to study the morphological changes associated with the Red blood cell leading to its sickling. The study done by [55] shows the multiple red blood cell deformation in the capillary with immersed boundary condition used for the interaction of the RBC and the hyper-elastic model used for the RBC membrane. In the capillary flow, the viscosity was more sensitive to the shear coefficient change of the cell membrane than the bending coefficient and dilation coefficient of the surface of the membrane while increased shear coefficient caused a drop in the pressure of the flow of blood through the capillaries resulting to constant rate of flux of the RBC. The study done by [56] showed the Single RBC deformation in a micro-vessel using numerical methods on a two-dimensional spring model representing the RBC membrane with the blood cells having different kinds of motion and shape deformation. The deformation of RBC with a spherical shape in a pair beam optical stretcher was calculated by [57] using a numerical method to impute the deformed morphology of the cell spherical shape from the distribution of photonics stress past a membrane that is

cellular. In the health care of children, with the prevalence of sickle cell disease (SCD) over the past decades, this disease continues to be associated with morbidity and premature mortality, with a 25 - to 30-year loss of life expectancy [58, 59]. Bone marrow transplant is presently the known cure for SCD. It requires the replacement of the disease stem cells in the bone marrow with healthy cells from a donor usually a relative. Others include the use of vaccinations, prophylactic antibiotics, pain medication, drugs to promote the formation of hemoglobin F, and blood transfusions. Early detection and treatment for complications is crucial for a sickle cell patient. Transfusion therapy reduces stroke and other complications with an increased risk of transfusion infection, reactions, and iron overload [60]. Hydroxyurea increases the total hemoglobin concentration, reduces vaso-occlusive complications of pain and acute chest syndrome, and attenuates mortality in adults [61]. However, less than thirty percent of patients are referred to take the drug but the majority of the adults do not respond to treatment. A small and un-quantified risk of latent transformation to leukemia with long-term use remains a concern [62]. Allogeneic HCT is currently another treatment therapy for SCD with encouraging results in young patients (less than 16 years of age) with SCD, with overall event-free survival (EFS) of approximately 85% and transplant-related mortality of less than 10% [63, 64, 65]. Recently, [66] carried out a study on mathematical modelling for improved blood flow in a sickle cell anemia patient with morphological effect. Other research on blood flow and their behavioral pattern in response to certain factors and conditions was studied by Omamoke et. al [67 – 73] .

Several approaches have been followed for the mathematical description and modeling of the SCD. In this study, we are proposing a mathematical model that describes the morphology of sickle cell disease. We are considering those factors that will lead to the loss in the shape of the normal red blood shape and its recovery. We are proposing a partial differential equation (PDE)

that will combine both the forces and the chemical composition that keep the RBC in the normal shape and to what extent these forces or the chemical composition acting on the red blood will cause it to regain its shape.

2 Mathematical Formulation




Figure 1 *Diagrammatic representation of the geometry of the Red Blood Cell Membrane.*

2.1 Mathematical Modeling of the Energy acting on the RBC Membrane.

There is a relationship between the acting force on the i th membrane and the energy of the red blood cells.

$$I = Fx \quad (1)$$

From Hooke's law

$$I = \frac{1}{2} kx^2 \quad (2)$$

The RBC elastic energy stored as a result of compression and stretching

$$x_{sk} = \left(\frac{d_i - d_o}{d_o} \right)^2 \quad (3)$$

$$I_{sk} = \frac{1}{2} k_{sk} \sum_{i=1}^n \left(\frac{d_i - d_o}{d_o} \right)^2 \quad (4)$$

Where $k_{sk} = 3 \times 10^{12} Nm$ is the Hooke's spring constant for compression and stretching.

The elastic energy stored in the RBC due to bending

$$x_b = \tan^2 \left(\frac{\theta_i}{2} \right) \quad (5)$$

$$I_b = \frac{1}{2} k_b \sum_{i=1}^n \tan^2 \left(\frac{\theta_i}{2} \right) \quad (6)$$

Elastic energy stored in the RBC due to the cross sectional area

$$x_{sA} = \left(\frac{A_i - A_o}{A_o} \right)^2 \quad (7)$$

$$I_{sA} = \frac{1}{2} k_{sA} \sum_{i=1}^n \left(\frac{A_i - A_o}{A_o} \right)^2 \quad (8)$$

Where $k_{sA} = 3 \times 10^{12} Nm$ is the Hooke's spring constant for compression and stretching, $A_o = \pi(2.8 \times 10^{-6})^2 \times 0.55m^2$ is the initial equivalent RBC membrane cross sectional area.

The total energy acting on the RBC membrane is expressed as

$$I = I_{sk} + I_b + I_{sA} \quad (9)$$

The forces acting on the i th membrane is expressed as

$$F_i = \frac{dI}{dx_i} \quad (10)$$

$$I = F_i x_i \quad (11)$$

This implies that the energy acting on the RBC membrane

$$I_{sk} + I_b + I_{sA} = F_i x_i \quad (12)$$

2.2 Mathematical Modeling of the Shape Geometry of the RBC Membrane.

$$x = r \cos \theta \quad (13)$$

$$y = r \sin \theta \quad (14)$$

$$x^2 + y^2 = r^2 \quad (15)$$

$$\cos^2 \theta + \sin^2 \theta = 1 \quad (16)$$

$$\cos^2 \theta + \sin^2 \theta = \frac{x^2 + y^2}{r^2} \quad (17)$$

$$L = \frac{e^{\frac{x^2 + y^2}{2r^2}}}{2\pi r^2} [74] \quad (18)$$

3 Governing equation

The governing equation in non-dimensional form for the chemical reaction effect on the RBC morphology and shape recovery is expressed below as

$$\rho A \mu \frac{\partial^2 \varphi}{\partial x^2} + EI \frac{\partial^4 \varphi}{\partial x^4} = 0 \quad (19)$$

The blood viscosity in relation to blood hematocrit non-dimensional form is expressed below as

$$\mu = \mu_p (1 + 2.5H) \quad (20)$$

$$E = \frac{\sigma}{\varepsilon} \quad (21)$$

The blood density in relation to cross sectional area of the blood cell in non-dimensional form is expressed below as

$$\rho = \frac{Q}{A} \quad (22)$$

The energy acting on the red blood cell RBC due to bending, cross sectional area, stretching and compression non-dimensional form is expressed as

$$I = Fx \quad (23)$$

The governing equation for the depth of the sickling of the RBC and shape recovery is expressed below as

$$\frac{\partial^2 \delta}{\partial t^2} + \lambda x \frac{\partial^2 \delta}{\partial x^2} = 0 \quad (24)$$

$$\lambda = \frac{\sigma Fx}{Q\mu\varepsilon} \quad (25)$$

4 First Method of Solution

The separation of variable method is used to solve the partial differential equation that models the dip, shear force and chemical composition for the memory and shape recovery.

$$\varphi = X_1 T_1 \quad (26)$$

$$\delta = X_2 T_2 \quad (27)$$

Boundary Conditions for the Erythrocytes

$$\varphi(0, t) = 0; \delta(0, t) = 0 \quad (28)$$

$$\varphi(L, t) = 0; \delta(L, t) = 0 \quad (29)$$

$$\varphi(x, 0) = 1; \delta(x, 0) = 1 \quad (30)$$

$$\varphi(x, \infty) = 0; \delta(x, \infty) = 0 \quad (31)$$

4.1 Solution to the Governing equation

$$T_1 = B_1 e^{\left(\frac{k}{\sqrt{\rho A}}\right)t} + B_2 e^{-\left(\frac{k}{\sqrt{\rho A}}\right)t} \quad (32)$$

$$X_1 = B_3 \cos\left(\sqrt[4]{\frac{k^2}{EI}}x\right) + B_4 \sin\left(\sqrt[4]{\frac{k^2}{EI}}x\right) \quad (33)$$

Substituting equation (32) and (33) into equation (26), the chemical reaction effect on RBC is expressed as

$$\varphi(x, t) = \left(B_1 e^{\left(\frac{k}{\sqrt{\rho A}}\right)t} + B_2 e^{-\left(\frac{k}{\sqrt{\rho A}}\right)t} \right) \left[(B_3) \cos\left(\sqrt[4]{\frac{k^2}{EI}}x\right) + (B_4) \sin\left(\sqrt[4]{\frac{k^2}{EI}}x\right) \right] \quad (34)$$

Applying the boundary conditions on equation (19), the chemical reaction effect on RBC growth is

$$\varphi(x, t) = \sum_{n=0}^{\infty} \left[\sin \frac{n\pi}{L} x \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right) \left(e^{-\left(\frac{n\pi}{L}\right)^2 \left(\frac{EI}{\rho A}\right)t} \right) \right] \quad (35)$$

$$\text{where } B_1 = 0, B_3 = 0, B_2 = 1, B_4 = \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right), L = \frac{e^{\frac{x^2+y^2}{2r^2}}}{2\pi r^2} \quad [74]$$

$$T_2 = C_1 e^{kt} + C_2 e^{-kt} \quad (36)$$

$$X_2 = C_3 \cos\left(\sqrt[4]{\frac{k^2}{\lambda}}x\right) + C_4 \sin\left(\sqrt[4]{\frac{k^2}{\lambda}}x\right) \quad (37)$$

Substituting equation (36) and (37) into equation (27), the depth/growth of RBC is expressed as

$$\delta(x, t) = (C_1 e^{kt} + C_2 e^{-kt}) \left[(C_3) \cos\left(\sqrt[4]{\frac{k^2}{\lambda}}x\right) + (C_4) \sin\left(\sqrt[4]{\frac{k^2}{\lambda}}x\right) \right] \quad (38)$$

Applying the boundary conditions on equation (24) the solution for the depth of RBC is

$$\delta(x, t) = \sum_{n=0}^{\infty} \left[\sin \frac{n\pi}{L} x \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right) \left(e^{-\left(\frac{n\pi}{L}\right)^2 (\sqrt{\lambda})t} \right) \right] \quad (39)$$

$$\text{Where } C_1 = 0, C_3 = 0, C_2 = 1, C_4 = \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right), L = \frac{e^{\frac{x^2+y^2}{2r^2}}}{2\pi r^2} \quad [74]$$

5 Second Method of Solution

From equation (19), we assume the solution to take the form

$$\varphi = \varphi_0 e^{i\omega t} \quad (40)$$

The expression for equation (19) will be given as

$$\rho A \mu \omega^2 \varphi_0 e^{i\omega t} + EI \frac{\partial^4 \varphi_0}{\partial x^4} e^{i\omega t} = 0 \quad (41)$$

Equation (41) is simplified and expressed as

$$\frac{\partial^4 \varphi_0}{\partial x^4} + \beta^4 \varphi_0 = 0 \quad (42)$$

$$\text{Where } \beta = \left(\frac{\rho A \mu \omega^2}{EI} \right)^{\frac{1}{4}} = \left(\frac{\mu k}{EI} \right)^{\frac{1}{4}} \quad (43)$$

$$\omega = \left(\frac{n\pi}{l} \right)^2 \sqrt{\frac{EI}{\rho A \mu}} \quad (44)$$

$$\varphi_0 = A_1 \cos(\beta y) + B_1 \sin(\beta y) + A_2 \cosh(\beta y) + B_2 \sinh(\beta y) \quad (45)$$

$$\varphi_0(y, t) = [A_1 \cos(\beta y) + B_1 \sin(\beta y) + A_2 \cosh(\beta y) + B_2 \sinh(\beta y)] e^{i\omega t} \quad (46)$$

Applying the boundary conditions in (28-29),

$$\varphi_0(l, t) = [A_1 \cos(\beta l) + B_1 \sin(\beta l) + A_2 \cosh(\beta l) + B_2 \sinh(\beta l)] e^{i\omega t} \quad (47)$$

The equation of the sickle cell growth model on the deformed RBC becomes

$$\varphi_0(x, t) = \sum_{n=0}^{\infty} \left[\sin \frac{n\pi}{L} x \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right) \left(e^{-\left(\frac{n\pi}{L} \right)^2 \left(\sqrt{\frac{EI}{\rho A}} \right) t} \right) \right] \quad (48)$$

$$\text{where } B_1 = 0, B_3 = 0, B_2 = 1, B_4 = \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right), L = \frac{x^2 + y^2}{2\pi r^2} \quad [74]$$

Applying the same method of solution on equation (24), the equation of the sickle cell depth/growth model on the deformed RBC is expressed as

$$\delta(x, t) = \sum_{n=0}^{\infty} \left[\sin \frac{n\pi}{L} x \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right) \left(e^{-\left(\frac{n\pi}{L} \right)^2 (\sqrt{\lambda}) t} \right) \right] \quad (49)$$

Where $C_1 = 0, C_3 = 0, C_2 = 1, C_4 = \left(\frac{2L}{n\pi} \left(1 - \cos \frac{n\pi}{L} \right) \right), L = \frac{x^2 + y^2}{2\pi r^2}$ [74]

6 Graphical Results and Discussion.




Figure 2 Changes in shape geometry with effect on the deformed RBC




Figure 3 Changes in cross sectional area with effect on the deformed RBC

From Figure 2.0 it was observed that an increase in the geometry of the RBC increases the RBC growth reducing the deformability and resulting in the restoration of the RBC. This is a result of the increased chemical reaction effect on the sickled RBC. Furthermore, the chemical reaction creates a stretching force which in turn affects the viscosity of the blood. The stretching force forces the sickle cell to regain its elasticity, improves the oxygen level of the blood, and reduces its stickiness helping it to recover its RBC shape depending on the amount of force applied which will affect the flow of the RBC. The increase in the shear force reduces the viscosity of the blood enabling improved flow with the shear force creating a correction on the sickled RBC reducing rigidity, stickiness, and breaking the frictional forces which afterwards help to reduce viscosity.

From Figure 3.0 it is observed that an increase in the cross-section area of the RBC causes an increase in the depth of the RBC due to the change in the radius of the RBC caused by the stretching force acting on the RBC. The long rods created in the hemoglobin to form a sickle shape are broken with the external forces acting on the sickle cell causing the shape to regain its

memory and recover its original shape. The sickling is reduced as a result of the parent heterozygote producing both mutated Hbs and NHA which does not develop the sickle cell disease causing it to become normal adult hemoglobin

7 Conclusion

The following conclusion was obtained from the research

1. An increase in chemical reaction increases the geometry of the RBC by increasing the stretching force, which in turn increases the growth of the RBC and reduces its deformability.
2. An increased cross-section area of the RBC increases the RBC depth due to the change in the RBC radius caused by the stretching force which breaks the long rods created in the hemoglobin which forms a sickle shape. This creates improved oxygenation of the blood and reduces the stickiness of the cell.

References

- [1] L. Paulings, H. A. Itano, S. J. Singer, I. C. Wells, Sickle cell anemia, a molecular disease Science 110 (1949) 543 – 548.
- [2] L. Gravitz, S. Pincock, Sickle-cell disease, Nature 515 (2014).
- [3] X. W. Tao, X. S. Wang, B. Cohen, G. Hongya 2010. Molecular Modeling of normal and sickle cell hemoglobin, New Jersey Institute of Technology, Newark, New Jersey 07102, USA 2McCoy School of Engineering, College of Science and Mathematics, Midwestern State University, Wichita Falls, Texas 76308, US.
- [4] B. R. Brooks, R. E. Bruccoleri, B. D. Olafson, D. J. States, S. Swaminathan, M. Karplus, CHARMM: A program for macromolecular energy, minimization, and dynamics calculations, J. Comp. Chem., vol. 4(2) pp. 187–217.
- [5] H. Lei, G. E. Karniadakis. Probing vaso-occlusion phenomena in sickle cell anemia via mesoscopic simulations, Proc Natl Acad Sci USA 110 11 326-330.

- [6] Y. C. Fung. *Biomechanics: Mechanical Properties of Living Tissues*, 2nd ed. Springer-Verlag, New York, 1993.
- [7] Omamoke, E., Funakpo, I., Olugbenro, O., Izah, S. C., Keneke, E. D., & Wilcox, B. K. (2023). Mathematical Modelling for Improved Blood Flow in a Sickle Cell Anaemia Patient with Morphological Effect, *American Journal of Applied Mathematics*, 11 (3), 40-51.
- [8] Omamoke, E. & Amos, E. (2023). Treatment and Slip Effect on MHD Blood Flow through a stenotic Artery: A mathematical Model, *Asian Research Journal of Mathematics*, 19 (6), 61-76.
- [9] Amos, E., Omamoke, E., Nwaigwe, C. (2022). MHD Pulsatile Blood Flow through an Inclined Stenosed Artery with Body Acceleration and Slip Effects, *International Journal of Theoretical and Applied Mathematics*, 8 (1), 1-13.
- [10] Omamoke, E. & Amos, E. (2022). Slip and Pulsatile Mhd Blood Flow Through An Inclined Stenosed Artery With Body Acceleration And Heat Source Effects, *IOSR Journal of Mathematics*, 18 (1), 1-13.
- [11] Amos, E., Omamoke, E., Nwaigwe, C. (2022). Chemical Reaction, Heat Source and Slip Effects on MHD Pulsatory Blood Flowing Past an Inclined Stenosed Artery Influenced by Body Acceleration, *International Journal of Mathematics Trend and Technology*, 68 (1), 1-23.
- [12] Omamoke, E., Amos, E. & Efere, J. (2020). Impact of Thermal Radiation and Heat Source on MHD blood flow with an inclined magnetic field in treating tumor and low blood pressure. *Asian Research Journal of Mathematics*, 16(9), 77-87, ISSN 2456-4778.
- [13] Omamoke, E. & Amos, E. (2020). Chemical Reaction, Radiation and Heat Source Effects on Unsteady MHD Blood Flow Over a Horizontal Porous Surface in the Presence of an Inclined Magnetic Field. *International Journal of Scientific and Engineering Research*, ISSN 2229-5518, Volume 11, Issue 4.
- [14] A. Perumbeti, T. Higashimoto, F. Urbinati, R. Franco, H. J. Meiselman, W. David P. Malik. A novel human gamma-globin gene vector for genetic correction of sickle cell anemia in a humanized sickle mouse model: critical determinants for successful correction, *BLOOD* 114 (2009) 6.
- [15] X. Li, H. Lei. (2015). Multiscale Modeling of Sickle Cell Anemia. e National Institutes of Health (NIH) under grants U01HL114476 and U01HL116323.
- [16] H. F. Bunn. Pathogenesis and treatment of sickle cell disease, *N. Engl. J. Med.* 337 (1997) 762–769.
- [17] G. A. Barabino, M. O. Platt, D. K. Kaul. Sickle Cell Biomechanics, *Annu Rev Biomed Eng.* 12 (2010) 345-367.
- [18] X. Li, M. Dao, G. Lykotrafitis, G. E. Karniadakis. Bio-mechanics and biorheology of red blood cells in sickle cell anemia. *J. Biomech.* PMC 2017.
- [19] S. C. Liu, L. H. Derick, Zhai, S. J. Palek. Uncoupling of the spectrin-based skeleton from the lipid bilayer in sickled red cells, *Science* 252 (1991) 574–576

- [20] D. K. Kaul, M. E. Fabry. In vivo studies of sickle red blood cells, *Microcirculation* 11 (2004) 153–165.
- [21] B. Modell, M. Darlison. Epidemiology of hemoglobin disorders and derived service indicators, *Bull World Health Organ*, 86 (2008) 480–487.
- [22] D. K. Kaul, M. E. Fabry, R. L. Nagel. Micro-vascular sites and characteristics of sickle cell adhesion to vascular endothelium in shear flow conditions: Pathophysiological implications, *Proc Natl Acad Sci USA* 86 (1989) 3356–3360
- [23] R. P. Hebbel. Beyond hemoglobin polymerization: the red blood cell membrane and sickle disease pathophysiology, *Blood*. 77 (1991) 214.
- [24] M. E. Fabry, R. L. Nagel. *Blood Cells* 8 (1989) 9-15.
- [25] R. P. Hebbel, M. A. Boogaerts, J. W. Eaton, M. H. Steinberg. *N. Engl. J. Med.* 302 (1980) 992-995.
- [26] D. K. Kaul. in *Genetically Abnormal Red Cells*, ed. Nagel, R. L. (CRC, Boca Raton, FL), 2 (1988) 161-176.
- [27] D. K. Kaul, R. L. Nagel, S. Baez. *Microvasc. Res.* 26 (1983) 170 - 181.
- [28] R. F. Eich. Mechanism of NO-induced oxidation of myoglobin and hemoglobin, *Biochemistry* 35 (1996) 6976–6983.
- [29] S. Herold, M. Exner, T. Nauser. Kinetic and mechanistic studies of the NO-mediated oxidation of oxymyoglobin and oxyhemoglobin, *Biochemistry* 40 (2001) 3385–3395.
- [30] Q. Gibson, F. J. W. Roughton. The kinetics and equilibria of the reactions of nitric oxide with sheep hemoglobin, *J. Physiol. London* 136 (1957) 507–526.
- [31] R. Cassoly, Q. Gibson. Conformation, co-operativity and ligand binding in human hemoglobin, *J. Mol. Biol.* 91 (1975) 301–313.
- [32] E. Carlsen, J. H. Comroe. The rate of uptake of carbon monoxide and of nitric oxide by normal human erythrocytes and experimentally produced Spherocytes, *J. Gen. Physiol.* 42 (1958) 83–107.
- [33] X. Liu. Diffusion-limited reaction of free nitric oxide with erythrocytes, *J. Biol. Chem.* 273 (1998) 18709–18713.
- [34] J. R. Lancaster. A tutorial on the diffusibility and reactivity of free nitric oxide, *Nitric Oxide* 1 (1997) 18–30.
- [35] M. W. Vaughn, L. Kuo, J. C. Liao. Effective diffusion distances of nitric oxide in the microcirculation, *Am. J. Physiol.* 274 (1998) H1705–H1714.
- [36] J. R. Hess, V. W. MacDonald, W. W. Brinkley. Systemic and pulmonary hypertension after resuscitation with cell-free hemoglobin, *J. Appl. Physiol.* 74 (1993) 1769–1778.

- [37] L. F. de-Figueiredo. Pulmonary hypertension and systemic vasoconstriction may offset the benefits of a cellular hemoglobin blood substitutes, *J. Trauma* 42 (1997) 847–856.
- [38] J. A. Ulatowski. Regional blood flow alterations after bovine fumaryl β crosslinked hemoglobin transfusion and nitric oxide synthase inhibition, *Crit. Care Med.* 24 (1996) 558–565.
- [39] D. H. Doherty. Rate of reaction with nitric oxide determines the hypertensive effect of cell-free hemoglobin, *Nature Biotechnology* 16 (1998) 672–676.
- [40] U. Pohl, D. Lamontagne. Impaired tissue perfusion after inhibition of endothelium-derived nitric oxide. *Basic Res. Cardiology.* 86 (1991) 97–105.
- [41] V. M. Ingram. Abnormal human hemoglobin: The comparison of normal human and sickle-cell hemoglobin by fingerprinting, *Biochim. Biophys. Acta.* 28 (1958) 539 – 545.
- [42] L. Huan, E. K. Goerge. Multiscale Modeling of Sickle Cell Anemia, *Modeling the Heart and the Circulatory System* (2015) 119 – 156.
- [43] F. A. Ferrone, J. Hofrichter, W. A. Eaton. Kinetics of sickle hemoglobin polymerization: studies using temperature-jump and laser photolysis techniques, *Journal of Molecular Biology* 183(4) (1985) 591 – 610.
- [44] F. A. Ferrone, J. Hofrichter, W. A. Eaton. Kinetics of sickle hemoglobin polymerization I. a double nucleation mechanism, *Journal of Molecular Biology*, 183(4) (1985) 611 – 631.
- [45] D. K. Kaul, M. E. Fabry, P. Windisch, S. Baez, R. L. Nagel. Erythrocytes in sickle cell anemia are heterogeneous in their rheological and hemodynamic characteristics, *Journal of clinical investigation* 72(1) (1983) 22 – 31.
- [46] M. R. Clark, N. Mohandas, S. B. Shoet. Deformability of deoxygenated irreversibly sickled cells, *Journal of Clinical investigation*, 65 (1980) 189 – 196.
- [47] E. Evans, N. Mohandas. Membrane associated sickle hemoglobin: a major determinant of sickle erythrocytes rigidity, *Blood* 70 (1987) 1443 – 1449.
- [48] T. Itoh, S. Chien, S. Usami.. Effects of hemoglobin concentration on deformability of individual sickle cells after de-oxygenation, *Blood* 85 (1995) 2245 – 2253.
- [49] J. M. Higgins, D. T. Eddington, S. N. Bhatia, L. Mahadevan. Sickle cell Vaso-occlusion and rescue in a microfluidic device. *Proceedings of the National Academy of Sciences* 104(51) (2007) 20,496 – 20,500.
- [50] D.K. Kaul, X. Liu. Rate of Deoxygenation Modulates Rheological behavior of Sickle Red Blood Cells at a given Mean Corpuscular Hemoglobin Concentration, *Clinical Hemorheology and Microcirculation* 21 (1999) 125-135.
- [51] D. K. Kaul, H. Xue. Rate of Deoxygenation and Rheologic behavior of Blood in Sickle Cell Anemia, *Blood* 77 (1991) 1353 – 1361.

- [52] K. Bridges, G. Barabino, C. Brugnara, M. Cho, G. Christoph, G. Dover, B. Ewestein, D. Golan, C. Guttman, J. Hofrichter, R. Mulkern, B. Zhang, W. Eaton. A Multiparameter Analysis of Sickle Erythrocytes in Patients undergoing Hydroxyurea Therapy, *Blood* 88 (1996) 4701 – 4710.
- [53] W. A. Eaton, J. Hofrichter. The biophysics of sickle cell hydroxyurea therapy, *Science* 268 (1995) 1142 – 1143.
- [54] O. S. Platt, S. H. Orkin, G. Dover, G. P. Beardsley, B. Miller, D. G. Nathan. Hydroxyurea enhances fetal hemoglobin production in sickle cell anemia, *J. Clin. Invest.* 74 (1984) 652 – 656.
- [55] G. Xiaobo, S. Kazuyasu, T. Shu, M. Yoichiro. The deformation Behaviour of Multiple Red Blood Cells in a Capillary, *Journal of Biomechanical Engineering*, 131 (2009).
- [56] H. P. G. Nayanajith, C. S. Suvash. Deformation of a Single Red Blood Cell in a Microvessel, *ANZIAM J.* 55 (2013) 64 – 79.
- [57] B. B. Paul, S. Yunlong, C. Yin-Quan, C. Arthur. Calculation of Spherical Red Blood Cell Deformation in a Dual- beam Optical Stretcher, *Optics Express*, 15(24) (2007).
- [58] O. S. Platt, D. J. Brambilla, W. F. Rosse. Mortality in sickle cell disease: life expectancy and risk factors for early death, *N Engl J Med.* 330 (23) 1994, 1639-1644.
- [59] D. R. Powars, L. S. Chan, A. Hiti, E. Ramicone, C. Johnson. Outcome of sickle cell anemia: a 4-decade observational study of 1056 patients, *Medicine (Baltimore)* 84 (2005) 363-376.
- [60] R. W. Hagar, E. P. Vichinsky. Major changes in SCD, *Adv Pediatr*, 47 (2000) 249-272.
- [61] M. H. Steinberg. Hydroxyurea treatment for SCD, *Scientific World Journal*, 2 (2002) 1706-1728.
- [62] O. S. Platt. Hydroxyurea for the treatment of sickle cell anemia, *N. Engl. J. Med.* 358 (2008) 1362-1369.
- [63] M. R. Abboud, S. M. Jackson, J. Barredo, J. Beatty, J. Laver. Bone marrow transplantation for sickle cell anemia, *Am J Pediatr Hematol Oncol* 16 (1994) 86-89.
- [64] M. C. Walters, M. Patience, W. Leisenring. Barriers to bone marrow transplantation for sickle cell anemia, *Biol Blood Marrow Transplant* 2(2) (1996) 100-104
- [65] M. C. Walters, M. Patience, W. Leisenring. Stable mixed hematopoietic chimerism after bone marrow transplantation for sickle cell anemia, *Biol Blood Marrow Transplant*, 7(12) (2001) 665-673.
- [66] Omamoke Ekakitie, Funakpo Isaac, Olugbenro Osinowo, Sylvester Chibueze Izah, Keneke Edwin Dauseye, et al. (2023). Mathematical Modelling for Improved Blood Flow in a Sickle Cell Anaemia Patient with Morphological Effect. *American Journal of Applied Mathematics*, 11(3), 40-51.

- [67] Omamoke, E. & Amos, E. (2023). Treatment and Slip Effect on MHD Blood Flow through a stenotic Artery: A mathematical Model, *Asian Research Journal of Mathematics*, 19 (6), 61-76
- [68] Amos, E., Omamoke, E., Nwaigwe, C. (2022). MHD Pulsatile Blood Flow through an Inclined Stenosed Artery with Body Acceleration and Slip Effects, *International Journal of Theoretical and Applied Mathematics*, 8 (1), 1-13.
- [69] Omamoke, E. & Amos, E. (2022). Slip and Pulsatile Mhd Blood Flow Through An Inclined Stenosed Artery With Body Acceleration And Heat Source Effects, *IOSR Journal of Mathematics*, 18 (1), 1-13.
- [70] Amos, E., Omamoke, E., Nwaigwe, C. (2022). Chemical Reaction, Heat Source and Slip Effects on MHD Pulsatory Blood Flowing Past an Inclined Stenosed Artery Influenced by Body Acceleration, *International Journal of Mathematics Trend and Technology*, 68 (1), 1-23.
- [71] Omamoke, E., Amos, E. & Isaac, F. (2021). Chemical Reaction, Heat Source and Soret Effects on MHD Pulsatile Flow over an Inclined Porous Plate with Thermal Radiation. *Journal of Association Mathematical Physics, JNAMP Vol. 61, page 77- 90.*
- [72] Omamoke, E., Amos, E. & Efere, J. (2020). Impact of Thermal Radiation and Heat Source on MHD blood flow with an inclined magnetic field in treating tumor and low blood pressure. *Asian Research Journal of Mathematics*, 16(9), 77-87, ISSN 2456-4778.
- [73] Omamoke, E. & Amos, E. (2020). Chemical Reaction, Radiation and Heat Source Effects on Unsteady MHD Blood Flow Over a Horizontal Porous Surface in the Presence of an Inclined Magnetic Field. *International Journal of Scientific and Engineering Research, ISSN 2229-5518, Volume 11, Issue 4.*
- [74] P. J. Vivek, L. Alfredo, P. J. Vinay, M. Carlos, T. W. Alexander, O. Daniel, Y. Ozlem, C. Pedro. Numerical Model for the Determination of Erythrocyte Mechanical Properties and Wall Shear Stress in Vivo from Intra-vital Microscopy, *Frontiers in Physiology* 10 (2019) 1562.