

Rootstocks ~~influenecs~~influence degree days requirement for producing quality wine of Cabernet Sauvignon grown in Pune region

ABSTRACT

The experiment was conducted to study the ~~rootstocks~~rootstocks' effect on the growth, quality, yield and -sensory evaluation of wine made from 'Cabernet Sauvignon' grapes. Among the rootstocks, pruning weight was significantly higher in 110R grafted vine. Days to bud sprout and days to veraison was early in Gravasec grafted rootstocks while days to harvest in Gravasec and SO4 rootstocks. Cabernet Sauvignon vines grafted on 110R rootstock recorded significantly higher yield than other rootstocks studied. Wine composition like volatile acid, total acids and ethanol was higher in 140RU; malic acids in 1103P and colour intensity in SO4 grafted vines. The wine sensory attributes were also positively influenced using different rootstocks, wine prepared from Cabernet Sauvignon grapes grafted on 1103P recorded the highest overall wine quality which was followed by Fercal rootstock.

Keywords: Cabernet Sauvignon, grape, rootstock, growth, yield, quality

1. Introduction

Grape (*Vitis vinifera* L.) is one of the important fruit crops widely cultivated in different regions (of what??). Though the grape ~~is~~ originated from temperate regions, it ~~is performing~~performs well under tropical ~~climate~~climates in the country where it grows as an evergreen vine without undergoing dormancy. In India, it is grown on an area of about 1.71 lakh hectares with an average ~~productivity~~production of 37.81 lakh metric ~~tonns~~tons and productivity of 22.10 MT/ha of grapes annually [1]. However, only about 2% of the total production of grapes is being used for juice and wine purpose [2]. The decline in yield due to the problems associated with soil and water salinity, chlorides in irrigation water and excess levels of sodium in soil and shortage of irrigation water in Maharashtra state alerted the situation. Since then, the use of rootstock to maintain the productivity of grapes under adverse ~~situation~~situations has gained ~~the~~ popularity. The choice of proper rootstock is becoming difficult due to availability of large number of rootstocks [3]. ~~Performance~~The performance of rootstock is different under different ~~condition~~conditions; hence it is necessary to evaluate rootstock best suited to the environment [4]. Rootstocks ~~isare~~ mostly utilized to make a better scion performance under ~~different conditions of various~~ cultivation condition in viticulture. However, its performance

mainly depends on scion cultivars, soil type and climatic conditions [5]. The selection of appropriate rootstock for the scion plays an important role in the growth, fruitfulness, and yield of grapevine scions [6].

In India, it is necessary to use appropriate rootstocks in grapes for profitable production against major abiotic stresses such as soil and water salinity, water scarcity, etc. [7]. There is no major issue of the effect of rootstocks on the fruit composition in table grapes except for berry size, TSS, acidity etc., but it is a major concern in wine grapes to produce good quality wines. Due to a lack of knowledge about the influence of rootstocks on fruit composition and other quality parameters (soluble solids, organic acids, and pH), most of the wine cultivars were grafted on Dogridge which is the choice for table grapes in India [8]. But many scientists concluded that Dogridge rootstock for wine grapes accumulates significantly higher concentrations of potassium in berries which ~~deteriorated~~-deteriorates the quality of wines in terms of high pH, poor ~~colour~~-color stability, and high organic acids when grown in warm regions [9,10].

As climate change continues to cause warmer and drier weather, it is having a significant impact on the growth, fruit composition, and early harvest. Grapevine yield and quality are heavily influenced by climate conditions and depend on complex interactions between temperatures, water availability, plant material and viticultural practices. Many growers have been able to improve yield and quality by using plant material and viticultural practices suited to their local climate. To adapt to higher temperatures, growers may need to change their planting material and modify their viticultural practices, such as adjusting their harvest dates [11].

Wine is one of the most popular beverages prepared from grapes through fermentation under the controlled conditions. It comprises phenolic compounds mainly classified ~~into~~-as flavonoids and non-flavonoids [12]. These compounds are considered to have antioxidant, anti-cancer and anti-inflammatory properties [13,14] and they are also responsible for some of the sensory properties like colour, aroma, flavor, bitterness and astringency in grapes and wine [15]. It is thus necessary to investigate how the rootstock is suitable for a given cultivar and location that affects the plant development, yield, and quality. The present study was therefore conducted to study the impact of rootstock on yield and quality in Cabernet Sauvignon wine grape variety.

2. Materials and Methods

2.1 Vineyard, Experiment Design, and Vine Management

The study was carried out over three years (2014-15, 2015-16 and 2016-17) in an experimental vineyard located in the ICAR-National Research Centre for Grapes, Pune, India (18.32° N

latitude, 73.85° E longitude and 559 m altitude). The cultivar 'Cabernet Sauvignon' grafted on eight different rootstocks (Fercal, Dogridge, SO4, 110R, Gravasec, 1103P, 101-14MGT, 140RU) were evaluated in a randomized block design with three replicates represented by five vines per replication. The grape vineyard was four years old, trained onto a mini-Y system of trellises spaced at 2.4×1.2 m accommodating about 3400 vines per hectare.

2.2 Determinations of berry physiochemical parameter

Pruning biomass ~~were-was~~ measured after fruit pruning (forward pruning) for selected vines and the average was calculated. Days taken for sprouting were recorded from the date of pruning to sprouting of first bud. The first sprouted bud with fully expanded leaf was taken as an indicator to calculate the days taken for sprouting. Days to veraison and days to harvest was calculated from date of fruit pruning for individual vines.

Harvesting was done about 145 days after forward pruning during the month of March. At harvest, soluble solids (Brix), titratable acidity (g L⁻¹ tartaric acid) and pH were measured using the juice of pressed berries (100 berries per treatment) collected. Soluble solids (°Brix) were determined using a handheld refractometer (ERMA, Japan) with temperature compensated to 20°C. The pH of pure juice of every sample was determined using a pH meter. Titratable acidity was determined by titration with 0.1 N NaOH to a phenolphthalein ~~end-point~~endpoint and expressed as g L⁻¹ [16]. Also, five vines were selected randomly from each rootstock. Juice recovery (%) was recorded by crushing 1 kg of berries. The observations on the number of berries/bunches, 100 berry weight (g), average bunch weight, yield per vine were recorded at the harvesting stage.

2.3 Growing Degree Days

Degree days requirement for maturity of Cabernet Sauvignon grapes was calculated, using following formula.

$$\text{Degree Days} = (\text{Maximum Temperature} - \text{Minimum Temperature})/2 - 10$$

2.4 Wine preparation and sensory evaluation

After harvest, grape bunches were washed with water to remove rotten and green berries and the de-stemming was done manually. The grape berries, were passed through a stainless-steel presser to prepare must. The must was inoculated with commercial yeast, culture (*Saccharomyces cerevisiae*) with viable cell count, *i.e.* 1.06 × 10⁸ mL⁻¹. Fermentation was carried out in 50 L capacity stainless steel tanks at 20-22 °C. During fermentation process, fermenting material was mixed, twice every day. The fermentation was completed by 13 days. The material like skin, seed and yeast lees were separated from the finished wine. The prepared

wine bottled properly and stored at 1-2 °C for further analysis. The sensory analysis of Cabernet Sauvignon wines was performed and overall quality at 1 to 10 rating scale, the means was calculated and data were expressed in graphical chart. The wine quality parameters (total acid, malic acid, pH, volatile acid, and ethanol) were recorded on Oeno Foss (FTIR based analyzer). The wine samples were drowned into microtube and centrifuged at 500 rpm for 5 minutes and the readings were recorded.

2.5 Statistical analysis

The experiment was conducted in Randomized Block Design (RBD) consisting of eight treatments as rootstocks which were replicated three times. Statistical analysis of data collected during studies was carried out by standard method of analysis of variance as described by [17] and data was analysed using Statistical Analysis System (SAS) software version 9.3. The standard error of mean ($S E_{m\pm}$) was worked out and the critical difference at 5 per cent level of significance was calculated wherever the results were found significant.

3. Results and Discussion

3.1 Growth parameters

Rootstocks non-significantly influences vigor of Cabernet Sauvignon, as indicated by values of pruning biomass (Table 1). The maximum pruning weight ~~were-was~~ recorded on 110R (1.00 kg) while minimum pruning weight ~~were-was~~ recorded on SO4 (0.62 kg). The difference in the pruned biomass among the rootstocks may be due to the difference in ~~the vigor of vinevine~~ vigor due to higher photo assimilation resulting in higher carbohydrate deposition [18]. The higher pruning weight might be attributed due to more efficiency for the absorption of water and minerals from the root system of rootstocks [19]. The previous study also reported that rootstocks influenced the vigour of the scion. Rootstock's effects on pruning weight may vary among different scion cultivars [20,10].

Days to bud sprouts was early on Gravasec among all the rootstocks (8.45 days) which was followed by SO4 (0.22) and Fercal (9.78) while the rootstock 140RU was late to sprout (11.67 days). Availability of stored food material that has helped to supply for early bud sprout, cultural practices, and temperature variation might be a reason for ~~variation-difference~~ in time taken for bud burst [21,22]. These results are in accordance with the earlier reports on the influence of rootstocks on bud break. [23]~~(No is not enough here write the name of author??)~~ reported that vines grafted on 110R took less time to achieve the maximum percentage of bud break as compared to those grafted on Freedom rootstocks.

The minimum days for veraison and harvest were recorded in Gravesec and SO4 rootstock grafted vines whereas, maximum days were recorded in Fercal and 110R grafted vines. The

interaction between stock and scion affects the root physiology which help in the proper uptake of water and minerals might be resulting into early veraison and harvest. Bunch load is also an important factor for early harvesting Cabernet Sauvignon grafted on 110R rootstock showed higher crop loads among the rootstocks which was responsible for late harvesting of its vines. Similar results were recorded by [24].

Table 1. Vegetative parameters in relation to different rootstocks in Cabernet Sauvignon (pooled means for three years)

3.2 Yield parameters

Rootstocks	Pruning weight (kg)	Days to bud sprouts	Days to veraison	Days to harvest
Fercal	0.85	9.78 ^d	106.56 ^a	149.00 ^a
Dogridge	0.64	11.44 ^{ab}	105.33 ^a	144.22 ^b
SO4	0.62	9.22 ^d	102.00 ^b	139.89 ^c
110R	1.00	10.78 ^{bc}	107.00 ^a	148.67 ^a
Gravasec	0.70	8.45 ^e	101.78 ^b	139.89 ^c
1103P	0.88	10.66 ^c	105.55 ^a	144.11 ^b
101.14MGT	0.70	11.66 ^a	105.44 ^a	144.11 ^b
140RU	0.76	11.67 ^a	105.44 ^a	144.1 ^b
SEm±	0.11	0.24	0.56	0.56
CD at 5%	0.33	0.72	1.70	1.70
<u>SigP value</u>	NS	**	**	**

(What do you mean by ** and NS, report cv% on table)

Number of bunches per vine, average bunch weight and total yield were significantly affected by rootstocks (Table 2). The maximum number of bunches were recorded on Cabernet Sauvignon grafted on 1103P (71.11) which was at par with SO4 (70.00) and 110R (68.45) rootstocks while the minimum number of bunches/vines were recorded on Dogridge (50.33) rootstock. This might be due to the higher carbohydrate reserves of the vine and proper accumulation of source reserve (I could not make sense of this sentence..Pleaser rewrite and is there any reference??). The highest average bunch weight was observed ~~in-on~~ vines grafted on to 101.14MGT (87.59 g) which was at par with Fercal (86.22 g) ~~which-were-was-as-the~~ lowest average bunch weight was recorded in Gravasec (72.03 g). The higher yield/vine was recorded on Cabernet Sauvignon grafted onto 110R (6.68 kg) followed by Fercal, 1103P and 140RU (6.08, 5.19 and 5.02 kg) rootstock, while lower yield was recorded on Cabernet Sauvignon grafted onto Gravasec (4.23 kg). The influence of rootstock on yield ~~observations~~ has been noted by many scientists [25,2,26]. Higher yield is directly related to more stored food material and high pruned biomass of vine (did you do any correlation study or there is any reference

for this sentence). Rootstocks vary in rooting distribution pattern and number of roots, which might affect the pruned biomass and yield components and also the yield to pruned biomass ratio [27]. Similar findings were reported by [28].

The maximum 100 berry weight was recorded in Fercal (95.93) whereas it was minimum in 1103P (73.97) rootstock. The maximum number of berries/bunches were recorded on Dogridge (106.34) rootstock while minimum number of berries/bunches were recorded on Gravasec (86.89) rootstock.

Table 2. Effect of rootstocks in relation to yield parameters in Cabernet Sauvignon (pooled means for three years)

Rootstocks	No of bunches/vine	Average bunch weight (g)	Yield /vine (kg)	100 berry weight (g)	No of berries/bunch
Fercal	60.44 ^c	86.22 ^{ab}	6.08 ^b	95.93	98.67
Dogridge	50.33 ^d	85.21 ^b	4.28 ^g	81.50	106.34
SO4	70.00 ^a	81.41 ^c	4.82 ^e	79.08	100.22
110R	68.45 ^{ab}	85.02 ^b	6.68 ^a	84.84	110.56
Gravasec	59.00 ^c	72.03 ^d	4.23 ^g	82.49	86.89
1103P	71.11 ^a	73.45 ^d	5.19 ^c	73.97	100.33
101.14MGT	51.44 ^d	87.59 ^a	4.60 ^f	86.78	102.33
140RU	62.11 ^{ab}	81.95 ^c	5.02 ^d	90.54	90.00
SEm±	2.48	0.63	0.04	7.34	13.65
CD at 5%	7.53	1.92	0.11	22.25	41.40
<u>SigP value</u>	**	**	**	NS	NS

(when 100 berry wt and # berries/ bunch are non significant how is it possible that av bunch wt is significantly different, please report the cv on all tables) (What do you mean by ** and NS)

3.3 Berry quality parameters

The basic biochemical composition of Cabernet Sauvignon grafted onto different rootstock varied as shown in the results (table??). The differences ~~for~~ in TSS among the rootstocks were non-significant (Table 3). The acidity varied from 5.64-6.33 g/lit. The minimum acidity was recorded on SO4 rootstock while the maximum was on 140RU rootstock. Acidity was decreased with the increase in TSS (how can you say this ? have you done regression or correlation analysis??). Similar results were found by [29] (rewrite this sentence what kind of similar result please write clearly in this sentence.. by 29 is not proper way of writing the sentence). The highest juice pH was recorded in Cabernet Sauvignon grafted on 1103P (3.69) which was at par with SO4, Fercal, Dogridge and Gravasec (3.68, 3.65 and 3.63) while, minimum ~~in~~ on 101.14MGT (3.54) rootstock. The pH value of the grape juice was not significantly affected by the rootstock (your result finding is different in Table 3??) [30]. The maximum juice recovery (64.08 %) was recorded on Dogridge rootstock while minimum juice

recovery was recorded on SO4 (53.42 %) rootstock. The volatile acids (where is volatile acid in the table??) in grape berries were higher in 1103P (0.13 g/L) while SO4, 110R and Gravasec recorded lower ~~concentration~~ concentrations (0.09 g/L). The maximum the potassium uptake efficiency of vines mainly depends on scion cultivars, rootstock used and soil conditions, some rootstock showed more potash uptake capabilities and increased potassium content of the berries which positively affects the biochemical contents of the berries [31]. [32] stated that the rootstock might be responsible for modifying the berry ripening by controlling various gene expressions (32 stated is not good way to start a sentence. Pls rewrite). The same results were reported by [23] and [33] (23 and 33 says is also not the right way to state in a sentence) in Fantasy Seedless and Manjari Naveen grapevines grafted on Dogridge rootstock, respectively.

Table 3. Influence of rootstocks on basic biochemical composition of Cabernet Sauvignon (pooled means for three years)

Rootstocks	TSS (°Brix)	Acidity (g/lit)	Juice pH	Juice recovery (%)	VA (g/lit)
Fercal	23.42	6.04	3.65 ^{ab}	57.27 ^e	0.10 ^{bcd}
Dogridge	22.87	6.06	3.63 ^{ab}	64.08 ^a	0.12 ^{ab}
SO4	23.16	5.64	3.68 ^a	53.42 ^f	0.09 ^{cd}
110R	23.92	6.32	3.57 ^{bc}	59.17 ^{cde}	0.09 ^d
Gravasec	23.63	5.92	3.63 ^{ab}	58.79 ^{de}	0.09 ^{cd}
1103P	23.20	6.01	3.69 ^a	60.24 ^{cd}	0.13 ^a
101.14MGT	22.70	5.98	3.54 ^c	61.42 ^{bc}	0.11 ^{abcd}
140RU	23.42	6.33	3.57 ^{bc}	62.99 ^{ab}	0.11 ^{abc}
SEm±	0.31	0.16	0.03	0.83	0.009
CD at 5%	0.93	0.49	0.09	2.52	0.026
<u>P value</u>	NS	NS	*	**	*

(What do you mean by ** and NS, report cv% on table)

3.4 Wine quality parameters

Significant differences were recorded among the different rootstocks for wine quality parameters studied (Table 4). The wine made from 140RU recorded the lowest pH (3.76) while the rootstock Fercal recorded the highest pH of 4.22 each, respectively. [34] (34 number starting a sentence is not right way) reported that pH value regulates the degradation of glucose and fructose as the lower the pH value, slow will be the degradation. It is also playing a modulating role in wine haze formation, which diminishes or overthrows the commercial value of wine [35]. The concentration of volatile acid was higher in wine made from 140RU (0.41 g/L) followed by 101.14MGT (0.29 g/L) while the rootstock Gravasec recorded the least volatile acids (0.17 g/L). Total acid was higher in 140RU and 110R (4.1 g/L) which was at par

with Dogridge (4.0 g/L) and least in Fercal (3.2 g/L). Volatile acid plays an important role in fermentation process as it delivers information about the degree of improper fermentation processes occurring during winemaking [36] while acids, ethanol and tannins are the primary factors that determine the wine aroma, taste and mouth feel in red wine [37].


The wine made from 140RU rootstock recorded the highest concentration of ethanol (13.6 %) which was at par with Dogridge (13.5 %), 1103P (13.3 %) and 110R (13.0 %) while the lower concentration of ethanol was recorded in SO4 (12.0 %) grafted vines. The concentration of ethanol (14-16%) was a fundamental requirement for the wine quality as it is linked to sugar content of grape berries, which affect the overall flavour of wine [38]. (You mean non of the rootstock was good enough for ethanol production??) However, it decreases astringency and increases the bitterness of wine [39] (but how with higher or lower concentration??). Malic acid concentration was higher in wine made from 1103P (2.2 g/L) followed by 110R (2.1 g/L), Dogridge and 140RU (2.0 g/L)

while it was less in Fercal and SO4 (1.1 g/L) rootstocks. Colour intensity was maximum in SO4 rootstock (3.497) while minimum was recorded in 110R rootstock (1.493). During the wine making process, malic acid influences fermentation. [40] reported that at high concentration of malic acid, all strains of *Saccharomyces* yeasts were positive ~~that-which~~ enhanced the rate of fermentation process consuming all the sugar. [41] reported the variation due to influence of grape cultivar on the taste and colour of wine. (41 initiated a sentence not the right way to start a sentence) [42] reported role of regional factors for the malic acid concentration in wine.

Table 4. Evaluation of rootstock in relation to wine quality of Cabernet Sauvignon

3.5 Degree days requirement of rootstocks

The data on degree days and days to harvest are presented in Figure 1. The maximum degree days were recorded in Fercal and 110R (1468.40) grafted vines followed by Dogridge, 1103P, 101.14 MGT and 140RU while, minimum degree days were reported in SO4 and Gravasec (1386.07). Minimum day to harvest was recorded in Fercal and 110R (144 days) grafted vines followed by Dogridge, 1103P, 101.14 MGT and 140RU (140.00) however, SO4 and Gravasec took maximum days to harvest (158 days). [43] reported that BRS Melodia grapevines required growing cycle of 138 days with a yield of 23.85 tons/ ha during the season 2013.




(it is not the right way to make line graph to compare varieties, also include error bar on your line graph)

Fig. 1. Growing degree day requirement and duration to harvest of different rootstock of Cabernet Sauvignon

The wine prepared from Cabernet Sauvignon grapes ~~were~~was significantly influenced by the use of different rootstocks (Fig. 2). In terms of overall quality, (how do you measured the overall quality...I cannot find in table and methodology section) wine prepared from Cabernet Sauvignon grapes grafted on 1103P rootstocks recorded the highest (8) overall wine quality which was followed by Fercal (7) and SO4 (6) rootstocks whereas the lowest ~~reading of~~ overall wine quality was recorded in wine prepared form Cabernet Sauvignon grapes grafted on 101.14

MGT (1) rootstock. Rootstocks significantly influenced the phenolic, biochemical and sensory parameters of the prepared wine [44,45]. There was very less research carried out which showed the rootstock had a positive effect on the wine sensory attributes. The aroma of Cabernet Sauvignon wine was improved when Cabernet Sauvignon grapevines were grafted on the Ruggeri rootstock, compared to those of Salt Creek [46]. Cabernet Sauvignon wine had recorded the highest rating scores when grafted on 161-49 C and 420A MGT rootstocks [47]. As these all experiments were conducted in different environments, soils and climatic conditions in addition vineyard management and wine making procedures might have influenced on the outcomes.



Varieties comparison with line graph is not the right way use bar graph with error bar

Rating scale – 1 to 10

Fig. 2. Influence of rootstocks on wine quality of Cabernet Sauvignon

4. Conclusion (this section seems result section rewritten. Please state the importance of your stud and way forward for future research in this section)

The results of the present study indicated that the growth, yield, chemical composition of berries and quality of wine prepared from Cabernet Sauvignon grapes varied with the rootstock used. 110R rootstock recorded a significantly higher yield than other rootstocks. The growth parameters i.e. days to early bud sprout and days to veraison were minimum with Gravasec and days to harvest in Gravasec and SO4 rootstocks. Berry quality i.e. acidity was higher with

140RU rootstock; juice recovery was higher with Dogridge rootstock. Wine composition parameters like volatile acid, total acids and ethanol were higher with 140RU rootstock; malic acid was higher with 1103P rootstock and colour intensity in SO4 rootstock. The wine sensory attributes were also positively influenced using different rootstocks, wine prepared from Cabernet Sauvignon grapes grafted on 1103P recorded the highest overall wine quality which was followed Fercal rootstock.

UNDER PEER REVIEW

REFERENCES

1. NHB. Annual report. https://nhb.gov.in/annual_report.aspx?. 2024.
2. Ausari, P.K., Gurjar PKS, Somkuwar, RG, Naruka, LS, Sharma AK, Gharate, PS. Effect of rootstocks on yield and wine quality of Sauvignon Blanc variety. *Plant archives*. 2024;24:1477-1482.
3. Loreti F, Massai R. State of the art on peach rootstocks and orchard systems. *Acta Hort*. 2006;713:253-268.
4. Shaffer RG. The Effect of Rootstock on the Performance of the *Vitis vinifera* Cultivars Pinot noir, Chardonnay, Pinot Gris and Merlot. A thesis submitted to Oregon State University, United States. 2002.
5. Cousin P. Using rootstocks to enhance grape and wine quality and vineyard sustainable practices. In: *Proc Texas Vitic Enol Res Symp*. 2009;10-12.
6. Satisha J, Prakash GS. The influence of water and gas exchange parameters on grafted grapevines under conditions of moisture stress. *S Afr J of Enol and Vitic*. 2006;27:40-45.
7. Singh M, Sharma JK. Effect of rootstocks on disease intensity of Perlette ~~Grape~~ *Grapevine*. *Haryana J Hortic Sci*. 2005;34:234-235.
8. Chadha KL, Shikhamany SD. The grapes improvement, production and postharvest management practices. Malhotra Publications, New Delhi. 1999.
9. Kodur S, Tisdall JM, Clingeffer PR, Walker RR. Regulation of berry quality parameters in 'Shiraz' grapevines through rootstocks. *Vitis*. 2013;52:125-128.
10. Wang Y, Chen WK, Gao XT, He L, Yang XH, He F, Duan CQ, Wang J. Rootstock-Mediated Effects on Cabernet Sauvignon Performance: Vine Growth, Berry Ripening, Flavonoids, and Aromatic Profiles. *Int J Mol Sci*. 2019;20:401.
11. Van Leeuwen C, Destrac-Irvine A, Dubernet M, Duchene E, Gowdy M, Marguerit E, Pieri P, Parker A, de Resseguier L, Ollat N. An update on the impact of climate change in viticulture and potential adaptations. *Agronomy*. 2019;9:514.
12. Garrido J, Borges F. Wine and grape polyphenols- a chemical perspective. *Food Research International*. 2013;54(2):1844-58.
13. Arranz S, Chiva-blanch G, Valderas-martinez P, Medina-remon A, Lamuelaraventos R M and Estruch R. Wine, beer, alcohol and polyphenols on cardiovascular disease and cancer. *Nutrients*. 2012;4(7):759-81.

14. Casas R, Chiva-blanch G, Urpi-sarda M, Llorach R, Rotches-ribalta M, Guille M, Estruch R. Differential effects of polyphenols and alcohol of red wine on the expression of adhesion molecules and inflammatory cytokines related to atherosclerosis: A randomized clinical trial. *American Journal of Clinic Nutrition*. 2012;95:326-334.
15. Del Rio D, Rodriguez-Mateos A, Spencer JPE, Tognolini M, Borges G, Crozier A. Dietary (Poly) phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxid Redox Sign*. 2013;18:1818-92.
16. Ryan JJ, Dupont JA. Identification and analysis of major acids from fruit juices and wines. *J Agr Food Chem*. 1973;21(1):45-49.
17. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*. ICAR Rev. Ed, 1995;97-156.
18. Somkuwar RG, Satisha J, Ramteke SD. Effect of different rootstocks on fruitfulness in Thompson Seedless (*Vitis vinifera* L.). *Asian J of Plant Sci*. 2006;5:150-152.
19. Mahmoud AA, Thanaa ME, Mohammed MH, Salah E, El- Shenawe and Ahmed Shehata. Performance of some Table Grape Cultivars Grafting on Different Rootstocks in El-Nubaria Region. *Asian J of Crop Sci*. 2015;7:256-266.
20. Shafer R, Sampaio TLB, Pinkerton J, Vasconcelos MC. Grapevine rootstocks for Oregon vineyards. Oregon State University, Oregon, EUA. 11p. 2004.
21. Buttrose MS, Hale CR. Effect of temperature on development of the grapevine inflorescence after bud burst. *Amer J Enol Vitic*. 1973;24(1):14-16.
22. Tomasi D, Pascarella G, Sivilotti P, Gardiman M. Grape Bud Burst: Thermal Heat requirement and Bud Antagonism. *Proc. Intl. WS on Grapevine. Acta Hort*. 2007;754.
23. Somkuwar RG, Taware PB, Bhande AM, Sharma J, Khan I. Influence of different rootstocks on growth, photosynthesis, biochemical composition and nutrient content in 'Fantasy Seedless' grapes. *Int J of Fruit Sci*. 2015; 00:1-16.
24. Donnell CM, Dry PR, Wample RL, Bastian S. Proceeding of the 2nd annual viticulture research conference, University of California, Davis. 2008.
25. Ollat N, Tandonnet JP, Bordenave L, Decroocq S, Geny L, Gaudillere JP, Fouquet R, Barrieu F, Hamdi S. Short and long term effects of three rootstocks on Cabernet Sauvignon vine behaviour and wine quality. *Acta Hort*. 2003;617:95-100.

26. Somkuwar RG, Sharma AK, Gobade N and Ausari PK. Rootstock influences yield, berry composition and wine quality in Sauvignon Blanc grown under semi-arid condition. *Journal of Advances in Biology and Biotechnology*. 2024;27(7):382-389.
27. Morano L, Kliwer WM. Root Distribution of Three Grapevine Rootstocks Grafted to Cabernet Sauvignon Grown on a Very Gravelly Clay Loam Soil in Oakville, California. *Am J Enol Vitic*. 1994;45:345-348.
28. Satisha J, Somkuwar RG, Sharma J, Upadhyay AK, Adsule PG. Influence of rootstocks on growth yield and fruit Composition of Thompson Seedless grapes grown in the Pune region of India. *S Afr J Enol Vitic*. 2010;31(1):1-8.
29. Thutte AS, Somkuwar RG, Dhemre JK, Gharate PS, Ausari PK and Kakade PK. Rootstock impact on quality attributes and shelf life of Crimson Seedless grapes. *International Journal of Advanced Biochemistry Research*. 2024;8(4):556-560.
30. Kodur S, Tisdall JM, Clingeffer PR, Walker RR. Regulation of berry quality parameters in 'Shiraz' grapevines through rootstocks. *Vitis*. 2013;52:125-128.
31. Garcia M, Gallego P, Daverede C and Ibrahim H. Effect of three rootstocks on grapevine (*Vitis vinifera* L.) cv. Negrette, grown hydroponically. I. Potassium, calcium and magnesium nutrition. 2001. *S. Afr. J. Enol. Vitic*. 22:101-103.
32. Corso M, Vannozzi A, Ziliotto F, Zouine M, Maza E, Nicolato T, Vitulo N, Meggio F, Valle G, Bouzayen M. Grapevine rootstocks differentially affect the rate of ripening and modulate auxin-related genes in Cabernet Sauvignon berries. *Front Plant Sci*. 2016;7:14.
33. Somkuwar RG, Samarth R, Ghule VS, Sharma AK. Crop load regulation to improve yield and quality of Manjari Naveen grape. *Indian Journal of Horticulture*. 2020;77(2): 381-383.
34. Pan W, Jussier D, Terrade N, Yada RY, deOrduna RM. Kinetics of sugars, organic acids and acetaldehyde during simultaneous yeast-bacterial fermentations of white wine at different pH values. *Food Research International*. 2011;44:660-666.
35. Lambri M, Dordoni RR, Giribaldi M, Violetta MR, Giuffrida MG. Effect of pH on the protein profile and heat stability of an Italian white wine. *Food Research International*. 2013;54:1781-1786.
36. Mateo E, Torija MJ, Masand A, Bartowsky EJ. Acetic acid bacteria isolated from grapes of South Australian vineyards. *Int J Food Microbiol*. 2014;178:98-106.

37. Scott CF, Harbertsonb JF, Heymann H. A full factorial study on the effect of tannins, acidity, and ethanol on the temporal perception of taste and mouthfeel in red wine. *Food Quality and Preference*. 2017;62:1-7.
38. Meillon S, Urbano C, Guillot G, Schlich P. Acceptability of partially dealcoholized wines measuring the impact of sensory and information cues on overall liking in real life settings. *Food Qual Prefer*. 2010;21(7):763-773.
39. Fontoin H, Saucier C, Teissedre PL, Glories Y. Effect of pH, ethanol and acidity on astringency and bitterness of grape seed tannin oligomers in model wine solution. *Food Qual Prefer*. 2008;19(3):286-291.
40. Bovo B, Nadai C, Vendramini C, Lemos Junior WJF, Carlot M, Skelin A, Giacomini A, Corich V. Aptitude of *Saccharomyces* yeasts to ferment unripe grapes harvested during cluster thinning for reducing alcohol content of wine. *Int J Food Microbiol*. 2016;236: 56-64.
41. Van Leeuw R, Kevers C, Pincemail J, Defraigne JO, Dommès J. Antioxidant capacity and phenolic composition of red wines from various grape varieties: Specificity of Pinot Noir. *Journal of Food Composition and Analysis*. 2014;36:40-50.
42. Zeravik J, Fohlerova Z, Milovanovic M, Kubesa O, Zeisbergerova M, Lacina K, Petrovic A, Glatz Z, Skladal R. Various instrumental approaches for determination of organic acids in wines. *Food Chemistry*. 2016;194:432-440.
43. Koyama R, Borges WFS, Colombo RC, Hussain I, Souza RTD, Roberto SR. Phenology and yield of the hybrid seedless grape 'BRS melodia' grown in an annual double cropping system in a subtropical area. *Horticulturae*. 2020;6(1),3.
44. Jogaiah S, Kitture AR, Sharma AK, Sharma J, Upadhyay AK, Somkuwar RG. Regulation of fruit and wine quality parameters of Cabernet Sauvignon grapevines (*Vitis vinifera* L.) by rootstocks in semiarid regions of India. *Vitis*. 2015;54:65-72.
45. Somkuwar RG, Kakade PB, Ausari PK, Bhor V, Sharma AK, and Karande P. Influence of varying rootstocks on biochemical properties and sensory characteristics of Cabernet Sauvignon wine produced under subtropical climate. *International Journal of Advanced Biochemistry Research*. 2024;8(5):379-386.
46. Bravdo B, Shoseyov O. Aroma studies on fruits and wine in Israel. *Acta Hort*. 2000; 526:399-405.
47. Silvilotti P, Zulini L, Peterlunger E, Petrucci C. Sensory properties of Cabernet Sauvignon wines as affected by rootstock and season. *Acta Horticulturae Leuven*. 2007; 654:443-448.

UNDER PEER REVIEW