

Review Article

Lokāt (*Eriobotrya japonica*; family: Rosaceae): A fruit with nutritional and medicinal properties, in the light of Unani Medicine and scientific studies

ABSTRACT

Eriobotrya japonica Lind. (Family: Rosaceae) named as *loquat*, is a subtropical fruit, which is well known medicinal plant originated in Japan and China. Various parts, like leaves, peels and fruits have been shown to possess various useful health benefits. In Unani medicine, it is vastly utilized in many illnesses, like fevers, nausea, de-arranged sanguinoshumour (diseases due to morbidity of blood), indigestion, liver diseases, vomiting, dysentery, wounds, inflammations etc. Loquat plant contains many active constituents, such as glycosides, flavonoids, polyphenolic compounds, tannin etc. and nutritional and mineral compounds like, carotenoids, vitamins, starch, amino acids, sugar and others. According to various pharmacological studies it is found that the plant has many biological effects like antitussive, anti-melanogenic, anti-diabetic, anti-inflammatory, antimicrobial, kidney protective, hepatoprotective and hypolipidemic activity. This review aims to shed light on the therapeutic applications of *loquat* based on both traditional Unani literature and scientific studies conducted on different parts of the plant.

Keywords: *Eriobotrya japonica* Lind.; loquat, Lokāt; Flavonoids; Unani Medicine; Hepatoprotective

1. INTRODUCTION

Eriobotrya japonica (Thunb.) Lindl. is a fruit tree of medium sized that belongs to the family Rosacea. It is usually named Loquat. It has been implanted for more than two thousand years, and is native to Japan and China but has recently **cultivated** commercially worldwide in over 30 countries, such as Japan, Turkey, Iraq, Spain, Italy, Syria, and others". [1,2] "The history of loquat cultivation is more than 2000 years old, dating from the Chinese Han dynasty (100 B.C.)" [3]. "It attains up to 6 meters or more in height, with thick and evergreen oval-oblong leaves. Fruit is yellow to orange, pear-shaped, with seeds 3 to 4 cm long, and sweet taste" [1,2]. "Among other fruits, unusually loquat flowers in early winter or autumn, and its fruit ripens in early spring or late winter" [4]. "Mainly its fruits are eaten but also have been utilized in jam, chutney, and jelly preparation". [5] Loquat is planted mostly for fruit production, and in the Unani medicine "Loquat" is commonly known as "*Lokāt*" and has also been utilized for various medicinal purposes like treatment of nausea, vomiting, indigestion, dysentery, liver diseases, wounds, inflammations etc. **Pharmacological studies have demonstrated that loquat exhibits various bioactive properties, including anti-inflammatory, diuretic, anti-tumor [6], hepatoprotective [7], and anti-HIV effects [8].** This review is compiled to gather the scattered information related to the phytochemicals, medicinal uses and scientific researches on Lokāt fruits and other parts of the tree.

2. METHODOLOGY

A comprehensive literature review was conducted by searching all available classical textbooks using key terms such as Lokāt, in the context of Unani medicine. Additionally, electronic databases including Google Scholar, Research Gate; Scopus and PubMed indexed journals were explored using keywords like *Eriobotrya japonica*, Lokāt, Loquat, Unani Medicine, etc. The search included both classical Unani terms and botanical nomenclature. Review articles and experimental studies were carefully considered for data collection and subsequent analysis. This meticulous approach aimed to gather relevant information from both traditional Unani sources and contemporary scientific literature, providing a comprehensive overview of the therapeutic applications and properties associated with *Eriobotrya japonica* in the context of Unani medicine and scientific researches.

3. Observations

3.1 Distribution

E. japonica Lindl. is widely found in subtropical regions of Japan, China, India, and the Mediterranean area [10]. In India it grows in Kashmir and Bengal. [9]

3.2 Botanical description:

"*E. japonica* Lindl. is a large, evergreen 5–10 meters (16–33 feet) tall but is often smaller, shrub or small tree with a rounded crown, a short trunk, and woolly new twigs. Flowers appear in the autumn or early winter, and the fruits are ripe at any time from early spring to early summer. The flowers are 2 cm (3/4 in) in diameter, white, with five petals, and produced in stiff panicles of three to ten flowers. They have a sweet, aroma. The color of loquat fruit shows a marked change from green to yellow during its developmental and maturation period and become yellow to deep orange when ripen. The fruit begins to ripen from spring to summer, depending on the temperature in the area. The leaves are dark green, simple, alternate, 10–25 cm long, tough, and leathery in texture, with a serrated margin". [11]

3.3 Taxonomic classification

Kingdom:	Plantae
Division:	Tracheophyta
Class:	Magnoliopsida
Order:	Rosales
Family:	Rosaceae
Genus:	<i>Eriobotrya</i>
Species:	<i>Eriobotrya japonica</i> (Thunb.)
Synonyms:	<i>Mespilus japonica</i> , <i>Photinia japonica</i> , <i>Folium eriobotryi</i> . [3]



Fig 1. Lokāt tree **a, b**, tree with flowers & dried leaves **c, d**, fruits **e, f** and seeds **g**.

3.4 Description in Unani Literature:

Lokāt is a tree introduced in India by English people. It reached to a height of *QalmiĀm* (mango tree). Leaves are big with dentate margins; fruits and flowers appear in clusters. The fruits are green when unripe and yellow when ripened. The size of the fruit is a pigeon's egg. Unripe fruits are citrus in taste, and ripe fruits are sweet in the taste. Fruits have 4-5 seeds which look like the seeds of *Sharīfa* (*Annona squamosa* L.) or *Khīrnī* (*Mimosopselengi* L.) seeds, but are larger in comparison. [9] Mainly its fruits and juice (*Āb-i-Lokat*) are recommended by the Unani physicians.

3.5 Mutarādifāt (vernacular names):

Urdu:	Lakhota
Hindi:	Lukat, Logat, Latku
English:	Loquat
Kashmiri:	Lokāt
Bengali:	Lakhot
Sanskrit:	Luttak
Kannad:	Lakkote
Marathi:	Lokat
Gujrati:	Logat
Tamil:	Alakota
Malayalam:	Nespali
English:	Loquat

3.6 Mizāj (temperament):

Cold and dry and some says cold and moist. [12]

3.7 Af'ālwaKhawās(actions and uses):

85 *Qāti'-i-Şafrā'* (anti-bilious), *MusakkinHiddat-i-Khūn*(blood heat moderator), *Dāfi' Qay'* (antiemetic), *Mufarriḥ*(exhilarant),
 86 *Dafi' Humma* [12,13], *Musakkin-i-Atash* (thirst quencher) [14].
 87

88 3.8 Traditional uses and dosage forms of Lokāt

89 Various parts of the plant are used to treat respiratory and digestive disorders and also to cure inflammations, wound and
 90 *Diabetes mellitus* etc. The uses are mentioned bellow:

91 3.8.1 Amrād-i-Ri'a (Respiratory diseases)

- 92 • For the treatment of cough, leaves, flowers and fruits of Lokāt are used in the form of decoction in a quantity of
 93 10-20 ml. [9]

94 3.8.2 Amrad-i-Nizām-i-Haḍm (Digestive disorders)

- 95 • Its fruit is taken to cure indigestion, vomiting, and dysentery.[9] The juice of Lokāt increases appetite and treats
 96 indigestion [9].
- 97 • The decoction of its leaves is useful for the treatment of liver diseases [9]

98 3.8.3 Inflammation and wounds

- 99 • Powder of its dried leaves, are applied locally to heal wounds. [9]

100 3.8.4 Dhayābītus (Diabetes mellitus)

- 101 • *Safūf-i-Sandal Dhayābītuswalais* taken with *Ab-i-Lokāt* for the treatment of *Diabetes*. [15]

102 3.9 Maḍarrat (toxicity, side effect or adverse effect, contraindication):

103 The fruit is contraindicated for the person having cold and phlegmatic temperament. [13]
 104

105 3.10 Musleh (Corrective):

106 To counter its side effects due to over use, *MirchSiyāh* (black piper) and *Namak*(salt) can be used as correctives. [13]
 107

108 3.11 Badal (Substitute or Alternative):

109 *Paniyāla (Flacourtiajangomas* (Lour.) Raeusch.), having similar properties, so recommended as a substitute of Lokāt. [13]
 110

111 3.12 Miqdār Khūrāk (Dosage):

112 2-4 pieces are taken therapeutically. [13]
 113

114 3.13 Compound formulations: In Unani medicine *Ab-i-Lokāt* is used for above mentioned medical conditions.
 115

116 3.14 Phytoconstituents:

117 In loquat, a broad range of phytochemical compounds like phenols, alkaloids, cardiac glycosides, flavonoids, mucilage,
 118 gums, and phytosterols have been reported, [16] that are responsible for various biological activities. [17] "By using high
 119 performance liquid chromatography, flavones like quercetin derivatives, hydroxycinnamic acid derivatives like chlorogenic
 120 acid along with other pcoumaric acid and caffeic acid derivatives, have been identified in the leaves, fruit, and flower of
 121 loquat". [18] The presence of bioactive and nutritional compounds in various parts of the plant is mentioned in **Table 1**.
 122

123 **Table 1.** Showing bioactive and nutritional compounds in different parts of the Lokāt tree.
 124

S.N	Parts of the plant	Bioactive compounds	Nutritional compounds
1	Fruit	In fruits presence of flavonoids, phenols, [19], caffeic acid, 4-o-caffeoylquinic acid, neochlorogenic acid, and chlorogenic acid, together with 4-hydroxybenzoic acid, protocatechuic acid, coumaric acid, ellagic acid and ferulic acid have been	The fruit contains sugars: levulose and sucrose; citric acid, tartaric acid, succinic acid, cryptoxanthin, β-carotene, neo-β carotene. The seeds contain amygdalin and fatty oil, [30], starch [19],

		reported [20].	Carbohydrates, protein, amino acids [16], Vitamins, carotenoid [19]. The kernel of loquat is rich with starch, minerals and proteins [21,22].
2	Kernel	The kernel of loquat is rich with tannins, [21,22], amygdalin, a cyanogenetic glycoside [23].	
3	Flowers	Flavonoids and phenolic compounds [24], oleanolic and ursolic acids are reported in loquat flowers. [20] 15 aromatic compounds are present in the flowers, and the most potent aromatic compound in fresh loquat is phenylacetaldehyde. Additionally, other aroma such as, hexanal, (E)-2-hexenal, hexanoic acid, and β -ionone are also important also have been reported. [25] Presence of amygdalin is also reported. [26] Flowers possess a higher content of quercetin and chlorogenic acid derivatives than new and old leaves. [18]	
4	Leaves	In the leaves of loquat, phenolic acids like p-coumaric, gallic, caffeic, ellagic acid, tormentic acid, and flavonoids like quercetin, epicatechin, and catechin have been identified [27,28] in addition to tannins, sesquiterpenes, triterpenes and megastigmane glycosides [24,29]. The isomeric pentacyclicoleanolic acid and ursolic acid are predominant triterpenoids found in <i>E. japonica</i> leaves [30]. Amygdalin, a cyanogenetic glycoside, also present in in considerable amounts [23]. The new leaves possess phenolic and caffeic acid derivatives [18]	
5	Stem	In stem four flavonoids, kaempferol 3-O- β -glucoside, quercetin, quercetin 3-O- α -rhamnoside, naringenin, and three triterpene acids, ursolic, corosolic, and oleanolic acids, were reported. [24]	
6	Stem bark	Stem bark contains catechin, β -sitosterol, β -sitosterol-3-O- β -D-glucopyranoside, oleanolic acid, cinchonainIIb, lyoniresinol, and lyoniresinol 2-a-O- β -Dxylopyranoside [1].	

125
126

3.15 Pharmacological activities of *Eriobotrya japonica*

127

3.15.1 Cytotoxic activity

128

“According to a study of loquat juice on cancer cell lines revealed that it contains polyphenolic compounds that stimulate glutathione-S-transferase enzymes (GSTs). These GSTs act as antioxidants, promoting cellular detoxification and apoptosis in cancer cells” [32]. Alwash (2017) reported that, “loquat fruit juice exhibited a pronounced anticancer effect on rhabdomyosarcoma (RD) and human cervical cancer (HeLa) cells”[2].

129

130

131

132

133

3.15.2 Anti-Diabetic activity

134

Shafi et al. reported (2019) that in “streptozotocin induced diabetic rat, the ethanol loquat fruit extract showed good anti-diabetic effect”. [33]. “Also, study done on maltose-loaded Sprague–Dawley rats, combination of green tea leaves and loquat leaves exhibited a reduction of blood levels of glucose and a corresponding decrease in serum secretion of insulin” [34].

135

136

137

138

139

3.15.3 Kidney protective effect

140

“*Diabetes mellitus* is a metabolic disease differentiated by hyperglycemia and several complications, such as neuropathy, retinopathy, angiopathy, nephropathy etc”. [35] Shafi et al. (2018) reported that, “ethanolic extract (50%) of seeds and

141

142 fruits of *Eriobotrya japonica* displayed renal effects in alloxan induced diabetic rats, which was evaluated by estimating
143 serum total proteins, serum creatinine and urea levels. The results showed a nonsignificant elevation in level of total
144 proteins and reduction in levels of serum urea, creatinine and also had effects on levels of serum glucose" [36].
145

146 **3.15.4 Hypolipidemic activity**

147 Shafi et al. (2019) also reported that "in streptozotocin induced diabetic rat, the ethanolic extract of fruit *Eriobotrya japonica*
148 showed significant hypolipidemic activity" [36]. "In high-fat diet mice, ethanolic extracts of loquat exhibited hypolipidemic
149 activity as it decreased white adipose tissue (WAT) weights (include visceral fat, mesenteric, peri-renal, and epididymal
150 WAT), body weight gain, hepatic tri-acylglycerol, and adipocyte size in the visceral depots" [37,38]. "Fermented tea
151 product (leaves mixture of both green tea and loquat) inhibited the synthesis of hepatic fatty acids and postprandial
152 hypertriacylglycerolemia by inhibition of pancreatic lipase, exhibiting anti-obesity action". [39] "The aqueous extract of
153 loquat leaves exhibited an anti-atherosclerotic effect in a hypercholesterolemic zebra fish model and in cellular assays"
154 [40].
155

156 **3.15.5 Anti-inflammatory activity**

157 Maher et al. study (2015) reported "the ethyl acetate-ethanol (1/2) extract and the dichloromethane methanol (0:1) fraction
158 of loquat suppressed the phospholipase A2 group IB (pG-IB) secreted in pig and the human secreted phospholipase A2
159 group IIA (hG-IIA)" [41]. "The N-butanol fraction of leaves of loquat demonstrated anti-inflammatory property by inhibiting
160 the expression of nitric oxide synthase and production of NO, also down-regulated cyclooxygenase-2 expression, pro-
161 inflammatory cytokine secretion such as interleukin-6 and tumor necrosis factor- α (TNF- α) in the LPS-activated murine
162 peritoneal macrophage model" [42]. Seong et al (2019) reported that "in lipopolysaccharide-induced RAW 264.7
163 macrophage cells, the ethanolic extract of loquat leaves showed an anti-inflammatory effect by suppressing TNF- α
164 production and NO expression" [43]. Zar et al. (2013) reported that "in lipopolysaccharide-induced RAW 264.7 cells,
165 aqueous extract of loquat leaves extracted by boiling for 15 minutes at 100°C, exerted an anti-inflammatory effect via
166 inhibiting PGE2 and COX-2 production. The new bioactive phenolic compounds in loquat tea may be responsible for its
167 anti-inflammatory potency" [44]. "Loquat tea water extract prepared by boiling it for 15 minutes at 100°C showed inhibitory
168 effects on the expression of TNF- α , interleukin-6, nitric synthase, and NO through the downregulation of pathways of the
169 TGF- β -activated kinase-mediated NF- κ B and MAPK" [45]. "This was observed in macrophage-like RAW 264.7 cells used
170 in the mouse paw edema model" [45,46].
171

172 **3.15.6 Antitussive and expectorant activity**

173 Wu et al. (2018) reported that "the ethanolic and aqueous extracts of loquat leaves showed expectorant and antitussive
174 effects. Aqueous extracts of growing leaves had a higher expectorant effect, which may be related to their higher flavonoid
175 content, such as quercetin, isoquercitrin, hyperoside, rutin, and others. Meanwhile, ethanolic extracts of fallen leaves
176 demonstrated better antitussive effects, which may be related to their higher triterpenoids content, such as tormentic acid,
177 corosolic acid, maslinic acid, ursolic acid, and others".
178 [46]
179

180 **3.15.7 Anti-melanogenic activity**

181 Seong et al. (2019) stated that ethanolic leaf extract of loquat showed anti-melanogenic activity because of its anti-
182 inflammatory and anti-oxidant activities. The ethanolic extract has a higher concentration of quercetin and other
183 polyphenols that limit the creation of melanin, it shows protection of human skin from oxidative stress and inflammation.
184 Since melanin is crucial in preventing the generation of UV-stimulated ROS, ethanolic extract can control melanin
185 formation. It also exhibits strong anti-inflammatory and antioxidant properties. [43,47] In B16 melanoma cells, the
186 methanolic leaf extract of *Eriobotrya japonica* exerted dose-dependant melanogenesis suppression
187 [43,47,48]. Furthermore, 70% and 30% loquat leaf ethanolic extracts inhibited mushroom tyrosinase for having whitening
188 effects [49].
189

190 **3.15.8 Hepatoprotective activity**

191 The study conducted by Shahat et al. (2018) shown that the methanolic extract (80%) of loquat leaves, as well as its
192 butanol, aqueous, and ethyl acetate fractions, had hepatoprotective effects in rats with hepatotoxicity induced by CCl4.
193 It significantly reduced biochemical parameter levels in rats like aspartate transaminase (AST), gamma-glutamyl
194 transferase, alanine aminotransferase (ALT), bilirubin, and alkaline phosphate levels but did not influence lipid profiles.
195 Administration of butanol and ethyl acetate fractions significantly suppressed CCl4-stimulated depletion of total protein
196 and the reduced levels of nonprotein sulfhydryl groups (NP-SH). [50].
197

198 **3.15.10 Antimicrobial activity**

199 Rashed et al. (2014) reported that the methanolic extract (80%) of stems of loquat extracted by
200 maceration process, demonstrated an antimicrobial effect against bacterial and fungal strains linked to the presence of
201 triterpenes and flavonoids. It was found to be more effective against *Candida albicans*, indicating that it can be used to
202 treat fungal infections and has no effect on other strains of bacteria or fungus. [24].

The presence of flavonoids [51], tannins [52], and kaempferol 3-O-β-glucoside [53] in the methanolic extract of loquat stems are responsible for its strong antibacterial and antioxidant properties.

3.15.11 Antiosteoporosis activity

Methanolic leaf extract of loquat showed antiosteoporotic effects in the model of ovariectomized mice [54]. Ursolic acid was isolated from loquat leaves and displayed an inhibitory effect on osteoclast differentiation. Ursolic acid was found to suppress the development of osteoclasts by targeting exportin 5 (XPO5), a nuclear exporter protein, to decrease osteoclast development. [55].

3.15.12 Antifibrosis activity

In a rat model of bleomycin-induced pulmonary fibrosis, triterpenic acids from an ethanolic extract of loquat leaves (prepared by cold technique, 2 hours) showed antifibrotic efficacy by reducing lung fibrosis and enhancing lung architecture. Rats with pulmonary fibrosis had lower levels of TGF-β1 and TNF-α production in their macrophages, both in terms of mRNA and protein.[56].

3.15.13 Antioxidant activity

Alwash (2017) reported that the fruit juice of *E. japonica* shown antioxidant properties [2]. By contributing their H atoms, the plant's juice can scavenge DPPH free radicals [57]. The phenolic compounds found in *E. japonica* fruit juice, particularly flavonoids, have the capacity to cause this reaction [17]. Their efficacy as antioxidants depends on the location and quantity of OH groups on the basic flavonoid structure; an increase in the number of hydroxyl groups is directly associated with an increase in antioxidant activity [1].

4. CONCLUSION

Lokāt (*Eriobotrya japonica*) emerges as a remarkable fruit with a rich profile of nutritional and medicinal properties, validated by both the ancient wisdom of Unani Medicine and contemporary scientific research. Traditionally, Unani practitioners have long valued Lokāt for its diverse therapeutic applications, including its potential to improve digestive health, support respiratory function, and regulate blood sugar levels. These traditional uses are increasingly supported by modern studies that highlight the fruit's abundance of essential nutrients, antioxidants, and bioactive compounds. Recent scientific investigations have shed light on loquat's capacity to combat oxidative stress, reduce inflammation, and provide protective effects against chronic diseases such as diabetes, cardiovascular ailments, and certain types of cancer. The convergence of Unani Medicine insights with cutting-edge research underscores Lokāt's potential as a functional food with significant health benefits. As the global interest in natural and holistic health solutions grows, Lokāt stands out as a valuable addition to the repertoire of nutraceuticals. Its integration into modern dietary practices could offer a natural means to enhance health and well-being. Future research should continue to explore and substantiate the therapeutic claims, ensuring a comprehensive understanding of this multifaceted fruit. In conclusion, Lokāt exemplifies the harmonious blend of traditional knowledge and modern science, reaffirming its place as a fruit with profound nutritional and medicinal promise. By embracing both historical perspectives and contemporary findings, we can fully appreciate and harness the potential of Lokāt for improving human health.

ACKNOWLEDGEMENTS

The authors are very thankful to Deputy Director RRIUM, Srinagar for providing facilities in the library of RRIUM, Srinagar.

COMPETING INTERESTS

Nil

AUTHORS' CONTRIBUTIONS

'Author A' designed the study, wrote the first draft of the manuscript. 'Author B' and C manage the literature search. All authors read and approved the final manuscript."

Disclaimer (Artificial intelligence)

Option 1:

261 Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc)
262 and text-to-image generators have been used during writing or editing of manuscripts.

263 Option 2:

264 Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during
265 writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI
266 technology and as well as all input prompts provided to the generative AI technology

267 Details of the AI usage are given below:

268 1. Grammar checker has been used only for checking and correcting the grammar.

269 2. No text to image generators have been used.

270 REFERENCES

- 271 1. Fouedjou RT, Nguenefack-Mbuyo EP, Ponou BK, Nguenefack TB, Barboni L and Taponjhou LA.
272 Antioxidant Activities and Chemical Constituents of Extracts from *Cordylinefruticosa* (L.) A. Chev.
273 (Agavaceae) and *Eriobotrya japonica* (Thunb) Lindl, (Rosaceae). *Pharmacologia*, 2016; 7 (2-3): 103-
274 113.
- 275 2. Alwash BMJ. Cytotoxic and Antioxidant Activity of Fruit Juice of *Eriobotrya Japonica* (Thunb.) Lind Plant
276 Cultivated in Iraq. *The Iraqi Journal of Agricultural Sciences*, 2017; 48(3): 892-898.
- 277 3. Chen SJ, Chen MM, Kang BB, Lin YY, Chen QQ. Effect of mechanical injury on postharvest physiology
278 of loquat fruit. *Journal of Fujian Agriculture and Forestry University* 2003; 33: 250-254.
- 279 4. Delfanian M, Kenari RE and Sahari MA. Antioxidant Activity of Loquat (*Eriobotrya japonica* Lindl.) Fruit
280 Peel and Pulp Extracts in Stabilization of Soybean Oil During Storage Conditions. *International Journal*
281 *of Food Properties*, 2015; 18: 2813–2824.
- 282 5. Koba K, Matsuoka A, Osada K, Huang YS. Effect of Loquat (*Eriobotrya japonica*) Extracts on LDL
283 Oxidation. *Food Chemistry*, 2007; 104(1): 308–316.
- 284 6. Shibata S. Chemistry and cancer preventing activities of ginseng saponins and some related
285 triterpenoid compounds. *Journal of Korean Medical Science* 2001; 16: S28–37.
- 286 7. Ma XH, Zhao YC, Yin L, Han DW, Ji CX. Studies on the effect of oleanolic acid on experimental liver
287 injury. *ActaPharmaceuticaSinica* 1982; 17: 93–97.
- 288 8. Kashiwada Y, Wang HK, Nagao T, Kitanaka S, Yasuda I, Fujioka T, et al. Anti-AIDS agents 30 Anti HIV
289 activity of oleanolic acid, pomolic acid, and structurally related triterpenoids. *Journal of Natural Products*
290 1998; 61: 1090–1095.
- 291 9. Kalam, MA. *Fawakeh Kashmir*. New Delhi; IdaraKitabusShifa. 2021
- 292 10. Lin S, Sharpe RH, Janick J, In: Jules, J. (Ed.), *Loquat: Botany and Horticulture*, *Horticultural Reviews*,
293 23. Wiley, New York, 1999, 234–276.
- 294 11. Kirtikar and Basu. *Indian Medicinal Plants*. Periodical Expert Book Agency, New Delhi. Vol II, 2012, p 989-
295 990
- 296 12. Ghani, HN. *KhazainulAdvia*. New Delhi; IdāraKitabusShifa. 2011
- 297 13. Hakim, MA. *BustanulMufradat*. New Delhi; IdāraKitabusShifa. 2011
- 298 14. Kabiruddin HM. *Makhzan al-Mufradat*. New Delhi: IdaraKitab al-Shifa. 2014, Pp-365-66
- 299 15. Khan A. Haziq. New Delhi: IdāraKitabusShifa. 2017. P-388-89.
- 300 16. Rajalakshmi P, Pugalanthi M, Subashini G, Miss. Kavitha and Vishnukumar S. Nutraceutical studies on
301 *Eriobotrya japonica* (Thunb.) Lindl. (Fruits & Seeds). *International Journal of Advances in Scientific*
302 *Research*, 2017; 3(04):44-48.
- 303 17. Matalka KZ, Abdulridha NA, Badr MM, Mansoor K, Qinna NA and Qadan F. *Eriobotrya japonica* Water
304 extract characterization: an inducer of interferongamma production mainly by the JAK-STAT Pathway.
305 *Molecules*, 2016; 21: 722.

- 311 18. Ahumada J, Fuentealba C, Olaeta JA, Undurraga P, Pedreschi R, Shetty K, Chirinos R, Campos D,
312 and Ranilla LG. Bioactive compounds of loquat (*Eriobotrya japonica* Lindl.) cv. Golden Nugget and
313 analysis of in vitro functionality for hyperglycemia management. *Cien. Inv. Agr.* 2017; 44(3): 272- 284.
- 314 19. Pareek S, Benkeblia N, Janick J, Cao S and Yahia EM .Postharvest physiology and technology of
315 loquat (*Eriobotrya japonica* Lindl.) fruit. *Journal of the Science of Food and Agriculture*, 2014;
316 94(8):1495-1504.
- 317 20. Xu H and Chen J. Commercial quality, major bioactive compound content and antioxidant capacity of
318 12 cultivars of loquat (*Eriobotrya japonica* Lindl.) fruits. *J. Sci. Food Agric.*2011; 91:1057–1063.
- 319 21. Fu XM, Kong WB, Peng G, Zhou JY, Azam M, Xu CJ, Grierson D, Chen KS. Plastid structure and
320 carotenogenic gene expression in red- and white-fleshed loquat (*Eriobotrya japonica*) fruits. *J. Exp. Bot.*
321 2012; 63:341–354.
- 322 22. Zhang WN, Zhao XY, Sun CD, Li X, Chen KS. Phenolic composition from different loquat (*Eriobotrya*
323 *japonica* Lindl.) cultivars grown in China and their antioxidant properties. *Molecules.* 2015; 20:542–555.
- 324 23. Zhuang YF. Determination of amygdalin content in *Eriobotrya japonica* leaves by high performance
325 liquid chromatography (HPLC). *Strait Pharmaceutical Journal* 2002; 14: 64–65.
- 326 24. Rashed KN and Butnariu M. Isolation and antimicrobial and antioxidant evaluation of bioactive
327 compounds from *Eriobotrya japonica* stems. *Adv. Pharmaceut. Bull.* 2014; 4 (1): 75-81.
- 328 25. Takahashi H, Sumitani H, Inada Y, Mori D, Nakano Y. Potent aroma volatiles in fresh loquat and its
329 canned product. *Nippon Shokuhin Kagaku KogakuKaishi* 2000; 47: 302-310.
- 330 26. Kwon HY, Hong SP, Hahn DH, Kim JH. Apoptosis induction of *Persicaria Semen* extract in human
331 promyelocytic leukemia (HL-60) cells. *Archives of Pharmacal Research* 2003; 26: 157–161.
- 332 27. Pande G and Akoh CC. Antioxidant capacity and lipid characterization of six Georgia-grown
333 pomegranate cultivars. *J Agric Food Chem.* 2009; 57(20):9427–36.
- 334 28. Fogo AS, Antonioli E, Calixto JB, and Campos AH. Tormentic acid reduces vascular smooth muscle
335 cell proliferation and survival. *European Journal of Pharmacology*, 2009; 615(1-3):50–54.
- 336 29. Nawrot-Hadzika I, Granicab S, Abela R, Czapor-Irzabekc H and Matkowskia A. Analysis of Antioxidant
337 Polyphenols in Loquat Leaves using HPLC-based Activity Profiling. *Natural Product Communications*,
338 2017; 12 (2):163-166.
- 339 30. Vasconcelos MA, Royo VA, Ferreira DS, Crotti AE, Andrade E, Silva ML, et al. In vivo analgesic and
340 anti-inflammatory activities of ursolic acid and oleanoic acid from *Miconia albicans* (Melastomataceae).
341 *Zeitschrift fur Naturforschung* 2006; 61: 477–482.
- 342 31. Park HJ, Yoon SH, Han LS, Zheng LT, Jung KH, Uhm YK, et al. Amygdalin inhibits genes related to cell
343 cycle in SNU-C4 human colon cancer cells. *World Journal of Gastroenterology* 2005; 11: 5156– 5161.
- 344 32. Soni A and Sosa S. Phytochemical analysis and free radical scavenging potential of herbal and
345 medicinal plant extracts. *JPP*, 2013; 2: 22-29.
- 346 33. Shafi S, Tabassum N. Effect Of Ethanolic Extract of Fruits Of *Eriobotrya Japonica* On Lipid Profile And
347 Body Weight In Streptozotocin Induced Diabetic Rats. *International Journal of Pharmaceutical Sciences*
348 *and Drug Research*, 2019; 11(1): 01- 06.
- 349 34. Tamaya K, Matsui T, Toshima A, Noguchi M, Ju Q, Miyata Y, Tanaka T, Tanaka K. Suppression of
350 blood glucose level by a new fermented tea obtained by tea-rolling processing of loquat (*Eriobotrya*
351 *japonica*) and green tea leaves in disaccharide-loaded Sprague–Dawley rats. *J. Sci. Food Agric.* 2010;
352 90:779–783.
- 353 35. Alberti KG, Zimmet PZ. Definition diagnosis and classification of diabetes mellitus and its complications.
354 Part I: diagnosis and classification of diabetes mellitus, provisional report of a WHO consultation.
355 *Diabetes Med.* 1998; 15:539-53.
- 356 36. Shafi S and Tabassum N. Phytochemical Screening And Renal Effects Of Ethanolic Extract Of
357 *Eriobotrya Japonica* Fruits And Seeds In Alloxan Induced Diabetic Rats. *International Journal of*
358 *Current Pharmaceutical Research*, 2018; 10(1): 3-7.

- 359 37. Shih CC, Lin CH, Wu JB. *Eriobotrya japonica* improves hyperlipidemia and reverses insulin resistance
360 in high-fat-fed mice. *Phytother. Res.* 2010; 24:1769–1780.
- 361 38. Shih CC, Ciou JL, Lin CH, Wu JB, Ho HY. Cell suspension culture of *Eriobotrya japonica* regulates the
362 diabetic and hyperlipidemic signs of high-fat-fed mice. *Molecules*, 2013; 18:2726–2753.
- 363 39. Tanaka K, Tamaru S, Nishizono S, Miyata Y, Tamaya K, Matsui T, Tanaka T, Echizen Y, Ikeda I.
364 Hypotriacylglycerolemic and antiobesity properties of a new fermented tea product obtained by tea-
365 rolling processing of third-crop green tea (*Camellia sinensis*) leaves and loquat (*Eriobotrya japonica*)
366 leaves. *Biosci. Biotechnol. Biochem.* 2010; 74:1606– 1612.
- 367 40. Kim JY, Hong JH, Jung HK, Jeong YS, Cho KH. Grape skin and loquat leaf extracts and acai puree
368 have potent anti-atherosclerotic and antidiabetic activity in vitro and in vivo in hypercholesterolemic
369 zebrafish. *Int. J. Mol. Med.* 2012; 30:606–614.
- 370 41. Maher K, Yassine BA, Sofiane B. Antiinflammatory and antioxidant properties of *Eriobotrya japonica*
371 leaves extracts. *African Health Sciences*, 2015; 15(2): 613-620.
- 372 42. Cha DS, Eun JS, Jeon H. Anti-inflammatory and antinociceptive properties of the leaves of *Eriobotrya*
373 *japonica*. *J. Ethnopharmacol.* 2011; 134: 305–312.
- 374 43. Seong N, Oh W, Kim I, Kim S, Seo J, Park C, Kim D, Ko J and Kim J. Efficacy and local irritation
375 evaluation of *Eriobotrya japonica* leaf ethanol extract. *Laboratory Animal Research*, 2019; 35:4.
- 376 44. Zar PPK, Sakao K, Hashimoto F, Morishita A, Fujii M, Wada K and Hou D. Antioxidant and anti-
377 inflammatory activities of loquat (*Eriobotrya japonica*) tea. *Functional Foods in Health and Disease*,
378 2013; 3(11):447-461.
- 379 45. Zar PPK, Morishita A, Hashimoto F, Sakao K, Fujii M, Wada K, Hou DX. Anti-inflammatory effects and
380 molecular mechanisms of loquat (*Eriobotrya japonica*) tea. *J. Funct. Foods.* 2014; 6:523–533.
- 381 46. Wu Y, Jian T, Lv H, Ding X, Zuo Y, Ren B, Chen J, Li W. Antitussive and expectorant properties of
382 growing and fallen leaves of loquat (*Eriobotrya japonica*). *Rev. bras. farmacogn.* 2018; 28(2):239-242.
- 383 47. Agar N and Young AR. Melanogenesis: a photoprotective response to DNA damage? *Mutat Res.* 2005;
384 571:121–32.
- 385 48. Tan H, Sonam T, Shimizu K. The potential of triterpenoids from loquat leaves (*Eriobotrya japonica*) for
386 prevention and treatment of skin disorder. *Int J Mol Sci.* 2017; 18(5):1030.
- 387 49. Park JO, Park JO, Joo CG. A study on whitening and anti-inflammatory effects of *Eriobotrya japonica*
388 leaf extracts with different extraction methods. *J SocCosmetSci Korea.* 2015; 41(2):151–7.
- 389 50. Shahat AA, Ullah R, Alqahtani AS, Alsaid MS, Husseiny HA, and Al Meanazel OTR. Hepatoprotective
390 Effect of *Eriobotrya japonica* Leaf Extract and Its Various Fractions against Carbon Tetra Chloride
391 Induced Hepatotoxicity in Rats. *Evidence-Based Complementary and Alternative Medicine*, 2018; 9:1-8.
- 392 51. Tapas AR, Sakarkar DM, Kakde RB. Flavonoids as Nutraceuticals: A review. *Trop J Pharm Res.* 2008;
393 7(3):1089–99.
- 394 52. Reddy MK, Gupta SK, Jacob MR, Khan SI, Ferreira D. Antioxidant, antimalarial and antimicrobial
395 activities of tannin-rich fractions, ellagitannins and phenolic acids from *Punicagranatum L.* *Planta Med.*
396 2007;73(5):461–7.
- 397 53. Skalicka-Wozniak K, Melliou E, Gortzi O, Glowniak K, Chinou IB. Chemical constituents of
398 *Lavateratrimestris L.*--antioxidant and antimicrobial activities. *Z Naturforsch C.* 2007; 62(11-12):797–
399 800.
- 400 54. Tan H, Furuta S, Nagata T, Ohnuki K, Akasaka T, Shirouchi B, Sato M., Kondo R, Shimizu K. Inhibitory
401 effects of the leaves of loquat (*Eriobotrya japonica*) on bone mineral density loss in ovariectomized
402 mice and osteoclast differentiation. *J. Agric. Food Chem.* 2014; 62:836–841.
- 403 55. Tan H, Zhao C, Zhu Q, Katakura Y, Tanaka H, Ohnuki K and Shimizu K. Ursolic Acid Isolated from the
404 Leaves of Loquat (*Eriobotrya japonica*) Inhibited Osteoclast Differentiation through Targeting Exportin
405 5. *J. Agric. Food Chem.* 2019; 67(12): 3333-3340.

- 406 56. Yang Y, Huang Y, Huang C, Lv XW, Liu L, Wang YY, Li J. Antifibrosis effects of triterpene acids of
407 *Eriobotrya japonica* (Thunb.) Lindl. leaf in a rat model of bleomycin-induced pulmonary fibrosis. *J.*
408 *Pharm. Pharmacol.* 2012;64: 1751–1760.
- 409 57. Shekhar T C and Anju G. Antioxidant activity by DPPH radical scavenging method of *Ageratum*
410 *conyzoides* Linn .leaves. *American Journal of Ethno medicine*, 2014; 1: 244-249.
- 411
- 412
- 413