

Influence of foliar spray with different doses of Zn (mention form of zinc) at various crop stages
Influence of foliar spray and stages of zinc application on seed on quality parameters of harvested radish seeds

Comment [A1]: Write clear title

Abstract

A laboratory experiment was conducted to analyze the influence of foliar spray and stages of zinc application on seed quality parameters of harvested radish seeds cv. Japanese White in laboratory of the Department of Seed Science and Technology, Dr Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan (HP). There were 12 treatment combinations of 4 zinc levels viz, 0g l^{-1} , 1g l^{-1} , 2g l^{-1} , 3g l^{-1} and 3 stages of application viz, Bolting, Bolting + 14 days after first application, Bolting + 14 days after first application+ 28 days after first application. Among treatment combinations Zn_2S_1 (2g l^{-1} at the time of bolting) was found superior in every aspect of quality attributes i.e. 1000 seed weight (14.36g), germination of harvested seeds (85.75%), seedling length (24.20cm), seedling dry weight (11.31mg), seedling vigour index-I (2,075.08) , seedling vigour index- II (966.77), electrical conductivity (0.017 dSm^{-1}), accelerated ageing test (78.25%) and cold test (81.00%).

Comment [A2]: Mention different factor instead 1st factor as stages and 2nd factor as treatment with different doses of Zn also mention the form of Zn used

Keywords: Radish, micronutrients, zinc, seed quality parameter

Introduction

Radish is a cruciferae family root crop that is grown in both tropical and temperate regions of the world. It is native to Europe and Asia. Radish seed output per unit area is low, however, it could be increased with better management practices such as foliar micronutrient supplementation. Micronutrients are essential for germination and seedling growth, from cell wall development to respiration, photosynthesis and chlorophyll generation. The administration of micronutrients accelerates the uptake of all plant nutrients while simultaneously generating a defence mechanism against insect - pests and diseases, hence enhancing growth and productivity (Anuprita et al., 2005). Micronutrients are required in minimal dose, even then lack of these might hinder growth and production. Foliar spray is the most recent type of crop nutrition, consisting of micronutrients given to leaves in liquid form. Foliar spray is 6 to 20 times more effective than soil application (Arif et al., 2006). Micronutrients such as zinc are critical for plant growth, development and biomass production because they are required for the creation of Indole Acetic Acid (IAA). It also aids in energy production, protein and growth regulator synthesis (Kumar et al., 2014), pollen development, sexual fertilization and germination (Cakmak, 2008), greater seed yield, and fruit number. It also improves growth, hormonebiosynthesis, starch creation, seed production and maturation (Brady and Weil, 2002). Zinc also improves the viability and vigour of seedlings. It is a necessary component of ribosomes and is critical to their structural integrity (Trivedi, 2013). Zinc plays numerous minor roles in plant growth, and a consistent and continuous supply is essential for optimal growth and output (Acquaah, 2002). The Zn ion is the form of zinc that is available to plants. Because of its adsorption with soil particles and less interaction with crop roots, foliar application of Zn^{2+} foliar spray is more effective than soil treatment (Wissuwa et al., 2008).

Comment [A3]: Mention the type of soil in which Zn is deficient and highlight the problem and objective clearly

Material and method

To study the effect of foliar application of zinc on quality of harvested seeds of radish, seed analysis was done at laboratory of the Department of Seed Science and Technology, Dr Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan (HP). After completion of field trial harvested seeds of radish cv. Japanese White were evaluated in laboratory for different seed quality parameters. In field trial, 12 different treatments were laid out by combining 4 different ~~doses levels~~ of zinc (mention the form

of Zn used) viz. Zn₁: 0g l⁻¹, Zn₂: 1g l⁻¹, Zn₃: 2g l⁻¹ and Zn₄: 3g l⁻¹ along with 3 different stages of application S₁: Bolting, S₂: Bolting + 14 days after first treatment, S₃: Bolting + 14 days after first application + 28 days after first application. Observations under laboratory conditions were recorded for 1000 seed weight, germination per cent, seedling length, seedling dry weight, seed vigour index-I, Seed vigour index-II, electrical conductivity, accelerated ageing test and cold test as per standard procedures. Experiments were conducted with 4 replications (ISTA, 2016) and results were statistically analyzed by using completely randomized block design (Panse and Sukhatme, 2000) following window-based computer application OPSTAT developed by Sheoran (2006).

Comment [A4]: Also do post hoc analysis for clearer result

Results and discussion

1000 seed weight

Data (Table 1) depicted that zinc levels, stages of foliar spray and their combinations had a significant influence on 1000 seed weight. Among the levels of zinc, 1000 seed weight was uncurvingly maximum (13.55 g) up to 2g l⁻¹ (Zn₃) and thereafter a significant reduction was observed when zinc was applied @ 3g l⁻¹ and minimum 1000 seed weight (12.28 g) was noticed in Zn₁ (Water spray). As far as stages of application were concerned, the highest 1000 seed weight (13.20 g) was observed with foliar application of zinc at S₁ (Bolting stage) and after that there was a linear decrease and lowest (12.99 g) 1000 seed weight was observed at S₃ (Bolting + 14 days + 28 days after first application). However, treatment combinations Zn₃S₁ (2g l⁻¹ foliar spray of zinc applied at bolting stage) produced maximum 1000 seed weight (14.36 g) among all interactions and minimum (12.28 g) was observed in treatment combination Zn₁S₂ (Water spray at bolting + 14 days after first application) and Zn₁S₃ (Water spray at bolting + 14 days + 28 days after first application) was at par with Zn₁S₁. The increase in 1000 seed weight could be attributed to zinc as it is responsible for improved seed filling due to its role in pollen germination, cell division, sugar transfer from source to sink and seed development Masuthiet *et al.*, (2009). Pariari *et al.*, (2009) in fenugreek and Lakshami *et al.*, (2017) in black gram support these findings. Zinc spray at bolting stage might have boosted the accumulation of assimilates in the seed, resulting in maximum 1000 seed weight as compared to later stages of zinc application.

Comment [A5]: Please do post hoc analysis which makes the result more clear if there is significant change in interaction there is no need for mentioning on individual factorial level

Germination percentage

In accordance with (Table 1), germination percentage of harvested seeds was significantly influenced by zinc levels, stages of application and their interaction. Among zinc levels, there was a significant increase in germination (80.92 %) up to Zn₃ (2g l⁻¹) and minimum (70.83 %) was recorded in Zn₁ (Water spray) (fig.1). However, in case of stages of application, maximum germination (80.62 %) was found during first stage of foliar application S₁ (Bolting stage) and foliar application at subsequent stages led to decrease in germination of the harvested seeds and minimum (74.19 %) was registered in S₃ (Bolting + 14 days + 28 days after first application). Whereas, in interaction, maximum germination (85.75 %) was observed in interaction Zn₃S₁ (2g l⁻¹ foliar spray at bolting stage) while minimum germination (70.75 %) was found in Zn₁S₂ (Water spray at bolting + 14 days after first application) and Zn₁S₃ (Water spray at bolting + 14 days + 28 days after first application) which was at par with Zn₁S₁. The increased germination of harvested seeds may be attributed to zinc's involvement in numerous catalytic activities and the breakdown of complex chemicals into their simplest form, such as glucose, amino acids and fatty acids. Zinc is also responsible for the aggregation of food molecules such as proteins and carbohydrates, as well as their rapid transport from source to growing seed, resulting in increased germination of harvested seeds (Shkolnik and Abdurashitov, 1958). Yoganand (2001) in bell pepper and Kiran (2006) in brinjal also noticed.

Seedling length (cm)

It is apparent from the data (Table 1) that there was a significant impact of zinc levels, stages of application and treatment combinations on seedling length. In case of zinc levels, maximum seedling length (22.13 cm) was observed in Zn₃ (2g l⁻¹) with a linear increase up to this level and at higher application of zinc (3g l⁻¹), seedling length was decreased significantly, whereas minimum seedling length (19.93 cm) was recorded in Zn₁ (Water spray). Among stages of application, maximum seedling length (21.75 cm) was noticed in S₁ (Bolting stage) while minimum (20.98 cm) was found in last stage of foliar application i.e.S₃ (Bolting + 14 days + 28 days after first application). Amongst treatment combinations, significantly maximum seedling length (24.20 cm) was registered in Zn₃S₁ (2g l⁻¹ zinc application at the time of bolting) and minimum (19.64) was obtained in Zn₁S₁ (Water spray at the time of bolting) being at par with Zn₁S₂ and Zn₁S₃. The increased seedling length could be attributed to higher metabolic activity of indole acetic acid and auxins (Krishnasamy, 2003), which could have resulted in rapid cell division and cell elongation, resulting in increased seedling length. The results coincide with the findings of Kiran (2006) in brinjal.

Seedling dry weight (mg)

According to data (Table 1), the zinc levels, stages of foliar application and their interaction significantly influenced the seedling dry weight. A continuous increase in the seedling dry weight (10.16 mg) was observed up to Zn₃ (2g l⁻¹) and there after a decrease was recorded in Zn₄ however, minimum seedling dry weight (9.16 mg) was recorded in Zn₁ (Water spray). In stages of application, maximum seedling dry weight (10.04 mg) was obtained in S₁ (Bolting stage) and afterwards, there was a sharp reduction in it having minimum seedling dry weight (9.46 mg) in S₃ (Bolting + 14 days + 28 days after first application). On the other hand, among the interactions, maximum seedling dry weight (11.31 mg) was recorded in Zn₃S₁ (2g l⁻¹ ZnSO₄ at the time of bolting) which was at par with Zn₂S₂ and Zn₄S₂ and minimum (9.12 mg) was observed in Zn₁S₃ (Water spray done at the time of bolting + 14 days + 28 days after first application) which was at par with Zn₁S₁ and Zn₁S₂ along with majority of other treatment combinations. Maximum seedling dry weight may be due to longer seedling length and lipid consumption via the glyoxylate cycle, which is the primary pathway that elevated the faster growth and seedling development to reach autotrophic stages sufficiently early and produce relatively more dry weight (Jayaraj, 1997). In brinjal, the results concur with the studies conducted by Kiran (2006).

Seed Vigour Index-I

Examination of the data (Table 1) revealed that seed vigour index-I was significantly affected by the zinc levels, stages of application and their interaction. Among different zinc levels, maximum SVI-I (1798.58) was observed in Zn₃ (2g l⁻¹) up to which there was uncurvingly increase in SVI-I and after that there was a decline in SVI-I and recording minimum (1411.02) in Zn₁ (Water spray). In stages of application, maximum SVI-I (1761.72) was obtained in S₁ (Bolting stage) and at later stages there was reduction in SVI-I with minimum (1557.37) being observed in S₃ (Bolting + 14 days + 28 days after first application). On the other hand, among treatment combinations, maximum SVI-I (2075.08) was recorded in Zn₃S₁ (2g l⁻¹ at the time of bolting) which was superior among all the combinations, however minimum (1394.25) was found in Zn₁S₁ (Water spray at the time of bolting) which was at par with Zn₁S₂ and Zn₁S₃. The findings are similar to that of Kiran (2006) in brinjal. The highest seed vigour index-I could be owing to the fact that SVI-I is a derived parameter as SV-I was determined by multiplying germination percentage and seedling length (cm), therefore as germination percentage and seedling length increased, so did the seed vigour index-I.

Table 1. Effect of foliar application of zinc, stages of application and their interaction on 1000 seed weight (g), germination % seeds, seedling length (cm), seedling dry weight(mg), seed vigour index-I in radish cv. Japanese White

Comment [A6]: Prepare more clear table


Factors	1000 Seed weight (g)	Germination %	Seedling length (cm)	Seedling weight (mg)	dry weight (mg)	Seed vigour index-I
Zn ₁	12.28	70.83(8.47)*	19.93	9.16		1,411.02
Zn ₂	13.19	78.17(8.89)	21.61	9.97		1,689.37
Zn ₃	13.55	80.92(9.05)	22.13	10.16		1,798.58
Zn ₄	13.31	78.75(8.93)	21.63	10.04		1,703.85
CD_{0.05}	0.06	0.07	0.47	0.04		40.00
Stages of application (S)						
S ₁	13.20	80.62(9.03)	21.75	10.04		1,761.72
S ₂	13.05	76.69(8.81)	21.24	9.99		1,633.03
S ₃	12.99	74.19(8.67)	20.98	9.46		1,557.37
CD_{0.05}	0.05	0.06	0.41	0.04		34.65
Interaction						
Zn × S						
Zn ₁ S ₁	12.29	71.00(8.48)*	19.64	9.18		1,394.25
Zn ₁ S ₂	12.28	70.75(8.47)	19.97	9.16		1,412.12
Zn ₁ S ₃	12.28	70.75(8.47)	20.17	9.12		1,426.71
Zn ₂ S ₁	12.68	82.75(9.15)	21.34	9.61		1,765.88
Zn ₂ S ₂	13.44	77.00(8.83)	22.23	10.83		1,712.62
Zn ₂ S ₃	13.45	74.75(8.70)	21.26	9.45		1,589.61
Zn ₃ S ₁	14.36	85.75(9.31)	24.20	11.31		2,075.08
Zn ₃ S ₂	13.26	81.50(9.08)	20.82	9.33		1,706.78
Zn ₃ S ₃	13.02	75.50(8.75)	21.37	9.83		1,613.89
Zn ₄ S ₁	13.47	83.00(9.16)	21.82	10.04		1,811.68
Zn ₄ S ₂	13.23	77.50(8.86)	21.95	10.64		1,700.58
Zn ₄ S ₃	13.22	75.75(8.76)	21.12	9.43		1,599.28
CD_{0.05}	0.10	0.12	0.82	0.72		69.29

*Figures in parenthesis represent the square root transformed values

<u>Crop stages</u>	<u>Zn dose</u>	<u>Parameter 1</u>	<u>Parmtr 2.</u>	<u>So on.....</u>
<u>S1</u>	<u>Z1</u>			
	<u>Z2</u>			
	<u>Z3</u>			
	<u>Z4</u>			
<u>S2</u>	<u>Z1</u>			
	<u>Z2</u>			
	<u>Z3</u>			
	<u>Z4</u>			
<u>s3</u>	<u>Same as above</u>			
<u>Factor s</u> <u>Factor Zn</u> <u>SX Zn</u>	<u>Mention C.d accordingly</u>			

Comment [A7]: Prepare table in this format include DMRT

UNDER PEER REVIEW



Comment [A8]: No need to make graph, or only make graph with clearer picture include standard error or Dmrt

Fig 1 :Germination percentage of harvested seeds

Seed Vigour Index- II

Analysis of the data (Table 2) depicted a significant difference in seed vigour index-II when zinc was sprayed at different levels and stages. A linear increase in SVI-II (822.64) was discernible with zinc up to 2g l^{-1} (Zn_3) and then there was a decline in SVI-II at Zn_4 and minimum value (648.49) was obtained in Zn_1 (Water spray). Similarly, among different stages of application, maximum SVI-II (810.78) was recorded in S_1 (Bolting stage) and then there was a decrease in SVI-II with every stage of zinc application with minimum SVI-II (701.90) in S_3 (Bolting + 14 days + 28 days after first application). However, in case of interactions, maximum (966.77) SVI-II was observed in Zn_3S_1 (2g l^{-1} at the time of bolting) and minimum (645.07) in observed in Zn_1S_3 (Water spray at bolting + 14 days + 28 days after first application) had statistical similarity with Zn_1S_1 and Zn_1S_2 . The highest seed vigour index-II could be because SVI-II is also a derived metric determined by multiplying germination (%) by seedling dry weight (mg). Thus, when the germination (%) and seedling dry weight increased, similar trend in SVI-II was obvious.

Electrical Conductivity (dS m^{-1})

In insight into the data (Table 2) showed that zinc levels, stages of application and their interaction had a significant influence on electrical conductivity. In case of different levels of zinc, minimum electrical conductivity (0.023 dS m^{-1}) was noticed in Zn_3 (2 g l^{-1}) and thereafter there was an increase in electrical conductivity with zinc applied @ 3 g l^{-1} while maximum EC (0.041 dS m^{-1}) was recorded in Zn_1 (Water spray). Regarding different stages of application, minimum EC (0.024 dS m^{-1}) was obtained in S_1 (Bolting stage) and after that EC of the seed was found to increase with every subsequent stage of application and maximum value (0.036 dS m^{-1}) was recorded in S_3 (Bolting + 14 days + 28 days after first application). Among different treatment combinations, minimum EC (0.017 dS m^{-1}) was registered in Zn_3S_1 (2 g l^{-1} at the time of bolting) whereas, maximum EC (0.041 dS m^{-1}) was recorded in Zn_1S_1 (Water spray at the time of bolting), Zn_1S_2 (Water spray at the time of bolting + 14 days first application) and Zn_1S_3 (Water spray at the time of bolting + 14 days + 28 days after first application). The electrical conductivity of seed is greatly dependent on seed vigour. The electrical conductivity of the seed is inversely proportional to its vigour, as seed vigour is greatest when electrical conductivity is lowest. The decrease in EC with zinc administration can be explained by the fact that this element is known to promote cell wall integrity, reducing solute leaking from the seed. In brinjal, the results concur with the findings of Kiran (2006).

Accelerated Ageing Test

From the data (Table 2), it is extrapolated that there were significant differences among different zinc levels, stages of application and their interaction for this parameter. Among zinc levels, maximum germination (74.33 %) for the parameter was obtained in Zn_3 (2 g l^{-1}) which was rectilinear maximum and minimum germination (70.17 %) was recorded in Zn_1 (Water spray). Whereas, in case of stages of application, maximum germination (74.93 %) was obtained in S_1 (Bolting stage) and thereafter there was a reduction in germination (%) of seed with every consecutive stage of application with minimum germination (71.31 %) being observed in S_3 (Bolting + 14 days + 28 days after first application). The improvement in germination (%) of the accelerated ageing test could be related to role of zinc increasing cell membrane integrity. As a result, the seed's vitality was increased, which lasted longer during the ageing process and resulted in a greater germination percentage after the accelerated ageing test. Artificial ageing damages DNA and mRNA, resulting in biochemical degradation of the reserved material in the seed, which increases seed vigour (Murthy et al., 2003).

Cold Test

An insight into the data (Table 2) revealed a significant effect of different levels of zinc, stages of application and their interaction on this parameter. Maximum germination (78.92 %) in cold test was recorded in Zn_3 (2 g l^{-1}) up to which there was linear increase and reduced as concentration of zinc application was increased to 3 g l^{-1} while minimum germination (70.67 %) was observed in Zn_1 (Water spray). Among stages of application, maximum germination (75.37 %) was obtained in S_1 (Bolting stage) and then, it declined with every subsequent stage reaching minimum (73.42 %) at S_3 (Bolting stage + 14 days + 28 days after first application). Amongst treatment combinations, maximum germination (81.00%) in cold test was found in Zn_3S_1 (2 g l^{-1} with foliar zinc at the time of bolting) and minimum germination (70.50%) was observed in Zn_1S_1 (Water spray at the time of bolting) and Zn_1S_3 (Water spray at the time of bolting + 14 days + 28 days after first application). The cold test was created to mimic harsh field conditions and assess the capacity of seeds to emerge. The germination percentage of Zn_3S_1 was higher because foliar application of zinc during the bolting stage might have increased photosynthates transport, available nutrients in seeds and cell membrane integrity, thereby preserving the structural inclination of macromolecules and ion transport systems, resulting in greater vigour and germination (Dang et al., 2010).

Table 2. Effect of foliar application of zinc, stages of application and their interaction on seed vigour index-II, electrical conductivity (dSm^{-1}), accelerated ageing test, cold test in radish cv. Japanese White

Factors	Seed vigour	Electrical	Accelerated ageing	Cold Test
---------	-------------	------------	--------------------	-----------

	index-II	Conductivity (dSm ⁻¹)	Test	
Zinc levels				
Zn ₁	648.49	0.041	70.17(8.44)*	70.66(8.47)*
Zn ₂	778.36	0.025	73.75(8.65)	74.75(8.70)
Zn ₃	822.64	0.023	74.33(8.68)	78.91(8.93)
Zn ₄	789.65	0.028	72.67(8.58)	73.41(8.62)
CD_{0.05}	12.81	0.001	0.05	0.08
Stages of application (S)				
S ₁	810.78	0.024	74.93(8.71)	75.37(8.74)
S ₂	766.67	0.029	71.94(8.54)	74.81(8.70)
S ₃	701.90	0.036	71.31(8.50)	73.12(8.61)
CD_{0.05}	11.10	0.001	0.04	0.07
Interaction				
Zn × S				
Zn ₁ S ₁	651.95	0.041	70.50(8.45)	70.50(8.46)
Zn ₁ S ₂	648.43	0.041	70.00(8.43)	71.00(8.48)
Zn ₁ S ₃	645.07	0.041	70.00(8.43)	70.50(8.46)
Zn ₂ S ₁	794.40	0.018	76.00(8.77)	76.75(8.82)
Zn ₂ S ₂	834.30	0.025	73.25(8.62)	74.00(8.66)
Zn ₂ S ₃	706.38	0.034	72.00(8.54)	73.50(8.63)
Zn ₃ S ₁	966.77	0.017	78.25(8.90)	81.00(9.05)
Zn ₃ S ₂	759.15	0.023	73.00(8.60)	80.75(9.04)
Zn ₃ S ₃	741.98	0.031	71.75(8.53)	75.00(8.71)
Zn ₄ S ₁	830.00	0.019	75.00(8.72)	73.25(8.62)
Zn ₄ S ₂	824.78	0.028	71.50(8.51)	73.50(8.63)
Zn ₄ S ₃	714.17	0.038	71.50(8.51)	73.50(8.63)
CD_{0.05}	22.19	0.001	0.08	0.14

*Figures in parenthesis represent the square root transformed value

Conclusion

Among different zinc levels, Zn₃(2g l⁻¹) was considerable superior to other levels for majority of seed quality parameters. In case of stages of application, S₁ (Bolting stage) showed its superiority above all other stages. In treatment combinations, Zn₃S₁ (2g l⁻¹ at the time of bolting) was superior over other treatment combinations for most of seed quality parameters namely, 1000 seed weight (g), germination percentage of harvested seeds, seedling length (cm), seedling dry weight (mg), seed vigour index-I, seed vigour index-II, accelerated ageing test, cold test.




Fig 2 : Foliar application of zinc

References

- Acquaah G 2002. Horticulture: Principles and Practices. 2nd ed. Upper Saddle River, USA: Prentice Hall. 822p.
- AnupritaH, Jadhav SR, Dalal RD and Rajeshwari P 2005. Effect of micronutrients on growth and flower production of gerbera under poly house condition. *Adv. Plant Sci.***18**: 755-758.
- ArifM, ChohanMA, Ali S, Gul, R and Khan S 2006. Response of wheat to foliar application of nutrients. *J. Agric. Biol. Sci.* **1**: 256-259.
- Brady NC and Weil RR2002. The Nature and Properties of Soils. 13th ed. Upper Saddle River, USA: Prentice Hall **95**: 393-394
- Cakmak I 2008. Enrichment of cereal grains with zinc: agronomic or genetic bio fortification. *Plant Soil***302**: 1-17.
- Dang HR, Li Y, Sun X, Zhang L Y 2010. Absorption, accumulation and distribution of zinc in highly-yielding winter wheat. *Agric. Sci. China* **9**: 965-973.
- ISTA 2016. International Seed Testing Association Manual. www.seedtesting.org (Assessed on 18th May, 2020)

Comment [A9]: Include recent papers from 2021 to 2024

- Jayaraj T 1997. Study of the effect of plant protection chemicals on seed quality in sesame(*Sesamum indicum* L.) cv. KRR 2 and TMV 3. M Sc Thesis, Tamil Nadu Agriculture University, Coimbatore. 76p.
- Kiran, J2006. Effect of fertilizer, biofertilizer and micronutrients on seed yield and quality of brinjal (*Solanum malongena*L.). M Sc Thesis, Department of Seed Science and Technology, University of Agricultural Sciences, Dharwad. 78p.
- Krishnasamy V 2003. Seed pelleting principles and practices. *ICAR short course on seed hardening and pelleting technologies for rainfed / garden land ecosystems*. Tamil Nadu Agriculture University, Coimbatore. 96p.
- Kumar H, Dogra P and Yadav V 2014. Effect of foliar application of N and Zn on growth and yield of cauliflower (*Brassica oleracea* var. *botrytis* L.) cv. Snowball-16. *Agric. Sustain. Dev.* **2**: 56-58.
- Lakshami EJ, Babu PVR, Reddy CP, Umamaheshwari P and Reddy AP K 2017. Effect of foliar application of secondary nutrients and zinc on growth and yield of black gram. *Int. J. Chem. Stud.***5**: 944-947.
- Masuthi D, Vyakaranahal BS and Deshpande V K 2009. Influence of pelleting with micronutrients and botanical on growth, seed yield and quality of vegetable cowpea. *Karnataka J. Agric. Sci.***22**: 898-900.
- Murthy UM, Kumar PP and Sun W Q 2003. Mechanisms of seed ageing under different storage conditions of *Vigna radiata* (L.) Wilczek: lipid peroxidation, sugar hydrolysis, maillard reactions and their relationship to glass state transition. *J. Exp. Bot.* **54**: 1057-1067.
- Panse V G and Sukhatme P V 2000. *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research, New Delhi, India. 157-165p.
- Pariari A, Khan S and Imam M N 2009. Influence of boron and zinc on increasing productivity of fenugreek seed (*Trigonella foenum graecum* L.) *J. Crop and Weed***5**: 57-58.
- Sheoran O P 2006. *Online Statistical Analysis Tool (OPSTAT)*. <http://14.139.166.opstat/> (Assessed on 3rd July, 2020).
- Shkolnik MI and Abdurashitov S A 1958. Influence of microelements on synthesis and translocation of carbohydrates. *Plant Physiol. USSR***5**: 393-399.
- Trivedi A P 2013. Effect of soil and foliar application of zinc and iron on the yield and quality of onion (*Allium cepa* L). *Bangladesh J. Agric. Res.***38**:41-48.
- Wissuwa M, Ismail AM and Graham R W 2008. Rice grain zinc concentrations as affected by genotype, native soil-zinc availability and zinc fertilization. *Plant Soil***306**: 37-48.
- Yoganand DK 2001. Effect of mother plant nutrition and growth regulators on plant growth, seed yield and quality of bell pepper cv. California Wonder. M Sc Thesis, University of Agricultural Sciences, Dharwad. 77p.