

# Original Research Article

## Genetic studies of heterosis and combining ability for foliar disease, yield and its contributing characters in faba bean (*Vicia Faba L.*)

### ABSTRACT

The present study was carried out in the farm of Sakha Agricultural Research Station, Kafr -El Sheikh, Egypt, during 2019/20, 2020 /21 seasons to produce seed hybrids ( $F_1$  seeds) which evaluated in 2021/22 season. Four parental genotypes namely; Almany, Giza 843, Misr 3 and Najeh were used as testers, while four material genotypes namely; Line 1, Line 2, Line 3 and Line 4 used as lines which crossed according to Line X tester mating system under insect free cages in the first and second seasons. In the third season the resulted  $F_1$ ,<sup>s</sup> sixteen crosses along with their parental genotypes were sown in a randomized complete blocks design with three replications during 2021/22 season. The data subjected to line x tester analysis. Highly significant of genotypes, parents and crosses were found for all traits. The magnitude of genotypes or any of its components i.e., parents, crosses and parents vs. crosses were several times large than corresponding mean squares of error. Mean squares of parent's vs. crosses as an indication to average heterosis over all crosses appeared to be highly significant heterotic effects for almost all traits. Partitioning the crosses mean squares to its components i.e., lines, testers and lines x testers revealed that, lines and testers mean squares seems to be not significant for all traits. While, lines x testers mean squares found to be highly significant for all traits. The ratio of  $\sigma^2_{gca}/\sigma^2_{sc}$  was less than unity for all traits, indicating that the non-additive gene effects were of greater important in the inheritance of these traits. The crosses; Line 1 x Tester 1, Line 1 x Tester 2, Line 2 x Tester 2, Line 3 x Tester 1, Line 3 x Tester 4, Line 4 x Tester 1 and Line 4 x Tester 2 had significant and /or highly significant mid and better parents heterotic effects for many important traits a result of over-dominance in approximately all studied traits. Tester 4 considered as good combiner parent for plant height, No of pods/plant, No of seeds /plant, seed yield /plant and curd protein percentage. The cross; Line 1 x Tester 2 expressed significant ( $\hat{S}_{ij}$ ) in favorable direction for rust disease reaction, flowering date, plant height and crude protein percentage. The cross; Line 2 x Tester 3 had significant ( $\hat{S}_{ij}$ ) in favorable direction for chocolate spot diseases reaction, plant height and No. of branches/plant and the cross; Line 4 x Tester 2 had significant ( $\hat{S}_{ij}$ ) in positive direction for No of branches/plant, No of pods/plant, No of seeds/plant, seed yield /plant and carbohydrate percentage. Broad sense heritability ranged from 24.6% for No. of seeds/plant to 77.90% for 100-seed weight. While, narrow- sense heritability ( $h^2$ ), ranged between 0.68% for 100-seed weight to 20.28% for chocolate spot diseases reaction. The predicted genetic advance ranged from 0.18 % for 100- seed weight to 6.92 % for chocolate spot diseases reaction.

Key words: Faba bean, heterosis, combining ability, heritability, genetic advance.

### INTRODUCTION

Faba bean is a major winter crop in the Mediterranean area, valued for its high protein content ranged from (24 – 34 %). It is an important crop for human animal nutrition in Egypt [1], as well as for improving soil fertility through nitrogen fixation. Also, play a key role as a break crop in cereal rotation system. In the 2023/24 season, Egypt cultivated approximately 130,000 feddan of faba bean, with an average yield of 10.9 ardab/feddan, resulting in total production of about 374,000 tons. However, the country's self-sufficiency rate is only 33.6 % [2]. To bridge the gap between production and

consumption, it is crucial to develop new high-yielding cultivars and implement proper culture practices.

Combining ability analysis is a useful tool for breeders to identify the most effective parent combinations that can be hybridized to take advantage of heterosis or create synthetic varieties. Bound [3] utilized the relative significance of general combining ability and specific combining ability effects to select parents for hybrid varieties. Induced mutagenesis can also be employment to increase the genetic diversity of quantitatively inherited traits.

The key factor in selecting appropriate breeding procedures is understanding the type of gene action of quantitative traits in faba bean to enhance seed yield and improve yield characteristics. Various genetic analysis methods such as half and full diallel matings, six-population models, and line x tester analysis have been utilized to determine the most effective breeding scheme. Simply selection superior genotype based on yield alone is ineffective due to the complexity of yield traits, and their interactions with the environment [4]. Successful breeding programs require genetic diversity within the plant collection to enable the section of desirable gene combinations [5].

A successful breeding program for faba bean genotypes depending on the genetic variation in the population. Reliable estimates of genetic and environmental variations are essential to determine heritability and predict genetic advance from selection. This study amide to assess variability, hetrosis, general and specific combining abilities, heritability, and genetic advance for foliar diseases, earliness and yield and yield components traits in sixteen faba bean crosses, via line x tester analysis.

#### MATERIALS AND METHODS

The present study was carried out under field condition in the farm of Sakha Agricultural Research Station, Kafr El-Sheikh, Egypt during the growing winter seasons of 2019/20, 2020/21 to produced hybrid seeds (F<sub>1</sub> seeds) which evaluated in 2021 /22 season. Four parental genotypes namely; Almany, Giza 843 Misr 3 and Najeh were used as testers, while four maternal genotypes namely; Line 1, Line 2, Line 3 and Line 4 used as lines were which crossed according to lines x testers mating system under insect free cages in the first season of (2019/20). In the second season (2020/21) the parental genotypes were sown again under wire cages and re-hybridized as done as in the first season to obtain more hybrid seeds. The resulted F<sub>1</sub>' s along with their parental genotypes were sown in a randomized complete blocks design with three replications during 2021/22 season. Flowers of F<sub>1</sub> plants grown under wire cage were manually tripped to ensure good pollination. Seeds were sown in single seeded hills, 20 cm apart, each plot consisted of one row for parents as well as their F<sub>1</sub>' s. The row was 3 meters long and 60 cm in between. The choice of parents was based on:

a- Genetic diversity, b- Differences in growth habit and disease reactions and c- Differences in yielding ability. The pedigree, disease reactions, agronomical characters and yielding level of parental genotypes are presented in Table (1).

**Table (1):** Names, pedigrees', disease reactions and agronomical characters of the parental faba bean genotypes used in this investigation.

Genotype	Pedigree	Agronomical characters	
		Flowering date	Characteristics
<u>Lines</u>			
Line 1 520/283/2015	Cross (Giza 843 x Misr 1)	Medium	Resistant to foliar disease and Tolerant to <i>Orobanche</i>
Line 2 313/453/2015	Cross (Giza 2 x Misr 1)	Early	Resistant to foliar disease and Tolerant to <i>Orobanche</i>
Line 3 249/230/2015	Cross (Giza 843 x Giza 2)	Medium	Resistant to foliar disease and Tolerant to <i>Orobanche</i>
Line 4 461/345/2015	Cross (Sakha 3 x Giza 2)	Medium	Moderate resistant to foliar disease and Tolerant to <i>Orobanche</i>
<u>Testers</u>		I	
Almany	Introduction from Germany	Medium	Resistant to foliar disease and high yield
Misr 3	L667 x (Cairo 241	Early	Tolerant to <i>Orobanche</i>

Giza 843	x Giza 461) 561/2076/85 x 461/845/83	Medium	Tolerant to <i>Orobanche</i>
Najhe	INRAT, Tunisia	Medium	Tolerant to <i>Orobanche</i>

Data were taken on the basis of individual plant as follows:

- |                                     |                           |                          |
|-------------------------------------|---------------------------|--------------------------|
| 1- Chocolate spot disease reaction. | 2- Rust disease reaction. | 3- Flowering date (day). |
| 4- Plant height (cm).               | 5- No of branches/plant.  | 6- No of pods /plant.    |
| 7- No. of seeds/plant.              | 8- 100-seed weight.       | 9- Seed yield / plant.   |
| 10- Crude Protein %.                | 11- Carbohydrates %.      | 12- Fibers %.            |

Reaction to foliar diseases was recorded on mid-February and mid-March for chocolate spot and rust diseases, respectively; according to the disease scales suggested by **Bernieret al [7]** as presented in Table (2).

**Table (2):** Rating scale for chocolate spot and rust diseases reactions.

Rate	Chocolate spot scale
1	Highly resistant, no disease symptom.
3	Resistant, few small discrete lesions.
5	Moderately resistant, some coalesced lesions with some defoliation.
7	Susceptible, large coalesced lesions, 50% defoliations, some dead plants.
9	Highly susceptible, extensive lesions on leaves, stems and pods, severe defoliation, heavy sporulation, death of more than 80% of plants.
Rust scale	
1	High resistant, no pustules or very small non- sporulation flecks
3	Resistant, few scattered pustules covering less than 1% of the leaf area, and few or no pustules on stem.
5	Moderately, resistant, pustules common on leaves covering 1-4% of leaf area, little defoliation and some pustules on stem.
7	Susceptible, pustules very common on leaves covering 4-8% of leaf area, some defoliation and many pustules on stem.
9	Highly susceptible, extensive pustules on leaves, petioles and stem covering 8—10% of leaf area, many dead leaves and several defoliation.

The statistical procedure was done according to the regular analysis of variance of randomized complete blocks design as outlines by **Cochran and Cox [8]**. The analysis of variance was done depending on the mean of the individual plant basis. The source of variation and degrees of freedom are shown in Table (3).

**Table (3):** Mean squares (MS) and expected mean of square (EMS)

S.O.V	d f	MS	EMS
Replication	r-1	M r	
Genotype (G)	(g-1)	MS g	$\sigma^2 e + r\sigma^2 g$
Error	(r-1) (g-1)	MS e	$\sigma^2 e$

Where: r is the number of replications; g is the number of genotypes;  $\sigma^2 g$  and  $\sigma^2 e$  refer to genotypic and error variances, respectively. The difference between any two means was tested according to the least significant difference (L.S.D) at both 5% and 1% level of significance as follows:

$$L.S.D : P \leq 0.05 = t_{0.05} (d.f) \times Sd$$

$$P \leq 0.01 = t_{0.01} (d.f) \times Sd$$

Where: r is the number of replications and

$MS_e$ : is the mean square of error

According to **Kempthorne [9]**, the variance of general and specific combining abilities may be computed from the covariance of full sib (F.S) and half sib (H.S) families.

Where:

$$\sigma^2 gca = Cov H.S = 1 + F/4 \times \sigma^2 A, \text{ with } F=1, \sigma^2 A = 2\sigma^2 gca$$

$$\sigma^2 sca = \{1 + F/2\}^2 \sigma^2 D, \text{ with } F=1, \sigma^2 D = \sigma^2 sca$$

Where F is the coefficient of inbreeding that ranged from 0 to 1 with open pollinated varieties to completely pure lines, respectively.

**Table (4): Line X tester analysis of variance including parents.**

S.O.V	df	MS	EMS
Replication	r-1		
Genotype (G)	(g-1)		
Parents	(P-1)		
Parent's vs. crosses	1		
Crosses	(C-1)		
Lines	(l-1)	M <sub>l</sub>	$\sigma^2_e + r \sigma^2_{2lt+r\sigma^2_l}$
Testers	(t-1)	M <sub>t</sub>	$\sigma^2_e + r \sigma^2_{lt+r\sigma^2_t}$
Lines x testers	(l-1) (t-1)	M <sub>lt</sub>	$\sigma^2_e + r \sigma^2_{lt}$
Error	(r-1) (g-1)	M <sub>e</sub>	$\sigma^2_e$

**Where:** the MS due to lines(M<sub>l</sub>) and testers (M<sub>t</sub>) were tested against MS due to lines x testers (M<sub>lxt</sub>) and the latter is, in turn, tested against MS due to error (M<sub>e</sub>) [9].

The genetical parameters; genotypic variance (V<sub>g</sub>), phenotypic variance (V<sub>ph</sub>) were computed according to the formula suggested by **Burton [10]**. For each character, PCV% and GCV% were computed based on the methods given by **Burton [10]**. Heritability in broad sense (H) was calculated according to **lush Jay [11]**. The range of heritability was categorized as suggested by **Johnson et al [12]**, where : low (< 30%), moderate (30-60%) and high (> 60%).

Narrow sense heritability (h<sup>2</sup>) estimates were calculated as **Acquaah [13]**. According to **Stansfield [14]**, the classification of narrow-sense heritability was as follows: Low (< 20%), moderate (20-50%) and high (> 50%). For each character, expected genetic advance (Ga) and predicted genetic advance (Ga%) were estimated in accordance with the methods illustrated by **Fehr [15]** assuming selection of the superior 5% of the genotype plants and the narrow-sense heritability (h<sup>2</sup>) was used to calculate the expected genetic advance (Ga). The range and frequency is as follows: low (less than 10), moderate (10-20) and high (more than 20).

#### **Seed Quality:**

All seed properties were carried out at Sakha Seed Technology Research section as Follow.

##### **1-Crude Protein (%)**

Tested seeds were ground to a fine powder to pass through 2 mm mesh and used to determine the crude protein percentage according to methods of **A.O.A.C [16]**.

**3-Carbohydrate contents:** were calculated by difference: [100-(protein + oil+ ash+fiber +moisture)]

**2-Fiber content:** was determined according to **A.O.A.C [16]**.

#### **RESULTS AND DISCUSSION**

##### **Analysis of variance:**

Highly significant mean squares of genotypes, parents and crosses were found for all studied traits (Table 5), except for parents mean square of No. of branches/plant. The magnitude of genotypes or any of its components i, e., parents, crosses and parents v s crosses were several times larger than corresponding mean squares of error. This would indicate the wide genetic variability between genotypes, therefore, it become statistically valid for the required diversity for the success of the planned crosses. Mean squares of parent vs. cross as an indication to average heterosis over all crosses appeared to be highly significant heterotic effects for all traits, except for crude protein percentage, carbohydrate percentage and fiber percentage. The latest three traits did not significant heterotic effects over all crosses. Partitioning the crosses mean squares to its three components i.e., lines, testers and lines x testers revealed that, lines and testers mean squares seems to be not significant for all the studied traits, except **testers** mean squares for plant **height and** carbohydrate %, and

where it's at least significant. Regarding to lines x testers mean squares, it's found to be highly significant for all studied traits in view, except, 100-seed weight, where it was only significant. Results presented in Table (5), revealed not significant  $\sigma^2_{gca}$  and  $\sigma^2_{sca}$  for all studied traits.  $\sigma^2_{gca}/\sigma^2_{sca}$  ratio was less than unity for all studied traits indicating the **non** additive types of gene action controlling these traits.

**Table (5): Analysis of variance for disease reaction, yield and its components of faba bean lines x testers crosses in the F<sub>1</sub> generation.**

S.O.V	d.f	Chocolate spot diseases reaction	Rust diseases reaction	Flowering date	Plant height (cm)	No. of branches / Plant	No. of pods/ plant	No. of seed/plant	Seed Yield/ plant	100-seed weight	crude protein (%)	Carbo-hydrate (%)	Fiber (%)
Rep	2	0.10	0.40	4.90	19.30	0.70*	3.80	40.00	55.20	41.10	0.70	1.10	0.30
Genotypes	23	2.10**	4.40**	53.1**	448.00**	3.30**	268.90**	2447.90**	1586.60**	130.20	22.3**	27.9**	3.50**
Parents (P)	7	3.30**	3.30**	28.20**	1171.70**	0.60	143.20**	1355.80**	975.8**	64.30**	26.4**	32.3**	6.40**
P vs C	1	8.50**	39.90**	156.30**	270.00**	34.50*	1560.50**	10534.00**	7560.80*	134.00**	14.00	13.60	3.50
Crosses (C)	15	1.20**	2.60**	57.8**	122.20**	2.40**	241.50**	2418.50**	1473.30**	160.80**	20.9**	26.8**	2.20**
Lines (L)	3	1.20	3.30	66.70	162.40	3.90	100.10	1017.30	993.50	32.20	23.90	13.00	0.50
Testers (T)	3	2.00	2.80	80.60	269.90*	0.40	333.00	4313.00	2020.30	104.30	44.50	88.9**	5.40
L x T	9	0.90**	2.30**	47.20**	59.50**	2.60**	258.10**	2254.1**	1450.9**	222.50*	12.0**	10.7**	1.70**
Error	46	0.30	0.30	9.40	11.40	0.20	9.70	830.00	213.00	87.90	0.60	0.60	0.40
$\sigma^2_{gca}$		0.009	0.010	0.367	2.176	0.007	0.577	5.709	0.778	2.143	0.308	0.559	0.018
$\sigma^2_{sca}$		0.210	0.690	12.617	16.036	0.802	82.81	723.70	412.6	44.86	3.81	3.367	0.424
$\sigma^2_{gca}/\sigma^2_{sca}$		0.043	0.014	0.029	0.136	0.009	0.007	0.008	0.002	0.048	0.808	0.166	0.042

\*and\*\* significant at 0.05 and 0.01 levels of probability, respectively.

### The mean performance:

The differences between mean performance of the parental lines and testers and their F<sub>1</sub> crosses of faba bean genotypes for studied traits are shown in Table 6. Highly significant difference between genotype was found the parental Line 1, Line 2, Tester 3 (Giza 843) and Tester 4 (Najeh) recorded the lowest values for chocolate spot and Rust diseases reaction. The parental Line 2 and Tester 2 recorded the lowest days of flowering date. While, Line 2, Tester 3 (Giza 843) and Tester 4 (Najeh) registered the best values of plant height. In addition to Line 1, Line 2, Tester 3 (Giza 843) and Tester 4 (Najeh) had the highest values of number of branches/plant, number of pods /plant, number of seeds/ plant, seed yield/ plant, 100-seed weight and crude protein%. For carbohydrate%, Line 2 and Tester 3 (Giza 843) gave the highest value. While, Line 2 and Tester 2 (Misr3) showed the highest values for fiber percentage.

The crosses; Line 1xTester 1, Line 1xTester 3, Line 2 x Tester 3 and Line 4xTester 1 were resistant for chocolate spot and rust diseases reaction where showed the lowest values of severity. In addition to the crosses; Line 1x Tester 2 Line 3xTester 2, Line 3xTester 3 and Line 4 xTester 1 were the earliest crosses of flowering date (40.00, 41.67 days, 43.33 and 43.33 respectively). The crosses; Line 1xTester 2 and Line 1xTester 4 had the highest values of plant height. For No. of branches/ plant, No. of pods/ plant, No. of seeds/ plant, seed yield/ plant and 100 -seed weight the crosses; Line 1x Tester 4, Line 2 xTester 2, Line 2xTester 4, Line 3xTester 1, Line 3xTester 3, Line 3xTester, Line 4xTester 2 and Line 4 x Tester 4) were the highest yielding crosses. Also, five crosses gave the desirable values of crude protein% ranged from (29.67% - 33.00%). On the other hand, seven crosses for carbohydrate % had the highest values ranged from (46.55% - 51.00%).

**Table (6). Mean performance of parents and their crosses of faba bean genotypes for studied traits.**

Crosses	Chocolate spot diseases reaction	Rust diseases reaction	Flowering date	Plant height (cm)	No. of branches/ Plant	No. of pods/ plant
Line 1	3.00	2.00	54.45	111.67	4.23	30.09
Line 2	4.00	3.00	45.11	128.33	3.86	25.53
Line 3	3.00	4.00	50.45	100.07	3.08	21.22

Line 4	5.00	5.00	50.00	117.38	2.91	16.65
Tester 1 (Almany)	5.00	4.00	53.33	110.50	3.10	20.30
Tester 2 (Misr 3)	4.17	3.43	48.33	112.49	3.09	25.30
Tester 3 (Giza 843)	3.33	2.80	53.33	155.00	3.67	38.55
Tester 4 (Najhe)	2.00	2.00	50.00	150.00	3.33	30.07
Line 1 x Tester 1	3.63	2.83	46.67	113.89	5.40	36.72
Line 1 x Tester 2	4.00	3.00	40.00	130.00	4.34	33.03
Line 1 x Tester 3	3.67	4.50	48.33	115.46	4.77	28.00
Line 1 x Tester 4	4.57	6.00	55.00	131.92	5.27	49.50
Line 2 x Tester 1	4.47	5.27	48.33	110.35	4.73	29.93
Line 2 x Tester 2	4.73	4.80	55.00	114.79	4.67	31.85
Line 2 x Tester 3	3.53	4.60	46.67	116.25	6.60	31.76
Line 2 x Tester 4	5.33	6.00	53.33	119.19	4.06	45.46
Line 3 x Tester 1	4.00	5.00	50.00	115.48	4.06	56.04
Line 3 x Tester 2	5.53	5.90	41.67	119.36	3.85	24.04
Line 3 x Tester 3	4.50	4.83	43.33	108.90	3.88	38.71
Line 3 x Tester 4	4.77	5.00	48.33	123.45	4.59	38.34
Line 4 x Tester 1	3.70	4.77	43.33	123.33	4.26	27.51
Line 4 x Tester 2	4.23	4.33	45.00	121.48	6.68	41.44
Line 4 x Tester 3	5.20	5.90	46.67	117.48	4.86	24.55
Line 4 x Tester 4	4.77	5.00	48.33	123.79	6.02	36.56
P LSD 0.05	0.41	0.95	3.44	7.77	1.03	2.78
P LSD 0.01	0.68	1.01	6.12	9.39	1.59	4.98
C LSD 0.05	1.09	0.83	5.89	5.97	0.59	6.15
C LSD 0.01	1.44	1.24	7.54	8.03	1.24	8.26
All LSD 0.05	0.89	0.82	5.03	5.55	0.77	5.12
All LSD 0.01	1.19	1.10	6.72	7.41	1.03	6.83

Table (6): Cont.

Crosses	No .of seed/ Plant	Seed Yield/ plant	100-seed weight	crude protein (%)	Carbo- hydrate (%)	Fiber (%)
Line 1	89.62	74.16	82.79	25.00	45.00	9.50
Line 2	67.88	57.06	84.05	22.27	50.67	12.50
Line 3	69.40	54.45	78.24	29.00	40.33	9.73
Line 4	47.75	33.52	70.17	24.73	44.00	8.60
Tester 1 (Almany)	52.46	40.20	76.05	26.00	46.00	10.20
Tester 2(Misr 3)	70.30	51.30	72.92	30.00	41.33	11.00
Tester 3(Giza 843)	112.41	88.32	78.83	29.67	44.00	8.77
Tester 4(Najhe)	89.00	69.13	77.60	30.00	41.67	7.93
Line 1 x Tester 1	99.51	81.56	82.21	27.00	46.55	9.07
Line 1 x Tester 2	87.17	73.14	83.96	33.00	42.00	10.00
Line 1 x Tester 3	81.20	63.18	77.73	28.67	45.17	7.97
Line 1 x Tester 4	147.59	121.08	82.01	31.00	42.67	10.00
Line 2 x Tester 1	84.48	72.39	85.69	23.67	51.00	10.00
Line 2 x Tester 2	91.08	83.84	91.89	25.67	44.00	9.47
Line 2 x Tester 3	82.66	63.84	77.33	27.00	46.00	8.00
Line 2 x Tester 4	143.28	82.11	58.07	30.00	42.00	9.00
Line 3 x Tester 1	154.46	129.58	84.02	27.25	47.33	10.33
Line 3 x Tester 2	64.70	50.12	77.78	25.53	45.00	8.67
Line 3 x Tester 3	109.77	85.48	77.92	27.00	46.33	8.33
Line 3 x Tester 4	110.29	96.89	87.86	32.67	38.20	11.00
Line 4 x Tester 1	74.59	59.03	79.22	28.00	46.60	9.60
Line 4 x Tester 2	113.75	88.72	77.99	25.67	47.00	9.67
Line 4 x Tester 3	60.48	51.29	84.80	26.50	47.67	9.00

<b>Line 4 x Tester 4</b>	<b>103.16</b>	<b>81.83</b>	<b>79.11</b>	<b>29.67</b>	<b>43.23</b>	<b>8.90</b>
<b>P LSD0.05</b>	<b>13.15</b>	<b>11.12</b>	<b>5.19</b>	<b>1.43</b>	<b>1.46</b>	<b>1.35</b>
<b>P LSD0.01</b>	<b>15.54</b>	<b>16.35</b>	<b>7.22</b>	<b>2.65</b>	<b>1.98</b>	<b>1.64</b>
<b>C LSD0.05</b>	<b>16.64</b>	<b>28.89</b>	<b>18.41</b>	<b>0.87</b>	<b>1.24</b>	<b>0.91</b>
<b>C LSD0.01</b>	<b>18.87</b>	<b>31.24</b>	<b>25.34</b>	<b>1.99</b>	<b>1.79</b>	<b>1.38</b>
<b>All LSD0.05</b>	<b>14.97</b>	<b>23.98</b>	<b>15.41</b>	<b>1.28</b>	<b>1.31</b>	<b>1.03</b>
<b>All LSD0.01</b>	<b>19.98</b>	<b>32.02</b>	<b>20.57</b>	<b>2.54</b>	<b>175</b>	<b>1.37</b>

### **Heterosis:**

It is a known fact that, the phenomena of heterosis (hybrid vigour) is of common occurrence in both cross and self- pollinated crops. Faba bean which is often a partially self-fertilized crop also show hybrid vigour when hybridization takes place between homozygous varieties.

The data illustrated in Table 7 show that, highly significant mid and better parents heterotic effects in the cross ; Line 1 x Tester 1 were found for rust disease reaction, flowering date, No. of branches/ plant , No. of pods /plant and No. of seeds/ plant in favourable direction a result of over - dominance as potence ratio pointed out in all cases , except rust disease reaction where partial dominance was controlled the heterotic effect in the trait of view.

Highly significant mid and better parents heterosis in the cross; line 1 x Tester 2 were detected for flowering date, plant height, No. of pods /plant, and protein % in favourable direction due to over – dominance.

Highly significant mid and better parents heterotic effects in the cross; Line 1 x Tester 4 were found for No. of branches / plant, No. of pods /plant, No. of seeds / plant and seed yield /plant in positive direction a result of over-dominance( $P > +1$ ). Highly significant mid and better parents heterosis in the cross; line 2 x Tester 2 were observed for No. of branches / plant, No. of pods /plant, No. of seeds / plant and seed yield / plant, except for No. of branches, No. of pods /plant and seed yield / plant , where the better parent heterosis were only significant , in all cases over – dominance was the main reason of heterotic effects ( $P > +1$ ).

Significant and / or highly significant mid and better parents heterosis were observed in the cross; Line 3 x Tester 1 for No. of branches / plant, No. of pods /plant, No. of seeds / plant, seed yield / plant and carbohydrate% in positive direction a result of over- dominance in all traits of view. Significant and / or highly significant mid and better parents heterosis were found in the cross; Line 3 x Tester 4 for No. of branches / plant, No. of pods /plant, No. of seeds / plant, seed yield / plant, crude protein % and carbohydrate% in positive direction due to over- dominance in the traits in consideration as potence ratio pointed out.

Highly significant mid and better parents heterosis were detected in the cross; Line 4 x Tester 1 for chocolate spot disease reaction and flowering date in negative direction, plant height; No. of branches / plant, No. of pods /plant, No. of seeds / plant and crude protein% in positive direction due to over- dominance( $P > +1$ ) in all traits in question . However, plant height was expressed only significant mid and better parents heterosis in positive direction and the data also shown highly significant mid and better parents heterotic effects in the cross; Line 4 x Tester 2 for No. of branches / plant; No. of pods /plant, No. of seeds / plant, seed yield / plant and carbohydrate %. Over- dominance was the main responsible of the heterotic effect in all traits mentioned, where potence ratio was exceeded plus one.

**Table (7): Heterotic effects relative to mid (M.P), potence ratios (P) and better (B.P) parent for disease reactions, yield and its components.**

Crosses	Chocolate spots disease reaction			Rust disease reaction			Flowering date (day)		
	M.P	p	B.P	M.P	p	B.P	M.P	p	B.P
Line 1 x Tester 1	-9.17	-0.37	21.11	-5.56**	-0.17	-29.17**	-13.41**	-12.90	-14.30**
Line 1 x Tester 2	11.63*	0.71	33.33*	10.43	0.40	-12.62	-22.17**	-3.72	-26.54**
Line 1 x Tester 3	15.79	3.00	22.22	87.50**	5.25	60.71**	-10.32*	-9.93	-11.24*
Line 1 x Tester 4	82.67**	4.13	128.33**	200.00**	∞	200.00**	5.31	1.25	1.00
Line 2 x Tester 1	-0.74	-0.07	11.67	50.48**	3.53	31.67**	-1.80	-0.22	-9.38
Line 2 x Tester 2	15.92	7.80	18.33	49.22**	7.31	39.81**	17.72*	5.13	13.79*
Line 2 x Tester 3	-3.64	-0.40	6.00	58.62**	17.00	53.33**	-5.19*	-0.62	-12.5*
Line 2 x Tester 4	77.78**	2.33	166.67**	140.00**	7.00	100.00**	12.15	2.36	6.67
Line 3 x Tester 1	0.00*	0.00	33.33*	25.00*	∞	25.00*	-3.64	-1.31	-6.25
Line 3 x Tester 2	54.42**	3.34	84.44**	58.74**	7.71	47.50**	-15.64**	-7.31	-17.4**
Line 3 x Tester 3	42.11**	8.00	50.00**	42.16*	2.39	20.83*	-16.49**	-5.93	-18.75**
Line 3 x Tester 4	90.67**	4.53	138.33**	66.67*	2.00	25.00*	-3.76	-8.46	-4.19
Line 4 x Tester 1	-26.00**	-∞	-26.00**	5.93	0.53	-4.67	-16.13**	-5.00	-18.75**
Line 4 x Tester 2	-7.64	-0.84	1.60	2.77	0.15	-13.33	-8.47	-5.00	-10.00
Line 4 x Tester 3	24.8**	1.24	56.00**	51.28*	1.82	18.00*	-9.68*	-3.00	-12.5*
Line 4 x Tester 4	36.19**	0.84	138.33**	42.86	1.00	0.00	-3.33	-∞	-3.33
L.S.D(0.05)	0.89		0.77	0.82		0.71	5.03		4.36
(0.01)	1.19		1.03	1.10		0.95	6.72		5.82

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

**Table (7):** Cont.

Crosses	Plant height(cm)			No of branches/plant			No of pods/plant		
	M.P	p	B.P	M.P	p	B.P	M.P	p	B.P
Line 1 x Tester 1	2.53	4.81	1.99	47.37**	3.06	27.62**	45.76**	2.36	22.05**
Line 1 x Tester 2	16.00**	43.73	15.57**	18.66	1.19	2.58	19.26**	2.23	9.77**
Line 1 x Tester 3	-13.40**	-0.82	-25.51**	20.68	2.88	12.60	-18.41	-1.49	-27.36
Line 1 x Tester 4	0.83**	0.06	-12.05**	39.29**	3.30	24.48**	64.58**	2331.04	64.54**
Line 2 x Tester 1	-7.59**	-1.02	-14.01**	35.96*	3.29	22.56*	30.59**	2.68	17.22**
Line 2 x Tester 2	-4.67**	-0.71	-10.55**	34.38*	3.08	20.90*	25.33*	55.43	24.76*
Line 2 x Tester 3	-17.94**	-1.91	-25.00**	75.38**	29.34	70.98**	-0.88*	-0.04	-17.62*
Line 2 x Tester 4	-14.36**	-1.84	-20.54**	12.88	1.76	5.18	63.52**	7.78	51.18**
Line 3 x Tester 1	9.69	1.96	4.51	31.34*	109.33	30.96*	169.93**	76.80	164.08**
Line 3 x Tester 2	12.31*	2.11	6.11*	24.87	363.28	24.79	3.35	0.38	-4.98
Line 3 x Tester 3	-14.61**	-0.68	-29.74**	15.06	1.74	5.88	29.52**	1.02	0.42**
Line 3 x Tester 4	-1.27**	-0.06	-17.70**	42.99**	10.94	37.59**	49.50**	2.87	27.5**
Line 4 x Tester 1	8.24*	2.73	5.07*	41.89**	13.15	37.51**	48.85**	4.95	35.48**
Line 4 x Tester 2	5.70	2.68	3.50	122.8**	41.40	116.38**	97.58**	4.74	63.83**
Line 4 x Tester 3	-13.74**	-0.99	-24.21**	47.70**	4.13	32.41**	-11.05**	-0.28	-36.31**
Line 4 x Tester 4	-7.41**	-0.61	-17.48**	92.95**	13.63	80.63**	56.49**	1.97	21.58**
L.S.D(0.05)	5.55		4.81	0.77		0.67	5.12		4.43
(0.01)	7.41		6.42	1.03		0.89	6.83		5.92

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

**Table (7):** Cont.

Crosses	No of seeds/plant	Seed yield/plant (g)	100-seed weight (g)
---------	-------------------	----------------------	---------------------

	M.P	p	B.P	M.P	p	B.P	M.P	p	B.P
Line 1 x Tester 1	40.08**	1.53	11.04**	42.63	1.44	9.98	3.52	0.83	-0.7
Line 1 x Tester 2	9.02*	0.75	-2.74*	16.59	0.91	-1.37	7.84	1.24	1.42
Line 1 x Tester 3	-19.62	-1.74	-27.76	-22.22*	-2.55	-28.46*	-3.81	-1.56	-6.1
Line 1 x Tester 4	65.26**	187.00	64.68**	69.00**	19.68	63.27**	2.27	0.70	-0.93
Line 2 x Tester 1	40.40**	3.15	24.46**	48.87	2.82	26.88	7.04	1.41	1.95
Line 2 x Tester 2	31.84**	18.18	29.57**	54.74*	10.30	46.94*	17.08	2.41	9.33
Line 2 x Tester 3	-8.30	-0.34	-26.46	-12.17*	-0.57	-27.71*	-5.05	-1.58	-8.00
Line 2 x Tester 4	82.66**	6.14	60.98**	30.14	3.15	18.78	-28.16**	-7.06	-30.91**
Line 3 x Tester 1	153.49**	11.04	122.56**	173.81**	11.55	137.98**	8.92	6.28	7.40
Line 3 x Tester 2	-7.37	-11.49	-7.96	-5.22	-1.75	-7.96	2.91	0.83	-0.59
Line 3 x Tester 3	20.76**	0.88	-2.34**	19.75	0.83	-3.21	-0.78	-2.04	-1.15
Line 3 x Tester 4	39.26**	3.17	23.92**	56.81*	4.78	40.15*	12.76	31.47	12.31
Line 4 x Tester 1	48.85**	10.40	42.17**	60.17	6.64	46.85	8.37	2.08	4.18
Line 4 x Tester 2	92.71**	4.85	61.81**	109.2**	5.21	72.93**	9.00	4.68	6.94
Line 4 x Tester 3	-24.47	-0.61	-46.19	-15.81**	-0.35	-41.93**	13.82	2.38	7.57
Line 4 x Tester 4	50.88**	1.69	15.92**	59.43	1.71	18.36	7.07	1.41	1.94
L.S.D(0.05)	14.97		12.97	23.98		20.77	15.41		13.34
(0.01)	19.98		17.31	32.02		27.73	20.57		17.81

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

**Table (7): Cont.**

Crosses	Crud protein %			Carbohydrate %			Fiber %		
	M.P	p	B.P	M.P	p	B.P	M.P	p	B.P
Line 1 x Tester 1	5.88	3.00	3.85	2.31	2.10	1.20	-7.95	-2.24	-11.11*
Line 1 x Tester 2	20.00**	2.20	10.00**	-2.7**	-0.64	-6.67**	-2.44	-0.33	-9.09
Line 1 x Tester 3	4.88	0.57	-3.37	1.50	1.33	0.37	-12.77**	-3.18	-16.14**
Line 1 x Tester 4	12.73	1.40	3.33	-1.54**	-0.40	-5.19**	14.72	1.64	5.26
Line 2 x Tester 1	-1.93**	-0.25	-8.97**	5.52	1.14	0.66	-11.89**	-1.17	-20.00**
Line 2 x Tester 2	-1.79**	-0.12	-14.44**	-4.35**	-0.43	-13.16**	-19.43**	-3.04	-24.27**
Line 2 x Tester 3	3.98**	0.28	-8.99**	-2.82**	-0.40	-9.21**	-24.76**	-1.41	-36.00**
Line 2 x Tester 4	14.80	1.00	0.01	-9.03**	-0.93	-17.11**	-11.91**	-0.53	-28.00**
Line 3 x Tester 1	-0.91**	-0.17	-6.03**	9.65*	1.47	2.90*	3.68	1.57	1.31
Line 3 x Tester 2	-13.45**	-7.93	-14.89**	10.20**	8.33	8.87**	-16.40**	-2.68	-21.21**
Line 3 x Tester 3	-7.95**	-7.00	-8.99**	9.88**	2.27	5.30**	-9.91**	-1.90	-14.38**
Line 3 x Tester 4	10.73**	6.33	8.89**	-6.83**	-4.20	-8.32**	24.53*	2.41	13.01*
Line 4 x Tester 1	10.38**	4.16	7.69**	3.55	1.60	1.30	2.13	0.25	-5.88
Line 4 x Tester 2	-6.21**	-0.65	-14.44**	10.16**	3.25	6.82**	-1.36	-0.11	-12.12*
Line 4 x Tester 3	-2.57**	-0.28	-10.67**	8.33**	∞	8.33**	3.65	3.80	2.66
Line 4 x Tester 4	8.40	0.87	-1.11	0.93	0.34	-1.74	7.66	1.90	3.49
L.S.D(0.05)	1.28		1.11	1.31		1.13	1.03		0.89
(0.01)	1.71		1.48	1.75		1.51	1.37		1.19

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

However, the crosses mention before could be used to evaluate faba bean genetic material of the present study through planned breeding program depending upon follow the bulk method in the advanced segregating generations, where according to this method non-additive

genetic variance reduced one fourth from generation to another and in the F<sub>5</sub> and F<sub>6</sub> generation this kind of genes approximately disappeared and additive genes would represented almost the genetic variance, hence, effective selection would be done with follow pedigree selection breeding method which allow to obtain some pure lines could be tested through yield trials.

Different values of heterosis might be due to the genetically diversity of the parents with non-allelic interactions, which increase or decrease the expression of heterosis[17]. Even in the absence of epistasis, multiple alleles at a locus could lead to either positive or negative heterosis[18]. These results are in good agreement with those obtained in faba bean by Attia et al [19], El-Hadyet al [20], El-Metwallyet al [21], Abo Mostafa et al [22], Haridy et al [23], Abo-Zaid et al [24], Abdalla et al [25], Abdalla et al [26], Bishnoi et al [27], Fouad [28], Heiba et al [29] and Ibrahime et al [30].

#### **Combining ability:**

The data presented in Table (6) revealed that, the ratios of  $\sigma^2_{gca}/\sigma^2_{sca}$  were less than unity for all studied traits; this might indicate that the non-additive gene effects were of greater importance in the inheritance of these traits. It could be concluded that, selection procedure based on the accumulation of additive would not be successful in improving these traits. To maximize selection advance, procedure which are known to be effective in shifting gene frequency when additive and non-additive genetic variances are involved would be preferred. However, in the case of predominant of non-additive genetic variance in the inheritance of a trait, the selection must be delayed to the later segregating generations and bulk method must be followed.

The data listed in Table (8) revealed the general combining ability effects ( $\hat{g}_i$ ) of the lines and testers used in the study for the studied traits. The data indicated that, Line 1 considered as good combiner parent for improving the tolerant to rust diseases with respect to the present materials due to its significant ( $\hat{g}_i$ ) in negative direction. Significant and/or highly significant ( $\hat{g}_i$ ) in positive direction for plant height and crude protein percentage were found in line 1 and tester 4. The parents; Line 3 and tester 4 showed highly significant ( $\hat{g}_i$ ) in positive direction for No. of pods/plant, no. of seeds/plant and seed yield /plant in the testers could be used in a suitable breeding program to improve seed yield. The parents line 4, tester 1 and tester 3; had significant ( $\hat{g}_i$ ) in positive direction for carbohydrates percentage. For tester parent 4 (Najeh) could be used in suitable breeding program to improve yielding potentiality and protein of the present material due to its at least significant( $\hat{g}_i$ ) for plant height, No. of pods and seeds/plant and seed yield /plant and crude protein percentage. GCA effects ( $\hat{g}_i$ ) was previously reported in faba bean by Drwish et al ([31], Ahmad [32], Haridy et al [23], Abo-Zaid et al [24], Abdalla et al [25], Abdalla et al [26], Bishnoi et al [27], Fouad [28], Heiba et al [29] and Ibrahime et al [30].

**Table (8) :** Estimates of parental lines and testers general combining ability effects for studied traits in the F<sub>1</sub> generation.

Genotypes	Chocolate spot diseases reaction	Rust diseases reaction	Flowering date	Plant height (cm)	No. of branches/ Plant	No. of pods/ plant	No. of seed/plant	Seed Yield/ plant	100- seed weight	crude protein (%)	Carbo- hydrate (%)	Fiber (%)
Line												
Line 1	-0.45	-0.78*	0.01	3.75*	0.07	0.97	3.36	4.48	1.01	1.9**	-0.95*	-0.05
Line 2	0.10	0.31	3.33*	-3.92*	0.14	-1.09	-0.14	-4.71	-2.23	-1.43**	0.70	-0.20
Line 3	0.29	0.33	-1.67	-2.27	-0.78*	3.44*	9.29*	10.26	1.42	0.09	-0.83*	0.27
Line 4	0.06	0.14	-1.67	2.45	0.58*	-3.32*	-12.52*	-10.04	-0.2	-0.56	1.08*	-0.02
LSD(gi-gj) 0.05 0.01	0.445 0.594	0.412 0.550	2.516 3.358	2.777 3.707	0.384 0.513	2.560 3.417	7.485 9.992	11.992 16.008	7.704 10.284	0.639 0.853	0.654 0.873	0.515 0.687
Tester												
Tester 1	-0.46	-0.39	-0.42	-3.31*	-0.26	1.71	2.75	5.39	2.31	-1.54**	2.82**	0.44
Tester 2	0.21	-0.35	-2.08	2.34	0.01	-3.25*	-11.34*	-6.3	2.43	-0.55	-0.55	0.14

Tester 3	-0.19	0.10	-1.25	-4.55*	0.15	-5.09*	-16.9**	-14.3*	-1.03	-0.73*	1.25*	-0.99*
Tester 4	0.44	0.64*	3.75*	5.51*	0.11	6.63**	25.57**	15.22*	-3.71	2.82**	-3.52**	0.41
LSD(gi-gj)	0.445	0.412	2.516	2.777	0.384	2.560	7.485	11.992	7.704	0.639	0.654	0.515
0.05	0.594	0.550	3.358	3.707	0.513	3.417	9.992	16.008	10.284	0.853	0.873	0.687

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

Specific combining ability effects ( $\hat{S}_{ij}$ ) were estimates of the sixteen crosses for the studied traits. The data presented in Table (9) showed that, the cross; Line 2 x Tester 3 had highly significant ( $\hat{S}_{ij}$ ) in favorable direction for chocolate spot disease reaction and crosses; Line 1 x Tester1, Line 1 x Tester 2 and Line 3 x Tester 4 had highly significant ( $\hat{S}_{ij}$ ) in favorable direction for rust disease reaction, indicating that the ability to use these crosses to improve the tolerant to the two diseases. While, cross; Line 1 x Tester 2 recorded negative significant ( $\hat{S}_{ij}$ ) for flowering date.

For plant height three crosses; Line 1 x Tester 2, Line 2 x Tester 3 and Line 4 x Tester1 exhibited significant positive ( $\hat{S}_{ij}$ ) and the crosses; Line 2 x Tester 3 and Line 4 x Tester 2 for No of branches/plant. The crosses; Line 1 x Tester 4 and Line 4 x Tester 2 possessed significant and highly significant ( $\hat{S}_{ij}$ ) in positive direction for No of pods /plant. The crosses; Line 1 x Tester 4, Line 2x Tester4, Line 3x Tester 1, Line 3x Tester 3 and Line 4x Tester 2 had significant and/ or highly significant ( $\hat{S}_{ij}$ ) for No of seeds /plant in positive direction.

**Table (9):** Estimates of specific combining ability effects for studied traits in the F<sub>1</sub> generation.

Crosses	No .of seeds/plant	Seed Yield/ plant	100-seed weight	Protein %	Carbohydrate %	Fiber %
Line 1 x Tester 1	0.13	-0.86*	-0.42	-5.62*	0.72	-1.80
Line 1 x Tester 2	-0.18	-0.73*	-5.42*	4.85*	-0.61	-0.54
Line 1 x Tester 3	-0.11	0.32	2.08	-2.81	-0.33	-3.73
Line 1 x Tester 4	0.16	1.28**	3.75	3.59	0.22	6.06*
Line 2 x Tester 1	0.41	0.49	-2.08	-1.49	-0.02	-6.53**
Line 2 x Tester 2	0.01	-0.02	6.25**	-2.69	-0.35	0.35
Line 2 x Tester 3	-0.79*	-0.67	-2.92	5.65*	1.44**	2.09
Line 2 x Tester 4	0.37	0.19	-1.25	-1.47	-1.06**	4.08
Line 3 x Tester 1	-0.24	0.21	4.58*	1.99	0.23	15.05**
Line 3 x Tester 2	0.62	1.07**	-2.08	0.22	-0.25	-11.99**
Line 3 x Tester 3	-0.01	-0.45	-1.25	-3.35	-0.36	4.51
Line 3 x Tester 4	-0.38	-0.83*	-1.25	1.14	0.38	-7.57**
Line 4 x Tester 1	-0.31	0.16	-2.08	5.12*	-0.93*	-6.72**
Line 4 x Tester 2	-0.45	-0.32	1.25	-2.37	1.22**	12.18**
Line 4 x Tester 3	0.91*	0.80*	2.08	0.50	-0.75	-2.58
Line 4 x Tester 4	-0.15	-0.64	-1.25	-3.25	0.46	-2.58
S.D(sij-skij) (0.05)	0.981	0.824	5.031	5.554	0.768	5.120
(0.01)	1.189	1.100	6.716	7.414	1.025	6.834

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

**Table (9):** Cont.

Crosses	No .of seeds/plant	Seed Yield/ plant	100-seed weight	Protein %	Carbohydrate %	Fiber %
---------	--------------------	-------------------	-----------------	-----------	----------------	---------

Line 1 x Tester 1	-7.10	-8.57	-1.58	-1.38*	-0.37	-0.63
Line 1 x Tester 2	-5.36	-5.30	0.05	3.63**	-1.55*	0.60
Line 1 x Tester 3	-5.69	-7.25	-2.72	-0.52	-0.17	-0.30
Line 1 x Tester 4	18.15*	21.12	4.24	-1.73**	2.09**	0.33
Line 2 x Tester 1	-18.64**	-8.54	5.13	-1.38*	2.43**	0.45
Line 2 x Tester 2	2.04	14.60	11.22	-0.37	-1.20*	0.21
Line 2 x Tester 3	-0.73	2.60	0.11	1.14	-1.00	-0.13
Line 2 x Tester 4	17.33*	-8.66	-16.47*	0.6	-0.23	-0.53
Line 3 x Tester 1	41.9**	33.67**	-0.19	0.68	0.29	0.31
Line 3 x Tester 2	-33.77**	-34.1**	-6.55	-2.03**	1.33*	-1.05*
Line 3 x Tester 3	16.95*	9.27	-2.94	-0.39	0.87	-0.26
Line 3 x Tester 4	-25.08**	-8.85	9.68	1.74**	-2.5**	1.00
Line 4 x Tester 1	-16.16*	-16.57	-3.37	2.08**	-2.35**	-0.13
Line 4 x Tester 2	37.09**	24.8*	-4.72	-1.24*	1.42*	0.24
Line 4 x Tester 3	-10.53	-4.62	5.55	-0.23	0.30	0.70
Line 4 x Tester 4	-10.40	-3.61	2.54	-0.61	0.63	-0.80
L.S.D(sij-skij) (0.05)	14.971	23.985	15.408	1.278	1.308	1.030
(0.01)	19.984	32.017	20.568	1.706	1.747	1.375

\*and \*\* refer to 0.05 and 0.01 levels of probability, respectively.

The crosses; Line 3 x Tester 1 and Line 4 x Tester 2 expressed significant ( $\hat{S}_{ij}$ ) in positive direction for seed yield /plant. The crosses; Line 1 x Tester 2, Line 3 x Tester 4 and Line 4 x Tester 1 had highly significant ( $\hat{S}_{ij}$ ) in positive direction for crude protein percentage. The crosses; Line 1 x Tester 4, Line 2 x Tester 1, Line 3 x Tester 2 and Line 4 x Tester 2 showed significant ( $\hat{S}_{ij}$ ) in favorable direction for carbohydrate percentage. However, The crosses; Line 1 x Tester 2, Line 1 x Tester 4, Line 3 x Tester 1 and Line 4 x Tester 2 had more significant traits for ( $\hat{S}_{ij}$ ). These crosses also had significant better parent heterosis for most traits, which might be performed that the source of both parameters i.e., heterosis and  $\hat{S}_{ij}$  is the non-additive genes. Indicating that these crosses could be used in breeding program to improve the reaction to chocolate spot disease and yielding ability. While, the cross; line 3 x tester 4 showed significant ( $\hat{S}_{ij}$ ) in favorable direction for fibers percentage. Similar trend are obtained by Abd El-Mohsen [33], El-Metwally et al [21], Ahmad [32], Haridy et al [23], Abo-Zaid et al [24], Bishnoi et al [27], Fouad [28], Heiba et al [29] and Ibrahim et al [30].

#### **IV- Heritability and genetic advance:**

The heritability is one of the most important parameter for determination the genetic behavior of a metric character. The heritability estimates is usually connected with breeding methods and many practical decisions about the techniques followed depends on its magnitude.

Phenotypic and genotypic coefficient of variation, heritability in broad and narrow-senses and expected and predicted genetic gain from selection are presented in Table (10). In this respect, phenotypic (PCV) and genotypic (GCV) coefficients of variation ranged from 8.4% to 34.33% and 6.83% to 26.75% for plant height and No. of branches/plant, respectively. Broad sense heritability (H) ranged from 24.6% for No. of seeds/plant to 77.90% for 100-seed weight. According to categorized suggested by Johanson [30], low (H) was detected for No. of seeds/plant, moderate values were noticed for No. of pods/plant, No. of branches/plant and seed yield /plant. The other traits had (H) values more than 60% which considered as high broad – sense heritability values. However, the values of (H) were considered as relatively low, which could be attributed to that the present genetic material is more affected by environmental conditions.

With respect to narrow- sense heritability ( $h^2$ ), the range is between 0.68% for 100-seed weight to 20.28% for chocolate spot diseases reaction. According to the classification of narrow-sense heritability by Stansfield [14], all estimates are considered as low (< 20%).

The low estimates of narrow- sense heritability in the present material could be expected results, because all studied traits were mainly controlled by non-additive gene effects as

mentioned before and the additive genes considered as low constitution of the genetic variance and subsequent lowering the narrow-sense heritability

The expected genetic advance (Ga) or predicted (Ga %) genetic gain upon selection are listed in Table (9). The predicted (Ga%) genetic gain upon selection ranged from 0.18 % for 100-seed weight to 6.92% for chocolate spot disease reaction, which considered as low values. This is an expected result due the low values of  $h^2$  and (PCV) which are the main component of (Ga) equation. These results are in the same line with those obtained by **El-Metwally et al [21], Abo Mostafa et al [22], Haridy et al [23], Abo-Zaid et al [24], Abdalla et al [25], Abdalla et al [26], Bishnoi et al [27], Fouad [28], Heiba et al [29] and Ibrahime et al [30].**

**Table (10): Means ( $\bar{X}$ ), Phenotypic (PCV) and genotypic (GCV) coefficient of variation, heritability (H) in broad-sense and narrow sense ( $h^2$ ), expected (Ga) and predicted genetic advance (Ga %) for all studied traits.**

Traits	$\bar{X}$	PCV	GCV(%)	Heritability		Genetic advance	
				H (%)	$h^2$ (%)	G a	Ga %
Chocolate-spot disease reaction	3.93	16.56	14.42	75.94	20.28	0.27	6.92
Rust disease reaction	4.59	16.37	14.12	74.33	0.71	0.01	0.24
Plant height(cm)	134.86	8.40	6.83	66.72	9.48	2.20	1.63
No .of branches/ plant	4.22	34.33	26.75	60.69	4.95	0.15	3.50
No .of pods/plant	22.56	30.87	24.01	60.49	1.06	0.15	0.67
No .of seeds/plant	65.16	24.93	12.36	24.60	3.37	1.14	1.73
Seed Yield/ plant	53.38	27.54	19.39	49.56	6.76	2.05	3.83
100-seed weight	80.94	12.67	11.12	77.90	0.68	0.14	0.18

## Conclusion

Overall, the study revealed significant genetic variability among the parental genotypes and their crosses for various important traits in faba bean. The results indicated the presence of both additive and non-additive gene effects in the inheritance of these traits. Tester 4 was identified as a good combiner parent for several traits, while the crosses Line 1 x Tester 1, Line 1 x Tester 3, Line 2 x Tester 3 and Line 4 x Tester 1 were resistant for chocolate spot and rust diseases reaction showed significant effects in favorable directions for disease resistance. Whilst the crosses; Line 1 x Tester 4, Line 2 x Tester 2, Line 2 x Tester 4, Line 3 x Tester 1, Line 3 x Tester 3, Line 3 x Tester, Line 4 x Tester 2 and Line 4 x Tester 4) were the highest yielding crosses. The heritability estimates varied among traits, with moderate to high values observed for some traits. The study provides valuable insights into the genetic control of important traits in faba bean and highlights the potential for developing high-yielding and disease-resistant hybrids through appropriate breeding strategies.

## Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

#### REFERENCES

1. Larralde, J. Estudio de algunos trastornos que se presentant en los animales tras la ingestion de semillas de Vicia faba L. Rev. Esp. Fisiol.1982;38: 345-351.
2. Eliw M. Economic analysis of supply response of broad beans crop in Egypt. New Valley Journal of Agricultural Science. 2021;1(1):10-25.
3. Bond, D. A. Combining ability of winter beans (*Vicia faba* L.) inbreeds. J. Agric. Sci. Cambridge University. 1967;68: 179-185.
4. Sindhu, P.S; Verma,M.; Cheema, H.S. and Sra, S.S. Genetic relationships among yield components in pigeonpea. Indian J. Agric. Sci. 1985;55(4):232-235.
5. Yassin, T.E. Genotypic and phenotypic variances and correlations in field beans (*Vicia faba* L.). J. Agric.Sci. Camb.1973;81 : 445-448.

6. Bernier, C.C.; S.B. Hanounik; M.M. Hussein and H.A. Mohamed. Field manual of common faba bean diseases in Niel Valley. Information Bulletin No. 3. ICARDA, P.O Box 5466, Aleppo, Syria; 1984.
7. Cochran, W.G. and G.M. Cox. Experimental Design. John Wiley and Sons. Inc. New York;1957.
8. Kempthorne, O. An introduction to genetic statistics. John Wiley and Sons. (eds) Ine .New York. 1957; 545pp.
9. Chaudhary, B.D and R.K. Singh. Biometrical Methods in Quantitative Genetic Analysis. At Kalyani Publishers, New Delhi-110002;1985.
10. Burton, G.W. Quantitative inheritance in grasses. Proceedings of the 6th International Grassland Congress, August 17-23, Pennsylvania College, USA.1952; pp. 277-283.
11. Lush, Jay. L. Intra-sire correlations or regressions of offspring on dam as a method of estimating heritability of characteristics. Proc. Amer. Soc. Anim. Prod. 1940; Pp.293-301.
12. Johanson, H.W.; Robinson and R.E. Comstoc. Estimates of genetic and environmental variability in soybean. Agron. J. 1955;47: 314-322.
13. Acquaah, G. Principles of plant genetics and breeding. Blackwell Publishing, Oxford.2007; P.128
14. Stansfield, W.D. Theory and problems of genetics. Mc. Grow Hills, Book Company;1991.
15. Fehr, W.R. Heterosis. In: Principle of cultivars. Development. Macmillan publishing company. A division of Macmillan Inc. New York.1987; pp.1:1-465
16. A.O.A.C. Official Methods of the Analysis. Association of Official Agricultural Methods. 15th Edition, Published by Association of Official Analytical Chemists, Arington, Virginia, U.S.A;1990.
17. Hyman, B.I. Interaction, heterosis and diallel crosses. Genetics,1957; 42:336-355.
18. Cress C.E. Heterosis of the hybrid related to gene frequency differences between two populations. Genetics. 1966;53:269-274.
19. Attia, Sabah M. and Salem Manal M. Analysis of yield and its components using diallel matings among five parents of faba bean. Egypt. J. Plant Breed. 2006;10(1):1-12.
20. El-Hady, M.M.;Sabah M. Attia; E.A.A. El-Emam, A.A.M.Ashrei and E.M.Rabie. Diallel mating among eight parents of faba bean (*Vicia faba* L) and performance of F<sub>1</sub> and F<sub>2</sub>. Egypt.J. of. Appl. Sci. 2008;23(5): 95-114.
21. El-Metwally, I.M., T.A. El-Shahawy and M.A. Ahmed. Effect of sowing dates and some broomrape control treatments on faba bean growth and yield. J. Applied Sci. Res. 2013; 9:197-204.
22. Abo Mostafa, R.A., Zeinab, E. G., Abbas, M.A., Gehan, G.A. and Sarhan,E.A. Combinedand genetic analysis for multiple-disease resistanceto chocolate spot and rust on faba bean yield. Inter.J. of Plant breed. and Genetics. 2014;51, 1-13.
23. Haridy, M.H.and El-Said, M.A.A. Estimation of genetic parameters using populations in faba bean (*Vicia faba* L.). J. Plant production, Mansoura Univ. 2016;7:1443-1447.

24. Abou-Zaid, G. G.; S. M. Mostafa and R. A. El-Refaey. Genotype x environment interaction effects on heritability and genetic advance for yield and its components of some faba bean genotypes. *J. Plant Production, Mansoura Univ.* 2017;8(6):665 -669.
25. Abdalla M.M.F, Shafik M.M, Sabah M. Attia, Hend A. Ghannam. Combining ability, heterosis and inbreeding effects in faba bean (*Vicia faba L.*). *Journal of Experimental Agriculture International.*2017a;15(5):1-13.
26. Abdalla M.M.F, Shafik M.M, Sabah Attia M, Hend A. Ghannam. Heterosis, GCA and SCA effects of diallel-cross among six faba bean (*Vicia faba L.*) genotypes. *Asian Research Journal of Agriculture*, 2017 b;4(4):1-10.
27. Bishnoi, S. K., Hooda, J. S. and Sharma, M. P. Analysis of gene effects for yield and yield components traits in faba bean (*Vicia faba L.*) genotypes. *The J. Anim. Plant Sci.* 2018;28:187-196.
28. Fouad, H.M. Six generation mean analysis using scaling and joint scaling tests in faba bean (*Vicia faba L.*). *Journal of Sustainable Sciences*, 2020;46 (1):1-11.
29. Heiba, H. E.; E. Mahgoub; A. Mahmoud; M. Ibrahim and E. M. B. Mahdy. Genetic improvement of faba bean (*Vicia faba L.*) genotypes selected for resistance to chocolate spot disease. *Agronomic Colombian.* 2022;40(2):186-197.
30. Ibrahim, M.A.; Gehan, G. Abo-Zaid and Salwa M. Mostafa. Estimation of some genetic parameters using six populations mean analysis in three faba bean crosses exposed to natural infection of foliar diseases. *Egyptian J. Agric. Res.* 2023;101(2):843-854.
31. Darwish, D.S.; M.M.F. Abdalla; E.A.A. El-Emam. Investigations on faba bean (*Vicia faba L.*) 19-diallel and triallel matings using five parents. *Proceed. Fourth PI. Breed. Conf. March 5, (Ismailia) Egypt J. Plant Breed.* 2005;9(1):197-208 Special Issue.
32. Ahmad, M.S.H. Studies on genetic variability, heritability and genetic advance in segregating generations of faba bean (*Vicia faba L.*). *Middle East J Agric. Res.* 2016;5(1):82-89.
33. Abd El-Mohsen, M.I. Heterosis and combining ability in faba bean for some quantitative characters. *Egypt. J. Plant Breed.* 2004;8:161-171.