

# ***Lactuca sativa* growth and yield as affected by cashew nut shells biochar, cattle manure and urea in northern and central Benin.**

## **ABSTRACT**

**Aims:** This study aims to determine the optimal combination of cashew nut shell biochar, cattle manure, and urea to enhance lettuce (*Lactuca sativa* L.) growth and yield in North and Central Benin.

**Place and Duration of Study:** Two agronomic trials were conducted: one in Tchaourou (Borgou department) in 2022 and another in Dassa (Colline department) in 2023.

**Methodology:** A Fisher design with three replications was used for the experimental design at each site, focusing on fertilization as the factor with seven treatments, including four reference treatments (control, producers' practice of organic fertilization = 40 t ha<sup>-1</sup> cattle manure; producers' practice of mineral fertilization = 100 kg ha<sup>-1</sup> urea; producers' practice of organo-mineral fertilization = 20 t ha<sup>-1</sup> cattle manure + 50 kg ha<sup>-1</sup> urea) and three treatments based on the reference treatments with biochar incorporation.

The experimental design on each site was a Fisher design with three replications, focusing on fertilization as the factor with seven treatments including four reference treatments (control, producers practice of organic fertilization = 40 t ha<sup>-1</sup> cattle manure; producers practice of mineral fertilization = 100 kg ha<sup>-1</sup> urea; producers practice of organo-mineral fertilization = 20 t ha<sup>-1</sup> cattle manure + 50 kg ha<sup>-1</sup> urea) and three treatments based on the reference treatments with biochar incorporation.

**Results:** The results demonstrated significant improvements in plant growth parameters and yield, with the application of biochar combined with cattle manure and urea or with cattle manure only. The obtained yields under applying of 10 t ha<sup>-1</sup> biochar + 20 t ha<sup>-1</sup> cattle manure and 10 t ha<sup>-1</sup> biochar + 25 t ha<sup>-1</sup> urea + 10 t ha<sup>-1</sup> cattle manure were at least respectively 165 and 318% higher than those of obtained under producers mineral fertilization practice on both sites.

**Conclusion:** The findings showed that integrating cashew nut shell biochar with organic and inorganic fertilizers can significantly enhance lettuce and other vegetable crop production while reducing the need for mineral fertilizers and dependence on cattle manure.

The findings showed that integrating cashew nut shell biochar with organic and inorganic fertilizers can significantly enhanced lettuce and thus vegetable crops production, while reducing the need on mineral fertilizers and dependence of cattle manure.

Formatted: Font: Not Bold

**Keywords:** Cashew nut shell biochar, cattle manure, urea, lettuce, Amendment, valorization, fertilization, vegetable crops

## **1. INTRODUCTION**

In sub-Saharan Africa, fruit and vegetable production has become a crucial agricultural sector due to their nutrition, and the significant economic income they generate [1,2]. This sector has the potential to drive agricultural and economic diversification, particularly for smallholders who can target local, regional, or export markets [3,4]. In Benin, gardening is a nationwide activity because it is essential for the population's nutritional balance and the improve agricultural producers' incomes [5,6]. Vegetable production is common in various ecologies, notably on plateaus, plains, valleys, lowlands, and along the coast [7].

Comment [U2]: You mention that "fruit and vegetable production has become a crucial agricultural sector due to their nutritional value and the significant economic income they generate." It could be beneficial to provide a specific example or statistic to highlight the extent of this economic impact or nutritional value, making the statement more concrete.

Among the primary leafy vegetables cultivated is lettuce (*Lactuca sativa*) [8], ~~which is widely consumed~~~~which is very consumed~~. According to the World Health Organization (WHO), the recommended vegetable intake per capita per day is 400 grams [9]. However, in Benin, vegetable consumption per adult is around 99.1 grams per day [10]. To meet this increasing demand, producers are resorting to intensive mineral fertilization [11]. Excessive use of chemical fertilizers exacerbates soil acidity and depletes organic matter, leading to reduced soil fertility and yields [12]. Therefore, appropriate fertilization management measures are necessary to restore and maintain soil fertility and ecological balance [13]. One promising solution is the use of organic fertilization. Various studies have reported that organic fertilizers are a good substitute for chemical fertilizers, improving soil fertility and plant development [14,15,16].

**Comment [U3]:** The sentence "To meet this increasing demand, producers are resorting to intensive mineral fertilization [11]" could benefit from specifying what demand you are referring to (e.g., "To meet the increasing demand for vegetables...").

In Benin, particularly in the North and Center, local organic resources are available in addition to animal waste (cattle manure, poultry droppings). Cashew nuts (*Anacardium occidentale*), the second-largest export product after cotton [17], generate substantial quantities of residues. This crop involves more than 200,000 actors who engage in production, processing, and marketing within different value chains [17]. With the increasing number of factories, there is an abundant production of shells (10,959 tonnes in 2013), which are challenging for processing units to manage [18]. To create added value, some processing units recycle cashew nut shells by extracting cashew nut shell liquid [19]. Additionally, the shells are stored in processing plants and used directly as fuel. Pyrolyzers convert cashew shells into gas and charcoal [18]. Studies by Godjo et al. [18] showed that the pyrolysis reactor could recover approximately 82% of the mass of treated shells as gas and produce about 18% as charcoal. However, they contain approximately 16% carbon, 3% to 7% ash, and 4% nitrogenous materials [20].

**Comment [U4]:** Your text transitions well between the different ideas, but the introduction of cashew nut shells as a resource could be more explicitly connected to the earlier discussion of fertilization challenges. For example, you could say, "Given the challenges of excessive mineral fertilization, exploring local organic resources like cashew nut shells offers a promising alternative."

Biochar, derived from the pyrolysis of organic waste under controlled conditions, has received increased global attention as an amendment to maintain a high stock of organic matter [21]. It gradually mineralizes in the soil because it contains a high recalcitrant C content in humid tropical soils [22]. Biochar significantly contributes to cation retention [23] and improves soil fertility by directly adding nutrients [21,24]. Thus, it offers better nutrient availability to plants, positively impacting their growth and development [25,26,27]. However, it needs to be combined with mineral or organic fertilizers to ensure satisfactory crop productivity [28,29,30]. Maccarthy et al.[29] showed that economic returns (Gross Margin, Gross Yield, and Benefit-Cost Ratio) improved when biochar was combined with inorganic fertilizers. Despite this potential, the adoption of biochar and organic/inorganic fertilization methods remains low among local farmers, as producers often prefer simpler methods [31].

**Comment [U5]:** It is important to briefly explain what recalcitrant C content is.

The main reason for this low adoption is the limited research focused on the valorization of cashew nut shells in soil fertilization or evaluating the potential of biochar in real-world crop fertilization. The central research question is: what is the influence of biochar from cashew nut shell associated or not to cattle manure, and or urea on lettuce growth and yield, in North and Central Benin?

**Comment [U6]:** "How does the incorporation of biochar derived from cashew nut shells, alone or in combination with cattle manure and/or urea, affect the growth performance and yield of lettuce (*Lactuca sativa* L.) in the diverse agroecological zones of North and Central Benin?"

The objective of this study was to evaluate the influence of cashew nut shell biochar, cattle manure, and urea on lettuce growth and yield, in North and Central Benin. We hypothesize that cashew nut shell biochar, integrated with or without organic fertilizers, can enhance productivity and improve soil fertility in market gardening, particularly lettuce cultivation, in North and Central Benin.

**Comment [U7]:** "The objective of this study was to evaluate the individual and combined effects of cashew nut shell biochar, cattle manure, and urea on lettuce growth and yield in North and Central Benin."

## 2. MATERIAL AND METHODS

### 2.1 Study area

This study was conducted in two agroecological zones in Benin. ~~The field trials were conducted during the same period~~~~The field trial was conducted at the same period~~ in the municipality of Tchaourou (Borgou department) in North Benin during 2022 (Site 1) and in the municipality of Dassa (Colline department) in Central Benin during 2023 (Site 2). The site 1 experiences a South Sudanian climate characterized by one rainy season and one dry season, with annual rainfall averaging between 1100 and 1200 millimeters and an average temperature of 26.8°C[32]. The main soil types in this area are weakly concreted tropical ferruginous types. In contrast, Site 2 is under a Sudano-Guinean climate with two rainy seasons and two dry seasons, often with unpredictable weather patterns. The annual number of rainy days ranges between 80 and 110, with annual rainfall averaging 907 millimeters and an average temperature of 29°C. The soils in this region are tropical ferruginous on a crystalline base, exhibiting highly variable characteristics[33]. The soil characteristics ~~on of the~~ two sites are presented on table 1. On both sites, the soils have a sandy-loamy texture, moderate content of P and very low content of potassium. They presented respectively neutral and moderately acid pH, very low and low nitrogen content, moderate and low carbon content, with rapid and slow decomposition of organic matter according to C/N ratio[34].

**Comment [U8]:** While you provide detailed soil characteristics, it might be useful to briefly discuss why these particular attributes (e.g., pH, nitrogen content, etc.) are relevant to your study. For example, "The sandy-loamy texture and low potassium content at both sites could affect nutrient availability and, consequently, the growth response of lettuce to different fertilization treatments."

**Table 1. Soil characteristics on site 1 and 2 (North and central Benin)**

Soil characteristics	Texture	pH	C (%)	N (%)	C:N	P mg/kg	K (meq/100g)
Site 1	Sandy-loam	6,89	1,14	0,035	32,57	15,45	0,59
Site 2	Sandy-loam	5,80	0,81	0,05	15,89	12,00	0,35

## 2.2 Plant Material and Organic Substrate Used

The plant material used in this study consisted of "EDEN" variety seed of lettuce (*Lactuca sativa* L.), which has a cycle length of approximately 51 days. The organic amendments used in this study were selected for their potential to enhance soil fertility and crop productivity. These included biochar derived from cashew nut shells, cattle manure, and the mineral fertilizer urea, which contains 46% nitrogen (see Table 2 for details). The organic amendments included biochar obtained from the pyrolysis of cashew nut shells, cattle manure and the mineral fertilizer urea containing 46% nitrogen (Tables 2).

**Comment [U9]:** The information about the lettuce variety "EDEN" is clear. However, it would be helpful to briefly mention why this variety was chosen, especially if it has specific traits that make it suitable for your study, such as adaptability to the local climate or its popularity among local farmers.

At the start and end of the experiment, representative composite samples were taken from the top 20 cm of soil in each replication. The laboratory analyses performed included:

- pH (water): measured by the potentiometric method[35];
- Total nitrogen: determined by the Kjeldahl method[36];
- Assimilable phosphorus: measured by the Bray I method[37];
- Exchangeable potassium: assessed by the Helmke and Sparks method[38];
- Organic matter: determined by incineration method of[39].

**Table 2. Chemical characteristics of organic substrates used at site 1 and 2**

Site	Fertiliser	C (%)	N (%)	C/N	P (mg/kg)	K (meq/100 g)	pH
Site 1	Cattle manure	15.72	0.93	16.9	1.47	5.87	8.01
	Biochar from hulls	37.04	0.61	60.72	0.51	2.10	9.81
Site 2	Cattle manure	19.26	1.33	25.6	0.43	1.39	8.51
	Biochar from hulls	34.40	0.58	59.3	0.33	2.02	10.75

## 2.3 Experimental Design and Management

The experimental design on each site was a Fisher design focusing on fertilization as the factor with seven treatments (T0 = control, PPFO = producers practice of organic fertilization = 40 t ha<sup>-1</sup> cattle manure, PPFM = producers practice of mineral fertilization = 100 kg ha<sup>-1</sup> urea, PPFOM = producers practice of organo-mineral fertilization = 20 t ha<sup>-1</sup> cattle manure + 50 kg ha<sup>-1</sup> urea, Biochar + FM = 10 t ha<sup>-1</sup> biochar + 50% of producers mineral fertilization rate = 10 t ha<sup>-1</sup> biochar + 50 kg ha<sup>-1</sup> urea, Biochar + FO = 10 t ha<sup>-1</sup> biochar + 50% of producers organic fertilization rate = 10 t ha<sup>-1</sup> biochar + 20 t ha<sup>-1</sup> cattle manure and Biochar + FM +FO = biochar + 25% of producers mineral fertilization rate + 25% of producers organic fertilization rate = 10 t ha<sup>-1</sup> biochar + 25 t ha<sup>-1</sup> urea +10 t ha<sup>-1</sup> cattle manure) including four (4) reference treatments (producers practice) and three replications (blocks) (Table 3).

Each experimental unit had an area of 2.4 m<sup>2</sup> (2 m x 1.2 m). The distance between the experimental plots was 0.5 m, and the distance between two blocks was 1 m. The sowing lines were arranged along the lengths of the experimental units with a spacing of 20 cm between lines and 20 cm between plants.

After soil weeding, the nursery was installed and the elementary plots were demarcated. Cattle manure and biochar were applied as basal fertilizers two weeks before transplanting to allow for initial soil conditioning and nutrient integration (Table 3). Cattle manure and biochar were applied to each elementary plot, according to the experimental design (Table 3). For this purpose, they were weighed and applied as basal fertilizer two weeks before transplanting on the elementary plots, which were watered twice a day (morning and evening) for one week before transplanting. The mineral fertilizer was applied in two equal fractions as maintenance fertilizer, one and three weeks after transplanting. Transplanting was done after 21 days of nursery. The plants were then watered twice a day (morning and evening), using two 14-liter watering

cans per plot. the Weeding and hoeing were carried out as needed. Phytosanitary treatment of the plants was carried out on the 15th and 30th day after transplanting; using neem oil, at a rate of 125 ml of neem oil diluted in 16 liters of water. Neem oil was selected for phytosanitary treatment due to its eco-friendly properties and effectiveness against common pests in the region.

**Table 3. Details of treatments included in the experimental design**

Fertilizers	Reference treatments				Treatments including biochar using			
	T0	PPFM	PPFO	PPFOM	Biochar + 50% PPFM	Biochar + 50% PPFO	Biochar + 25% PPFM + 25% PPFOM	
Urea	-	100 kg ha-1	-	50 kg ha-1	50 kg ha-1	-	25 kg ha-1	
Cattle manure	-	-	40 t ha-1	20 t ha-1		20 t ha-1	10 t ha-1	
Biochar	-	-	-	-	10 t ha-1	10 t ha-1	10 t ha-1	

## 2.4 Growth and Yield Parameters Measurements

The growth data collected during the trial were: the height of the plants measured from the collar to the apex using a graduated ruler; and the number of leaves emitted by counting all the leaves emitted by the plant. These two data were collected from one week after transplanting and weekly on six (06) plants chosen at random from the useful surface area. Harvest took place six weeks after transplanting by pulling up of plants on each elementary plot. these plants were washed with water to remove sand from the roots and leaves and then weighed using an electronic balance. To calculate the average weight of plants and yield of fresh biomass, the number of plants harvested from each elementary plot were counted. The average mass of harvested plants (P) per treatment was estimated by the following formula:

$$m \text{ (g/plant)} = M/N$$

Where M = total mass of plants harvested from a plot (g); N = number of plants harvested from an elementary plot.

The fresh biomass yield (R) was determined by the following formula:  $R \text{ (t/ha)} = m \cdot D \cdot 1000$

Where m = average mass of harvested plant (kg/plant) and D = lettuce density per hectare

## 2.5 Statistical Analysis

For Data analysis, the factors involved in experimentation were considered differently depending on their nature. Treatments (fertilization modalities) and Time were considered as fixed factors, and replication (Block) as a random factor. For growth data, a linear mixed-effect model was performed on plant height and a generalized linear mixed-effect model of the fish type was performed on number of leaves counted on each plant. This was done to assess the effect of treatments and time on plant height and number of leaves. For yield data, a linear mixed-effect model was performed to evaluate treatments effect. All Analyses were performed using R 4.3.3. statistical software.

## 3. RESULTS

### 3.1 Influence of Cashew Nut Shells Biochar, Cattle Manure and Urea on Lettuce Growth

The table 4 ~~shows~~ showed the results of linear mixed model analysis on growth data ~~realized on growth data~~. The treatments significantly influenced plant height at both sites over time (p < 0.05). Additionally, treatment and time had a distinctly significant effect on the number of leaves produced at both Site 1 and Site 2 (Table 4). It appears that treatments influenced significantly (p < 0.05) the plants height at the two sites, depending on time. Treatment and time had a distinctly significant effect on number of leaves emitted on both Site 1 and 2 (Table 4).


To this end, we noted an increase in leaves number over time until four weeks after transplanting (T4) on both sites. Indeed at site 1 and 2, the highest number of leaves were recorded under treatments Biochar + FO + FM ( $6 \pm 2$  and  $9 \pm$

3.39 leaves respectively), PPFO ( $6 \pm 2$  and  $10 \pm 3.77$ ), PPFOM ( $5 \pm 2$  and  $9 \pm 3.54$ ), Biochar + FO ( $5 \pm 2$  and  $9 \pm 2.87$ ) and Biochar + FM ( $5 \pm 2$  and  $9 \pm 2.91$  leaves respectively), while the lowest number of leaves were recorded under PPFM ( $4 \pm 2$  and  $6 \pm 2.08$ ) and control treatment ( $4 \pm 2$  and  $7 \pm 2.38$ ) (Figure 1).


Like leaves number, the plants height also increased significantly over time. Thus, at four months after transplanting (T4) we note at Site 1 a highest height under treatments PPFO ( $8.56 \pm 0.54$  cm), Biochar + FO + FM ( $8.34 \pm 0.54$  cm) and Biochar + FO ( $8.15 \pm 0.54$  cm); while at site 2, the highest height was observed under treatments PPFOM ( $19.75 \pm 0.69$  cm), PPFO ( $19.59 \pm 0.69$  cm), Biochar + FO + FM ( $19.12 \pm 0.69$  cm), followed by treatment Biochar + FO. ( $18.73 \pm 0.69$  cm) (Figure 2).

**Table 4. Linear mixed model outputs on growth data**

Treatments	Leaves number		Heigth (cm)	
	Site 1	Site 2	Site 1	Site 2
Biochar + FM	$5 \pm 2$ a	$9 \pm 2.91$ a	$5.18 \pm 2.03$ a	$11.66 \pm 3.77$ b
Biochar + FO	$5 \pm 2$ a	$9 \pm 2.87$ a	$5.36 \pm 2.28$ a	$13.65 \pm 4.01$ a
Biochar + FO + FM	$6 \pm 2$ a	$9 \pm 3.39$ a	$5.90 \pm 2.30$ a	$13.12 \pm 5.04$ ab
PPFM	$4 \pm 2$ b	$6 \pm 2.08$ b	$4.13 \pm 1.76$ b	$7.12 \pm 2.30$ c
PPFO	$6 \pm 2$ a	$10 \pm 3.77$ a	$5.67 \pm 2.26$ a	$14.51 \pm 4.32$ a
PPFOM	$5 \pm 2$ a	$9 \pm 3.54$ a	$5.24 \pm 1.85$ a	$13.62 \pm 4.73$ a
T0	$4 \pm 2$ b	$7 \pm 2.38$ b	$3.70 \pm 1.92$ b	$7.25 \pm 2.28$ c
T1	$3 \pm 1$ c	$6 \pm 1.08$ a	$3.45 \pm 0.94$ d	$7.14 \pm 1.94$ d
T2	$5 \pm 4$ b	$7 \pm 1.39$ b	$4.36 \pm 1.34$ c	$10.33 \pm 3.05$ c
T3	$5 \pm 2$ b	$9 \pm 1.93$ c	$5.41 \pm 1.96$ b	$12.50 \pm 3.95$ b
T4	$7 \pm 3$ a	$12 \pm 3.10$ d	$6.88 \pm 2.49$ a	$16.27 \pm 4.60$ a
Source of variation	Pr>(F/Chisq)	Pr>(F/Chisq)	Pr>(F/Chisq)	Pr>(F/Chisq)
Treatment	$< 2.2e-16$ ***	$< 2.2e-16$ ***	0.001248 **	934e-08 ***
Time	$6.726e-13$ ***	$< 2.2e-16$ ***	$< 2.2e-16$ ***	$7.811e-07$ ***
Treatment:Time	0.4254	0.9999	0.010149 *	$5.980e-07$ ***
Block	-	-	0.15782	0.20398
Treatment:Block	-	-	0.03199 *	0.08876
Time:Block	-	-	1.00000	0.80839
Treatment:Time:Block	-	-	$1.73e-09$ ***	$1.993e-06$ ***



**Figure 1. Influence of treatments and time on leaves number**



**Figure 2. Effect of treatment: time on plants height**

### 3.2 Influence of Cashew Nut Shells Biochar, Cattle Manure and Urea on Lettuce Yield

The table 5 presents the influence of treatments on lettuce yield. It shows that lettuce yield varied significantly depending on treatment ( $p < 0.05$ ). Thus, the best yields at the first site were obtained from plants having received the treatment PPFO ( $10.49 \pm 4.62$  t/ha), followed by Biochar + FO + FM ( $8.25 \pm 4.41$  t/ha). On site 2, the best yields are also obtained from plants having received the PPFO treatment ( $40.96 \pm 11.59$  t/ha) and Biochar + FO + FM ( $34.57 \pm 5.94$ ), followed by PPFOM ( $30.01 \pm 8.73$  t/ha) and Biochar + FO ( $27.15 \pm 1.25$  t/ha) treatments. The lowest yields were observed under the T0 and PPFM treatments at the two sites with respectively  $2.27 \pm 0.79$  t/ha,  $3.11 \pm 1.83$  t/ha on site 1 and  $6.48 \pm 3.84$  t/ha,  $3.78 \pm 1.38$  t/ha on site 2 (Table 5).

**Table 5. Effect of treatments on lettuce yield**

Treatments	Yield (t/ha)	
	Site 1	Site 2
Biochar + FM	$6.52 \pm 2.64$ ab	$16.25 \pm 3.32$ bc
Biochar + FO	$6.72 \pm 1.72$ ab	$27.15 \pm 1.25$ b
Biochar + FO + FM	$8.25 \pm 4.41$ ab	$34.57 \pm 5.94$ ab
PPFM	$3.11 \pm 1.83$ b	$6.48 \pm 3.84$ c
PPFO	$10.49 \pm 4.62$ a	$40.96 \pm 11.59$ a
PPFOM	$6.80 \pm 2.15$ ab	$30.01 \pm 8.73$ b
T0	$2.27 \pm 0.79$ b	$3.78 \pm 1.38$ c
Source of variation	Pr>(F/Chisq)	Pr>(F/Chisq)
Treatment	0.002912 **	2.22e-05 ***
1   Block	0.09621	0.1896
1   Treatment:Block	1.00000	1.00000

## 4. DISCUSSION

This study examined the impact of cashew nut shell biochar, cattle manure, and urea on growth and yield of lettuce (*Lactuca sativa*) in North and Central Benin. The results demonstrate significant effects of fertilizations modalities on lettuce growth and yield, highlighting the potential of organic and integrated fertilization strategies to enhance sustainable vegetable production.

### 4.1 Growth Parameters

The treatments influenced key growth parameters such as number of leaves and plant height. The treatments: Biochar + FO ( $10 \text{ t ha}^{-1}$  biochar +  $20 \text{ t ha}^{-1}$  cattle manure) and Biochar + FO + FM ( $10 \text{ t ha}^{-1}$  biochar +  $25 \text{ t ha}^{-1}$  urea +  $10 \text{ t ha}^{-1}$  cattle manure) consistently yielded the highest growth metrics across both sites, indicating that biochar, when combined with organic fertilizers, or both organic and mineral fertilizers can significantly boost lettuce growth. These results can be explained by the fact that biochar improved soil nutrient availability, especially that of nitrogen [26,40]. Indeed, Biochar can enhance nutrient retention and release rates, improving soil nutrient use efficiency [41, 42]. Biochar can enhance nutrient retention and release rates, improving soil nutrient use efficiency [41, 42]. ~~biochar can improve nutrient fixation and release rates, thereby improving soil nutrient use efficiency [41,42].~~ Nutrient enrichment of biochar allows the penetration of available nutrients into the micro and nanopores of the biochar and they are released slowly according to the synchrony between nutrient supply and plant demand [43,44]. The slow-release mechanism based on crop physiological needs increases nitrogen use efficiency (NUE), reduces leaching losses and increases nutrient bioavailability, resulting in increased crop productivity [45,46,47,48,49]. Moreover, manure stimulates soil microbial activity and further promotes nutrient release and transformation [50]. Liang et al.[13]observed in their study, an improvement in soil nutrient use efficiency and a more favorable soil environment for rice growth by the combined application of biochar and manure. Our results corroborate those of Sarmiento et al. [51], who also obtained a significant increase in growth parameters on the lettuce, by combining application of biochar and goat manure. This finding also aligns with finding of Glaser et al. [24] and Agegnehu et al. [28], who have reported improved plant growth due to the enhanced nutrient availability and soil structure associated with biochar amendments.

**Comment [U10]:** The discussion could benefit from a more explicit connection between the results and the underlying mechanisms. For example, you mention that biochar improved soil nutrient availability, especially nitrogen, but it would be helpful to directly link how this led to the observed growth improvements in lettuce.

### 4.2 Yield

Lettuce yield varied significantly depending on treatments, with the PPFO (40 t ha<sup>-1</sup> cattle manure) and Biochar + FO + FM (10 t ha<sup>-1</sup> biochar + 25 t ha<sup>-1</sup> urea +10 t ha<sup>-1</sup> cattle manure) treatments (followed by PPFOM: 20 t ha<sup>-1</sup> cattle manure + 50 kg ha<sup>-1</sup> urea and Biochar + FO: 10 t ha<sup>-1</sup> biochar + 20 t ha<sup>-1</sup> cattle manure) producing the highest yields. These results suggest that integrating biochar with organic and mineral fertilizers can substantially enhance crop production. The superior yields from these treatments can be attributed to the synergistic effects of improved soil fertility, increased nutrient retention, and better water holding capacity provided by the biochar [21,23]. Indeed, biochar mixed with minerals or organic amendments provides more available nutrients in the soil and increases nutrient absorption by plants [13,43,46]. Raj et al.[52]observed an increase in the availability of soil nutrients (total N, P, and K) after applying biochar combined with bovine urine. Pandit et al. [53] also showed an increase in available P up to 105 mg kg<sup>-1</sup> after applying nutrient-enriched biochar (co-composted biochar) compared to non-enriched biochar (38 mg kg<sup>-1</sup>). Several studies have reported significant positive effects of available P and K in the soil on crop productivity [46,54,55,57]. Our results align with those of Raj et al. [52], who demonstrated that biochar enriched with bovine urine increases corn yield, unlike biochar added separately, even when both received the same amount of organic and inorganic supplements. Similarly, Schmidt et al. [44] observed that biochar mixed with cow urine led to a pumpkin yield of 82.6 t·ha<sup>-1</sup>, an increase of over 300% compared to the treatment where only urine was applied, and an 85% increase compared to the treatment with biochar alone. Similar observations were noted by Sarmiento et al. [51], who evaluated the application of goat manure and biochar on lettuce yield in Brazil; and Lele et al. [58], who assessed the effect of mineral fertilizer combined with biochar on corn and cassava yield. These results showed that biochar improves the efficiency of fertilizer use [59]. Likewise, biochar co-applied with inorganic NPK improves soil quality, leading to better growth and yield of crops such as cowpea and rice [29,60]. Additionally, biochar mixed with minerals or organic amendments effectively reduces a significant portion of nitrogen losses to the environment [44,49,52]. High nitrogen losses from agricultural areas lead to reduced yields and add to economic and environmental costs [61]. Numerous studies have shown that biochar mixed with minerals or organic amendments leads to increased land use and crop yields while significantly reducing nitrogen losses through leaching and emissions [46,49,62]. Tan et al.[63]found that cumulative N<sub>2</sub>O emissions decreased by 33.7% when rice husk biochar + NPK fertilizer was applied to soils compared to the treatment where NPK fertilizer was applied without biochar.

#### 4. CONCLUSION

The study demonstrated that using cashew nut shell biochar, cattle manure, and urea as soil amendments can significantly improve growth and yield of lettuce cultivation in North and Central Benin. Among the biochar treatments, those combining biochar, cattle manure, and urea Among the treatments including biochar, the treatments which combines biochar, cattle manure, and urea (10 t ha<sup>-1</sup> biochar + 25 t ha<sup>-1</sup> urea +10 t ha<sup>-1</sup> cattle manure) and biochar and cattle manure (10 t ha<sup>-1</sup> biochar + 20 t ha<sup>-1</sup> cattle manure), consistently resulted in a good yield across both sites, comparatively to producers' mineral fertilization practice. This combination also enhanced plant growth parameters such as height and leaf number, thereby contributing to sustainable agricultural practices. Given the abundant availability of cashew nut shells in the region, promoting the adoption of their derived biochar in addition to cattle manure, by reducing mineral fertilizers rate and dependence of cattle manure, could lead to more sustainable and profitable lettuce production systems. Future research should focus on economical profitability and impact of these practices on soil fertility to further validate these findings.

#### CONSENT (WHEREEVER APPLICABLE)

Not applicable

#### ETHICAL APPROVAL (WHEREEVER APPLICABLE)

Not applicable

#### REFERENCES

- 1.FAO. Transforming food systems for affordable healthy diets. Rome: FAO; 2020. 287 p. (The state of food security and nutrition in the world).
- 2.Martin T, Saidi M, Komlan FA, Simon S, Kasina M, Vidogbéna F, et al. Des filets anti insectes pour protéger les cultures maraichères en Afrique Subsaharienne: une technologie rentable et adaptée aux conditions climatiques. In 2014.French.

3. Ekesi S, Chabi-Olaye A, Subramanian S, Borgemeister C. Horticultural pest management and the African economy: successes, challenges and opportunities in a changing global environment. In 2009. p. 165-83.
4. Weinberger K, Lumpkin TA. Diversification into horticulture and poverty reduction: a research agenda. *World development*. 2007;35(8):1464-80.
5. Adjovi ISM, Adjovi C, Ayi-Fanou L, Ayandji P, Sanni A. Les pratiques d'utilisation des pesticides par les maraichers au Bénin: perspectives sociologiques. *Global Journal of Arts, Humanities and Social Sciences*. 2020;8(2):22-45. French.
6. Kpéra GN, Segnon AC, Saïdou A, Mensah GA, Aarts N, van der Zijpp AJ. Towards sustainable vegetable production around agro-pastoral dams in Northern Benin: current situation, challenges and research avenues for sustainable production and integrated dam management. *Agriculture & food security*. 2017;6:1-13.
7. MAEP. Plan Stratégique de Relance du Secteur Agricole (PSRSA). 2011. French.
8. Kouglénou SD, Jerrold Agbankpé A, Béhanzin JG, Victorien Dougnon T, Aniambossou AV, Baba-Moussa L, et al. Microbiological Safety of Leafy Vegetables Produced at Houeyiho and Sèmè-Kpodji Vegetable Farms in Southern Benin: Risk Factors for *Campylobacter* spp. *International journal of food science*. 2019;2019(1):8942608.
9. World Health Organization. Carbohydrate intake for adults and children: WHO guideline. World Health Organization. 2023.
10. Kouévi TA, Adé CK, N'Danikou S, Mongbo RL, Komlan C, Dagnon GN, et al. Preferences of market gardeners for traditional vegetables and associated factors in urban areas of southern Benin. 2023;11(5). *Future of Food: Journal on Food, Agriculture and Society*. Available: <https://kobra.uni-kassel.de/handle/123456789/15363>
11. Avadí A, Hodomihou NR, Amadji GL, Feder F. LCA and nutritional assessment of southern Benin market vegetable gardening across the production continuum. *Int J Life Cycle Assess*. 2021;26(10):1977-97.
12. Liu F, Wang X, Chi Q, Tian M. Spatial variations in soil organic carbon, nitrogen, phosphorus contents and controlling factors across the "Three Rivers" regions of southwest China. *Science of The Total Environment*. 2021;794:148795.
13. Liang D, Ning Y, Ji C, Zhang Y, Wu H, Ma H, et al. Biochar and Manure Co-Application Increases Rice Yield in Low Productive Acid Soil by Increasing Soil pH, Organic Carbon, and Nutrient Retention and Availability. *Plants*. 2024;13(7):973.
14. Ahmad M, Ishaq M, Shah WA, Adnan M, Fahad S, Saleem MH, et al. Managing phosphorus availability from organic and inorganic sources for optimum wheat production in calcareous soils. *Sustainability*. 2022;14(13):7669.
15. Thibault C, Lecompte F. Gestion de la fertilité des sols en cultures légumières et maraichères. 2018. French.
16. Zhou Z, Zhang S, Jiang N, Xiu W, Zhao J, Yang D. Effects of organic fertilizer incorporation practices on crops yield, soil quality, and soil fauna feeding activity in the wheat-maize rotation system. *Frontiers in Environmental Science*. 2022;10:1058071.
17. MAEP. Plan Stratégique de Développement du Secteur Agricole (PSDSA): Orientations stratégiques 2025 Plan National d'Investissements Agricoles et de Sécurité Alimentaire et Nutritionnelle PNIASAN 2017 - 2021. 2017;135. French.
18. Godjo T, Tagutchou JP, Naquin P, Gourdon R. Valorisation des coques d'anacarde par pyrolyse au Bénin. *Environnement, Ingénierie & Développement*. 2015. French.
19. Kyei SK, Eke WI, Nagre RD, Mensah I, Akaranta O. A comprehensive review on waste valorization of cashew nutshell liquid: Sustainable development and industrial applications. *Cleaner Waste Systems*. 2023;6:100116.
20. Samb C, Biteye M, Faye E, Diarra R, Diaw N, Thiam M, et al. Caractérisation agromorphologique, chimique et biochimique des noix brutes de trois provenances d'anacardiens (*Anacardium occidentale* L.) au Sud du Sénégal. *Agronomie Africaine*. 2021;33(2):177-89. French.
21. Janczak D, Malińska K, Czekala W, Cáceres R, Lewicki A, Dach J. Biochar to reduce ammonia emissions in gaseous and liquid phase during composting of poultry manure with wheat straw. *Waste Management*. 2017;66:36-45.
22. Frimpong KA, Amoakwah E, Osei BA, Arthur E. Changes in soil chemical properties and lettuce yield response following incorporation of biochar and cow dung to highly weathered acidic soils. *Journal of Organic Agriculture and Environment*. 2016;4(1):28-39.
23. Liang B, Lehmann J, Solomon D, Kinyangi J, Grossman J, O'Neill B, et al. Black carbon increases cation exchange capacity in soils. *Soil science society of America journal*. 2006;70(5):1719-30.
24. Glaser B, Lehmann J, Zech W. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal—a review. *Biology and fertility of soils*. 2002;35:219-30.
25. Hossain MZ, Bahar MM, Sarkar B, Donne SW, Ok YS, Palansooriya KN, et al. Biochar and its importance on nutrient dynamics in soil and plant. *Biochar*. 2020;2:379-420.
26. Sahin O, Taskin M, Kaya E, Atakol O, Emir E, Inal A, et al. Effect of acid modification of biochar on nutrient availability and maize growth in a calcareous soil. *Soil Use and Management*. 2017;33(3):447-56.
27. Taskin MB, Kadioglu YK, Sahin O, Inal A, Gunes A. Effect of acid modified biochar on the growth and essential and non-essential element content of bean, chickpea, soybean, and maize grown in calcareous soil. *Communications in Soil Science and Plant Analysis*. 2019;50(13):1604-13.

28. Agegnehu G, Srivastava AK, Bird MI. The role of biochar and biochar-compost in improving soil quality and crop performance: A review. *Applied soil ecology*. 2017;119:156-70.
29. MacCarthy DS, Darko E, Nartey EK, Adiku SGK, Tettey A. Integrating Biochar and Inorganic Fertilizer Improves Productivity and Profitability of Irrigated Rice in Ghana, West Africa. *Agronomy*. 2020;10(6):904.
30. Tahery S, Munroe P, Marjo CE, Rawal A, Horvat J, Mohammed M, et al. A comparison between the characteristics of a biochar-NPK granule and a commercial NPK granule for application in the soil. *Science of the Total Environment*. 2022;832:155021.
31. Qian S, Zhou X, Fu Y, Song B, Yan H, Chen Z, et al. Biochar-compost as a new option for soil improvement: Application in various problem soils. *Science of The Total Environment*. 2023;870:162024.
32. Kora O, Guidibi E. Monographie de la Commune de Tchaourou. Bénin: Afrique Conseil. 2006. French.
33. INSAE. Cahier des villages et quartiers de ville: Département de l'Atacora. Institut National de la Statistique et de l'Analyse Economique (INSAE). 2016. French.
34. Hazelton P, Murphy B. Interpreting soil test results: What do all the numbers mean? CSIRO publishing. 2016.
35. Jackson ML. Soil chemical analysis: advanced course: a manual of methods useful for instruction and research in soil chemistry, physical chemistry of soils, soil fertility, and soil genesis. UW-Madison Libraries parallel press. 2005.
36. Fleck A. The Determination of Organic Nitrogen. *Proceedings of the Association of Clinical Biochemists*. 1967;4(7):212-5.
37. Bray RH, Kurtz LT. Determination of total, organic, and available forms of phosphorus in soils. *Soil science*. 1945;59(1):39-46.
38. Helmke PA, Sparks DL. Lithium, sodium, potassium, rubidium, and cesium. *Methods of soil analysis: Part 3 chemical methods*. 1996;5:551-74.
39. Bell JS. On the einstein podolsky rosen paradox. *Physics Physique Fizika*. 1964;1(3):195.
40. Zhu X, Li C, Li J, Xie B, Lü J, Li Y. Thermal treatment of biochar in the air/nitrogen atmosphere for developed mesoporosity and enhanced adsorption to tetracycline. *Bioresource Technology*. 2018;263:475-82.
41. Liu Y, Li H, Hu T, Mahmoud A, Li J, Zhu R, et al. A quantitative review of the effects of biochar application on rice yield and nitrogen use efficiency in paddy fields: A meta-analysis. *Science of the Total Environment*. 2022;830:154792.
42. Spokas K, Novak J, Masiello C, Johnson M, Colosky E, Ippolito J, et al. Physical disintegration of biochar: an overlooked process. *Environmental Science & Technology Letters*. 2014;1(8):326-32.
43. Schmidt H, Pandit BH, Cornelissen G, Kammann CI. Biochar-based fertilization with liquid nutrient enrichment: 21 field trials covering 13 crop species in Nepal. *Land Degradation & Development*. 2017;28(8):2324-42.
44. Schmidt HP, Pandit BH, Martinsen V, Cornelissen G, Conte P, Kammann CI. Fourfold increase in pumpkin yield in response to low-dosage root zone application of urine-enhanced biochar to a fertile tropical soil. *Agriculture*. 2015;5(3):723-41.
45. Borchard N, Schirrmann M, Cayuela ML, Kammann C, Wrage-Mönnig N, Estavillo JM, et al. Biochar, soil and land-use interactions that reduce nitrate leaching and N<sub>2</sub>O emissions: a meta-analysis. *Science of the Total Environment*. 2019;651:2354-64.
46. Kammann CI, Schmidt HP, Messerschmidt N, Linsel S, Steffens D, Müller C, et al. Plant growth improvement mediated by nitrate capture in co-composted biochar. *Scientific reports*. 2015;5(1):11080.
47. Kizito S, Luo H, Lu J, Bah H, Dong R, Wu S. Role of nutrient-enriched biochar as a soil amendment during maize growth: Exploring practical alternatives to recycle agricultural residuals and to reduce chemical fertilizer demand. *Sustainability*. 2019;11(11):3211.
48. Munera-Echeverri JL, Martinsen V, Strand LT, Zivanovic V, Cornelissen G, Mulder J. Cation exchange capacity of biochar: An urgent method modification. *Science of the total environment*. 2018;642:190-7.
49. Puga AP, Grutzmacher P, Cerri CEP, Ribeirinho VS, de Andrade CA. Biochar-based nitrogen fertilizers: Greenhouse gas emissions, use efficiency, and maize yield in tropical soils. *Science of the Total Environment*. 2020;704:135375.
50. Iqbal A, Liang H, McBride SG, Yuan P, Ali I, Zaman M, et al. Manure applications combined with chemical fertilizer improves soil functionality, microbial biomass and rice production in a paddy field. *Agronomy Journal*. 2022;114(2):1431-46.
51. Sarmiento JJA, Costa CC, Dantas MV, Lopes KP, De Macedo IC, Bomfim SMP, et al. Productivity of lettuce under organic fertilization. *Journal of Agricultural Science*. 2019;11(1):333-43.
52. Pandit NR, Sipkhan P, Sharma SS, Dawadi D, Vista SP, Raut P. Cattle-Urine-Enriched Biochar Enhances Soil Fertility, Nutrient Uptake, and Yield of Maize in a Low-Productive Soil. *Nitrogen*. 2024;5(1):16-27.
53. Pandit NR, Schmidt HP, Mulder J, Hale SE, Husson O, Cornelissen G. Nutrient effect of various composting methods with and without biochar on soil fertility and maize growth. *Archives of Agronomy and Soil Science*. 2020.
54. Arif M, Ali S, Ilyas M, Riaz M, Akhtar K, Ali K, et al. Enhancing phosphorus availability, soil organic carbon, maize productivity and farm profitability through biochar and organic-inorganic fertilizers in an irrigated maize agroecosystem under semi-arid climate. *Soil Use and Management*. 2021;37(1):104-19.

55. Gautam DK, Bajracharya RM, Sitaula BK. Effects of Biochar and Farm Yard Manure on Soil Properties and Crop Growth in an Agroforestry System in the Himalaya. *SAR*. 2017;6(4):74.
56. Martinsen V, Mulder J, Shitumbanuma V, Sparrevik M, Børresen T, Cornelissen G. Farmer-led maize biochar trials: Effect on crop yield and soil nutrients under conservation farming. *Journal of Plant Nutrition and Soil Science*. 2014;177(5):681-95.
57. Yamato M, Okimori Y, Wibowo IF, Anshori S, Ogawa M. Effects of the application of charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. *Soil science and plant nutrition*. 2006;52(4):489-95.
58. Lele B, Lejoly J, Kachaka C. Effet de l'application d'engrais minéral complet NPK et de biochar sur les performances de la culture pure du manioc et de l'association manioc-maïs dans les conditions du plateau des Batéké en République Démocratique du Congo (RDC). *Tropicicultura*. 2016;34(1):45-55. French.
59. Abewa A, Yitafaru B, Selassie YG, Amare T. The role of biochar on acid soil reclamation and yield of Teff (*Eragrostis tef* [Zucc] Trotter) in Northwestern Ethiopia. *Journal of Agricultural Science*. 2014;6(1):1.
60. Phares CA, Atiah K, Frimpong KA, Danquah A, Asare AT, Aggor-Woananu S. Application of biochar and inorganic phosphorus fertilizer influenced rhizosphere soil characteristics, nodule formation and phytoconstituents of cowpea grown on tropical soil. *Heliyon*. 2020;6(10).
61. Ortuzar-Iragorri A, Castellón A, Besga G, Aizpurua A, Fuertes-Mendizabal T, Estavillo JM. Nitrogen losses: gaseous and leached nitrogen balance. *Global wheat production*. 2018;79-98.
62. Cayuela ML, Van Zwieten L, Singh B, Jeffery S, Roig A, Sánchez-Monedero M. Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis. *Agriculture, Ecosystems & Environment*. 2014;191:5-16.
63. Tan G, Wang H, Xu N, Liu H, Zhai L. Biochar amendment with fertilizers increases peanut N uptake, alleviates soil N<sub>2</sub>O emissions without affecting NH<sub>3</sub> volatilization in field experiments. *Environmental Science and Pollution Research*. 2018;25:8817-26.