

Review Article

Impact of organic amendments on Entomopathogenic Nematodes-an overview

Abstract:

Organic amendments are applied in the field in order to increase the organic matter content of soil for sustainable and healthy soil. Healthy soil enhances crop yield. Build-up of insect pest populations under organic production is problematic which not only reduces yield but also lowering crop quality. Integrated Pest Management (IPM) strategies that incorporate cultural management, biological management, use of resistant varieties are viable options for reduction of pest population. Among the biocontrol agents, Entomopathogenic nematodes (EPNs) are efficacious against various insect pests in a variety of habitats. However, the effectiveness of EPNs against insect pests is largely dependent on the method of application and nematode survival under field conditions. Cultural management practice such as usage of organic amendments can alter the soil balance that enables their occurrence and activity. Understanding the interactions is essential to reveal ways to enhance the potential of EPN as biocontrol agents in a particular soil type, environmental condition and agricultural scenario. This review provides brief overviews of impact of sawdust on EPNs from the standpoint of pest management.

Key words: Organic agriculture, Pests, Cultural management, Biological management, Organic amendments, sawdust, Entomopathogenic nematodes (EPNs).

Introduction

Agricultural production relies on healthy soils that guarantee essential soil functions such as carbon, nutrient, and water cycling. Traditionally, soil quality has been characterized by the presence of nutrients and water and their availability to the crop (Patzelet *al.*, 2000). The development of agricultural practices have largely enhanced crop yield; but protecting the crops against herbivores and diseases is problematic (Oerke, 2005). Insect pests destroy 18-26% of the world's total crop production annually, with an estimated cost of more than \$470 billion (Kergunteuilet *al.*, 2016). Attempts to control this problem with chemical pesticides have resulted in significant environmental damage, harm to beneficial insects, humans and other animals.

Alternative to chemical pesticides include IPM strategies that incorporate cultural practices like crop rotation, cover crops and organic amendments; use of resistant varieties; biocontrol agents including parasitoid wasps, entomopathogenic fungi, Bt toxins, entomopathogenic nematodes (EPNs) etc. Among all the biocontrol agents, EPNs in the families Steinernematidae and Heterorhabditidae are efficacious against numerous insect pests in a variety of habitats. There are no negative impacts of EPNs on the environment, human health and non-target organisms when compared to pesticides (Ehlers, 1996). Their potential to recycle in their host populations could be further employed for long-term pest management. However, in the agro-ecosystem, EPNs are affected by various abiotic soil properties such as soil texture, moisture, temperature, and soil organic matter, which might be drastically altered by agricultural management practices (Jaffuel et al., 2016). Farmers add soil amendments such as compost to increase organic matter content, improve soil aeration, soil porosity, and soil water holding capacity. EPNs in many agricultural soils are mixed with soil amendments (Uluget al., 2014). Application in the field by using compost as a carrier might protect the nematodes against UV damage, buffer temperature extremes and promote contact between the nematodes and the pest insects (Georgis et al., 2006).

Entomopathogenic Nematodes (EPNs) in the soil environment

As the soil entomopathogens are intimately associated with their environment, any disruption of the soil structure (i.e. tillage) could cause a change in activity or survival of pathogens. Soil physical and chemical characteristics play an important role in the abundance, distribution and structure of nematode communities. EPNs occur in soils of natural and agricultural areas worldwide (Gaugler 1988). Environmental variables, soil characteristics and anthropogenic activities may affect their diversity and occurrence. They occur in their non-feeding stage, called infective juvenile (IJ), capable of locating suitable hosts (Stock, 2015). IJs invade insect bodies through the mouth, anus, spiracles, or cuticle and release their bacterial symbionts, γ -Proteobacteria (*Xenorhabdus* spp. bacteria for *Steinernema* spp. and *Photorhabdus* spp. bacteria for *Heterorhabditis* spp.) into the insect's hemolymph (Boemare, 2002). The bacteria release toxins to kill the insect, though sometimes are aided by anti-immune agents produced by the nematodes themselves, and proliferate in the cadaver. The IJs consume the bacteria, complete their development, and reproduce. After one or more generations of nematodes inside the cadaver, new IJs leave to seek new hosts in the soil (Kaya, 1990). As the soil is their native habitat, EPNs have a marked potential to control soil insects where there is protection from UV

radiation and rapid desiccation (Campos-Herrera *et al.*,2008). The effectiveness of EPNs against insect pests is largely dependent on the method of application and nematode survival under field conditions (Peters, 1996). Inside the soil matrix, EPNs are exposed to a broad biodiversity of soil organisms, potential insect host, predators, microbial pathogens, free-living nematodes as competitors, and chemical attractants and repellents released by plants and other soil organisms (Lewis *et al.*, 2015). EPNs use soil cracks and openings caused by root growth to move through the soil matrix. EPNs exploit volatile organic compounds emitted by roots in response to herbivory to locate damaged root systems and thereby insect hosts. EPNs foraging behavior is therefore greatly influenced by root architecture and their effectiveness negatively correlated with root complexity (Demartaet *et al.*, 2014). All these factors affect the survivorships, reproduction, host searching capacity and movement of EPN in many agricultural soils (Ishibashi and Kondo,1986; Helmberger *et al.*,2017). For survival, IJs must cope with soil and environmental conditions, overcome competitors for hosts, and avoid predators and other soil organisms (Stuart *et al.*, 2015). When applied, many EPNs do not persist in soil longer than a few weeks or months (Ebssa and Koppenhofer, 2011), some native EPN populations, adapted to a particular environment, have been shown to persist over many years (Shields *et al.*, 2020). Presence of hosts within the field can provide a chance for EPN to reproduce and increase persistence of EPN up to two years in the field (Susurluket *et al.*,2011).

Agricultural management practices: Use of compost

Agricultural management practices can modify soil properties in ways that may disrupt or enhance the abundance and activity of beneficial organisms in the soil. Agricultural practices such as irrigation, tillage and fertilization may affect the viability, reproduction and insecticidal activities of EPN. Soil water holding capacity is one of the most vital soil factors in crop production. The water holding capacity of the soil can be enhanced by increasing soil organic matter content due to its effects on soil texture, porosity, and bulk density. Furthermore, compost can be used to improve crop production, promote plant growth, and suppress, prevent and control soil-borne plant pathogens (Herren *et al.*,2018). A large diversity of microorganisms has the potential to protect plants against pests and diseases if applied in an appropriate manner.The application of compost enriched with antagonists or beneficial microorganisms holds great promise in agricultural and horticultural settings since compost itself inherently infers benefits to soil and plant growth. Compost can increase the N and P availability in soil, as well as the

activity and the size of its microbial population. Furthermore, it can be used to improve crop production, plant growth and suppress diseases that are caused by soil-borne plant pathogens. Beneficial effects of compost on soil quality or plant growth (e.g. increased nutrient capacity, disease suppression) are dependent on the quality of the organic matter, presumably including its inhabiting biology and its biochemical stability. Using compost or a compost and mulch mix improved plant growth and yield (Burkhard *et al.*,2009), provided pest control, and benefitted natural enemies (Renkema *et al.*,2016). Learning more about the factors that determine the prevalence of the organisms that are directly associated with EPN, can provide additional information on how farming practices might contribute to this important feature of soil health. Soils with high levels of natural organic matter and minimal soil disturbance sustain larger numbers of soil microorganisms, including EPN and associate organisms (Susurluk , 2008). Diverse interactions between EPNs and other members of the soil community exist, from plants and fungi to arthropods and annelids (Blanco-Pérez *et al.*,2020) . These interactions influence EPN performance under field conditions, specifically, if they contribute to the variability limiting EPN efficacy and wide-scale adoption. Soil texture, temperature, moisture, agronomic practices, soil antagonists, and host resources are major factors affecting nematode persistence in the soil environment (Lawrence et al.,2006), and could be responsible for the differences in detection of entomopathogenic nematodes among different habitats. Agricultural management affects both soil quality variables and EPN distribution. Soil conditions could also be manipulated so that nematodes are established and sustained for prolonged periods after inundative releases. If soil organic content does increase both water holding capacity and water retention then possibly it may also extend the time over which an initial application of the IJs of EPNs would be efficacious as biocontrol agents(Shapiro et al., 1999).

Various studies have measured the effect of different organic and (physical status, nutrient content, nutrient mineralization rate, and decomposition products) inorganic fertilizers on nematode fitness. The effect of inorganic fertilizers varied according to their composition, the EPN species exposed, and the duration of its exposure to the used fertilizer (Kaya 1990). The addition of compost was shown to increase the overall nematode presence, while modifying species composition, favouring bacterivorous taxa and decreasing the ratio of fungivorous nematodes (Sikora *etal.*,2005). Composts may also improve the virulence of and pest mortality rates caused by EPNs compared to certain soil types (Khumalo *et al.*, 2021), although

immaturely-cured compost may negatively affect EPN virulence (Herren *et al.*, 2018). In general, application of EPNs into composts, mulches, or other soil amendments rather than soil may protect EPNs from UV damage and buffer them from temperature and moisture extremes while promoting EPN movement and contact with pests (Georgis *et al.*,2006). Plant bodies contain various amounts of cellulose, chitin, flavonoids, lignin, phenols, ricin, saponins, tannins, terpenoids, etc., which significantly affect the growth of microorganisms and thus affect the resulting cooperating soil microbiome during the decomposition. Compost improves soil quality due to increased organic matter content and soil microbial populations, and it has also shown that it improves plant resistance to insect pests (Kurzemann *et al.*,2020).However, the accumulation of compost in soils can affect the presence and abundance of the nematode community when applied regularly over time (Puissant *et al.*,2021). Compost can be used as a carrier for field application of EPN (Hoitink and Boehm,1999; Herren *et al.*, 2018; Georgis *et al.*, 2006) use in a wide array of agricultural and horticultural systems (Lacey *et al.*, 2015).Application of organic manure may encourage nematode establishment and recycling. Bednarek and Gaugler (1997) observed that organic manure resulted in increased densities of native population of *Steinernema feltiae* as influenced by host population under field condition. Campos-Herrera *et al.*, (2015;2019,2021) found a positive effect of organic soil management on EPN populations. Application in the field by using compost as a carrier might protect the nematodes against UV damage, buffer temperature extremes and promote contact between the nematodes and the insect(Georgis *et al.*,2006).

Effect of Sawdust as organic amendment

Sawdust are organic waste of the saw-milling, pulp, paper, and wood processing industries. When used as mulch, sawdust mulch enhances soil structure, improves aeration in heavy soils, enhances water absorption and penetration, and conserves moisture and reduce evaporation and ultimately the distribution of soil elements within the plant growth profile. Sawdust is a carbonaceous organic substance with a high carbon to nitrogen ratio 300:1. Application of sawdust mulch has been found to increase soil oxygen diffusion rate, promote a more uniform soil temperature, reduce surface crusting and soil bulk density, and improve aeration porosity and soil moisture content. Sawdust mulch is a viable and sustainable soil management practice (Abiodun *etal.*,2022). Ground covers by applying sawdust can significantly affect soil temperature and moisture in the top 12cm of soil, which may directly influence pathogen activity

or survival. Type of tillage appeared to be the primary factor affecting soil entomopathogens whereas conservation tillage apparently conserves entomopathogens and alternate hosts (Duncan *et al.*,2007).

A compost and woodchip/sawdust mulch caused mortality of Japanese beetle, *Popillia japonica* population in blueberry with application of *H.bacteriophora* when temperatures are optimal to mulches (Renkema and Parent,2021) .

Deol *et al.*,(2011) used a peat and bark based potting medium to deliver EPN to the soil via the use of infected host cadavers, concluding that this application method extended nematode survival and virulence to 180 days (at 25°C) in comparison with 60 days through aqueous application. The use of compost as a carrier to deliver EPN directly to the soil could therefore be useful for agricultural settings.

Heterorhabditiszealandica together with pine chips, wheat straw, pine wood shavings, blackwood and applewood chips were evaluated to control diapausing codling moth, *Cydia pomonella* larvae; after application it was observed that IJs of *H. zealandica* were having < 10% persistence(de Waal et al.,2011).. The type of compost used (mulch soil amendment) based on its feedstock composition, the concentration of nutrients, and organic matter mixed with a specific type of soil texture had a significant effect on the tolerance of *H.bacteriophora* to different dehydration periods. Using Petri dish assays, the mulch soil amendment resulted in the highest *G.mellonella* larval mortality rate as compared to different organic compost application concentrations (mass:mass). Virulence and efficacy decreased over 7 days of progressive dehydration. Infectivity and survivorship of IJs of *H.bacteriophora* was improved by the use of organic compost as the soil undergoes progressive moisture loss (Khumalo *et al.*, 2021).

Conclusion and future perspectives

There is increasing interest in identifying the factors that define EPN population dynamics in agro-ecosystems. To improve the efficacy of EPNs as biocontrol agent in the field, knowledge about the kinds of impact, which the physical nature of the agricultural soil environment may have on nematode survival, persistence, and host location, is important. Lack of predictability is partly a function of poor nematode persistence in the soil. Around 10% of the initial inoculum of EPNs may be present in the soil seven days after application(Curran,1993). Many agricultural practices expose EPNs to extreme conditions (temperature,UVlight etc.) that significantly reduce their biocontrol potential, even reaching the extinction of the natural populations. Their

identification can allow us establishing the best practices to favor suitable ecological scenarios to enhance their activity. Improper selection of nematode species and strains for the climatic conditions at the release site as well as for the insect community present have been cited as possible factors affecting field performance (Koppenhofer *et al.*,2020). Identifying the abiotic and biotic factors responsible for the persistence of endemic nematode populations, particularly the host community, may improve the matching of species and strains with the conditions of the habitat in which nematodes are released (Brust, 1991). Given the drawbacks of the current application methods and the low field survival rates of EPN, there is a need for more efficient delivery methods for EPN into the field either through application of compost like sawdust to increase their persistence (Shapiro-Ilan and Dolinski,2015). As a first step towards the development of new biological control methods for sustainable agriculture, a comprehensive inventory of selected beneficial organisms of a given agroecosystem is required, together with application method as well as knowledge about the factors that might determine their abundance and survival.

References

- Abiodun FO., Isola JO, Smart MO. 2022. Influence of sawdust mulch on soil properties, growth and yield performance of okra *Abelmoschus esculentus* (L.) Moench in an alfisol. *FUTY Journal of the Environment*. 16(3):12-22.
- Bednarek A. ,Gaugler R. 1997. Compatibility of soil amendments with entomopathogenic nematodes. *Journal of Nematology*, 29 (2): 220-227
- Blanco-Pérez R, Sáenz-Romo MG , Vicente-Díez I , Ibáñez-Pascual S , Martínez-Villar E , Marco-Mancebón VS, Pérez-Moreno I, Campos-Herrera R. 2020. Impact of vineyard ground cover management on the occurrence and activity of entomopathogenic nematodes and associated soil organisms. *Agriculture, Ecosystems and Environment*. 301 (2020) 10702. <https://doi.org/10.1016/j.agee.2020.107028>
- Boemare N., 2002. Biology, taxonomy and systematic of *Photorhabdus* and *Xenorhabdus*. In: *Entomopathogenic Nematology*. ed. R.Gaugler. Wallingford, UK: CABI Publishing pp.35-56. [Doi.org/ 10.1079/ 9780851995670.0035](https://doi.org/10.1079/9780851995670.0035)
- Brust GE. 1991. Augmentation of an endemic entomogenous nematode by agroecosystem manipulation for the control of a soil pest. *Agriculture, Ecosystems & Environment*. 36(3-4): 175-184. [https:// doi.org/ 10.1016/ 0167-8809\(91\)90014-O](https://doi.org/10.1016/0167-8809(91)90014-O).

- Burkhard NE., Lynch DH., Percival DC., Sharifi M. 2009. Organic mulch impact on vegetation dynamics and productivity of highbush blueberry under organic production. *HortScience*. 44:1-9.
- Campos-Herrera R., Gómez-Ros JM, Escuer M, Cuadra L, Barrios L, Gutiérrez C, 2008 Diversity, occurrence, and life characteristics of natural entomopathogenic nematode populations from La Rioja (Northern Spain) under different agricultural management and their relationships with soil factors. *Soil Biology and Biochemistry*. 40(6): 1474-1484. <https://doi.org/10.1016/j.soilbio.2008.01.002>.
- Campos-Herrera R., Piedra-Buena, A. Escuer M., Montalba B., Gutierrez C. 2010. Effect of seasonality and agricultural practices on occurrence of entomopathogenic nematodes and soil characteristics in La Rioja (Northern Spain). *Pedobiologia*, 53: 253-258.
- Campos-Herrera, R., Blanco-Pérez, R., Bueno-Pallero, F.A., Duarte, A., Nolasco, G., Sommer, R.J., Rodríguez Martín, J.A., 2019. Vegetation drives assemblages of entomopathogenic nematodes and other soil organisms: evidence from the Algarve, Portugal. *Soil Biol. Biochem.* 128, 150-163. <https://doi.org/10.1016/j.soilbio.2018.10.019>
- Campos-Herrera, R., El-Borai, F.E., Duncan, L.W., 2015. Modifying soil to enhance biological control of belowground dwelling insects in citrus groves under organic agriculture in Florida. *Biol. Control* 84: 53-63. <https://doi.org/10.1016/j.biocontrol.2015.02.002>.
- Campos-Herrera, R., Vicente-Díez, I., Blanco-Pérez, R., Chelkha, M., Gonzalez-Trujillo, M., del, M., Puelles, M., Cepulyte-Rakauskiene, R., Pou, A., 2021. Positioning entomopathogenic nematodes for the future viticulture: exploring their use against biotic threats and as bioindicators of soil health. *Turk. J. Zool.* 45, 335-346. <https://doi.org/10.3906/zoo-2106-40>
- Curran J. 1993. Post-application biology of entomopathogenic nematodes in soil. pp.67-78. In: Bedding RA., Akhurst R. and Kaya HK (Eds.) *Nematodes and the Biological Control of Insects*. CSIRO Publications, East Melbourne, Australia.
- de Waal, J.Y., Malan, A.P., Addison, M.F., 2011. Evaluating mulches together with *Heterorhabditis zealandica* (Rhabditida: Heterorhabditidae) for the control of diapausing codling moth larvae, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *Biocontrol Sci. Technol.* 21: 255-270. <https://doi.org/10.1080/09583157.2010.540749>.

- Demarta L, Hibbard BE, Bohn MO, Hiltbold I. 2014. The role of root architecture in foraging behavior of entomopathogenic nematodes. *J Invertebr Pathol.* 122:32-39.
- Deol YS., Jagdale GB., Canas L., Grewal PS. 2011. Delivery of entomopathogenic nematodes directly through commercial growing media via the inclusion of infected host cadavers: a novel approach. *Biol. Control.* 58(1):60-67. doi.org /10. 1016 /j. biocontrol. 2011.03.002
- Duncan LW., Graham JH., Zellers J., Bright DB., Dunn DC., El-Borai FE., Porazinska DL. 2007. Food web responses to augmenting the entomopathogenic nematodes in bare and animal manure-mulched soil. *J. Nematol.* 39: 176-189.
- EbssaL., Koppenhofer AM. 2011. Efficacy and persistence of entomopathogenic nematodes for black cutworm control in turfgrass. *Biocontrol Science and Technology.* 21:779-796
- Ehlers RU. 1996. Current and future use of nematodes in biocontrol: Practice and commercial aspects in regard to regulatory policies. *Biocontrol Science and Technology*, 6: 303-316.
- Gaugler R. 1988. Ecological considerations in the biological control of soil-inhabiting insects with entomopathogenic nematodes. *Agriculture, ecosystems & environment.* 24: 351-360.
- Georgis, R., Koppenhöfer, A. M., Lacey, L. A., Bélair, G., Duncan, L. W., Grewal, P. S., Samish, M., Tan, L., Torr, P. and van Tol, R. W. H. M. 2006. Successes and failures in the use of parasitic nematodes for pest control. *Biological Control* 38:103-23.
- Helmberger, M.S., Shields, E.J., Wickings, K.G., 2017. Ecology of belowground biological control: entomopathogenic nematode interactions with soil biota. *Appl. Soil Ecol.* 121, 201–213. <https://doi.org/10.1016/j.apsoil.2017.10.013>.
- Herren, G. L., Binnemans, I., Joos, L., Viaene, N., Ehlers, R. U., Vandecasteele, B., Bert, W. and Steel, H. 2018. Compost as a carrier medium for entomopathogenic nematodes - the influence of compost maturity on their virulence and survival. *Biological Control* 125:29-38.
- Hoitink HAJ; Boehm MJ. 1999. Biocontrol within the context of soil microbial communities: A substrate-dependent phenomenon. *Annual Review of Phytopathology.* 37:427-446.
- Hummel, R.L., Walgenbach, J.F., Barbercheck, M.E., Kennedy, G.G., Hoyt, G.D., Arellano, C., 2002. Effects of production practices on soil-borne entomopathogens in Western North Carolina vegetable systems. *Environ. Entomol.* 31, 84–91. <https://doi.org/10.1603/0046-225X-31.1.84>.

- Ishibashi, N., Kondo, E., 1986. *Steinernema feltiae* (DD-136) and *S. glaseri*: Persistence in soil and bark compost and their influence on native nematodes. *J. Invertebr. Pathol.*18: 310-316
- Jaffuel, G., Mader, P., Blanco-Pérez, R., Chiriboga, X., Fliessbach, A., Turlings, T.C.J., Campos-Herrera, R., 2016. Prevalence and activity of entomopathogenic nematodes and their antagonists in soils that are subject to different agricultural practices. *Agric. Ecosyst. Environ.* 230: 329-340. <https://doi.org/10.1016/j.agee.2016.06.009>
- Kaya H.K. 1990. Soil Ecology. In: Gaugler R, and Kaya HK. (eds). *Entomopathogenic Nematodes in Biological Control*. CRC Press, Boca Raton, FL, pp. 93-111.
- Kergunteuil A, Bakhtiari M, Formenti L, Xiao Z, Defosse E et al. 2016. Biological control beneath the feet: a review of crop protection against insect root herbivores. *Insects*:7:e70.
- Khumalo NN., Lephoto TE., Gray VM. 2021. The effect of organic compost and soil texture on the survival and infectivity of entomopathogenic nematode species. *Archives of Phytopathology and Plant Protection*.54(17-18):1443-1455. <https://doi.org/10.1080/03235408.2021.1914369>.
- Koppenhofer AM., Shapiro-Ilan D., Hiltbold I. 2020. Entomopathogenic nematodes in sustainable food production. *Front. Sustain. Food Syst.* Sec. Crop Biology and Sustainability.4, doi.org/10.3389/fsufs.2020.00125
- Kurzemann FR., Plieger U., Probst M., Spiegel H., Sanden T., Ros M., Insam H. 2020. Long-term fertilization affects soil microbiota, improves yield and benefits soil. *Agronomy*.10,1664.
- Lawrence JL., Hoy CW., Grewal PS. 2006. Spatial and temporal distribution of endemic entomopathogenic nematodes in a heterogeneous vegetable production landscape. *Biological Control*.37(3):247-255. doi.org/10.1016/j.biocontrol.2006.02.002
- Lewis, E.E., Hazir, S., Hodson, A., Gulcu, B., 2015. Trophic relationships of entomopathogenic nematodes in agricultural habitats. In: Campos-Herrera, R. (Ed.), *Nematode Pathogenesis of Insects and Other Pests: Ecology and Applied Technologies for Sustainable Plant and Crop Protection*. Springer International Publishing, AG Switzerland, pp. 139–163. https://doi.org/10.1007/978-3-319-18266-7_5
- Oerke EC. 2005. Crop losses to pests. *J Agr Sci*.144: 31-43. [doi.org.10.1017/S0021859605005708](https://doi.org/10.1017/S0021859605005708).

- Patzel N., Sticher H., Karlen DL. 2000. Soil fertility-phenomenon and concept. *J. Plant Nutr. Soil Sci.* 163, 129-142.
- Peters A. 1996. The natural host range of *Steinernema* and *Heterorhabditis* spp. and their impact on insect populations. *Biocontrol Science and technology*, 6 (3): 389-402.
- Puissant J., Villenave C., Chauvin C., Plassard C., Blanchart E., Trap J. 2021. Quantification of the global impact of agricultural practices on soil nematodes: A meta-analysis. *Soil Biology and Biochemistry*. 161, 108383. doi.org/ 10.1016/j.soilbio.2021.108383
- Renkema JM., Lynch DH., Cutler GC., MacKenzie K., Walde SJ. 2016. Organic mulches in highbush blueberries alter beetle (Coleoptera) community composition and improve functional group abundance and diversity. *Agric. For. Entomol.* 18:119-127.
- Renkema JM., Parent JP. 2021. Mulches used in highbush blueberry and entomopathogenic nematodes affect mortality rates of third-instar *Popillia japonica*. *Insects*. 12, 907. https://doi.org/10.3390/insects12100907
- Shapiro D.I., L.C. Lewis, J.J. Obrycki., M. Abbas 1999. Effects of fertilizers on suppression of black cutworm (*Agrostis ipsilon*) damage with *Steinernema carpocapsae*. Supplement to the *Journal of Nematology*, 31 (4S): 690-693.
- Shapiro-Ilan D., Dolinski C. 2015. Entomopathogenic nematode application technology. In: Campos-Herrera R. ed. *Nematode pathogenesis of insects and other pests-ecology and applied technologies for sustainable plant and crop protection*. Springer International Publishing. pp. 231-534.
- Shields EJ., Testa AM. 2020. Multi-year biological control of black vine weevil, *Otiorynchus sulcatus*, with persistent entomopathogenic nematodes. *The Great Lakes Entomologist*. 53(2):119-125. doi.org/10.22543/0090-0222. 2377
- Sikora RA., Bridge J., Starr JL., 2005. Management practices: An overview of integrated nematode management technologies. In: *Plant Parasitic nematodes in subtropical and tropical Agriculture*, 2nd ed: Luc M., Sikora RA., Bridge J. Eds; CAB International: Wallingford, UK, pp. 793-825.
- Susurluk I.A., N.A. Kumral, U. Bilgili & E. Acikgoz 2011. Control of a new turf pest, *Dorcadion pseudopreissi* (Coleoptera: Cerambycidae), by the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Journal of Pest Science*, 84: 321-326.

Susurluk I.A. 2008. Effects of various agricultural practices on persistence of the inundative applied entomopathogenic nematodes, *Heterorhabditisbacteriophora* and *Steinernema feltiae* in the field. Russian Journal of Nematology, 16 (1): 23-32

Ulug D., Hazir S., Kaya HK., Lewis EE.2014.Natural enemies of natural enemies: The potential top-down impact of predators on entomopathogenic nematode populations.Ecological Entomology. 39(4). doi.10. 1111 /een.12121

UNDER PEER REVIEW