

## **Review Article**

# **Interplay of light intensity and nutrients on ginger responses related to growth, yield and quality of ginger in various cropping systems: a retrospective analysis**

### **Abstract**

Ginger is an important crop belonging to the Zingiberaceae family. It is grown for its rhizomes, which are vitally important in the food and spice sectors as well as in the pharmaceutical and perfumery industries as a therapeutic herb. Its cultivation has been greatly expanded in several nations, particularly India, as it is a lucrative crop. The majority of ginger is produced and exported from India. Since ginger can withstand partial shade, it is a good choice for intercropping in several cropping systems. The effect of light intensity, as well as how they interact with nutrients to affect crop performance in terms of growth and development, production and quality is crucial for achieving maximum yield. However, the relationship between the growing environment and nutrient managements which affect the elements that govern its growth, development and quality is not well understood. It is crucial to comprehend the proper amount of nutrients to apply in an intercropping system in order to minimise wastage since the intercrop may potentially interact with the main crop for nutrients and other inputs. Therefore, it is worthwhile to undertake the current literature study to learn how ginger responds to the stimulus of nutrients and varying light intensities in diverse cropping systems. From this, it can be inferred that ginger grown under partial shade in various cropping systems, can be a very profitable spice crop with superior yield and quality rhizomes.

**Keywords:** *Ginger, cropping system, light intensity, nutrients, interaction, growth*

**Comment [A1]:** And specially for agroforestry systems

## **1. INTRODUCTION**

Ginger (*Zingiber officinale* Roscoe) is an important spice crop grown throughout the world. It is well known for the medicinal values imparted by unique phytochemical compounds. There are more than 400 different chemical compounds present in ginger. Major constituents of rhizome include carbohydrates (50–70%), lipids (3–8%), terpenes (zingiberene,  $\beta$ -bisabolene,  $\alpha$ -farnesene,  $\beta$ -sesquiphellandrene, and  $\alpha$ -curcumene), and phenolic compounds (gingerol, paradols, and shogaol) [1]. In India, rhizomes of ginger are used for cooking with vegetables and for adding aroma and flavour to food. In some of the countries like Thailand, leaves of ginger are also used as salad. It is also used as a carminative relieving flatulence [2]. It is a traditional medication that has been used to treat a wide range of ailments, including sickness and confinement in women, as well as headaches, motion sickness, nausea, dyspepsia, and a reduction in colic and gas.

Increasing attention in ginger as spice, food ingredient, medicine and industrial component led to increased cultivation in many countries. Most of the ginger (78.3 %) is grown in Asian countries like India, China, Japan, Thailand, Indonesia and Taiwan; about 19.8 percent is cultivated in some African countries like Nigeria, Mauritius, Sierra Leone, Ghana; 1.7 per cent in some parts of North and South America and only 0.3 % in Oceania. India is the largest producer and exporter of the ginger; it produces 1,844,000 tonnes of ginger from an area of 1,72,000 hectare [3]. Most of the ginger is produced in Assam, Maharashtra, West Bengal, Gujarat, Kerala and North-Eastern states.

Ginger can tolerate partial shade conditions and therefore it is well suited as intercrop in agricultural, horticultural and agroforestry systems, although it is also grown as a stand-alone crop. In southern states of India, it is grown under plantation crops such as coconut and arecanut. Growing ginger under the shade of fruit crops like guava, aonla, citrus etc. is a common practice in sub-Himalayan region. In sub-tropical areas, it is frequently interplanted in forest trees such as between rows of *Populus*, *Ailanthus*, *Luceana*, etc. [4]. The purpose of planting vegetable crops in these production systems is to utilize every inch of land as the available land for agriculture is decreasing at an alarming rate. In silviculture system a lot of land remain unutilized and a large proportion of land goes wasted. Moreover, there is less utilization of inputs like fertilizers and irrigation water. It is, however, possible to grow more crops on the same piece of land in a given amount of time by using the land wisely in terms of time and space.

Intercrop selection is a crucial idea in agricultural, horticultural, and agroforestry systems. The intercrop should be selected in such a way that it should be complementary with the main crop, there should not be any competition for light, water and nutrients between the crops and most importantly the cropping system should be profitable [5]. Ginger is one of the best crops which fits into this system because of its tolerance to partial shade and high economic value of ginger rhizomes which are in continuous demand in global trade. It is important to note that light intensity plays a crucial role in the photosynthetic process as well as plant growth and development. Also, the intercrop may potentially interact with the main crop for nutrients and other inputs, it is important to understand the appropriate amount of nutrients to apply in an intercropping system to prevent waste. There are conflicting reports on the increase and decline of ginger yields when grown in the open as well as in shades with varying light intensity [6, 7, 8, 9, 10, 11, 12, 13]. Therefore, to ascertain how ginger responds to the stimulus

Comment [A2]: Add Space

Comment [A3]: However, it is

Comment [A4]: planning

Comment [A5]: wastage of fertilizers

Comment [A6]: are several

Comment [A7]: performances

of nutrients, and various intensities of light in different cropping systems it is worthwhile to conduct the current literature study.

## 2. GINGER UNDER DIFFERENT CROPPING SYSTEMS

The foremost objective of agriculturists and farmers is to enhance crop productivity, which can be done by increasing production per unit area with the expansion of cultivated land. However, the arable land is decreasing, so there is a need to analyse the role of multiple cropping as a method of crop production to enhance the agricultural production [14]. The success of multiple cropping system lies in the fact that the intercrop uses space and labour more efficiently [15]. Moreover, well-consumption of climatic factors, higher and more stable yields under varying environmental conditions, soil conservation, and the extent of inputs and outputs and their contribution to the stabilisation of household food supply [16] are some of the other factors contributing to its increased trend. An intercrop combination is considered best when it provides greater yield per unit area and is more competent to use resources than would otherwise be the case if each crop was grown as a monoculture [17]. Ginger, being a long-duration crop and having partial tolerance to shade, is well suited for multiple cropping in agriculture, horticulture and agroforestry systems. Intercrop of ginger in larger spacing of poplar (5 × 4m) resulted in increased plant height, a greater number of tillers and leaves per plant. Additionally, the intercropped ginger had greater length and width of rhizome and yield per plant than monoculture [18]. Shade plays an important role in productivity of ginger, while growing as an intercrop in an agroforestry system. It was observed that ginger grows well with mango at 50±5% shade and yielded 2.45-fold better than the least yield i.e., 5.07 t/ha in ginger-guava cultivation with 70±5% shade [19]. In another study based on intercropping of ginger in different fruit trees, the yield of ginger was highest in coconut + guava-based system with 70-80 % PAR [20]. Intercropping of high fertilizer requirement crops can lead to reduced yields of both crops as compared to sole cropping. The reduction in the number of ears of sweet maize by 17.35 %, ear weight by 8.07 % and decreased number of ginger branches (5.12 to 6.47 %) and rhizome weight (5.64 to 16.43 %) was also reported [21]. However, higher comprehensive economic benefits were reported in ginger-sweet maize intercropping than sole cropping. The qualities of the soil and its microbial microecological environment are altered over time by the continuous cultivation of one crop on the same piece of land, which limits the development of the medicinal plant business by lowering the chemical components of medicinal plants. The intercropping of ginger with *Pogostemoncablin* improved the abundance and diversity of soil microbes and community structure by enhancing the number of dominant bacteria and activities of soil enzymes, which ultimately led to an increase in the active ingredient in *Pogostemoncablin* [22]. Similarly, Table 1 shows how intercropped ginger performed with different crops and its effect on various aspects of crop production.

**Table 1: Features of intercropping ginger in different crops**

S.No.	Intercrop/ Mixed crop	Effect of intercropping	Location	Reference
1	Sapota- Jatropha	Higher fresh rhizome yield and increased morphological traits like number of fingers, plant height, length	Gujarat, India	[23]

Comment [A8]: add space

Comment [A9]: needed/requiring

Comment [A10]: However,

Comment [A11]: SL.

Comment [A12]: References

		and width of rhizome.		
2	Maize at varying row spacing	Higher ginger yield, gross return and margin and benefit cost ratio when maize was planted on 3 <sup>rd</sup> , 4 <sup>th</sup> and 5 <sup>th</sup> ginger row with 100 % ginger	Chapainawab ganj, Bangladesh	[24]
3	<i>Paulownia elongata</i>	Ginger gave high yields as compared to maize and beans	Buckingham, China	[25]
4	Two bamboo species: <i>Dendrocalamus asper</i> <i>Dendrocalamus miltonii</i> )	Increase in yield by 10.5–15.6% as compared to open field Maximum returns (3,595.67 US\$ ha <sup>-1</sup> yr <sup>-1</sup> ) when ginger grown under the canopy of <i>D. asper</i>	Himachal Pradesh, India	[26]
5	Spinach, cauliflower, bitter gourd	Sole crop of ginger lead to highest yield The benefit cost ratio of Ginger + spinach was best	Jharkhand, India	[27]
6	Poplar	Increase in yield, plant height, number of tillers/plants, rhizome length and width at 5 X 4 m spacing of poplar.	Himachal Pradesh, India	[18]
7	Sweet maize	Decreased number of branches and rhizome weight per plant in intercropping and Increased income in intercropping	Jinan, China	[21]
8	Cowpea, soybean, mungbean, lablab	Enhanced growth, rhizome yield and other yield traits in intercropping with legume crops; highest yield was attained when ginger was intercropped with mung bean; yield was maximum when one row of ginger was planted with two rows of legume crops	Abuja, Nigeria	[28]
9	Ginger + Maize-Soybean	Increased total land output (2% higher than ginger sole cropping) and increased potential economic return by 6%.	Pokhara, Nepal	[29]
10	<i>Pogostemon cablin</i>	Improved soil microbial abundance, diversity and community structure Modification of soil physical and chemical properties	Guangzhou, China	[22]
11	Maize, French bean, Soybean, pumpkin, groundnut	Ginger + maize + French bean + pumpkin system recorded 15.20%, higher than sole ginger crop yield The intercropping of ginger + groundnut recorded highest benefit cost ratio (3.42)	Meghalaya, India	[30]
12	Maize	At 24074 maize plants/ha, the production of intercropped ginger (250000 plants/ha) was noticeably comparable to the yield of ginger grown as a solitary crop. At this plant population, the production of maize	Zaria, Nigeria	[31]

Comment [A13]:

Comment [A14]: structure,

		grains was likewise maximum.		
13	Teak	Intercropping of pruned teak and application of fertilizer in ginger increased the net weight of ginger	Karangduwet, Indonesia	[32]
14	Pigeon pea and Maize	Increased yield and reduces rhizome rot incidence	Assam, India	[33]

### 3. EFFECT OF SHADE AND LIGHT INTENSITIES ON GROWTH AND YIELD OF GINGER

Light intensity is a crucial element in plant growth and development. It is defined as the amount of visible light emitted in one unit of time per unit of solid angle. It is also the main component of photosynthesis and food production by plants. Accordingly, increasing light intensity will increase the photosynthetic capacity and also have a positive impact on plant growth [34, 35]. Since ginger prefers shade, the amount of light that it receives in a cropping system will affect its growth characteristics and yield. Numerous research studies demonstrated an increase in ginger yield and contributing attributes with various levels of photosynthetic active radiation (PAR) under natural and artificial systems of shade [6, 7, 8, 9, 10]. In contrast to these studies, it was found that solitary cropping of ginger at 29200 lux was considerably more productive than intercropping with cashew plantations at 25774 lux in terms of fresh weight of rhizome, quantity of secondary rhizomes, and clump size [13]. According to Wilson and Ovid [36], ginger intercropped with okra (*Abelmoschus esculentus* L.), corn, and peas had more plant height, number of tillers, and rhizome yield than corn alone and in the absence of shade. Studying the impact of shade levels on ginger growth and yield, it was reported that plant height and tiller count were higher in 50% shade levels, while fresh rhizome weight, dry weight, and total yield were higher in 30% shade levels [37]. Ginger can also be successfully cultivated under the partial shade of multipurpose trees in a forest. Another study [38] reported higher rhizome weight and yield under the partial shade of guava trees (180 cm × 180 cm) than planting with the dense shade of gamar trees (90 cm × 90 cm). Moreover, it is also observed that ginger grown in shade remained greener and physiologically active for longer periods of time [4]. Pruning is an important practice in fruit crops to maintain a balance between reproductive and vegetative growth. Pruning of branches will lead to decreased interception of PAR by the fruit crops, allowing more light to reach the intercrop. Ginger, while growing as an intercrop with oranges, responded well to pruning, and authors reported an increase of 70 and 50 percent in growth and yield, respectively with pruning compared to unpruned orange trees [39]. Similarly, in an agroforestry system, pruning of teak trees resulted in increase in rhizome yield of *Zingiber officinale* var. *amarum* [32]. Comparing the efficiency of light authors reported that green light was less efficient wavelength and it can only lead to increase in plant height as compared to white light treatment [40]. Later, it was found that supplementing green light under strong sunlight had a positive impact on the growth characteristics of ginger, leading to a higher number of shoots, increased crown density, and greater crown volume [41]. Intercropping ginger with pigeon pea and maize led to attenuation of total PAR ( $1688.1 \mu \text{mol s}^{-1} \text{m}^{-2}$ ) by 79.5 and 69.2%, respectively, leading to decreased rhizome rot incidence and increased yield due to unfavourable soil and air temperatures and soil moisture for the spread of disease [33].

**Comment [A15]:** It would be more interesting if the data were presented in tabular form like the previous table.

#### 4. EFFECT OF LIGHT INTENSITY ON BIOACTIVE COMPOUNDS AND QUALITY OF GINGER

Besides being used as a spice and flavouring compound, ginger is widely grown as a medicinal herb throughout the world, especially in Asia. The species *Z. officinale* is known for its wide range of bioactive compounds and phytochemical antioxidants, which have many health-promoting benefits [42]. Similar to other herbs that are used as flavouring agents in food, ginger is also a good source of phenolic and flavonoid compounds[43]. Quality control becomes an important issue while growing ginger for medicinal purposes, the altered concentrations of the bioactive compounds can be harmful to consumers. The concentration of phenolics and bioactive compounds in plants is highly influenced by environmental conditions such as light intensity, temperature, and other biotic and abiotic factors [43, 44]. Furthermore, the concentration of phenolics and flavonoids may vary among different plant parts while grown in different light intensities. The flavonoid content was higher in the leaves in 60 percent shade, while phenolics were favouring the rhizomes in zero percent shade. Caffeic acid was only detected in ginger grown under zero per cent shade, while tannic acid only accumulated in the leaves of ginger grown under 60% shade level [45].The synthesis of phenolic compounds and flavonoids in ginger, viz., quercetin, rutin, catechin, epicatechin, and naringenin, was high in plants grown at low light intensities. The antioxidant activities in leaves also increased with increasing flavonoid concentration at low light intensities ( $310 \mu\text{mol m}^{-2}\text{s}^{-1}$ ). The ferric reducing activity of the rhizomes was higher than that of the leaves in  $310 \mu\text{mol m}^{-2}\text{s}^{-1}$  of sun light[43]. The growth and development of ginger are also affected by the function of the green light.The chlorophyll metabolism gene was upregulated, and the amount of chlorophyll-a was increased when green light was added with intense sunshine. The expression of photosynthetic genes and the rate of photosynthesis were both efficiently boosted by the enhancement of green light radiation [41].

In India, ginger is either eaten raw or dried, and in the importing nations, it is used to make oleoresin and essential oils. Dry ginger is necessary to command a higher price in the market, and ginger's essential oils, oleoresin, starch, protein, and dietary fibre are crucial qualities [46]. These traits are highly influenced by light intensity. According to a study, the dry ginger yield, non-volatile ether extract, starch content and crude fibre content increased at 20 and 40 percent shade as compared to 60 and 80 percent shade and volatile oil content increased with the increase in shade level[47].

#### 5. INTERACTION OF NUTRIENTS AND SHADE AFFECTS GINGER'S GROWTH

The availability of nutrients is an essential factor in the growth and development of ginger, thus choosing the right fertilizer type and rate of application to get the best yield is important. In *Zingiberaceae*, which includes plants like ginger (*Z. officinale*) and turmeric (*Curcuma longa*) increasing fertilizer rates greatly increases the rhizome production [47, 48]. Ginger is grouped as an exhaustive crop which responds greatly to manure application [49].Ginger prefers shade, and the relationship between shade and fertiliser type and its rate of application impacts the crop's development and output.The interaction between shade levels and NPK fertilizers in *Zingiber zerumbet* (L.), reported an increase in yield with the increase in NPK fertilizer doses and shade levels [37]. The best plant growth and maximum rhizome yield was produced at 30 % shade level and N: P:

**Comment [A16]:** It would be more interesting if authors would quantify an approximate fertilizer dose suitable for intercropping.

Kat 120:140:230 kg/ha/year. Ginger cultivated in shaded conditions can result in increased N and K uptake, necessitating a 150% increase in fertiliser dose[50]. Therefore, it was recommended that a fertiliser dose of 150 kg N: 100 kg P<sub>2</sub>O<sub>5</sub>: 100 kg K<sub>2</sub>O ha<sup>-1</sup> be used to increase ginger yield when grown under coconut trees [47]. Crop growth and productivity in an agro-forestry system are regulated by the interaction of crop canopy, soil nutrient content, soil temperature and moisture, etc. It was revealed that dense canopy and increased leaf litter of bamboo species (*D. asper*), and application of FYM improved the soil's physical, chemical, and moisture qualities, which in turn led to better development of ginger and an increase in rhizome production[26]. Similar to this, when teak and ginger were intercropped, pruning of teak increased light transmission, which together with fertilizer application resulted in heavier clumps of ginger as opposed to fertilized and non-fertilized treatments under open conditions [7, 32]. The grain legume crops like cowpea, soybean, mung bean and lablab as a sole crop or as an intercrop improve the nutrient status of the soil. Ginger intercropped with legume crops lead to enhanced rhizome growth and yield variables as compared to sole crops. However, growth variables of ginger can be reduced due to intercrop competition and reduced PAR due to excess shading brought about by the crop canopy. A ginger-to-legume planting ratio higher or lower than 1:2 may lead to under or overutilization of growth resources such as soil nutrients, light, and moisture by the component crops in the mixture[28]. Another study discovered a substantial interaction effect between light intensity and vermicompost on ginger's fresh rhizome yield. According to reports, vermicompost alone produced yields comparable to those of recommended dose under normal lighting conditions [51]. The fact that ginger accumulates more dry matter and absorbs more nutrients in the shade can be highlighted here. This suggests that if ginger is grown in an intercropped scenario with low light intensities, the fertilizer dose may need to be increased to get the best yield.

## 6. MECHANISMS INVOLVED IN GINGER'S REACTION TO LIGHT AND NUTRIENTS

Light and nutrients are the most crucial components for successful plant growth when environmental considerations in agriculture are taken into account. Photosynthesis, a complicated process for producing biomass that involves the synthesis of pigments, the Calvin cycle, and electron transport, requires functional light [52]. Along with influencing plant growth and yield potential, light quality and intensity also have an impact on the architecture and mechanism of the photosynthetic apparatus[53, 54]). A certain amount of light is required by plants for growth; light levels that are too high or too low may affect photosynthesis adversely. Light also affects the environmental conditions like soil and air temperature, humidity, carbon dioxide concentrations around the plant [55]. In addition to light, balanced nutrients must also be provided to meet the needs of plant for growth and development, as deficiencies in nitrogen, phosphorus, potassium and other essential nutrients can hinder plant growth and development, photosynthesis, and reduce leaf area[56, 57, 58]). Finding a balance between lighting and fertilization based on regional circumstances is crucial for greater quality and higher yields. For instance, different nitrogen rates, as well as red-blue LED light intensities, had an impact on lettuce development. Higher light intensities and lower nitrogen contents enhanced photosynthesis and lettuce output[59]. The following is a description of some potential reasons that could explain how nutrients, light, and their interactions affect ginger's growth and quality.

### 6.1) Probable causes of the varying effects of light on ginger growth and yield

Comment [A17]: .

Ginger, being a shade-loving plant, can carry out its physiological processes appropriately at reduced light intensities[60]; however, the effectiveness of relative light intensity may vary with the geographical region, season, day length, intensity of light, humidity, temperature and cultivar. Ginger as an intercrop intercepted less PAR with a lower reflection coefficient as compared to a sole crop, indicating efficient use of PAR (Figure 1). Here, the higher rhizome yield and increased value for other morphological parameters of the plant may be attributed to the low light intensity and shade-loving nature of the crop [6, 9, 18, 61] reported that ginger grows more rapidly when light transmission is reduced and air temperatures and relative humidity (RH) are raised. As a matter of fact, relative humidity drops as the temperature rises if no additional moisture is supplied to the air[62, 63, 64]). Therefore, the inadequate combination of light, temperature, and humidity may be the cause of the lower growth parameters in open field conditions. Additionally, high light intensity may cause damage to shade-loving plants by reducing the leaf area due to high radiation level causing damage to photosynthetic pigment [37]. On the other side, more photosynthates accumulated under shadowed conditions due to higher leaf area values, which in turn enhanced bulking rate, net assimilation rate, and crop growth rate, increasing rhizome output. In addition, increased weight of rhizomes results from early bulking and progressive accumulation of photosynthates even during the latter phases of plant growth [65]. Similarly, increasing the shade level was reported to produce a higher biomass in turmeric [66, 67]. Growing ginger as intercrop with crops like maize and pigeon pea provided congenial microclimate which led to optimum growth and high yield of ginger and reduced incidence of rhizome rot disease [33]. Contrastingly, there are reports of increased rhizome yield in open conditions as compared to shading. Authors have suggested high light interception, less competition for nutrient and moisture in sole crop [11, 12, 13] and competition for light, source and sink relationship in plant and mobilisation of assimilates from the source to primary and secondary rhizomes for decreased yield in partial shade [68, 69]. The allelopathic nature of *Tectonagrandis*, competition for essential nutrients and inadequate light intensity for the poor tiller production by ginger grown in the teak was also reported [7].


## 6.2) Reasons for differential effect of light intensity on bioactive compounds and ginger quality

During growth and development, plants have to interact with the biotic and abiotic factors of their ecological environment. Among abiotic factors, viz., water, light, temperature, soil, etc., light is the most crucial of all; extremes of it cause stress, which leads to variation in the synthesis of secondary metabolites[70]. The phenolic and flavonoid content of the same plant may vary depending on the ecological type it is domesticated into. It has been reported in ginger and many other crops that low light levels are ideal for producing the most flavonoids, but high light levels were ideal for phenolic acid production [45, 71, 72, 73]. This is because UV and blue light at zero per cent shade level inhibits the activity of the enzyme chalcone synthase, which is crucial for the flavonoid pathway [74, 75].

**Figure 1. Flowchart showing factors responsible for improved performance of ginger under shade**

Comment [A18]: .

Comment [A19]: It would be better added this to paragraph 4.



While the increased phenolic compounds in high light intensity is due to increased activity of Phenylalanine ammonialyase (PAL) enzyme under high light intensity and UV (Fig.1)[76]. Additionally, different shade levels had a substantial impact on the synthesis, accumulation, and partitioning of flavonoids and phenolic acids in different parts of ginger and different genotypes had a variable concentration of flavonoids and phenolic acids. The strong positive relationships between total flavonoids contents and antioxidant activity was also reported in other studies[43, 77]. Therefore, antioxidant activities were also higher at low light intensities due to increased free radical scavenging power, which rose along with the increase in total flavonoids, suggesting that flavonoids had a greater impact on free radical scavenging than phenolics[43]. The antioxidant activities are in general higher in leaves than the rhizomes in most of the studies [42, 78, 79]). The most abundant flavonoid in ginger is quercetin and apigenin. Recent studies on the potential carcinogenicity of quercetin have also looked at its potential as a cancer preventative. The synthesis of quercetin content in leaves and rhizomes increased in low light intensity, suggesting that medical power of ginger can be improved by growing it in low light intensity.

## 7. CONCLUSION

Ginger is able to effectively utilise low light intensities, suggesting that the growth performance of ginger can be improved under partially shaded conditions. However, it is well known that too much shade can lead to reduced yields. Here, it is crucial to emphasise that the efficiency of relative light intensity may vary depending on the geographical region, season, length of the day, light intensity, humidity, temperature, and cultivar. Therefore, with the proper management of light intensity and nutrition, ginger can be accommodated as an intercrop in agricultural, horticultural, and agroforestry systems. The tall crops of these systems can modify the microclimate by providing shade during sprouting and rhizome development stages. Additionally, by including ginger as an intercrop in various systems, farmers' income per unit area can be increased.

Comment [A20]: were

Comment [A21]: Conclude the suitable and acceptable percentage of shade for ginger production.

Comment [A22]: Please conclude an approximate fertilizer dose for ginger in intercropping/mixed cropping systems.

Comment [A23]: Nevertheless

## REFERENCES

1. Grzanna R, Lindmark L, Frondoza CG (2005) Ginger—an herbal medicinal product with broad anti-inflammatory actions. *J Med Food*, 8: 125-132. [10.1089/jmf.2005.8.125](https://doi.org/10.1089/jmf.2005.8.125)
2. Kikuzaki H, Nakatani N (1993) Antioxidant effects of some ginger constituents. *J Food Sci*, 58: 1407-1410. <https://doi.org/10.1111/j.1365-2621.1993.tb06194.x>
3. FAO, 2020. Food and Agriculture Organization of the United Nations. [www.faostat.fao.org](http://www.faostat.fao.org).
4. Pandey V, Shukla S, Saxena J, Khan S, Kumar R (2018) Ginger (*Zingiberofficinale*Rosc.) productivity in tree based agro forestry system. *JOJ Horticulture & Arboriculture*, 1: 555568. [10.19080/JOJHA.2018.01.555568](https://doi.org/10.19080/JOJHA.2018.01.555568)
5. Maitra S, Hossain A, Brestic M, Skalicky M, Ondrisik P, Gitari H, Brahmachari K, Shankar T, Bhadra P, Palai, JB, Jena J (2021) Intercropping—A low input agricultural strategy for food and environmental security. *Agronomy*, 11: 343. <https://doi.org/10.3390/agronomy11020343>
6. Kumar RD, Sreenivasulu GB, Prashanth SJ, Jayaprakashnarayan RP, Nataraj SK, Hegde NK (2010) Performance of ginger in tamarind plantation (as intercrop) compared to sole cropping (ginger). *Int J Agric Sci*, 6; 193-195.
7. Oladele AT, Popoola L, Jimoh SO (2012) Effects of teak canopy cover and NPK fertilizer application on growth of ginger in agroforestry trial, Ile-Ife, Nigeria. *JASR*, 12: 147-157.
8. Babu MS, Kumar BP, Swami DV, Krishna KU, Emmanuel N (2019) Impact of shade net condition on growth, rhizome and yield characters of ginger. *J. Pharmacogn. Phytochem.* 8: 3481-3485.
9. Aly MM, El Sawy A, El Gendy RA (2019) Comparative study of different shading types on growth and yield of ginger plants. *Middle East J*, 8:1264-1270. <https://doi.org/10.36632/mejar/2019.8.4.28>
10. Hossain S, Rahman MS, Kona KN, Bari MS, Akter N, Ali MM (2020) Growth Performance of Two Ginger (*Zingiberofficinale* Roscoe) Varieties under Different Agroforestry Systems in Bangladesh. *Asian Plant Res J*, 3: 1-10. [10.9734/aprj/2019/v3i3-430074](https://doi.org/10.9734/aprj/2019/v3i3-430074)
11. Hegde NK, Sulikeri GS, Rethnum BP (2000) Distribution of Photosynthetically Active Radiation (PAR) and performance of ginger under arecanut shade. In *Spices and aromatic plants: challenges and opportunities in the new century. Contributory papers. Centennial conference on spices and aromatic plants, Calicut, Kerala, India, 20-23 September, 2000: 107-112*. Indian Society for Spices.
12. Kumar RD (2005) Intercrop studies in tamarind (*Tamarindusindica* L.) plantation. Dissertation, University of Agricultural Sciences, Dharwad, Karnataka, India.
13. Vikram HC, Hegde NK (2014) Performance of ginger in cashew plantation (as intercrop) compared to sole cropping. *Asian J Hort*, 9: 187-189.
14. Midmore DJ (1993) Agronomic modification of resource use and intercrop productivity. *Field Crops Res*, 34: 357-380. [https://doi.org/10.1016/0378-4290\(93\)90122-4](https://doi.org/10.1016/0378-4290(93)90122-4)
15. Baldy C, Stigter CJ (1997) Agrometeorology of multiple cropping in warm climates. INRA, Paris.
16. Beets WC (1982) Multiple cropping and tropical farming system, grower. *London, Britain, and West views press, Colorado, USA. 156p*.
17. Inal A, Gunes A, Zhang F, Cakmak (2007) Peanut/maize intercropping induced changes in rhizosphere and nutrient concentrations in shoots. *Plant PhysiolBiochem*, 45: 350-356. [10.1016/j.plaphy.2007.03.016](https://doi.org/10.1016/j.plaphy.2007.03.016)

18. Jaswal SC, Mishra VK, Verma KS (1993) Intercropping ginger and turmeric with poplar (*Populusdeltoides* 'G-3'Marsh.). *Agroforsyst*, 22: 111-117. <https://doi.org/10.1007/BF00705140>
19. Amin MR, Ikbal TMT, Miah MMU, Hakim MA, Amanullah ASM (2010) Performance of ginger under agroforestry system. *Bangladesh Res. Pub. J*, 4: 208-217.
20. Bhuiyan MM, Roy S, Sharma PC, Rashid MH, Bala P (2012) Impact of multistoreyedagro-forestry systems on growth and yield of turmeric and ginger at Mymensingh, Bangladesh. *Crop Prod*, 1:19-23. <http://www.escijournals.net/EJCP>
21. Qingsheng REN, Chengyong LI (2021) Effect of Intercropping Ginger and Sweet Maize on Growth, Yield and Benefits. *Asian Agric Res*, 13: 45-53. [10.22004/ag.econ.316971](https://doi.org/10.22004/ag.econ.316971)
22. Zeng J, Liu J, Lu C, Ou X, Luo K, Li C, He M, Zhang H, Yan H (2020) Intercropping with turmeric or ginger reduce the continuous cropping obstacles that affect *Pogostemoncablin* (patchouli). *Frontiers Microbiol*, 11: 579719. <https://doi.org/10.3389/fmicb.2020.579719>
23. Pandey SBS, Pandey M, Jadeja DB, Tandel MB, Nayak D (2017) Growth and yield of ginger (*Zingiberofficinale* L.) under Sapota-Jatropha based agroforestry systems in south Gujarat. *J. Pharmacogn. Phytochem*, 6: 247-251.
24. Islam MK, Nayeem IKA, Uddin MZ, Rahman MM, Hossain MM (2021) Productivity of intercropping maize with ginger at varying row spacing. *J Biosci Agric Res*, 28: 2374-2380. <https://doi.org/10.18801/jbar.280221.288>
25. Newman SM, Bennett K, Wu Y (1997) Performance of maize, beans and ginger as intercrops in Paulownia plantations in China. *AgroforSysts*, 39: 23-30. <https://doi.org/10.1023/A:1005938310106>
26. Garima, Bhardwaj DR, Thakur CL, Kaushal R, Sharma P, Kumar D, Kumari Y (2021) Bamboo-based agroforestry system effects on soil fertility: Ginger performance in the bamboo subcanopy in the Himalayas (India). *Agron J*, 113: 2832-2845. <https://doi.org/10.1002/agj2.20684>
27. Kumar A, Singh RK, Bharti N and Krishna G (2018) Performance and profitability study of different ginger-based cropping system in Bokaro district of Jharkhand. *Int J CurrMicrobiol App Sci*, 7: 2934-2939.
28. Nwaogu EN, Muogbo PC (2015) Effect of ginger-grain legume cropping system and spatial arrangement on soil fertility management and yield of intercropped ginger in the guinea savanna of Nigeria. *Int Res J Agric Sci Soil Sci*, 5: 1-7. <http://dx.doi.org/10.14303/irjas.2014.076>
29. Chapagain T, Pudasaini R, Ghimire B, Gurung K, Choi K, Rai L, Magar S, Bishnu BK, Raizada MN (2018) Intercropping of maize, millet, mustard, wheat and ginger increased land productivity and potential economic returns for smallholder terrace farmers in Nepal. *Field Crops Res*, 227: 91-101. <https://doi.org/10.1016/j.fcr.2018.07.016>
30. Rymbai H, Das A, Mohapatra KP, Talang HD, Nongbri B, Law I (2021) Ginger (*Zingiberofficinale*) based intercropping systems for enhancing productivity and income—a farmers' participatory approach. *Indian J Agric Sci*, 91: 956-60. <https://doi.org/10.56093/ijas.v91i7.115024>
31. Lyocks SWJ, Tanimu J, Dauji LZ (2013) Growth and yield parameters of ginger as influenced by varying populations of maize intercrop. *J Agric Crop Res*, 1: 24-29.

32. Sudomo A, Sebastian GE, Perdana A, Prameswari D, Roshetko JM (2019) Intercropping of *Zingiberofficinale* Var. *Amarumon* teak silviculture in Karangduwet, Paliyan, GunungKidul Yogyakarta. In *IOP Conference Series: Earth Environ Sci*, 250: 012104. 10.1088/1755-1315/250/1/012104
33. Neog P, Goswami RK, Saikia D, Gogoi N, Kalita MK (2022) Response of modified microclimates on growth, yield, and incidence of rhizome rot disease of ginger in Assam, India. *International J Biometeorol*, 66: 719-730. <https://doi.org/10.1007/s00484-021-02231-9>
34. Abrams MD, Mostoller SA (1995) Gas exchange, leaf structure and nitrogen in contrasting successional tree species growing in open and understory sites during a drought. *Tree physiol.* 15: 361-370. <https://doi.org/10.1093/treephys/15.6.361>
35. Makino A, Sato T, Nakano H, Mae T (1997) Leaf photosynthesis, plant growth and nitrogen allocation in rice under different irradiances. *Planta*, 203: 390-398. <https://doi.org/10.1007/s004250050205>
36. Wilson H, Ovid A (1993) Growth and yield responses of ginger (*Zingiberofficinale* Roscoe) as affected by shade and fertilizer applications. *J Plant Nutr*, 16: 1539-1545. <https://doi.org/10.1080/01904169309364631>
37. Goh S, Lee Abdullah T, Hassan SA, Stanslas J (2018) Breaking dormancy and effects of shade level and NPK fertilizer rates on yield of *Zingiber zerumbet* (L.) Smith (Lempoyang). *Agriculture*, 8: 198. <https://doi.org/10.3390/agriculture8120198>
38. Aslam Ali M, Jamaluddin M, Mujibur Rahman GM (2005) Ginger cultivation under multipurpose tree species in the hill forest. *Korean J Soil Sci Fert*, 38: 218-221. <https://www.researchgate.net/publication/264137143>
39. Tabin T, Balasubramanian D, Arunachalam A (2015) Influence of Canopy Pruning on Orange Growth and Rhizome Yield of Intercrop Ginger under Agri-Horticulture System. *Indian J Hill Farming*, 28: 1. <https://ebook.icar.gov.in/index.php/IJHF/article/view/51196>
40. Zhang R, Xu K, Dong C, Li Y, Lv J (2008) Effects of light quality on growth and light utilization characteristics in ginger. *Acta Hortic Sin* 35: 673–680. <https://doi.org/10.3724/SP.J.1005.2008.01083>
41. Cao B, Lv X, Chen Z, Xu K (2021) Supplementing green light under strong sunlight improves growth and functional ingredients of ginger (*Zingiberofficinale*Rosc.) in summer. *Ind Crops Prod*, 167: 113527. <https://doi.org/10.1016/j.indcrop.2021.113527>
42. Chan EWC, Lim YY, Wong LF, Lianto FS, Wong SK, Lim KK, Joe CE, Lim TY (2008) Antioxidant and tyrosinase inhibition properties of leaves and rhizomes of ginger species. *Food Chem*, 109: 477-483. <https://doi.org/10.1016/j.foodchem.2008.02.016>
43. Ghasemzadeh A, Jaafar HZ, Rahmat A, Wahab PEM, Halim MRA (2010) Effect of different light intensities on total phenolics and flavonoids synthesis and anti-oxidant activities in young ginger varieties (*Zingiberofficinale* Roscoe). *Int J Mol Sci*, 11: 3885-3897. <https://doi.org/10.3390/ijms11103885>
44. Southwell IA, Bourke CA (2001) Seasonal variation in hypericin content of *Hypericum perforatum* L. (St. John's wort). *Phytochem*, 56: 437-441. [https://doi.org/10.1016/S0031-9422\(00\)00411-8](https://doi.org/10.1016/S0031-9422(00)00411-8)

45. Ghasemzadeh A, Ghasemzadeh, N (2011) Effects of shading on synthesis and accumulation of polyphenolic compounds in ginger (*Zingiberofficinale* Roscoe) varieties. J Med Plant Res, 5: 2435-2442. <http://www.academicjournals.org/JMPR>
46. Govindarajan VS, Connell DW (1983) Ginger—chemistry, technology, and quality evaluation: part 1. Crit Rev Food Sci Nutr, 17: 1-96. 10.1080/10408398209527343
47. Ajithkumar K, Jayachandran BK (2001) Effect of major nutrients on yield of ginger (*Zingiberofficinale* Rose.) intercropped in coconut garden. J Spices Aromat Crops, 10: 17-23.
48. Akhter S, Noor S, Islam MS, Masud MM, Talukder MR, Hossain MM (2013) Effect of potassium fertilization on the yield and quality of ginger (*Zingiberofficinale*) grown on a K deficient terrace soil of level Barind Tract (AEZ 25) in Northern Bangladesh. Research Findings, e-*ifc*, (35).
49. Susharna P K, Jose A I (1994) Nutrition of ginger. In: Chadha K L & Rethinam P (Eds.) Advances in Horticulture Vol. 9, Plantation and Spices Crops Part I, (pp 490-497). Malhotra Publishing House, New Delhi
50. Ancy J, Jayachandran BK (1996) Nutrient requirement of ginger (*Zingiberofficinale*R.) under shade. Indian Cocoa, Arecanut and Spices Jpomal. 20: 15- 116.
51. Vastrad NV, Hegde RV, Giritammanavar VA (2006) Effect of Light intensity and Vermicompost on Yield of Ginger. Karnataka J Agric Sci, 19: 941-945
52. Wang L, Liang W, Xing J, Tan F, Chen Y, Huang L, Cheng CL, Chen W (2013) Dynamics of chloroplast proteome in salt-stressed mangrove *Kandeliacandel* (L.) Druce. J Proteome res, 12: 5124-5136. 10.1021/pr4006469
53. He B, Chen Y, Zhang H, Xia C, Zhang Q, Li W (2018) The effect of colored plastic films on the photosynthetic characteristics and content of active ingredients of *Dysosmaversipellis*. Hortic Environ Biote, 59: 519-528. 10.1007/s13580-018-0061-8
54. Yang F, Fan Y, Wu X, Cheng Y, Liu Q, Feng L, Chen J, Wang Z, Wang X, Yong T, Liu W (2018) Auxin-to-gibberellin ratio as a signal for light intensity and quality in regulating soybean growth and matter partitioning. Front Plant Sci, 9: 56. <https://doi.org/10.3389/fpls.2018.00056>
55. Song R, Kelman D, Johns KL, Wright AD (2012) Correlation between leaf age, shade levels, and characteristic beneficial natural constituents of tea (*Camellia sinensis*) grown in Hawaii. Food Chem, 133: 707-714. <https://doi.org/10.1016/j.foodchem.2012.01.078>
56. Mu X, Chen Y (2021) The physiological response of photosynthesis to nitrogen deficiency. Plant PhysiolBiochem, 158: 76-82. <https://doi.org/10.1016/j.plaphy.2020.11.019>
57. Meng X, Chen WW, Wang YY, Huang ZR, Ye X, Chen LS, Yang LT (2021) Effects of phosphorus deficiency on the absorption of mineral nutrients, photosynthetic system performance and antioxidant metabolism in *Citrus grandis*. PLoS One, 16: e0246944. <https://doi.org/10.1371/journal.pone.0246944>
58. Tabaglio V, Boselli R, Fiorini A, Ganimede C, Beccari P, Santelli S, Nervo G (2020) Reducing nitrate accumulation and fertilizer use in lettuce with modified intermittent Nutrient Film Technique (NFT) system. Agronomy, 10: 1208. <https://doi.org/10.3390/agronomy10081208>
59. Fu Y, Li H, Yu J, Liu H, Cao Z, Manukovsky NS, Liu H (2017) Interaction effects of light intensity and nitrogen concentration on growth, photosynthetic characteristics and quality of lettuce (*Lactuca sativa* L. Var. *youmaica*). Sci hortic, 214: 51-57. 10.1016/j.scienta.2016.11.020

60. Minoru T, Hori Y (1969) Studies on photosynthesis of vegetable crops I. photosynthesis of growing plant vegetables in relation to light intensity. Bull In: Phil, 139-140.
61. Balasimha D (1989) Light penetration patterns through arecanut canopy and leaf physiological characteristics of intercrops. J. Plantation Crops, 16: 61-67.
62. Hemm MR, Rider SD, Ogas J, Murry DJ, Chapple C (2004) Light induces phenylpropanoid metabolism in Arabidopsis roots. Plant J, 38: 765-778. <https://doi.org/10.1111/j.1365-313X.2004.02089.x>
63. Briskin, D.P. and Gawienowski, M.C., 2001. Differential effects of light and nitrogen on production of hypericins and leaf glands in *Hypericum perforatum*. Plant PhysiolBiochem, 39: 1075-1081. [https://doi.org/10.1016/S0981-9428\(01\)01326-2](https://doi.org/10.1016/S0981-9428(01)01326-2)
64. Kurata H, Matsumura S, Furusaki S (1997) Light irradiation causes physiological and metabolic changes for purine alkaloid production by a *Coffea arabica* cell suspension culture. Plant science, 123: 197-203. [https://doi.org/10.1016/S0168-9452\(96\)04588-8](https://doi.org/10.1016/S0168-9452(96)04588-8)
65. Sangeetha KS, Subramanian S, Marthandan V (2016) Physiological Parameters and Yield Characters of Ginger Genotypes (*Zingiberofficinale*Rosc.) under Coconut Ecosystem. IJBSM, 7: 939-942. [10.23910/IJBSM/2016.7.4.1630](https://doi.org/10.23910/IJBSM/2016.7.4.1630)
66. Srikrishnah S, Sutharsan, S (2015) Effect of different shade levels on growth and tuber yield of turmeric (*Curcuma longa* L.) in the batticaloa district of Sri Lanka. Am Eurasian J Agric Environ Sci, 15: 813-816. [10.5829/idosi.aejaes.2015.15.5.12642](https://doi.org/10.5829/idosi.aejaes.2015.15.5.12642)
67. Amzad Hossain M, Akamine H, Ishimine Y, Teruya R, Aniya Y, Yamawaki K (2009) Effects of relative light intensity on the growth, yield and curcumin content of turmeric (*Curcuma longa* L.) in Okinawa, Japan. Plant Prod Sci, 12: 29-36. <https://doi.org/10.1626/pp.s.12.29>
68. Singh VB, Kar PL (1991) Effect of planting materials on the productivity of turmeric variety Lakadong. Indian Cocoa, Arecanut and Spice Journal, 14:153-154
69. Vastrad NV, Sulikeri GS, Hegde RV (2002) Effects of light intensity and vermicompost on the yield of ginger (*Zingiberofficinale*Rosc). J ApplHort, 4: 25-26. [10.37855/jah.2002.v04i01.07](https://doi.org/10.37855/jah.2002.v04i01.07)
70. Verma N, Shukla S (2015) Impact of various factors responsible for fluctuation in plant secondary metabolites. J Med Aromat, 2: 105-113. <https://doi.org/10.1016/j.jarmap.2015.09.002>
71. Fonseca JM, Rushing JW, Rajapakse NC, Thomas RL, Riley MB (2006) Potential implications of medicinal plant production in controlled environments: the case of feverfew (*Tanacetum parthenium*). HortScience, 41: 531-535. <https://doi.org/10.21273/HORTSCI.41.3.531>
72. Mosaleeyanon K, Zobayed SMA, Afreen F, Kozai T (2005) Relationships between net photosynthetic rate and secondary metabolite contents in St. John's wort. Plant Sci, 169: 523-531. <https://doi.org/10.1016/j.plantsci.2005.05.002>
73. Bergquist SÅ, Gertsson UE, Nordmark LY, Olsson, ME (2007) Effects of shade nettings, sowing time and storage on baby spinach flavonoids. J Sci Food Agric J, 87: 2464-2471. <https://doi.org/10.1002/jsfa.2956>
74. Logemann E, Tavernaro A, Schulz W, Somssich IE, Hahlbrock K (2000) UV light selectively conduces supply pathways from primary metabolism and flavonoid secondary product formation in parsley. PNAS, 97: 1903-1907. <https://doi.org/10.1073/pnas.97.4.1903>

75. Loyall L, Uchida K, Braun S, Furuya M, Frohnmeyer H (2000). Glutathione and a UV light-induced glutathione S-transferase are involved in signaling to chalcone synthase in cell cultures. *Plant Cell*, 12: 1939-1950. [10.1105/tpc.12.10.1939](https://doi.org/10.1105/tpc.12.10.1939)
76. Kumari R, Singh S, Agrawal B (2009) Effects of supplemental ultraviolet-B radiation on growth and physiology of *Acorus calamus* L. (sweet flag). *Acta BiolCracoviensia, Ser. Bot*, 51: 19-27.
77. Oktay M, Gülçin İ, Küfrevioğlu Öİ (2003) Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *LWT-Food Sci Technol*, 36: 263-271. [https://doi.org/10.1016/S0023-6438\(02\)00226-8](https://doi.org/10.1016/S0023-6438(02)00226-8)
78. Agnani H, Menut C, Bessi re, JM (2004) Aromatic plants of tropical central Africa. Part XLIX: Chemical composition of essential oils of the leaf and rhizome of *Aframomum giganteum* K. Schum from Gabon. *Flavour Fragr J*, 19: 205-209. <https://doi.org/10.1002/ffj.1403>
79. Elzaawely AA, Xuan TD, Tawata S (2007) Essential oils, kava pyrones and phenolic compounds from leaves and rhizomes of *Alpinia zerumbet* (Pers.) BL Burtt. & RM Sm. and their antioxidant activity. *Food Chem*, 103: 486-494. <https://doi.org/10.1016/j.foodchem.2006.08.025>

UNDER PEER REVIEW