

Review Article

Egg Parasitoids of Hemipteran Insects: A special account on Scelionidae (Platygastroidea: Hymenoptera)

Abstract: Parasitoids are considered as one of the key biotic factors regulating the population dynamics of other insects in the natural settings. Egg parasitoids have been used successfully for many decades as inundative and augmentative biological control agents against a wide range of economically important agricultural pests. In this review parasitoids in the family Scelionidae, especially those attacking the eggs of hemipteran insects were discussed. Information on six genera of scelionids namely *Trissolcus* Ashmead, *Telenomus* Haliday, *Gryon* Haliday, *Idris* Förster, *Protelenomus* Kieffer, *Hadronotus* Förster and *Psix* Kozlov as the egg parasitoids of hemipteran insects has been gathered. Among these six genera, several species belonging to *Trissolcus* Ashmead and *Telenomus* Haliday have been extensively studied and exploited for the biological control of hemipteran pests. The main target hosts of these egg parasitoids found to be the heteropteran pests in the families Pentatomidae, Scutelleridae, Urostylididae and Coreidae. Release of these parasitoids have provided successful control of several pests in soybean, cowpea, rice and squash especially for the control of invasive stink bugs. Further, it was found that, the effective utilization of these parasitoids has the greatest potential to manage several agriculturally important pests in the ever-changing agricultural ecosystems where chemical free pest management is the main priority. A small effort is made to pool the information from the existing literature on the parasitoid species involved and the host insects associated with them to understand their diversity and biocontrol potential.

Introduction

Maintaining the health of crops and protecting them from insect pests is a major concern for the farmers [1]. Insect pests belonging to diverse taxa with different feeding habits cause various kinds of injury to crop plants. Among these, the sucking insects belonging to order Hemiptera causes serious injury to plants directly by sucking the sap, and indirectly by transmitting plant diseases leading to severe yield losses. For instance, Pea aphid, *Acyrtosiphon pisum* (Aphididae: Hemiptera) poses a serious threat to commercial pulse production as it can cause direct damage to crops by extracting sap from leaves, stems, and

pods. Additionally, it serves as a vector for transmitting over 30 plant viruses, including cucumber mosaic virus, beet yellows virus, pea enation mosaic virus, and bean leafroll virus [2]. In Okra, leafhoppers and whiteflies causes 54-66% loss of yield [3]. After pod borers, the pigeon pea pod sucking bug, *Clavigralla gibbosa* (Coreidae: Hemiptera), has emerged as a significant menace to pigeonpea quality grain production [4]. Typically, the damage inflicted on grain yield by this bug range from 25 to 40 % [5]. The occurrence of the mirid bug, *Nesidiocoris cruentatus* (Hemiptera: Miridae) has recently been recorded on tender leaves and young fruits across various regions of eastern Uttar Pradesh on bottle gourd [6]. In recent years, there has been a notable surge in the global movement of invasive exotic species from one region to another. Among economically significant arthropods, whiteflies and mealybugs play a substantial role in this invasion. In India, there are 469 species of whiteflies belonging to 71 genera known to feed on a wide range of agricultural, horticultural, and forestry crop plants, including 8 invasive species [7]. Despite various control measures available, farmers primarily rely on systemic chemical pesticides to control these insect pests. However, the unrealistic and indiscriminate use of synthetic pesticides over the years has led to outbreaks of both insect, and non-insect pests and sever environmental problems. Farmers, both in developed and underdeveloped countries, solely depend on synthetic insecticides, collectively spending an estimated 9 billion US Dollars annually [8].

Stink bugs (Hemiptera: Pentatomidae) are serious pests of a wide variety of agricultural crops and are often found on a succession of host plants throughout the season. Several hymenopteran egg parasitoids are known to attack Pentatomidae, including Scelionidae (*Gryon* Haliday, *Telenomus* Haliday, *Trissolcus* Ashmead), Eupelmid (*Anastatus* Motschulsky) and Encyrtidae (*Ooencyrtus* Ashmead). The eupelmids and encyrtids associated with pentatomids are generalists and include members that are facultative hyperparasitoids [9]. In contrast, scelionids in the genus *Trissolcus* and several species in the genus *Telenomus* are strictly associated with Pentatomidae, cause high levels of parasitism, and are used as biological control agents for native and invasive pentatomid species [10]. Attributes like high searching ability, high reproductive rates and lack of hyperparasitoids place them as highly potential and promising biocontrol agents [10].

The superfamily Platygastroidea is the third largest group after Ichneumonoidea and Chalcidoidea in the Parasitic Hymenoptera [11]. The Scelionidae (exclusively attacking the eggs of insects) and Platygastriidae (parasitizing both insect eggs and immature stages of

Cecidomyiidae and sternorrhynchous Hemiptera) of Platygastroidea are rich in species and are of immense importance as biocontrol agents. The subfamily Telenominae (Scelionidae) has high potential to control several agricultural pests belonging to Lepidoptera and Hemiptera. *Trissolcus basalis* (Wollaston) which attacks the green stink bug *Nezaravirudula* (L.) eggs are being used widely in biological control programmes for its management [12]. Scelionid parasitoids of hemipteran eggs are the subject of active study, driven by the economic damage caused by a variety of bug pests, and by recent works that accelerate further advancement [13]. Biological control of other pentatomid pests has focussed on the use of egg parasitoids and the study of their biology [14]. In the Indian subcontinent, egg parasitoids from several genera in the family Scelionidae have been reported, among them *Telenomus* Haliday, *Trissolcus* Ashmead and *Gryon* Haliday are important [15]. In this review, we tried to pool some vital information regarding the Scelionid egg parasitoids of agriculturally important hemipteran pests.

Diversity of Hemipteran egg parasitoids in the superfamily Platygastroidea

The superfamily Platygastroidea (Hymenoptera: Proctotrupomorpha) is a diverse taxon of wasps that are parasitoids of nine orders of insects as well as spiders [16]. Superfamily Platygastroidea comprises seven extant families: Geoscelionidae, Janzenellidae, Neuroscelionidae, Nixoniiidae, Platygastriidae, Scelionidae and Sparasionidae [16]. Among these families, except Platygastriidae and Scelionidae, the information on host range, the species composition of remaining five families are scanty. From the available information it is understood that these five families attack mainly the eggs of Orthoptera. Austin *et al.* (2005) [11] suggested that the most likely ancestral host for Platygastroidea are the eggs of Orthoptera, however, individual groups of platygastroids have undergone subsequent changes to attack different arthropod groups. The Scelionidae and Platygastriidae are the two major families of Platygastroidea, known for their diverse species composition and better understanding on their host range of which many are hemipteran insects. For instance, 26 species of Scelionids belonging to 4 genera (*Trissolcus*, *Telenomus*, *Psixand* *Gryon*) recorded as the egg parasitoids of *Nezaraviridula* [17]. The Scelionidae comprises more than 4000 species in 176 genera [18] with the greatest diversity of hosts in the superfamily and most biological data in general. The host taxa include: Odonata, Orthoptera, Mantodea, Embiidina, Hemiptera, Coleoptera, Neuropteran, Lepidoptera and Diptera) as well as spiders [19]. The subfamily Telenominae, particularly the genera *Trissolcus* Ashmead and *Telenomus* Haliday,

have been extensively studied because of their role as biological control agents of their hosts especially hemipterans. The genera *Gryon* Haliday and *Telenomus* have been reported to use several groups of hosts (e.g., *Telenomus* species are parasitoids of the eggs of Hemiptera, Lepidoptera, Diptera, and Neuroptera), but according to Taekulet *al.* (2014) [20], the ground-plan hosts of both *Gryon* and *Telenomus* are likely to be Hemiptera. The Platygasteridae comprises 69 genera [21]. This is strikingly the most species rich family characterized by their minute size and reduction in the wing venation, number of antennomeres, number of metasomal segments and overall body sculpture. As per the available data, the most species of Platygasteridae are larval or egg-larval parasitoids of Cecidomyiidae (gall midges). The subfamily Sceliotrachelinae is more biologically diverse and has been reared from the eggs of Coleoptera and auchenorrhynchous Hemiptera, the nymphs of sternorrhynchous Hemiptera.

There are six genera of scelionids recorded as the egg parasitoids of hemipteran insects. A detailed list of the documented species and the host insects they associated with is presented in the Table 1. The genus *Trissolcus* Ashmead is a cosmopolitan genus with at least 180 described species of egg-parasitoid wasps associated with stink bugs (Pentatomidae, Scutelleridae, Urostylididae), many of which are important insect pests [22]. During the past decade, the Asian fauna of *Trissolcus* has received increased attention, driven largely by the search for biological control agents to manage two invasive stink bugs of global significance: *Halyomorpha halys* (Stål) and *Bagrada hilaris* (Burmeister) (Pentatomidae). The genus *Telenomus* Haliday is by far the largest genus in the subfamily Telenominae and includes more than 600 species reported worldwide. Species of *Telenomus* share the hosts of *Trissolcus* (Pentatomidae, Scutelleridae, Urostylididae), but also attack a wider range of Heteroptera, as well as Auchenorrhyncha, Lepidoptera, Diptera and Neuroptera. *Protelenomus* Kieffer, another genus in this subfamily, is known to parasitize coreid bugs [23][24]. *Protelenomusanoplocnemidis*, *P. flavicornis* and *P. areolatus* Rajmohana are the only previously described species in this genus [25], however Venakumari *et al.* (2019) [24] described six new species and added and revised the genus *Protelenomus*. The *Hadronotus* Förster another genus in the family Scelionidae and the species of *Hadronotus* are known to parasitize the eggs of hemipteran pests, including pentatomids such as *Piezodorushybnieri*, *Dolycoris baccarum*, *Nezara antennata* and *Halyomorpha halys* [26].

Table 1. List of Scelionid parasitoids and the hemipteran insect hosts they associated.

Parasitoid species	Hosts	Reference
--------------------	-------	-----------

		s
<i>Trissolcus edessae</i> (Fouts)	<i>Halyomorphahalys</i> , <i>Euschistus</i> <i>servus</i> ,	[27]
<i>Trissolcus euschisti</i>	<i>Chinaviahilare</i>	
	<i>Euschistus</i> <i>conspersus</i>	[28]
	<i>Edessa</i> <i>meditabunda</i>	[29]
	<i>Chinaviahilare</i> , <i>Brochymena</i> <i>quadripustulata</i>	[30]
<i>Trissolcus kozlovi</i>	<i>Halyomorphahalys</i> ,	[31]
<i>Trissolcus semistriatus</i> (Nees von Esenbeck)	<i>Eurydema</i> sp. (Scutelleridae)	[22]
<i>Trissolcus podisi</i>	<i>Podisus</i> <i>maculiventris</i> Say	[32]
	<i>Halyomorphahalys</i> (Stal) (Pentatomidae) , <i>Euschistus</i> <i>servus</i> (Say), <i>Chinaviahilare</i> (Say)	[27]
<i>Trissolcus cultratus</i> (Mayr)	<i>Hippotiscus</i> <i>dorsalis</i> Stål (Pentatomidae) , <i>Urochelaluteovaria</i> Distant(Urostylididae)	[22]
<i>Trissolcus elasmuchae</i> (Watanabe)	<i>Nipheelongata</i> (Dallas) (Pentatomidae)	
<i>Trissolcus latusulcus</i> (Crawford)	<i>Poecilocoris</i> <i>latus</i> Dallas (Scutelleridae)	
<i>Trissolcus yamagishii</i> Ryu	<i>Nipheelongata</i> (Dallas) (Pentatomidae)	
<i>Trissolcus japonicus</i> (Ashmead)	<i>Erthesina</i> <i>fullo</i> (Thunberg) and <i>Rhaphigaster</i> <i>nebulosa</i> (Poda) (Pentatomidae)	
	<i>Halyomorphahalys</i>	[33, 34]
	<i>Halyomorphahalys</i>	[31]
<i>Trissolcus grandis</i>	<i>Eurygaster</i> <i>integriceps</i> ,	[35]
	<i>Eurygaster</i> spp. <i>Eurygaster</i> <i>integriceps</i>	[36]
<i>Trissolcus basalis</i> (Wollaston)	<i>Nezaraviridula</i>	[17, 37, 38, 39]
	<i>Halyomorphahalys</i>	[31]
<i>Trissolcus simoni</i>	<i>Eurydemaventrals</i>	[40]
<i>Trissolcus simoni</i>	<i>Eurydemaventrals</i>	[41]
<i>Trissolcus brochymenae</i>	<i>Murgantia</i> <i>histrionica</i>	
<i>Trissolcus utahensis</i>	<i>Euschistus</i> <i>conspersus</i>	[28]
<i>Trissolcus elimatus</i>	<i>Edessa</i> <i>meditabunda</i> (F.)(Pentatomidae)	[29]
<i>Trissolcus urichi</i> (Crawford)	<i>Euschistus</i> <i>heros</i> and <i>Chinavia</i> <i>ubica</i>	[42]
	<i>Edessa</i> <i>meditabunda</i>	[29]
	<i>Oebalus</i> <i>poecilus</i> ,	[43]
	<i>Tibracalimbativentris</i>	
	<i>Piezodorus</i> <i>guildinii</i> ,	[44]
	<i>Nezaraviridula</i> ,	
	<i>Dichelops</i> sp.,	
	<i>Euschistus</i> <i>heros</i> ,	
	<i>Piezodorus</i> <i>guildinii</i>	
<i>Trissolcus plautiae</i> (Watanabe)	<i>Plautia</i> <i>stali</i> Scott (Pentatomidae)	[45]
<i>Trissolcus mitsukurii</i> (Ashmead)	<i>Nezaraviridula</i> , <i>Scotinopharalurida</i> , <i>Ensarcoris</i> sp.	[46]
	<i>Nezara</i> <i>antennata</i>	[46][47]
	<i>Gonopsis</i> <i>affinis</i> , <i>Lagnotmus</i>	[47]

	<i>elongatus</i> , <i>Dolycoris baccarum</i> L., <i>Piezodorus hybneri</i> ,	
	<i>Halyomorpha halys</i>	[31][48]
	Pentatomidae	[22]
<i>Trissolcus hyalinipennis</i>	<i>Bagradahilaris</i> (Pentatomidae)	[49]
<i>Telenomus nakagawai</i> Watanabe	<i>Nezaraviridula</i> , <i>Scotinopharalurida</i>	[46]
	<i>Nezara antennata</i>	[47]
<i>Telenomus triptus</i>	<i>Piezodorus hybneri</i>	[50]
	<i>Scotinopharacoarctata</i>	[51]
<i>Telenomus truncatus</i> (Nees von Esenbeck)	Pentatomidae	[52]
<i>Telenomus gifuensis</i> Ashmead	<i>Scotinopharalurida</i> , <i>D. baccarum</i> ,	[47]
	<i>Eusarcoris parvus</i> , <i>Eusarcoris guttiger</i>	[53]
	<i>Nezaraviridula</i>	[46]
	<i>Nezara antennata</i> , <i>Anacanthocoris concoloratus</i> , <i>Riptortus clavatus</i> , <i>Scotinopharalurida</i> (Pentatomidae)	[53]
<i>Telenomus perplexus</i> Girault	<i>Brochymena</i> sp	[54]
<i>Telenomus podisi</i> Ashmead	<i>Oebalus poecilus</i> ,	[43]
	<i>Euschistus heros</i> ,	
	<i>Tibracalimbativentris</i>	
	<i>Murgantia histrionica</i>	[55]
	<i>Euschistus heros</i>	[43][56][57]
	<i>Euschistus conspersus</i>	[28]
	<i>Euschistus servus</i>	[27]
	<i>Halyomorpha halys</i>	[27]
	<i>Piezodorus guildinii</i> ,	[59]
	<i>Euschistus</i> spp., <i>Nezaraviridula</i>	
	<i>Dichelops</i> sp., <i>Piezodorus guildinii</i> ,	[44]
	<i>Nezaraviridula</i> ,	
	<i>Euschistus heros</i>	
	<i>Dichelops melacanthus</i> , <i>Euschistus heros</i> ,	[60]
	<i>Podisus nigrispinus</i>	
<i>Telenomus turesis</i>	<i>Palomena prasina</i> L. (Hemiptera: Pentatomidae)	[61]
	<i>Halyomorpha halys</i>	[31]
<i>Telenomus ashmeadi</i>	<i>Pentatomaligata</i> , <i>P. sayi</i>	[62]
	<i>Euschistus fissilis</i> ,	[54]
	<i>E. tristigmus</i>	
	<i>Piezodorus guildinii</i> ,	[59]
	<i>Euschistus</i> spp., <i>Nezaraviridula</i> ,	
	<i>Podisus maculiventris</i> (Say), <i>Chinavia hilaris</i>	
<i>Telenomus calvus</i>	<i>Euschistus conspersus</i>	[28]
	<i>Podisus neglectus</i>	[63]
	<i>Podisus maculiventris</i>	[64, 65, 66]

<i>Telenomus cuspis</i> Rajmohana and Srikumar	<i>Helopeltis antonii</i> (Miridae)	[67]
<i>Gryon homoeoceri</i> Nix.	<i>Amblypeltanianihotis</i> Blote, <i>Dasynuspiperis</i> China	[68, 69]
<i>Gryon sp</i> (Scelionidae)	<i>Amblypeltalutescenspapuensis</i>	[70]
	<i>Edessa mediatubunda</i> (F.)(Pentatomidae)	[29]
	<i>Nilaparvatalugens</i>	[71]
<i>Gryon aetherium</i> Talamas	<i>Bagrada hilaris</i> (Burmeister) (Pentatomidae)	[72, 73]
<i>Gryon ancinla</i> Kozlov and Lê	<i>Acanthocoris scaber</i> (L.) (Coreidae)	[74]
<i>Gryon clavigrallae</i> Mineo	<i>Clavigralla spp.</i> (Coreidae)	[75]
<i>Gryon gonikopalense</i>	<i>Bagradahilaris</i>	[49]
<i>Idris elba</i> Talamas	<i>Bagradahilaris</i>	[76]
<i>Protelenomus</i> sp	<i>Anoplocnemisphasiana</i> (Coreidae)	[77]
<i>Protelenomusflavicornis</i>	<i>Anoplocnemisphasiana</i> (Coreidae)	[23] [24]
<i>Protelenomusgajadanta</i>	<i>Pseudothertapsdevastans</i> (Coreidae)	[24]
<i>Protelenomusanoplocnemidis</i>	<i>Anoplocnemiscurvipes</i> (Coreidae)	
<i>Hadronotus pubescens</i> (Motschoulsky)	<i>Riptortus pedestris</i> (Fab.) (Alydidae)	[78]
<i>Hadronotuspennsylvanicus</i> (Ashmead)	<i>Anasa tristis</i> and <i>Leptoglossus</i> sp (Coreidae)	[79]
<i>Psixabnormis</i> Kozlov and Lê, <i>Psixsaccharicola</i> (Mani), <i>Psixsunithae</i> , <i>Psixlacunatus</i> , <i>Psixconfluus</i> Johnson and Masner, <i>Psix robustus</i> Rajmohana, <i>Psixstriaticeps</i> (Dodd), <i>Psixvariosus</i> Johnson and Masner	Several pentatomids and coreids	[80]

Gryon Haliday, 1833 is another genus within the family Scelionidae. It is the largest genera in Scelioninae with 332 species known in the world [81]. This genus is notable for its role as an egg parasitoid, primarily targeting the eggs of various hemipteran pests, mainly Pentatomidae, Reduviidae, and Coreidae [82]. Recent studies have identified several new species within this genus, such as *Gryon ancinla* and *Gryon aetherium*, which have been reported in regions like Vietnam, China, and Pakistan, respectively [22, 73]. Similarly, another genus *Idris* Förster, 1856 is also include several egg parasitoids, however majority of them are parasitic on spiders. Few species like *Idriselba*Talamas is parasitic on eggs of *Bagradahilaris*[76]. *Hadronotus* is a genus of wasps in the Scelionidae family that includes about 217 species. Some species of *Hadronotus* include. *Hadronotuspennsylvanicus* and *Hadronotuspubescens* which are egg parasitoids of bugs belongs to the heteropteran families Coreidae and Alydidae[7] [83]. The parasitoids in the genus *Psix*Kozlov, 1976 have been documented to parasitize the eggs of various hemipteran pests, contributing to the control of populations that can cause significant agricultural damage [80].

Host parasitisation potential and use of Scelionid parasitoids in biological control of hemipteran pests: few important case reports

Both Scelionidae and Platygasteridae contain candidates for classical biological control programs, however Scelionidae contain most egg parasitoids of hemipteran pests, and many of them have been introduced for control of invasive pests. A clear example is *Trissolcus basalis*, which has been recorded in six zoogeographical regions. *Trissolcus basalis* is the representative example which is widely introduced into many countries for control of *Nezaraviridula* (L., 1758) (Pentatomidae: Hemiptera) throughout the world [37]. The parasitoid–host association of *Trissolcus basalis* and *N. viridula* has become a favored model system in ecological, behavioral, and physiological research on insects [11]. Some other species of Scelionidae including, *Trissolcus mitsukurii*, *Trissolcus colemani* and *Telenomus chloropus*, have also been widely transported to control *N. viridula*. Attempts were also made to use species of Scelionidae in the framework of classical biological control.

The Neotropical Brown Stink Bug, *Euschistus heros* (Pentatomidae), the most prevalent soybean pest, presents serious management issues due to its high frequency and abundance of occurrence [84]. Egg parasitoids are widely used in augmentative biological control and can be regarded the most important stink bug biocontrol agents [85]. Among the numerous species of egg parasitoids that can be used in augmentative biological control of *E. heros*, *Telenomus podisi* is notable for its high parasitism and control efficacy against its hosts [60]. Bueno *et al.* (2020) [56] reported that release of *T. podisi* in the field increased *E. heros* egg parasitism to 70% and 50%, in 2017/2018 and 2018/2019, respectively. Thus, they suggested that *T. podisi* can be efficiently used to control stink bug eggs in the form of encapsulated pupa or unprotected, with similar efficacy. Garipey *et al.* (2019) [30] reported five species of scelionids parasitizing eleven pentatomid species. 70% of all the eggs masses examined were parasitized by parasitoids. *Telenomus podisi* Ashmead being the dominant parasitoid and was detected in all host species. *Trissolcus euschisti* Ashmead was detected in many host eggs, but was significantly more prevalent in *Chinavia hilaris* (Say) and *Brochymena quadripustulata* (Fabricius). *Trissolcus brochymenae* Ashmead and *Trissolcus thyantae* Ashmead were recorded sporadically. Parasitism of *H. halys* was 55%, and this species was significantly less likely to be parasitized than native pentatomids. The scelionid species composition of *H. halys* consisted of *Te. podisi*, *Tr. euschisti* and *Tr. thyantae*.

Although these species cannot develop in fresh *H. halys* eggs, we demonstrate that parasitoids attempt to exploit this host under field conditions.

Across the United States, the squash bug, *Anasa tristis* (De Geer) (Coreidae: Hemiptera), is a serious pest of cucurbit crops. Boyle *et al.* (2023) [79] reported that before the release of egg parasitoid, *Hadronotuspennsylvanicus* (Ashmead) (Scelionidae: Hymenoptera), there was very little parasitisation which was about <21% in 2020 and <8% in 2021. However, in both the years, the percentage of *A. tristis* eggs parasitized within 2 weeks of release of *H. pennsylvanicus* was significantly greater at release sites (~60%) than at no-release sites (~14%). High rates of *H. pennsylvanicus* parasitism (>72%) were further observed at release sites 4, 6, 8, and 10 weeks following parasitoid deployment. Other Scelionidae species have also been reported to have successfully released egg parasitoids. For example, *Gryon muscaeformis* released in hazel boosted parasitism rates on *Gonocerusacuteangulatus* (Heteroptera: Coreidae) eggs [86]. Similar results were seen when *Telenomus gifuensis* was released early in the season. This resulted in higher parasitism rates on *Scotinopharalurida* (Pentatomidae) eggs than in untreated areas [53]. *Gryon aetherium* has been evaluated for its effectiveness against the invasive stink bug *Bagradahilaris*, a significant pest of crops. Research indicates that *Gryon* parasitoids can significantly reduce pest populations by targeting their eggs, thus preventing the emergence of adult pests [73]. In Brazil, a total of 23 species of egg parasitoids targeting stink bugs in soybean crops have been identified. Among these, *Trissolcus basalis* and *Telenomus podisi* (Ashmead) exhibit higher rates of parasitism compared to other species and are frequently utilized for the control of stink bugs [87].

Clavigralla spp. (Coreidae) are the serious sucking pests of pulse crops. They lay eggs in clusters. The field collections of eggs in clusters at International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) by Shanower *et al.* (1996) [75] reported 69-100% eggs parasitized by that *Gryon clavigrallae* Mineo (Scelionidae). Overall, more than 39 % of *Clavigralla* spp. eggs were parasitized by *G. clavigrallae*. The percentage of egg clusters parasitized and the percentage of eggs parasitized in a cluster were positively correlated with the size of the egg cluster. The percentage of eggs and egg clusters parasitized by *G. clavigrallae* increased through the season. A survey carried out annually from 2010 to 2013 at the Directorate of Cashew Research in Puttur, Karnataka, India, identified three species of egg parasitoids: *Telenomus* sp. (Scelionidae), *Chaetostricha* sp.

(Trichogrammatidae), and *Erthymelushelopeltidis*(Mymaridae) associated with *Helopeltis* spp. *Telenomus* sp. was the most prevalent species, observed throughout the year except in March and April, with peak parasitism occurring during the monsoon months of June and July, ranging from 6.89% to 28.21% [67]. Thus, the long-term benefits of a more environmentally friendly pest control technique, such as the employment of *T. podisi*, may outweigh the risk of a slight decrease in productivity. For example, the organic market typically offers higher pricing, which may compensate for minor yield losses. Biological management is required for organic crops because synthetic pesticides are not permitted. Furthermore, organic farming has showed environmental benefits at the farm level, increasing the use of biological control methods.

Conclusion

This review emphasizes the significant role of egg parasitoids, particularly those from the Scelionidae family, in controlling population of hemipteran pests. Several genera of these parasitoids are specialized in targeting the eggs of various hemipteran species, especially the stick bugs in the families Pentatomidae and Coreidae, which are major pests in the agricultural ecosystems. The ecological significance of these parasitoids as natural pest control agents is of immense importance, further, using them as biological control agents has provided viable option for safer pest management.

Future studies

Understanding the complex interactions between egg parasitoids and their hemipteran hosts will be crucial. Future studies should investigate the dynamics of host-parasitoid relationships, including factors influencing parasitism rates, host specificity, and the impact of environmental variables on these interactions. Given that the environmental conditionals greatly influence the performance of these parasitoids, future studies must focus on how climate change affect their efficiency. Studies like those examining the development of *Gryon aetherium* under varying temperature could provide insights into their adaptability and potential range expansion in response to changing climates. Further, investigating optimal release strategies and timing, as well as the impact on non-target species, will be essential for successful implementation in pest management programs.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

1. Kataria R, Kumar D. Occurrence and infestation level of sucking pests: Aphids on various host plants in agricultural fields of Vadodara, Gujarat (India). *International Journal of Scientific and Research Publications*. 2012;2(7).
2. Paudel S, Bechinski EJ, Stokes BS, Pappu HR, Eigenbrode SD. Deriving economic models for pea aphid (Hemiptera: Aphididae) as a direct pest and a virus vector on commercial lentils. *Journal of Economic Entomology*. 2018; 111:2225–2232.
3. Rai AB, Halder J, Kodandara MH. Emerging insect pest problems in vegetable crops and their management in India: An appraisal. *Pest Management in Horticultural Ecosystems*. 2014;20(2):113–122.
4. Chakravarty S, Agnihotri M, Kumar L, Ahmad MA. Influence of weather parameters on natural egg parasitization of pod bug, *Clavigralla gibbosa* Spinola (Hemiptera: Coreidae) on pigeon pea. *The Bioscan*. 2016;11(3):1759–1763.
5. Gopali JB, Sharma OP, Yelshetty S, Rachappa V. Effect of insecticides and biorationals against pod bug (*Clavigralla gibbosa*) in pigeonpea. *Indian Journal of Agricultural Sciences*. 2013;83(5):582–585.
6. Halder J, Kushwaha D, Deb D, Rai AB. Effect of meteorological parameters on abundance of mirid bug, *Nesidiocoris cruentatus* (Ballard) (Hemiptera: Miridae): An emerging insect pest of bottle gourd. *Journal of Agrometeorology*. 2017;19(2):180–182.
7. Sundararaj R, Krishnan S, Sumalatha BV. Invasion and expansion of exotic whiteflies (Hemiptera: Aleyrodidae) in India and their economic importance. *Phytoparasitica*. 2021; 49:851–863.
8. Cork A, Kamal NQ, Alam SN, Choudhury JS, Talekar NS. Pheromone and their applications to insect pest control. *Bangladesh Journal of Entomology*. 2003;13(2):1–13.
9. Roversi PF, Maltese M, Simoni S. *Graphosomalineatum* (Hemiptera: Pentatomidae): a suitable host for mass rearing *Ooencyrtustelenomicida* (Hymenoptera: Encyrtidae). *International Journal of Pest Management*. 2017;1–9.
10. Orr DB. Scelionid wasps as biological control agents: a review. *Florida Entomologist*. 1988; 71:506–528.
11. Austin AD, Johnson NF, Dowton M. Systematics, evolution, and biology of scelionid and platygastriid wasps. *Annual Review of Entomology*. 2005; 50:553–82.
12. Powell JE, Shepard M. Biology of Australian and United States strains of *Trissolcus basalis*, a parasitoid of the green vegetable bug *Nezaraviridula*. *Australian Journal of Ecology*. 1982; 7:181–186.
13. Yan CJ, Talamas E, Lahey Z, Chen HY. *Protelenomus* Kieffer is a derived lineage of *Trissolcus* Ashmead (Hymenoptera, Scelionidae), with comments on the evolution of phoresy in Scelionidae. *Journal of Hymenoptera Research*. 2022; 94:121–137. <https://doi.org/10.3897/jhr.94.95961>

14. Forouzan M, Shirazi J, Safavi A, Rezaei M. Oviposition model of *Trissolcus basalis* Wholaston (Hym.: Scelionidae) on sunn pest eggs. JAST. 2015; 17:551–560.
15. Rajmohana K. A checklist of the Scelionidae (Hymenoptera: Platygastroidea) of India. Zoos' Print Journal. 2006; 21:2506–2513.
16. Chen H, Lahey Z, Talamas EJ, Valerio AA, Popovici OA, Musetti L, Klompen H, Polaszek A, Masner L, Austin AD, Johnson NF. An integrated phylogenetic reassessment of the parasitoid superfamily Platygastroidea (Hymenoptera: Proctotrupomorpha) results in a revised familial classification. Systematic Entomology. 2021;46(4):1088–113.
17. Jones WA. World review of the parasitoids of the southern green stink bug, *Nezaraviridula* (L.) (Heteroptera: Pentatomidae). Annals of the Entomological Society of America. 1988;81(2):262–73.
18. Johnson NF. Catalog of world species of Proctotrupeoidea, exclusive of Platygastriidae (Hymenoptera). Memoirs of the American Entomological Institute. 1992; 51:1–825.
19. Muesebeck CFW. Superfamily Proctotrupeoidea. In: Krombein KV, Hurd PD, Smith DR, editors. Catalog of Hymenoptera in America north of Mexico. Washington (DC): Smithsonian Institution Press; 1979. p. 1121–1186.
20. Taekul C, Valerio AA, Austin AD, Klompen H, Johnson NF. Molecular phylogeny of telenomine egg parasitoids (Hymenoptera: Platygastriidae s.l.: Telenominae): evolution of host shifts and implications for classification. Systematic Entomology. 2014; 39:24–35.
21. Vlug HJ. Catalogue of the Platygastriidae (Platygastroidea) of the world (Insecta: Hymenoptera). Hymenopterorum Catalogus. 1995; 19:1–168.
22. Chen H, Talamas EJ, Pang H. Notes on the hosts of *Trissolcus* Ashmead (Hymenoptera: Scelionidae) from China. Biodiversity Data Journal. 2020a;8: e53786. <https://doi.org/10.3897/BDJ.8.e53786>
23. Veenakumari K, Mohanraj P. Redescription of *Protelenomus flavicornis* Kieffer (Platygastroidea: Platygastriidae) from India. Entomofauna, Zeitschrift für Entomologie. 2015; 36:305–312.
24. Veenakumari K, Notton DG, Polaszek A. World Revision of the Genus *Protelenomus* Kieffer (Hymenoptera: Scelionidae: Telenominae). Annales Zoologici. 2019;69(2):381–406. <https://doi.org/10.3161/00034541ANZ2019.69.2.006>
25. Rajmohana K. On a new species of *Protelenomus* Kieffer, 1906 (Hymenoptera-Platygastriidae-Telenominae) from Arunachal Pradesh, being the first record of the genus from India. Colemania. 2013; 38:1–7.
26. Zhang YZ, Li W and Huang DW (2005) A Taxonomic study of Chinese species of *Ooencyrtus* (Insecta: Hymenoptera: Encyrtidae). *Zoological Studies* 44: 347–360.
27. Ogburn EC, Bessin R, Dieckhoff C, Dobson R, Grieshop M, Hoelmer KA, Walgenbach JF. Natural enemy impact on eggs of the invasive brown marmorated stink bug, *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae), in organic agroecosystems: a regional assessment. Biological Control. 2016; 101:39–51.
28. Krupke CH, Brunner JF. Parasitoids of the consperse stink bug (Hemiptera: Pentatomidae) in north central Washington and attractiveness of a host-produced pheromone component. Journal of Entomological Science. 2003;38(1):84–92.
29. Golin V, Loiacono MS, Margaría CB, Aquino DA. Natural incidence of egg parasitoids of *Edessa meditabunda* (F.) (Hemiptera: Pentatomidae) on *Crotalaria*

- spectabilis* in Campo Novo do Parecis, MT, Brazil. *Neotropical Entomology*. 2011; 40:617–618.
30. Garipey TD, Bruin A, Konopka J, Scott Dupree C, Fraser H, Bon MC, Talamas E. A modified DNA barcode approach to define trophic interactions between native and exotic pentatomids and their parasitoids. *Molecular Ecology*. 2019;28(2):456–470.
 31. Moraglio ST, Tortorici F, Pansa MG, Castelli G, Pontini M, Scovero S, Visentin S, Tavella L. A 3-year survey on parasitism of *Halyomorpha halys* by egg parasitoids in northern Italy. *Journal of Pest Science*. 2020; 93:183–194.
 32. Riley CV, Howard LO. *Insect Life*. U. S. Dep. Agric., Div. Ent. 1891; IV:124.
 33. Bertoldi V, Rondoni G, Brodeur J, Conti E. An egg parasitoid efficiently exploits cues from a coevolved host but not those from a novel host. *Frontiers in Physiology*. 2019; 10:746.
 34. Holthouse MC, Schumm ZR, Talamas EJ, Spears LR, Alston DG. Surveys in northern Utah for egg parasitoids of *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae) detect *Trissolcus japonicus* (Ashmead) (Hymenoptera: Scelionidae). *Biodiversity Data Journal*. 2020;8: e53363. <https://doi.org/10.3897/BDJ.8.e53363>.
 35. Buleza VV, Mikheev AV. Localisation in the host's body of substances stimulating host-seeking by the egg parasite *Trissolcus grandis* (Thom.). *KhemoretseptsiyaNasekomykh*. 1979; 4:95–100.
 36. Trissi AN, El Bouhssini M, Ibrahim J, Abdulhai M, Parker BL, Reid W, El-Haramein FJ. Effect of egg parasitoid density on the population suppression of Sunn Pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae), and its resulting impact on bread wheat grain quality. *Journal of Pest Science*. 2006; 79:83–87.
 37. Waterhouse DF. *Biological Control of Insect Pests: Southeast Asian Prospects*. Canberra: Australian Centre for International Agricultural Research; 1998. 548 p.
 38. Colazza BS, McElfresh JS, Millar JG. Identification of volatile synomones, induced by *Nezaraviridula* feeding and oviposition on bean spp., that attract the egg parasitoid: *Trissolcus basalis*. *Journal of Chemical Ecology*. 2004;30(5):945–964.
 39. Tillman PG. Parasitism and predation of stink bug (Heteroptera: Pentatomidae) eggs in Georgia corn fields. *Environmental Entomology*. 2010;39(4):1184–1194.
 40. Colazza S, Bin F. Risposta di *Trissolcus simoni* (Mayr) (Hym.: Scelionidae) and *Eurydemaventrata* Klt. (Het.: Pentatomidae) in olfattometro. In: *Proceedings of the Atti XV Congresso Nazionale Italiano di Entomologia*, pp. 833–840, L'Aquila, Italy; 1988.
 41. Conti E, Salerno G, Bin F, Vinson SB. The role of host semiochemicals in parasitoid specificity: a case study with *Trissolcus brochymenae* and *Trissolcus simoni* on pentatomid bugs. *Biological Control*. 2004;29(3):435–444. <https://doi.org/10.1016/j.biocontrol.2003.08.009>
 42. Sousa KKA, Silva NNP, Querino RB, Silva PHS, Grazia J. Diversity, seasonality, and egg parasitism of hemipteran (Coreidae and Pentatomidae) from a cowpea crop in northeastern Brazil. *Florida Entomologist*. 2019;102(1):29–35.
 43. Silva NN, Sousa KK, Silva PHS, Querino RB. New records of egg parasitoids of stink bugs (Hemiptera: Pentatomidae) on rice in Piauí, Brazil: rate parasitism, incidence and seasonality. *Entomological Communications*. 2021;3:ec03020–ec03020.

44. de Almeida Paz-Neto A, Querino RB, Margaría CB. Egg parasitoids of stink bugs (Hemiptera: Coreidae and Pentatomidae) on soybean and cowpea in Brazil. *Florida Entomologist*. 2015;929–932.
45. Matsuo K, Hirose Y, Johnson N. A taxonomic issue of two species of *Trissolcus* (Hymenoptera: Platygastridae) parasitic on eggs of the brown-winged green bug, *Plautia stali* (Hemiptera: Pentatomidae): resurrection of *T. plautiae*, a cryptic species of *T. japonicus* revealed by morphology, reproductive isolation and molecular evidence. *Applied Entomology and Zoology*. 2014;49(3):385–394. <https://doi.org/10.1007/s13355-014-0260-4>
46. Hokyo N, Kiritani K. Two species of egg parasites of the southern green stink bug, *Nezaraviridula*. *Japanese Journal of Applied Entomology and Zoology*. 1963; 7:214–227.
47. Yasumatsu K, Watanabe C. A Tentative Catalogue of Insect Natural Enemies of Injurious Insects in Japan. Part 2: Host Parasite-Predator Catalogue. Japan: Faculty of Agriculture, Kyushu University; 1964. p. 7–8.
48. Zapponi L, Zhang YZ, Li W, Huang DW. A Taxonomic study of Chinese species of *Ooencyrtus* (Insecta: Hymenoptera: Encyrtidae). *Zoological Studies*. 2005; 44:347–360.
49. Tofangsazi N, Hogg BN, Hougardy E, Stokes K, Pratt PD. Host searching behavior of *Gryon gonikopalense* and *Trissolcus hyalinipennis* (Hymenoptera: Scelionidae), two candidate biological control agents for *Bagradahilaris* (Hemiptera: Pentatomidae). *Biological Control*. 2020; 151:104397.
50. Higuchi H. Seasonal prevalence of egg parasitoids attacking *Piezodorushybneri* (Heteroptera: Pentatomidae) on soybeans. *Applied Entomology and Zoology*. 1993;28(3):347–352.
51. Arida GS, Shepard BM, Perez VA. Parasitization of the Malayan black bug (MBB) by five species of egg parasitoids.
52. Tortorici F, Orrù B, Timokhov AV, Bout A, Bon MC, Tavella L, Talamas EJ. *Telenomus* Haliday (Hymenoptera, Scelionidae) parasitizing Pentatomidae (Hemiptera) in the Palearctic region. *Journal of Hymenoptera Research*. 2024; 97:591–620.
53. Hidaka T. Biological investigation on *Telenomus gifuensis* Ashmead (Hym. Scelionidae), an egg parasite of *Scotinopharalurida* Burmeister (Hem. Pentatomidae) in Japan. *Acta Hymenopterorum*. 1958; 1:75–93.
54. Girault. *Entomological News*. 1904;336.
55. McPherson JE, Bundy CS, Kuhar TP. *Murgantia histrionica* (Hahn), in *Invasive Stink Bugs and Related Species (Pentatomidae): Biology, Higher Systematics, Semiochemistry, and Management*, ed. by JE McPherson. Boca Raton (FL): CRC Press; 2018. p. 333–350.
56. Bueno AF, Braz ÉC, Favetti BM, de Barros França-Neto J, Silva GV. Release of the egg parasitoid *Telenomus podisi* to manage the Neotropical Brown Stink Bug, *Euschistus heros*, in soybean production. *Crop Protection*. 2020; 137:105310.
57. Parra LM, Carvalho JR, Hoback WW, Oliveira RCD. Optimizing mass rearing of the egg parasitoid, *Telenomus podisi*, for control of the brown stink bug, *Euschistus heros*. *Insects*. 2023;14(5):435.

58. Cornelius ML, Dieckhoff C, Vinyard BT, Hoelmer KA. Parasitism and predation on sentinel egg masses of the brown marmorated stink bug (Hemiptera: Pentatomidae) in three vegetable crops: importance of dissections for evaluating the impact of native parasitoids on an exotic pest. *Environmental Entomology*. 2016; nvw134.
59. Moonga MN, Kamminga K, Davis JA. Status of stink bug (Hemiptera: Pentatomidae) egg parasitoids in soybeans in Louisiana. *Environmental Entomology*. 2018;47(6):1459–1464.
60. Queiroz AP, Taguti EA, Bueno AF, Grande MLM, Costa CO, Moraglio. Host preferences of *Telenomus podisi* (Hymenoptera: Scelionidae): parasitism on eggs of *Dichelops melacanthus*, *Euschistus heros*, and *Podisus nigrispinus* (Hemiptera: Pentatomidae). *Neotropical Entomology*. 2018; 47:543–552.
61. Ozdemir IO, Tuncer C, Tortorici F, Ozer G. Egg parasitoids of green shield bug, *Palomena prasina* L. (Hemiptera: Pentatomidae) in hazelnut orchards of Turkey. *Biocontrol Science and Technology*. 2023;33(1):61–75. DOI: 10.1080/09583157.2022.2158308.
62. Morrill AW. Description of a new species of *Telenomus* with observations on its habits and life history. *The American Naturalist*. 1907;41(487):417–430.
63. Aldrich JR. Chemical communication in true bugs and exploitation by parasitoids and commensals. In: Carde RT, Bell WJ, editors. *Chemical Ecology of Insects II*. London: Chapman & Hall; 1995. p. 318–63.
64. Aldrich JR, Kochansky JP, Abrams CB. Attractant for a beneficial insect and its parasitoids: pheromone of a predatory spined soldier bug, *Podisus maculiventris* (Hemiptera: Pentatomidae). *Environmental Entomology*. 1984; 13:1031–6.
65. Orr DB, Russin JS, Boethel DJ. Reproductive biology and behavior of *Telenomus calvus* (Hymenoptera: Scelionidae), a phoretic egg parasitoid of *Podisus maculiventris* (Hemiptera: Pentatomidae). *Canadian Entomologist*. 1986; 118:1063–1072.
66. Bruni R, Sant'Ana J, Aldrich JR, Bin F. Influence of host pheromone on egg parasitism by scelionid wasps: comparison of phoretic and nonphoretic parasitoids. *Journal of Insect Behavior*. 2000;13(2):165–73.
67. Srikumar KK, Bhat PS, Vanitha K. Phenology and Parasitization Behaviour of *Telenomus cuspis* (Hymenoptera: Platygasteridae) an Egg Parasitoid of *Helopeltis antonii* (Hemiptera: Miridae) in Cashew. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.* 2015;85:437–442. <https://doi.org/10.1007/s40011-014-0365-0>
68. Phillips JS. A search for parasites of dasynine bugs in the Netherlands Indies. *Transactions of the Royal Entomological Society of London*. 1941; 91:119–144.
69. Kalshoven LG. *Pests of Crops in Indonesia*. 1950;1065 pp.
70. Greve JVS, Ismay JW, editors. *Crop Insect Survey of Papua New Guinea from July 1st 1969 to December 31st, 1978*.
71. Manjunath TM, Rai PS, Gowda G. Parasites and predators of *Nilaparvatalugens* in India. *Pans*. 1978;24(3):265–269.
72. Rojas-Gálvez NR, Talamas E, Albornoz MV, Flores MF, Barros-Parada W, Bout A. *Gryon aetherium* Talamas (Hymenoptera, Scelionidae): Parasitoid of *Bagradahilaris* (Burmeister) (Hemiptera, Pentatomidae) Adventive in Chile. *Journal of Hymenoptera Research*. 2021; 87:493–501.

73. Hogg BN, Grettenberger IM, Borkent CJ. Parasitism by *Gryon aetherium* (Hymenoptera: Scelionidae) on *Bagradahilaris* (Hemiptera: Pentatomidae) eggs in northcentral California. *Journal of Economic Entomology*. 2023;116(5):1540–1550. <https://doi.org/10.1093/jee/toad172>.
74. Chen H, Talamas EJ, Bon MC, Moore MR. *Gryon ancinla* Kozlov & Lê (Hymenoptera: Scelionidae): host association, expanded distribution, redescription and a new synonymy. *Biodiversity Data Journal*. 2020b;8. doi: 10.3897/BDJ.8.e47687
75. Shanower TG, Anitha V, Bhagwat VR, Dreyer H. Parasitism of *Clavigralla* spp. (Hemiptera: Coreidae) eggs by *Gryon clavigrallae* Mineo (Hymenoptera: Scelionidae). *Journal of Biological Control*. 1996; 10:1–7.
76. Lomeli-Flores JR, Rodríguez-Rodríguez SE, Rodríguez-Levya E, González-Hernández H, Garipey TD, Talamas EJ. Field studies and molecular forensics identify a new association: *Idris elba* Talamas, sp. nov. parasitizes the eggs of *Bagradahilaris* (Burmeister). *Journal of Hymenoptera Research*. 2019; 73:125–141. <https://doi.org/10.3897/jhr.73.38025>
77. Kohno K. Phoresy by an egg parasitoid, *Protelenomus* sp. (Hymenoptera: Scelionidae), on the coreid bug *Anoplocnemisphasiana* (Heteroptera: Coreidae).
78. Raju MR, Arifunnahar M, Mostafiz MM, Alim MA, Hossain MA, Talamas EJ. *Hadronotus pubescens* (Motschoulsky) (Hymenoptera, Scelionidae): Redescription, biological attributes, and parasitism on eggs of *Riptortus pedestris* (Fab.) (Hemiptera, Alydidae). *Journal of Hymenoptera Research*. 2022; 94:139–161.
79. Boyle SM, Salom S, Schultz P, Lopez L, Weber DC, Kuhar TP. Augmentative biological control for squash bug (Hemiptera: Coreidae) using the egg parasitoid, *Hadronotus pennsylvanicus* (Hymenoptera: Scelionidae). *Environmental Entomology*. 2023;52(5):779–86.
80. Singh LRK, Johnson NF, Ramamurthy VV. New species of the genus *Psix* Kozlov & Lê (Hymenoptera: Platygastriidae). *Zootaxa*. 2012;3530(1):35–42.
81. Johnson NF. Hymenoptera Online (HOL). [Internet]. 2019. Available from: <http://hol.osu.edu/> [last accessed on 26/09/2019].
82. Masner L. A revision of *Gryon* Haliday in North America (Hymenoptera: Proctotrupeoidea: Scelionidae). *Canadian Entomologist*. 1983; 115:123–174. <https://doi.org/10.4039/Ent115123-2>.
83. Boyle SM, Cornelius M, Talamas EJ, Straser RK, Wilson H, Daane KM, Weber DC, Kuhar TP. *Hadronotus pennsylvanicus* (Hymenoptera: Scelionidae): a biological control agent for pestiferous leaf-footed bugs (Hemiptera: Coreidae). *Journal of Integrated Pest Management*. 2024;15(1):15.
84. Panizzi AR. History and contemporary perspectives of the integrated pest management of soybean in Brazil. *Neotropical Entomology*. 2013; 42:119–127.
85. Laumann RA, Moraes MCB, Silva JPD, Vieira AMC, Silveira SD, Borges M. Egg parasitoid wasps as natural enemies of the neotropical stink bug *Dichelops melacanthus*. *Pesquisa Agropecuária Brasileira*. 2010; 45:442–449.
86. Mineo G, Lucido G. Difesa del nocciolodagliartropodidannosi. XI. Prove di controllo biologico del *Gonocerus acuteangulatus* (Goeze) con lanci in-oculativi di alcuni suoiparassitoidi: *Gryon muscaeformis* (Nees), *Trissolcus flavipes* (Thoms.),

Ooencyrtus gonoceri Vigg. Bollettino dell'Istituto di Entomologia Agraria e Osservatorio Fitopatologico di Palermo. 1976; 9:109–128.

87. Bueno AF, Sosa-Gómez D, Corrêa-Ferreira B, Moscardi F, Bueno R. Inimigos naturais das pragas da soja. In: Hoffman-Campo CB, Corrêa-Ferreira BS, Moscardi F, editors. Soja: Manejo Integrado de Insetos e Outros Artrópodes-Praga. Brazil: Embrapa Soja; 2012. p. 493–629.

UNDER PEER REVIEW