

Original Research Article

Assessment of Early blight disease resistant Tomato F₁ hybrid ArkaAbhed in Karimnagar District of Telangana, India

ABSTRACT

Tomato is the major Solanaceous vegetable crop grown in Karimnagar District of Telangana. *Alternariatomatophila* and *Alternariasolani* cause early blight disease and severe yield losses to the tomato growers in Karimnagar District. Early blight is the major problem in Karimnagar from June to September months causing severe crop losses to the tomato farmers. Usually farmers are spraying different fungicides based on Copper to control early blight disease for private hybrids. It controls 25-35% of the disease and increased cost of cultivation. Hence, early blight disease resistant hybrid ArkaAbhed was tested against private hybrid in KVK, Karimnagar operational area. Data on disease incidence was collected at 15, 30, 60, 90, 120 DAP in all the hybrids. The data revealed that Arkaabhed recorded significantly very lower incidence of early blight disease than private hybrid during all crop growth stages. It recorded 1.5% early blight incidence whereas private hybrid recorded 32.4%. Reduced number of sprayings to control early blight disease were also significantly low i.e. two sprays were carried out whereas in private hybrid seven sprays were done during in rainy period. According to yield ArkaAbhed recorded 68.3t/ha whereas it was 58.6t/ha in private hybrid. The peduncle in green in colour for long days and taste also same for all the days in storage period for 8-12 days in ArkaAbhed. Average fruit weight is 86 gr to 112 grams were recorded.

Key Words: Early blight in tomato, ArkaAbhed, Disease resistant tomato

INTRODUCTION

One of the most well-liked and extensively consumed vegetable crops worldwide is the tomato (*Solanumlycopersicum* L.), and a high-quality yield is a necessary need for its commercial success worldwide. Fourth most grown crop worldwide is the tomato. In terms of acreage and production, India comes in second place globally, right after China. With an annual yield of 20.51 million tonnes and a productivity of 21.13 t/ha, tomatoes are produced on 0.81 million hectares in India (NHB, 2018–19). Carotenoids, especially the unsaturated alkylic component lycopene, which lowers the risk of cancer and heart disease, are abundant in it. Tomato fruits are thus in great demand all year round. Due to severe selection and inbreeding during domestication and development, cultivated tomatoes have a limited genetic diversity, making them more vulnerable to disease outbreaks throughout the growing season (Zhang et al. 2002).

One of the most terrible tomato diseases, early blight can cause up to 78% yield and loss of yield (Datar and Mayee 1981 and Bessadat et al. 2014). Early blight on foliage, collar rot on seedlings' basal stem, stem lesions on adult plants, and tomato fruit rot are all indications of *Alternariasolani* infection. The symptoms of early blight manifest on the plant's aerial parts, including the leaves, stem, and fruits. Brown, spherical spots that may develop up to half an inch in diameter are seen on leaves. After sporulation, larger patches with concentric rings emerge, which develops characteristic bull eye-shaped patches or "target board" symptoms due to

sporulation pattern. The Alternaria blight has been managed with a variety of fungicides; nevertheless, fungicide treatment is neither ecologically nor economically viable. According to Kemmitt et al. (2002), fungicides are first sprayed 1-2 days after transplanting and thereafter need to be routinely administered at intervals of 7-10 days for successful control. This increases the expense of production and pollutes the environment. The most cost-effective and long-lasting method of controlling early blight is the development of resistant cultivars. Due to its devastating nature, early blight draws a lot of attention. The tomato plant (*Solanumlycopersicum* L. [Peralta et al. 2005, syn. *L. esculentum* Mill.]) is especially susceptible to this disease, which in extreme circumstances can result in total defoliation, in areas with high humidity, heavy rainfall, and relatively high temperatures between 24 and 29 degree Celsius. In semi-arid areas with regular, long-lasting nighttime dews, epidemics can also happen (Rotem and Reichert 1964).

Therefore, the goal of the current study was to pinpoint tomato resistance sources against early blight disease. University, which would offer a more extensive genetic base to support the emergence of resistant cultivars.

MATERIALS AND METHODS

The present study was carried out by the KrishiVigyan Kendra during *Kharif* season from 2021 to 2023 at the farmers' fields of different villages of the Karimnagar district of Telangana. In total 323 Ha of tomato fields of Karimnagar District. Selected farmers based on crops growing in the village and conducted training programmes to the famers. A group of cooperative farmers were identified based on their participation and feedback received during the preliminary survey and interactive FPO meetings

The seeds of ArkaAbhed bought from IIHR, Bengaluru. We had grown nursery in KVK Jammikunta, the seedlings were transplanted during the 1st week of July in the years 2020-21, 2021-22, and 2022-23, right into the fields of Karimnagar farmers. For this on-farm trial, ArkaAbhed and the hybrid PHS-448 check hybrid were transplanted at same days. The seedlings used in the transplantation were between 33 days old and were spaced at 75 x 45 cm intervals. During the land preparation process, 6 tonnes per hectare (t/ha) of Vermicompost was incorporated into the soil. Further more, a combination of NPK fertilizers, The nutrient requirement for hybrids is 200:250:250 kg of NPK per ha. 75 % of P (187.5 kg P which comes to 1172 kg of superphosphate) is applied as basal. The remaining quantity of 200:62.5:250 kg of NPK per ha applied through fertigation. Every day irrigation had given for one hour. Along with this, water soluble fertilizers have to be given.

This dose was split and given once in 3 days for the entire crop period through fertigation as detailed below.

ArkaAbhed characters: It is a multiple disease resistant hybrid *i.e.* resistant to late blight, early blight, leaf curl virus and bacterial wilt with a crop duration of 140-145 days and yield potential of 70-75t/ha. It is suitable for cultivation during *Kharif*, *Rabi* and Summer.

Table No.1 :Disease rating scale (Henfling 1987)

Mean severity (%)	Limits (%)	Symptoms
0	0	No symptoms
2.5	Tr < 5	Up to 10 lesions per plant
10	5 -15	Lesions easily seen at closer distance. Maximum

		foliage area affected up to 20 leaflets.
25	15 - 35	About 25 % of foliage is covered with lesions
50	35 - < 65	Lower leaves are dead. About half the foliage area is destroyed
75	65 - < 85	About 75 % leaf area destroyed; field appears neither brown nor green
90	85 - < 95	Only top leaves are green. Many stems have large lesions
97.5	95 - < 100	All plants in a spot are brown-colored. A few top leaves still have some green areas. Most stems have lesions
100	100	All leaves and stems dead

Plant protection measures were implemented in both the check and trial plots to accurately observe the growth and yield parameters of the tomato hybrids. Tomato F1 hybrid ArkaAbhed (H-397) is renowned for its resistance to multiple diseases, including tomato leaf curl disease (ToLCV), bacterial wilt (BW), early blight, and late blight. ArkaAbhed tomato plants exhibit a semi-determinate growth habit and possess dark green foliage. The fruits of ArkaAbhed are firm, oblate round, and of medium-large size, typically weighing between 90-100 grams. Key characteristics of ArkaAbhed include its resistance to late blight, early blight, leaf curl virus, and bacterial wilt, a crop duration of 140-145 days, and a remarkable yield potential of 70-75 tonnes per hectare. This hybrid is suitable for cultivation during the Kharif, Rabi, and summer seasons.

The fruits of ArkaAbhed are characterized by their oblate to high round shape, large size (ranging from 90-97 grams), deep red coloration, and firm texture. On the other hand, PHS-448 is a private hybrid variety with plants that exhibit a growth pattern as semi-determinate. These plants feature robust foliage cover and vigor. The time to the first harvest for PHS-448 is typically 62-67 days. The fruits produced by this hybrid are flat and round, possessing green coloration with a very firm structure and a distinctly acidic taste. Their color matures to a deep red, and the average fruit weight falls within the range of 100-120 grams. Additionally, PHS-448 exhibits intermediate resistance to tomato leaf curl virus. The seeds for the tomato hybrids ArkaAbhed were obtained from the Indian Institute of Horticulture Research (IIHR) in Bengaluru.

A field experiment was conducted across various locations in Thimapur, Mogilipalem villages within the Karimnagar during the Rabi season for three consecutive years: 2021-22, 2022-23 and 2023-24. The experiment utilized hybrids ArkaAbhed and PHS-448 Hybrid as a reference or check hybrid. The experimental design followed a randomized block layout with five replications. Various traits, early blight symptoms, virus infestation, including plant height, days to first flowering, days required for 50% flowering, days to the initial harvest, number of fruits per plant, average fruit weight (in grams), fruit yield per individual plant, yield per hectare, net returns per hectare, and the benefit-cost ratio, were meticulously recorded and subsequently subjected to statistical analysis using the methodology outlined by Panse and Sukhatme (1985).

RESULTS AND DISCUSSION

“Data on Early blight incidence was collected from vegetative stage to fruiting stage in all the three treatments in all the locations. There is significant difference among tomato hybrid ArkaAbhed and private hybrid with respect to Early blight incidence during all growth stages. During vegetative stage of the crop, Early blight incidence was observed very less in tomato hybrid ArkaAbhed whereas in PHS448 and private hybrid, half of the foliage *i.e.* lower leaves are dried due to early blight disease. During flowering stage, 2-4 % of early blight incidence was observed in ArkaAbhed in heavy rainy days only whereas private hybrid 35-45% early blight incidence was observed through the

crop period. During this stage about 60-75% of leaf area was blighted in the susceptible Private hybrid. During fruiting stage 36-40 % of early blight incidence was observed in private hybrid respectively whereas in ArkaAbhed 0 % (very minute observations) of early and late blight incidence was observed. This might be attributed to high disease resistance and favorable environmental conditions for spread of the disease. And late blight disease also observed in private and but it is very less in Arkaabhed triple resistant hybrid” [26].

Table No2. Demonstration and Check results given below:

Treatments	Yield(t/ha)	Costofcultivation (Rs/ha)	Grossincome (Rs/ha)	Netincome (Rs/ha)	B:C ratio
ArkaAbhed	68.3	118900.00	409800.00	290900	1:2.4
Privatehybrid	58.6	124300.00	351600.00	227300.00	1:1.8

Table No.3 Disease incidence particulars

S.No	Particulars	Arkaabhed	Private hybrid PHS -448
1.	Seed rate /Ha	80gr	80gr
2.	Seed cost	2400.00	4800.00
3.	Land preparation	12000.00	9000.00
4	Cost of mulching +trellis+labour	56000.00	56000.00
5	Fertilizers and pesticides	26500.00	31200.00
6	Weeding labour charges	6000.00	9800.00
7	Fruit picking charges	16000.00	13500.00
8	Cost of cultivation	118900.00	124300.00
9	ToLCV, BW & EB	Very less	observed more
10.	Yield (t / Ha)	68.3	58.6
11	Average Market rate /Kg	6	6
12	Gross returns	409800.00	351600.00
13	Net returns	290900.00	227300.00
14	C:B Ratio	1:2.4	1:1.8

Table No.4:Disease incidence particulars

Sl.No	Particulars	ArkaAbhed	Farmers Plot(Check)
1	Early blight symptoms	2% observed	35 %observed
2	Late blight symptoms	3% observed	40 %observed
3	Viral symptoms	Not observed	Not observed

Table5:Meanseveritypercentageof early blight amongtwohybridsduringdifferentstages.

Cropstage	Treatment	Different Percentages average
Vegetative stage (30DAP)	ArkaAbhed ^a	0
	Privatehybrid ^b	10
Flowering stage(60DAP)	ArkaAbhed ^a	0
	Privatehybrid ^b	35-40
Fruitingstage (90DAP)	ArkaAbhed ^a	0.02
	Privatehybrid ^b	40-65
Total	ArkaAbhed ^a	0.03
	Privatehybrid ^b	45

CONCLUSION

ArkaAbhed performed very well in farmers fields with good fruit quality, longer crop duration, high yield and observed disease resistance for early blight under Karimnagar field conditions which are highly conducive to early blight and late blight development in rainy season in Karimnagar District. If farmers manage early blight disease in early stages from 10th day after transplanting and in flowering period, they will get good yields and good income.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. **Not used**

2.

3.

References

1. P Adhikari, Y Oh, DR Panthee - journal of molecular sciences (2017) discussed in Current status of **early blight** resistance in **tomato**
2. R Chaerani, RE Voorrips -Journal of general plant pathology (2006) Tomato Early blight (*Alternaria solani*) the pathogen ,genetics and breeding for resistance
3. Xie, Y Shao, X Li, Y He -Scientific reports(2015)- Detection of early blight and late blight diseases on tomato
4. CK Roy, N Akter, MKI Sarkar, MU Pk... - ... Microbiology Journal(2019)-Control of early blight of tomato caused by and screening of tomato varieties against the pathogen
5. Robin,E.,&Choen,Y.(2004).Oospores associated with tomato seed may lead to seed borne
i. transmission of *Phytophthora infestans*.
6. *Phytoparasitica*, 32: 237-245. Saleem, M. Y., Akhtar, K.P., Iqbal, Q., Asghar, M., Hameed, A., &
a. Shoaib, M. (2016). Development of tomato hybrids with multiple disease tolerance.
b. *Pakistan Journal of Botany*, 48(2):771-778.
7. Sadashiva, A. T., Hanson, P., Reddy, M. K., Ravishankar, K. V., Prasad, M., Prasanna, H. C., & Bhat, L.
(2017). Breeding tomato (*Solanum lycopersicum* L.) for resistance to biotic and abiotic stresses. *Journal of Horticultural Sciences*, 12(2): 91-105.

8. Shattock, R. (2002). *Phytophthora infestans*: populations, pathogenicity and phenylamides. *Pest Management Science*, 58: 944-950.
9. Sunitha, C. H., Chinnam Naidu, D., Raja Kumar, N., BhagyaLakshimi, K. and ChittiBabu, G. (2020).
 - a. Assessment of tomato (*Solanum lycopersicon* L.) hybrids for performance and adoptability at
 - b. Srikakulam District, Andhra Pradesh. *International Journal of Ecology and Environmental Sciences*, 2(3): 317-319.
10. Akhtar, K. P., Saleem, M. Y., Asghar, M., Ahmad, M., & Sarwar, N. (2010). Resistance of *Solanum*
 - a. species to Cucumber mosaic virus subgroup IA and its vector *Myzus persicae*. *European Journal of Plant Pathology*, 128(4): 435-450.
11. Azam, F., & Shah, S. J. (2003). Exploring the role of farmer led management practices on various
 - a. and cucumber diseases in Peshawar and Dragai areas of NWFP. In: *Final Reports on PHF funded research projects*. pp. 1-75.
12. Chen, Y., & Halterman, D. A. (2011). Phenotypic characterization of potato late blight resistance mediated by the broad-spectrum resistance gene *RB*. *Phytopathology*, 101: 263–270.
13. Hardham, A. R., Jones, D. A., & Takemoto, D. (2007). Cytoskeleton and cell wall function in
 - a. resistance. *Current Opinion in Plant Biology*, 10: 342–348.
14. Henfling, J. W. (1987). Late blight of potato. Technical Bulletin No. 4. International Potato Centre Lima,
 - a. Peru P.25.
15. Pragma Adhikari, Yeonyee Oh, and Dilip R. Panthee. *Int. J. Mol. Sci.* 2017, 18(10), 2019. Current Status of
16. Early Blight Resistance in Tomato, Allelopathy and microclimatic modification of intercropping
17. With marigold on tomato early blight disease development
18. Horsfield, A., Wicks, T., Davies, K. *et al. Australasian Plant Pathol.* 39, 368–375 (2010)
19. Bio-genetic analysis of resistance in tomato to early blight disease, *Alternaria alternate*
20. C. Singh, Rajesh Kumar*, Major Singh, Ashutosh Rai, M.C. Singh** and Mathura Rai
21. Effect of fungicide use strategies on the control of early blight (*Alternaria solani*) and potato yield
22. Identification of resistant sources against early blight disease of tomato
23. D.C. Batista, M.A. Lima, F. Haddad, L.A. Maffia, E.S.G. Mizubuti
24. Validation of decision support systems for tomato early blight and potato late blight, under
25. Brazilian conditions
26. Ravuri PL, Deva S, Kumar MR. Assessment of Late Blight Resistant Tomato Hybrid Arka Abhed In *Biological Forum—An International Journal* 2021 (Vol. 13, No. 3, pp. 613-616).