

Original Research Article

ECONOMICS OF THE INDIVIDUAL CROPS IN MAIZE AND GROUNDNUT CROPPING SYSTEMS GROWN UNDER DIFFERENT IRRIGATION SYSTEMS WITH VARIED IRRIGATION AND NITROGEN LEVELS

Profitability of Maize and Groundnut crop production under Different Irrigation Systems with Varied Irrigation and Nitrogen Level Application in Northern India

ABSTRACT

An experiment was conducted to study the effects of irrigation systems, irrigation₁ and N levels on the economics of maize and groundnut during 2021-22 and 2022-23 at the College farm, College of Agriculture, Rajendranagar, PJTSAU, Hyderabad. The experiment consisted of 18 treatment combinations (3 irrigation systems as the main plot, 3 irrigation levels as the sub-plot, and 2 nitrogen levels as the sub-sub-plot) in a split plot design replicated thrice. The experiment revealed that among irrigation systems, irrigation₁ and N levels in both crops (maize and groundnut) significantly higher gross returns, net returns₁ and BCR were recorded with M₂ (sub surface sub-surface irrigation system), S₁ (1.2 Epan; IW/CPE) and N₁ (100 per cent percent RDN). In both crops, the interaction between irrigation system and irrigation levels (M x S) revealed significantly higher returns and BCR with sub-surface drip irrigation levels @ 1.2 Epan & 0.9 Epan ratios (M₂S₁ and M₂S₂) followed by surface drip irrigation levels @ 1.2 Epan & 0.9 Epan ratios (M₁S₁ and M₁S₂), surface irrigation @ 1.2 IW/CPE ratio (M₃S₁) whereas surface. In contrast, surface irrigation @ 0.6 IW/CPE ratio (M₃S₃) resulted in least in the least returns and BCR during both the years. The interaction effects of irrigation systems and N levels (M x N), irrigation₁ and N levels (S x N) were non-significant in the groundnut crop whereas they were significant in maize crop crops. The M₂N₁ recorded significantly higher mean gross returns (₹. 1,46,852 ha⁻¹), net returns (₹. 90,960 ha⁻¹) and BCR (2.62) whereas M₃N₂ recorded the lowest mean gross return (₹. 97,052 ha⁻¹), net return (₹. 45,305 ha⁻¹) and BCR (1.87). Among S x N interaction effects, the S₁N₁ (1.2 Epan; IW/CPE with 100 per cent percent RDN) resulted in significantly higher gross returns (₹. 1,44,385 ha⁻¹), net returns (₹. 89,256 ha⁻¹) and and BCR (2.61) followed by S₂N₁ while. In contrast, S₃N₂ resulted in significantly least returns and BCR. The interaction effects of irrigation systems, irrigation₁ and N levels (M x S x N) was were non-significant in the groundnut crop crops while showed showing a significant impact on the economics of maize. The sub-surface irrigation system (M₂) with 1.2 Epan (S₁) and 100 per cent percent RDN (N₁) recorded significantly higher mean gross returns (₹. 1,58,749 ha⁻¹), net return (₹. 1,10,243 ha⁻¹) and BCR (2.82) followed by M₂S₂N₁ (₹. 1,56,106 ha⁻¹, ₹. 1,14,542 ha⁻¹ and 2.79), M₁S₁N₁ and M₁S₂N₁ while M₃S₃N₂ (surface irrigation system with 0.6 IW/CPE and 75 per cent RDN) recorded least mean gross returns (₹. 65,253 ha⁻¹), net returns (₹. 14,106 ha⁻¹) and BCR (1.28) thus making it the least remunerative among all treatment combinations studied.

Keywords: Maize, groundnut, irrigation systems, net returns, gross returns and BCR

1. INTRODUCTION

Maize, the ~~fastest growing~~ fastest-growing cash crop, is ~~growing~~ increasing throughout the world and has the highest production among all ~~the~~ cereals. It is ~~the~~ preferred staple food for 900 million poor, 120 -140 million poor farm families, and about one-third of all malnourished children globally (Murdia *et al.* 2016). Maize is the third most important food grain following wheat and rice for ~~the~~ Indian population and is cultivated over ~~a 9.6 million hectare~~ 9.6-million-hectare area with an annual production and productivity of 28.7 million tonnes and 3.0 t ha⁻¹ (India Stat, 2023). ~~Similarly~~ Similarly, groundnut is considered ~~as~~ the most important oilseed crop in India with a cultivated area of 4.8 million ~~hectare~~ hectares with an annual production and productivity of 9.95 million tonnes and 2.06 t ha⁻¹. Both maize and groundnut crops ~~ranks~~ rank 4th and 1st in acreage but 6th and 2nd in production globally which ~~clearly~~ indicates that the yield potential of maize and groundnut crops are inevitably smaller than ~~the~~ productivity of USA (10.7 & 4.3 t ha⁻¹), China (6.3 & 3.9 t ha⁻¹) and Argentine (7.5 & 3.5 t ha⁻¹) because achieving maximum yield potential requires near perfect management of crop and soil factors which are lacking in farm fields in India (Sreekanth *et al.* 2017).

In Telangana, ~~an 8.5 million hectare~~ 8.5-million-hectare area is gross cropped employing 66.13% of state workers which contributed 15 & 19% in real and nominal terms to the state gross value addition (GVA) in the triennium ending 2021-22 (MOSPI, 2021). Similarly, 36% of the state's land is ~~doubled cropped~~ double-cropped, 64.6% ~~of~~ farmers are marginal and 23.7% ~~of~~ farmers are small. Telangana is rice and cotton centric contributing 52% of state crop value of output (VOO), however, the cereal contribution has shrunk from 44% in 2014-15 to 18% in 2019-20 ~~despite the fact that even though~~ apart from gram, pulses do not appear to be profitable for Telangana farmers as Telangana agricultural households earns about ₹. 4,917 per month as ~~compare~~ compared to Punjab's crop income of ₹.12,597 per month which indicates that the higher agricultural growth in Telangana has not translated into ~~high income~~ high-income levels for farmers (Saini *et al.*, 2023). On this consideration, there is an imminent need to improve cultural practices like fertilizer application along with ~~the~~ proper method of irrigation as water and nutrients are the basic imperative inputs in crop production due to its diverse needs (Sezen *et al.*, 2011) and on this basis, micro irrigation along with fertigation provides the scope for maximizing the crop productivity and monetary returns per unit area and input (Jain *et al.*, 2021). Selection of appropriate crops especially when micro irrigation systems are resorted is a critical decision to be made, considering the initial costs involved. The significance of the selection is more paramount under sandy loam soils of Telangana wherein increasing the cropping intensity is more challenging due to its innate lower productivity and low water & ~~nutrient holding~~ nutrient-holding capacity nature.

With productive farmland facing increased pressure from ~~the~~ growing population, cereal-legume rotations have been proposed as an effective mean to increase the productivity of nutrient-depleted and low ~~water holding~~ water-holding capacity soils in low-external input systems (Buerkert *et al.*, 2001) and to ensure nutritional security. The resource use efficiency, ~~and~~ gross and net returns can be increased by fertigation wherein the ~~plants~~ plants' nutrients are applied with irrigation water, mainly through micro irrigation systems (Jat *et al.*, 2011). Fertigation is relatively a recent innovative method by which fertilizers are applied along with irrigation water through ~~a~~ drip system to enhance fertilizer-use efficiency besides increasing ~~the~~ crop yields and monetary returns. Studies ~~shows~~ show that ~~drip irrigated~~ drip-irrigated maize and groundnut had greater yield, market grade, and gross revenue compared with conventional irrigated regimes (Sorensen and Lamb, 2009). However, information on ~~the~~ economics ~~for~~ of this intensive cropping system is limited. Particularly under

different irrigation systems with varied irrigation and nitrogen levels. Therefore, the present experiment on economics in ~~maize-groundnut~~maize-groundnut crop ~~system systems~~ with varied irrigation ~~systems~~systems, irrigation levels, and nitrogen levels was conducted.

2. MATERIAL AND METHODS

The current experiment was conducted during ~~the~~ *rabi* and summer seasons of 2021-22 and 2022-23 at College ~~farm~~Farm, College of Agriculture, Rajendranagar. The soil of ~~the~~ experimental site was sandy clay loam in texture, moderately alkaline in reaction (pH 7.90), low in available nitrogen ($213.58 \text{ kg ha}^{-1}$), moderately high in phosphorus (25.32 kg ha^{-1}), and medium in potassium content ($180.54 \text{ kg ha}^{-1}$). The moisture content at field capacity and permanent wilting point were 18.44 and 7.88% respectively. The bulk density was 1.41 Mg cm^{-3} . The experiment consisted of three irrigation systems as main plots *viz.*, M₁ - surface drip irrigation system, M₂ - sub-surface drip irrigation system, M₃ - surface irrigation system, three irrigation levels as sub-plot *viz.*, S₁-1.2 Epan; IW/CPE, S₂-0.9 Epan; IW/CPE, S₃-0.6 Epan; IW/CPE and two nitrogen levels as sub-sub-plot *viz.*, N₁-100 ~~per cent~~percent RDN, N₂- 75 ~~per cent~~percent RDN in split plot design replicated thrice. Maize and groundnut varieties KMNH-4010141 and Leepakshi constituted the experimental material. The gross and net plot sizes were 6.0 m x 4.8 m and 4.8 m x 3.6 m, respectively. A complete drip system was installed by Netafim Irrigation Limited. The water distribution system consisted of a ~~main line~~mainline and eighteen sub-mains, each having ~~a~~ control valve for water regulation. Irrigation water from manifolds flowed into 16 mm dripper lines laid out on the ground surface at 0.60 m apart with ~~a~~ spacing of 0.40 m between two inline emitters delivering 2 L hr^{-1} in surface drip irrigation system while in ~~sub surface~~subsurface irrigation system the dripper lines were laid out 15 cm below the soil surface with the same spacing and specification as in surface drip irrigation system. Control valves were fixed separately to each treatment plot to facilitate controlling the water flow as per treatments. ~~Water-A~~ water meter was fixed at the head control unit to quantify the amount of water delivered in each irrigation treatment. Scheduling of irrigation in M₁ and M₂ ~~were was~~ fixed ~~on daily basis~~daily for maize and groundnut crops based on daily evaporation data recorded from (USWB open pan Evaporimeter) obtained from the ACRC, ARI, Rajendranagar, Hyderabad. In ~~the~~ surface irrigation system (M₃) the ~~sub-treatment~~sub-treatment plots were leveled manually, ~~and~~ ridge & furrow and flat-bed land configurations were maintained for maize and groundnut crops during both years of ~~the~~ experiment. In maize and groundnut crops the irrigation was scheduled based on ~~a~~ climatological approach when CPE reached ~~50 and 60 mm~~50- and 60-mm depth respectively.

Recommended doses of N, P₂O₅, and K₂O were 240:80:80 and 30:40:50 kg ha⁻¹ NPK for maize and groundnut respectively, applied in the form of urea, single super phosphate, and ~~sulphate~~sulfate of potassium for maize while urea, single super phosphate, and muriate of potash for groundnut. In both surface and sub-surface drip irrigation systems (M₁ and M₂) a comprehensive fertigation schedule was adopted for maize crop which was already developed by PJTSAU based on crop growth stages and their uptake patterns (Table 1) while ~~in~~ groundnut nitrogen was fertigated in the form of urea in four splits at ~~one week~~one-week interval after proper crop establishment. In ~~the~~ surface irrigation system (M₃) for maize crop, 1/3rd N, full dose of P₂O₅ and K₂O were applied as basal while remaining 2/3rd N was applied in two splits; 1/3rd N at knee height stage and remaining 1/3rd at tasseling depending on irrigation levels in both years whereas in groundnut crop under surface irrigation

system, 2/3rd N, full dose of P₂O₅ and K₂O were applied as basal dose while remaining 1/3rd N was applied at 30 DAS.

What about the fertigation schedule for rabi Groundnut

Table 1. Fertigation schedule for *rabi* maize.

Crop growth stage	Nutrient dose (kg ha ⁻¹ day ⁻¹)	
	Urea	SOP
After sowing; 20 days (10 – 30 DAS)	3.22	1.48
Grand growth period; 25 days (31-55 DAS)	8.66	2.48
Reproductive stage; 20 days (56 – 75 DAS)	8.16	1.98
Kernel development stage; 25 days (76 – 90 DAS)	4.95	1.98

3. RESULTS AND DISCUSSION

3.1 Effects of irrigation systems, irrigation_i and N levels on economics in maize and groundnut.

3.1.1 Effects of irrigation systems

A scrutiny of data (Table 2 and 3) indicated that the mean gross returns (₹. 1,40,507 and 1,91,409 ha⁻¹), net returns (₹. 85,006 and 1,39,683 ha⁻¹) and BCR (2.53 and 3.70) in sub-surface drip irrigation system (M₂) were significantly highest in both maize and groundnut crops when compared to surface drip irrigation (M₁) and surface irrigation systems (M₃) while the returns and BCR under M₃ remained significantly inferior to M₁ and M₂ in both crops during both the seasons and in means. In maize crop, the M₂ recorded 13.12% & 40.94%, 23.1% & 78.7%, 12.4% & 32.5% increased mean gross returns, net returns, and BCR when compared to M₁ and M₃ (Table 2) while in groundnut crop with same irrigation system (M₂) an increased (8.8% & 47.3%, 12.1% & 71.2% and 8.1% & 37.9%) mean gross returns, net returns and BCR was recorded over two other irrigation systems (M₁ and M₃) (Table 3). The higher returns and BCR in drip irrigation systems could be attributed to higher economic yields, lesser cost of cultivation, and efficient use of fertilizers. Current results are also in similarity with Sivanappan, (1978) who worked out the economics of drip irrigation and reported that drip irrigation gave an additional amount of ₹. 10,000 year⁻¹ on a small farm where the available water was not sufficient to irrigate by surface method. Pawar *et al.* (2015) also reported that the drip irrigation system resulted in 27% higher gross returns (₹. 1,13,480 ha⁻¹) over than surface irrigation system (₹. 89,302 ha⁻¹). Joshi *et al* (2015) also observed higher net returns and BCR in drip irrigation over surface irrigation systems. These findings are also supported by Shruti and Aladakatti (2017) who observed furrow method of irrigation recorded the least net returns (₹. 58,317 ha⁻¹) as well as the lowest BCR (2.18) when compared with the drip irrigation system.

3.1.2 Effects of irrigation levels

Data presented in Table-Tables 2 and 3 indicated that, among the three irrigation levels, 1.2 Epan; IW/CPE (S₁) recorded significantly higher gross returns, net returns, and BCR over the other two irrigation levels (S₂ and S₃) in both crops (maize and groundnut) during both the seasons and in mean-while the lowest returns

and BCR was recorded under irrigation scheduled at 0.6 Epan; IW/CPE (S₃). In [the](#) maize crop an overall 5.83% & 44.89%, 9.31% & 100.75%, 4.96% & 42.95%, and in [the](#) groundnut crop 3.64% & 53.3%, 4.7% & 88.2%, 2.7% & 50.9% increased mean gross, net returns and BCR was witnessed with S₁ over two other irrigation levels (S₂ & S₃) (Table 2 and 3). The reason for higher returns and BCR with S₁ (1.2 Epan; IW/CPE) could be attributed to favorable soil moisture conditions maintained throughout the crop growth period which enhanced the photosynthetic rate, biomass accumulation, and partition into economic parts. The lowest returns and BCR under S₃ (0.6 Epan; IW/CPE) might be [due to the fact that because](#) moisture was not sufficient for [the](#) crop to absorb the nutrients efficiently, as water is [a](#) medium for nutrient absorption, which resulted in reduced leaf area, photosynthesis, biomass production and consequently lesser economic yield. Similar findings were also reported by Shivakumar *et al.* (2011), Sharan (2012), and Bibe *et al.* (2017).

3.1.3 Effects of Nitrogen levels

Among N levels significantly higher mean gross returns (₹. 1,27,180 and ₹.1,72,190 ha⁻¹), net returns (₹. 72,524 and ₹.1,21,650 ha⁻¹) and BCR (2.31 and 3.39) was observed with N₁ (100 [per cent](#) RDN) in both maize and groundnut crops and hence proved economically profitable over N₂ (75 [per cent](#) RDN) which resulted in comparatively lesser mean gross returns (₹. 1,15,756 and ₹.1,59,336 ha⁻¹), net returns (₹. 61,879 and ₹. 1,08,892 ha⁻¹) and BCR (2.14 and 3.14) (Table 2 and 3). The [mean%mean](#) increase with N₁ over N₂ was 9.87%, 17.2% & 8.23% in maize and 8.1%, 11.7%, and 7.9% in groundnut crop in [term terms](#) of mean gross returns, net returns and BCR. The application of 100 [per cent](#) RDN (N₁) was [in accordance withby](#) the [plants plants'](#) nutrient needs resulting in higher [economics economic](#) yields. The higher returns and BCR in N₁ [is are](#) attributed to [the](#) higher availability of N in the soil solution, which resulted in higher absorption and improved crop growth and ultimately final yield, which resulted in higher economic returns. Similarly, Fanish *et al.* (2011) and Sharan (2012) also reported higher economic returns under increased levels of fertilizers.

3.2 Interaction effects

The interaction effects of different treatments on [the](#) economics of maize and groundnut are depicted in ([Table Tables](#) 4, 5, 6, and 7). The interaction between irrigation systems and irrigation levels (M x S), irrigation systems and N levels (M x N), irrigation levels and N levels (S x N), and the interaction effect within irrigation systems, irrigation levels and N levels (M x S x N) was significant in maize crop while in groundnut all the interactions were non-significant except for the interaction between irrigation systems and irrigation levels (M x S) during both the years.

3.2.1 Interaction between different irrigation systems and irrigation levels

Whenever maize and groundnut crops are grown with limited water specific methodologies/systems shall be adopted to ensure maximum returns. In [the](#) current study in case of interaction between [the](#) irrigation system and irrigation levels (M x S), the sub-surface drip irrigation levels with either 1.2 or 0.9 Epan (M₂S₁ and M₂S₂) resulted in higher gross returns, net returns, and BCR followed by surface drip irrigation levels with 1.2 or 0.9 Epan (M₁S₁ and M₁S₂) over surface flood irrigation with 1.2, 0.9 or 0.6 IW/CPE ratios (M₃S₁, M₃S₂ or M₃S₃) (Table 4). The mean gross returns (₹. 1,51,594 and ₹. 2,13,539 ha⁻¹), net returns (₹. 95,760 and ₹. 1,61,479 ha⁻¹) & BCR (2.71 and 4.10) were significantly higher in sub-surface drip irrigation system with 1.2 Epan (S₁) and decreased in order: M₂S₂ > M₁S₁ > M₁S₂ > M₃S₁ > M₂S₃ > M₃S₂ > M₁S₃ >

M₃S₃ in both maize and groundnut crops. An overall 6.3, 15.2, 53.8, 1.4, 25.7, 25.6, 36.4 & 125.5% increased mean gross returns and 10.0, 25.5, 19.0, 2.04, 46.3, 41.2, 61.9 & 510.0% increased mean net returns were observed in maize crop whereas in groundnut crop an overall 5.03, 6.4, 72.5, 3.7, 37.9, 39.4, 48.9 & 129.1 % increased mean gross returns and 6.5, 8.1, 121.9, 4.8, 56.1, 55.1, 69.7 & 255.5% increased mean net returns were recorded with M₂S₁ over M₁S₁, M₁S₂, M₁S₃, M₂S₂, M₂S₃, M₃S₁, M₃S₂ and M₃S₃ respectively. Similarly, M₂S₂ registered 4.8, 13.6, 51.6, 23.9, 23.7, 34.4 and 122.2% increased mean gross returns and 7.8, 23.0, 14.5, 43.4, 38.3, 58.7 and 497.7% increased net returns in maize and 1.3, 2.6, 66.3, 33.0, 34.4, 43.6 & 120.9% increased mean gross returns and 1.7, 3.2, 111.8, 49.0, 48.0, 61.9 & 239.3 % increased net returns in groundnut over M₁S₁, M₁S₂, M₁S₃, M₂S₃, M₃S₁, M₃S₂ and M₃S₃ respectively. The maximum returns and BCR returns in sub-surface drip irrigation ~~system-systems~~ with 1.2 and 0.9 Epan ~~is-are~~ because of maximum economic yield under ~~the~~ same irrigation system and levels. These findings acquire reasonable support from Aladakatti *et al.* (2012) who reported better monetary returns (₹. 34,025 ha⁻¹) and BCR (1.93) under drip irrigation at 1.0 Epan over surface irrigation system (₹. 29,556 ha⁻¹ and 1.79).

3.2.2 Interaction between ~~the different irrigation system-systems~~ and N levels

The interaction effects of irrigation systems and N levels were non-significant in groundnut ~~crop-crops~~ however in maize ~~crop-crops~~ a significant interaction was observed. Across irrigation systems and N levels, the M₂N₁ had significantly higher mean gross returns (₹. 1,46,852 ha⁻¹), net returns (₹. 90,960 ha⁻¹) and BCR (2.62) whereas M₃N₂ recorded lowermost mean gross returns (₹. 97,052 ha⁻¹), net returns (₹. 45,305 ha⁻¹) and BCR (1.87) (Table 5). An overall 10.9, 26.5, 9.45, 43.5 and 51.3% increased mean gross returns and 18.4, 48.4, 15.1, 82.6 and 100.7% mean net returns ~~was-were~~ observed with M₂N₁ over M₁N₁, M₁N₂, M₂N₂, M₃N₁ and M₃N₂ respectively while M₂N₂ registered 1.3, 15.6, 31.1 and 38.2% increased mean gross returns and 2.9, 29.0, 58.7 and 74.5% increased net returns over M₁N₁, M₁N₂, M₃N₁ and M₃N₂ respectively. Similar results were also reported by Selvakumar, (2006) who observed that drip irrigation at 1.0 Epan recorded ~~an additional net returns additional net returns~~ (₹. 1,23,679) and BCR (3:30) in chilli followed by 0.8 Epan with 100% RDF registering an additional net return of (₹. 1,19,488) and BCR over surface irrigation system. These results findings are in further accordance ~~to-with~~ the findings of Senthilkumar (2000), Ramaprabha Nalini (1999) in groundnut and Suresh Kumar (2000) in capsicum under ~~a~~ drip irrigation system.

3.2.3 Interaction between different irrigation levels and N levels

The interaction effects of irrigation and N levels were non-significant in groundnut ~~crop-crops~~ however in maize ~~crop-crops~~ the interaction effects between irrigation levels and N levels showed that S₁N₁ (1.2 Epan; IW/CPE with 100 ~~per-centpercent~~ RDN) resulted in significantly higher gross returns (₹. 1,44,385 ha⁻¹), net returns (₹. 89,256 ha⁻¹) and BCR (2.61) followed by S₂N₁ which obtained ₹. 1,38,194 ha⁻¹ gross returns, ₹. 83,564 ha⁻¹ net returns and 2.52 BCR (Table 6). The S₁N₁ registered 9.2, 4.5, 17.2, 45.9 & 75.1% increased mean gross returns and 14.7, 6.8, 28.8, 99.5 & 131.9% increased mean net returns over S₁N₂, S₂N₁, S₂N₂, S₃N₁ and S₃N₂ respectively while S₂N₁ registered 4.5, 12.2, 39.6 & 50.3% increased mean gross returns and 7.4, 20.6, 86.7 & 117.1% increased net returns over S₁N₂, S₂N₂, S₃N₁ and S₃N₂ respectively. Similar results were also observed by Sharan (2012) who recorded significantly higher net returns and BCR with drip irrigation level with 1.2 Epan (₹. 57,266 ha⁻¹ and 2.6) followed

by 1.0 Epan (₹. 54,331 ha⁻¹ and 2.6) and 0.8 Epan (₹.39655 ha⁻¹ and 2.2) and lowest with surface irrigation level with 1.0 IW/CPE. (₹.19489 ha⁻¹ and 1.5). These findings are in further accordance ~~to~~ with the findings of Vishwanatha *et al.* (2000) and Ramah *et al.* (2010).

3.2.4 Interaction between different irrigation systems, irrigation levels, and N levels

The interaction effects of irrigation systems (M), irrigation levels (S) and N levels (N) showed a significant impact on the economics of maize and revealed that sub-surface irrigation system (M₂) with 1.2 Epan (S₁) and 100 ~~per cent~~ percent RDN (N₁) recorded significantly higher mean gross returns (₹. 1,58,749 ha⁻¹), net returns (₹. 1,10,243 ha⁻¹) and BCR (2.82) followed by M₂S₂N₁ which resulted in ₹. 1,56,106 ha⁻¹, ₹. 1,14,542 ha⁻¹ and 2.79 mean gross returns, net returns, and BCR. In later treatments M₁S₁N₁ (surface drip irrigation system with 1.2 Epan and 100 ~~per cent~~ percent RDN) and M₁S₂N₁ (surface drip irrigation system with 0.9 Epan and 100 ~~per cent~~ percent RDN) resulted in comparatively higher gross returns (₹. 1,49,705 and 1,45,378 ha⁻¹), net returns (₹. 93,820 and 89,792 ha⁻¹) and BCR (2.68 and 2.61), however, M₁S₂N₁ remained statistically at par with M₂S₁N₂ (sub-surface drip irrigation system with 1.2 Epan and 75 ~~per cent~~ percent RDN) during both the years (2021-22 and 2022-23) whereas the surface irrigation system with 0.6 IW/CPE and 75 ~~per cent~~ percent RDN (M₃S₃N₂) recorded least gross returns (₹. 65,253 ha⁻¹), net returns (₹. 14,106 ha⁻¹) and BCR (1.28) thus making it the least remunerative among all treatment combinations studied (Table 7). Higher gross returns, net returns, and BCR in the current study were due to higher crop yields, and efficient use of fertilizers & irrigation water under a sub-surface irrigation system (M₂). These findings are also supported by Shruti and Aladakatti (2017) who observed higher net returns (₹. 134262 ha⁻¹) and improved BCR (3.25) under drip fertigation combination of 1.0 Epan with 100 ~~per cent~~ percent RDNK over other ~~treatment treatments~~ combinations such as 1.0 Epan with 75 ~~per cent~~ percent RDNK.

4. CONCLUSION

~~On the basis of~~ Based on the results obtained in the present investigation it can be concluded that among different irrigation systems with varied irrigation and N levels, the implementation of a sub-surface drip irrigation system, 1.2 & 0.9 Epan ratios and 100 ~~per cent~~ percent RDN could result in ~~significantly significant~~ higher mean gross returns, net returns, and BCR in both maize and groundnut crops while satisfactory economics returns couldn't be achieved with surface irrigation system with 0.9 and 0.6 IW/CPE ratios and 75 ~~per cent~~ percent RDN in both crops.

Net return = Total Revenue – Total cost (Total Variable Cost)

Where is your Total cost to earn Net Return? Total Variable cost is costs include the labor cost, land rent cost, fertilizer cost, seed cost, oxen or tractor cost, etc please include the variable costs to get a net return. output did not get without incurring the cost(employing inputs)

Table 2. Effects of irrigation systems, irrigation, and N levels on [the economies profitability](#) of maize (2021-22 and 2022-23)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost-cost-benefit-cost ratio (BCR)		
	2021-22	2022-23	Mean	2021-22	2022-23	Mean	2021-22	2022-23	Mean
Main plots (M - Irrigation systems)									
M ₁ (Surface drip irrigation)	116347	132058	124202	62659	75421	69040	2.16	2.33	2.25
M ₂ (Sub-surface drip irrigation)	128364	152651	140507	74337	95674	85006	2.37	2.68	2.53
M ₃ (Surface furrow irrigation)	94585	104805	99695	43923	51193	47558	1.86	1.95	1.91
S. Em±	1066	336	-	1066	336	-	0.020	0.006	-
C.D. (P = 0.05)	4186	1319	-	4186	1319	-	0.078	0.025	-
Subplots (S - Irrigation levels)									
S ₁ (1.2 Epan; IW/CPE)	128023	148554	138288	74758	92340	83549	2.40	2.64	2.52
S ₂ (0.9 Epan; IW/CPE)	120598	140751	130674	67834	85036	76435	2.28	2.52	2.40
S ₃ (0.6 Epan; IW/CPE)	90674	100209	95442	38327	44912	41619	1.72	1.80	1.76
S. Em±	1613	1000	-	1613	1000	-	0.030	0.018	-
C.D. (P = 0.05)	4972	3082	-	4972	3082	-	0.092	0.055	-
Sub-subplots (N - Nitrogen levels)									
N ₁ (100 per cent RDN)	118356	136005	127180	65174	79873	72524	2.22	2.41	2.31
N ₂ (75 per cent RDN)	107841	123671	115756	55439	68319	61879	2.05	2.23	2.14
S. Em±	761	764	-	761	764	-	0.014	0.013	-
C.D. (P = 0.05)	2260	2271	-	2260	2271	-	0.042	0.040	-

Table 3. Effects of irrigation systems, irrigation, and N levels on [the economicsprofitability](#) of groundnut (2022 and 2023)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost ratio (BCR)		
	2022	2023	Mean	2022	2023	Mean	2022	2023	Mean
Main plots (M - Irrigation systems)									
M ₁ (Surface drip irrigation)	174885	177000	175943	124872	124238	124555	3.49	3.35	3.42
M ₂ (Sub-surface drip irrigation)	190969	191849	191409	140618	138747	139682	3.79	3.61	3.70
M ₃ (Surface flood irrigation)	128212	131663	129937	81225	81926	81575	2.72	2.64	2.68
S. Em±	1787	3150	-	1787	3150	-	0.036	0.062	-
C.D. (P = 0.05)	7017	12368	-	7017	12368	-	0.141	0.243	-
Subplots (S - Irrigation levels)									
S ₁ (1.2 Epan; IW/CPE)	189425	190627	190026	139836	138288	139062	3.81	3.63	3.72
S ₂ (0.9 Epan; IW/CPE)	182463	184210	183337	133374	132371	132872	3.70	3.54	3.62
S ₃ (0.6 Epan; IW/CPE)	122177	125675	123926	73505	74252	73878	2.50	2.43	2.46
S. Em±	1572	2284	-	1572	2284	-	0.031	0.045	-
C.D. (P = 0.05)	4845	7040	-	4845	7040	-	0.097	0.139	-
Sub-subplots (N - Nitrogen levels)									
N ₁ (100 per cent RDN)	171598	172782	172190	122433	120867	121650	3.47	3.31	3.39
N ₂ (75 per cent RDN)	157780	160893	159336	108711	109074	108892	3.20	3.09	3.14
S. Em±	1319	1684	-	1319	1684	-	0.027	0.033	-
C.D. (P = 0.05)	3919	5005	-	3919	5005	-	0.080	0.098	-

Table 4. Interaction effects of irrigation systems and irrigation levels on [the profitability economics](#) of maize and groundnut (2021-22 and 2022-23)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost ratio (BCR)		
	2021-22	2022-23	Mean	2021-22	2022-23	Mean	2021-22	2022-23	Mean
Maize									
M ₁ S ₁	131020	154096	142558	76999	97125	87062	2.42	2.70	2.56
M ₁ S ₂	121932	141096	131514	68212	84425	76318	2.27	2.49	2.38
M ₁ S ₃	96088	100982	98535	42767	44711	43739	1.80	1.79	1.80
M ₂ S ₁	139071	164118	151594	84711	106808	95760	2.56	2.86	2.71
M ₂ S ₂	134102	164641	149371	80042	107631	93837	2.48	2.89	2.68
M ₂ S ₃	111917	129194	120556	58258	72584	65421	2.09	2.28	2.18
M ₃ S ₁	113977	127450	120713	62565	73088	67826	2.22	2.34	2.28
M ₃ S ₂	105760	116515	111137	55248	63053	59150	2.09	2.18	2.14
M ₃ S ₃	64018	70452	67235	13956	17440	15698	1.28	1.33	1.30
Irrigation systems means at the same or different levels of irrigation levels.									
S. Em±	2403	1434	-	2403	1434	-	0.045	0.026	-
C.D. (P = 0.05)	7769	4485	-	7769	4485	-	0.144	0.081	-
Irrigation levels means-mean at the same or different levels of irrigation systems.									
S. Em±	2795	1732	-	2795	1732	-	0.052	0.031	-
C.D. (P = 0.05)	8612	5339	-	8612	5339	-	0.160	0.096	-
Groundnut									
M ₁ S ₁	202986	203622	203304	152640	150526	151583	4.03	3.83	3.93
M ₁ S ₂	199460	202010	200735	149414	149214	149314	3.99	3.83	3.91
M ₁ S ₃	122209	125369	123789	72563	72973	72768	2.46	2.39	2.43
M ₂ S ₁	213052	214026	213539	162367	160591	161479	4.20	4.01	4.10
M ₂ S ₂	206035	205745	205890	155650	152610	154130	4.09	3.87	3.98
M ₂ S ₃	153821	155776	154798	103836	103041	103438	3.08	2.95	3.02
M ₃ S ₁	152237	154235	153236	104500	103748	104124	3.19	3.05	3.12
M ₃ S ₂	141896	144876	143386	95059	95289	95174	3.03	2.92	2.98
M ₃ S ₃	90502	95880	93191	44115	46743	45429	1.95	1.95	1.95
Irrigation systems means at the same or different levels of irrigation levels.									
S. Em±	2558	3924	-	2558	3924	-	0.051	0.077	-
C.D. (P = 0.05)	8729	13708	-	8729	13708	-	0.175	0.270	-
Irrigation levels means at the same or different levels of irrigation systems.									
S. Em±	2723	3957	-	2723	3957	-	0.054	0.078	-
C.D. (P = 0.05)	8391	12193	-	8391	12193	-	0.168	0.241	-

Table 5. Interaction effects of irrigation systems and N levels on [the profitability economics](#) of maize (2021-22 and 2022-23)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost ratio (BCR)		
	2021-22	2022-23	Mean	2021-22	2022-23	Mean	2021-22	2022-23	Mean
M ₁ N ₁	123409	141294	132351	69331	84267	76799	2.28	2.48	2.38
M ₁ N ₂	109285	122822	116053	55987	66574	61281	2.05	2.18	2.12
M ₂ N ₁	135139	158565	146852	80722	101198	90960	2.48	2.76	2.62
M ₂ N ₂	121589	146737	134163	67952	90150	79051	2.27	2.59	2.43
M ₃ N ₁	96519	108157	102338	45467	54156	49812	1.89	2.00	1.94
M ₃ N ₂	92650	101453	97052	42378	48231	45305	1.84	1.90	1.87
Irrigation systems means at the same or different levels of nitrogen levels.									
S.Em±	1416	995	-	1416	995	-	0.026	0.018	-
C.D. (P = 0.05)	4974	3063	-	4974	3063	-	0.093	0.055	-
Nitrogen levels means-mean at the same or different levels of irrigation systems.									
S.Em±	1317	1324	-	1317	1324	-	0.025	0.023	-
C.D. (P = 0.05)	3915	3933	-	3915	3933	-	0.073	0.069	-

Table 6. Interaction effects of irrigation and N levels on [the profitability economics](#) of maize (2021-22 and 2022-23)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost ratio (BCR)		
	2021-22	2022-23	Mean	2021-22	2022-23	Mean	2021-22	2022-23	Mean
S ₁ N ₁	135012	153759	144385	81357	97155	89256	2.51	2.71	2.61
S ₁ N ₂	121033	143350	132192	68159	87526	77843	2.29	2.56	2.42
S ₂ N ₁	127238	149149	138194	74084	93045	83564	2.39	2.65	2.52
S ₂ N ₂	113958	132352	123155	61584	77028	69306	2.17	2.39	2.28
S ₃ N ₁	92817	105109	98963	40079	49421	44750	1.75	1.88	1.81
S ₃ N ₂	88532	95310	91921	36574	40402	38488	1.69	1.73	1.71
Irrigation levels means-mean at the same or different levels of nitrogen levels.									
S. Em±	1863	1370	-	1863	1370	-	0.035	0.024	-
C.D. (P = 0.05)	5690	4151	-	5690	4151	-	0.106	0.074	-
Nitrogen levels means-mean at the same or different levels of irrigation levels.									
S. Em±	1317	1324	-	1317	1324	-	0.025	0.023	-
C.D. (P = 0.05)	3915	3933	-	3915	3933	-	0.073	0.069	-

Table 7. Interaction effects of irrigation systems, irrigation levels, and N levels on the economics of maize (2021-22 and 2022-23)

Treatment	Gross returns (₹. ha ⁻¹)			Net returns (₹. ha ⁻¹)			Benefit-cost ratio (BCR)		
	2021-22	2022-23	Mean	2021-22	2022-23	Mean	2021-22	2022-23	Mean
M ₁ S ₁ N ₁	138438	160973	149705	84027	103612	93820	2.54	2.81	2.68
M ₁ S ₁ N ₂	123601	147219	135410	69970	90638	80304	2.30	2.60	2.45
M ₁ S ₂ N ₁	133970	156786	145378	79859	99725	89792	2.48	2.75	2.61
M ₁ S ₂ N ₂	109895	125407	117651	56564	69126	62845	2.06	2.23	2.14
M ₁ S ₃ N ₁	97819	106124	101971	44108	49463	46785	1.82	1.87	1.85
M ₁ S ₃ N ₂	94358	95840	95099	41427	39959	40693	1.78	1.72	1.75
M ₂ S ₁ N ₁	149554	167943	158749	94804	110243	102524	2.73	2.91	2.82
M ₂ S ₁ N ₂	128588	160292	144440	74619	103372	88995	2.38	2.82	2.60
M ₂ S ₂ N ₁	140269	171942	156106	85820	114542	100181	2.58	3.00	2.79
M ₂ S ₂ N ₂	127935	157339	142637	74265	100719	87492	2.38	2.78	2.58
M ₂ S ₃ N ₁	115592	135809	125701	61542	78809	70176	2.14	2.38	2.26
M ₂ S ₃ N ₂	108243	122580	115411	54973	66360	60666	2.03	2.18	2.11
M ₃ S ₁ N ₁	117042	132360	124701	65241	77609	71425	2.26	2.42	2.34
M ₃ S ₁ N ₂	110911	122539	116725	59889	68567	64228	2.17	2.27	2.22
M ₃ S ₂ N ₁	107476	118719	113097	56574	64867	60720	2.11	2.20	2.16
M ₃ S ₂ N ₂	104044	114311	109177	53922	61239	57580	2.08	2.15	2.11
M ₃ S ₃ N ₁	65040	73393	69216	14588	19991	17290	1.29	1.37	1.33
M ₃ S ₃ N ₂	62996	67510	65253	13324	14888	14106	1.27	1.28	1.28
Nitrogen levels means-mean at the same combination of irrigation systems and irrigation levels									
S. Em±	2282	2293	-	2282	2293	-	0.043	0.040	-
C.D. (P = 0.05)	6780	6813	-	6780	6813	-	0.127	0.120	-
Irrigation levels means-mean at the same combination of irrigation system and levels of nitrogen									
S. Em±	3227	2373	-	3227	2373	-	0.060	0.042	-
C.D. (P = 0.05)	9855	7190	-	9855	7190	-	0.183	0.128	-
Irrigation systems means at the same combination of irrigation levels and levels of nitrogen									
S. Em±	2991	2178	-	2991	2178	-	0.056	0.039	-
C.D. (P = 0.05)	9442	6622	-	9442	6622	-	0.176	0.118	-

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Aladakatti, Y.R., Hallikeri, S.S., Nandagavi, R.A., Shivamurthy, D., & Malik Rehan. (2012). Precision irrigation and fertigation to enhance the productivity and economic returns of bt cotton in vertisols. *Proceedings of Agro-Informatics and Precision Agriculture*, 341-343.
2. Bharathraj, H.R., Joshi, M., & Vishaka, G.V. (2015). Effect of surface fertigation on nutrient uptake, fertilizer use efficiency, and economics of interspecific hybrid Bt cotton. *Universal Journal of Agricultural Research*, 3, 46–48.
3. Bibe, S.M., Jadhav, K.T., & Chavan, A.S. (2017). Response of irrigation and fertigation management on growth and yield of maize. *International Journal of Current Microbiology and Applied Science*, 6(11), 4054-4060.
4. Buerkert, A., Bagayoko, M., Alvey, S., & Bationo, A. (2001). Causes of legume rotation effects in increasing cereal yields across the Sudanian, Sahelian, and Guinean ~~zone-zones~~ of West Africa. In: *Plant nutrition, food security, and sustainability of agro-ecosystems through basic and applied research. Developments in Plant and Soil Science. edited Edited-Edited* by Horst, W. 972–973. Kluwer Academic Publishers, Dordrecht, The Netherlands.
5. ~~Fanish~~Finish, S.A, Muthukrishnan, P., & Santhi, P. (2011). Effect of drip fertigation on field crops- A review. *Agricultural Review*, 32 (1), 14-25.
6. Jain, N.K., Yadav, R.S., & Jat, R.A. (2021). Drip fertigation influences the yield, nutrient uptake, and soil properties of ~~peanut-peanuts~~ (*Arachis hypogaea*). *Indian Journal of Agricultural Sciences*, 91(2), 258–262
7. Jat, R.A., Wani, S.P., Sahrawat, K.L., Singh, P., & Dhaka. B. L. (2011). Fertigation in vegetable crops for higher productivity and resource use efficiency. *Indian Journal of Fertilisers*, 7(3), 22–37.
8. Joshi, M., Bharath-Raj, H.R., & Vishaka, G.V. (2015). Effect of surface fertigation on nutrient uptake, fertilizer use efficiency, and economics of inter-specific hybrid Bt Cotton. *Universal Journal of Agricultural Research*, 3(3), 46-48.
9. Ministry of Statistics and Programme Implementation. (2021). Situation assessment of agricultural households and holdings of households in rural India, 2019. Ministry of Statistics and Programme Implementation. Government of India.
10. Murdia, L.K., Wadhwani, R., Wadhawan, N., Bajpai, P., & Shekhawat, S. (2016). Maize Utilization in India: An Overview. *American Journal of Food and Nutrition*, 4(6), 169-176.
11. Pawar, N., Bishnoi, D.K., Singh, M., & Dhillon, A. (2015). Comparative economic analysis of drip irrigation vis-a-vis flood irrigation system on productivity of Bt. ~~cotton-Cotton~~ in Haryana. *Agricultural Science Digest-A Research Journal*, 35(4), 300-303.
12. Ramah, K., Santhi, P., & Ponnuswamy, K. (2010). Economic viability of drip fertigation in maize (*Zea mays*) based cropping system. *Madras Agricultural Journal*, 97(1-3), 12-16.

13. Ramaprabha N.R. (1999) Influence of micro sprinkler irrigation on the performance of groundnut. M. Sc. (Ag.) Thesis. Tamil Nadu Agricultural University Coimbatore, India.
14. Saini S., Khatri, P., & Kumari, R.V. (2023). Agricultural Transformation in Telangana: Understanding Drivers of Growth and Planning Ahead. Arcus Research Report 3, New Delhi.
15. Selvakumar, T. (2006). Performance evaluation of drip fertigation on growth, yield, and water use in hybrid ~~chilli~~chili-chili (*Capsicum annuum* L.). PhD Thesis. Tamil Nadu Agricultural University, Coimbatore, India.
16. Senthilkumar. 2000. Effect of ~~micro-sprinkler~~micro-sprinkler irrigation and fertigation on yield and quality of groundnut. M.Sc. (Ag.) Thesis. Tamil Nadu Agricultural University, Coimbatore, India.
17. Sharan, B. (2012). Performance of sweet corn hybrid under different levels of irrigation and nitrogen applied through the drip system. M.Sc. (Ag.) Thesis. Acharya N G Ranga Agricultural University, Hyderabad, India.
18. Shivakumar, H.K., Ramachandrappa, B.K., Nanjappa, H.V., & Mudulagiriappa. (2011). Effect of ~~phenophase-based~~phenophase-based irrigation schedules on growth, yield and quality of baby corn (*Zea mays* L.). Agricultural Sciences, 2 (3), 267-272.
19. Shruti, M.Y., & Aladakatti, Y.R. (2017). Effect of drip irrigation and fertigation on yield, economics and water use efficiency of intra-hirsutum Bt cotton. Journal of Farm Science, 30, 185–189.
20. Sivanappan, R.K. (1978). Economics of drip irrigation method in small and marginal farms. Madras Agricultural Journal, 65(12), 809-813.
21. Sorensen, R.B., & M.C. Lamb. (2009). Peanut yield, market grade, and economics with two surface drip lateral spacings. Peanut Science, 36(1), 85–91.
22. Sreekanth, M., Hakeem, A.H., Quadri, J.A.P., & Irfath, R. (2017). Low productivity of Indian agriculture with special reference ~~on-to~~on-to cereals. Journal of Pharmacognosy and Phytochemistry, 6(5), 239-243.
23. Suresh Kumar, P. (2000). Performance evaluation of drip fertigation system with ~~water-solubles~~soluble soluble soluble water-soluble fertilizers on water, fertilizer use and yield in hybrid capsicum. M.Sc. (Ag.) Thesis. Tamil Nadu Agricultural University, Coimbatore, India.
24. Vishwanatha, G.B., Ramachanrappa, B.K., & Nanjappa, H.V. (2000). Effect of drip irrigation and methods of planting on root and shoot biomass, tasseling-silking interval, yield and economics of sweet corn (*Zea mays* L.). Mysore Journal of Agricultural Sciences, 34, 134-141.
25. Yazar, A., Gokcel, F., & Sezen, M.S. (2009). Corn yield response to partial ~~root~~zone zone zone root-zone drying and deficit irrigation strategies applied with drip system. Plant and Soil Environment, 55(11), 494–503.