

# Production potential of diversified intercropping system in paired row rice in summer fallows

## ABSTRACT

Globally diversification of rice based cropping system builds ecosystem resilience and food security. A field experiment was carried out during summer season of 2023-24 at Integrated farming systems research station, Karamana, Thiruvananthapuram to study the production potential of diversified intercrops in paired row planted rice in summer fallows of double cropped lowland rice fields. The field experiment was laid out in a randomized, completely block design with 10 treatments and three replications. The treatments were T<sub>1</sub>: Paired row of Rice + Grain Cowpea, T<sub>2</sub>: Paired row of Rice + Finger millet, T<sub>3</sub>: Paired row of Rice + Foxtail millet, T<sub>4</sub>: Paired row of Rice + Proso millet, T<sub>5</sub>: Sole crop of Paired row Rice, T<sub>6</sub>: Sole crop of Grain Cowpea, T<sub>7</sub>: Sole crop of Finger millet, T<sub>8</sub>: Sole crop of Foxtail millet, T<sub>9</sub>: Sole crop of Proso millet, T<sub>10</sub>:Fallow. Among the treatments, the higher number of tillers m<sup>-2</sup> and leaf area index were in the sole crop of paired row rice (T<sub>5</sub>). Among the yield attributes, rice had the highest number of productive tillers m<sup>-2</sup>, grain yield and straw yield in the sole crop treatment(T<sub>5</sub>). Sole crop treatments performed best in terms of growth and yield attributes than their intercropping treatments. The highest yield (5.85 t ha<sup>-1</sup>) was observed in the sole crop of rice (T<sub>5</sub>), which was superior to all other treatments. The yields recorded in T<sub>1</sub> (paired row of rice + grain cowpea) and T<sub>4</sub> (paired row of rice + proso millet) were statistically at par and significantly higher than T<sub>3</sub> (paired row of rice + foxtail millet). Paired row of rice + grain cowpea (T<sub>1</sub>) recorded the highest monetary advantage index of ₹ 55820 ha<sup>-1</sup>.The analyzed data showed that T<sub>5</sub>(sole crop of paired row rice) resulted in an 8 to 15 per cent grain yield increase over intercropping treatments. It can be concluded from the study that among the intercrops, grain cowpea can be recommended for diversification in paired row planting of rice in summer fallows for higher productivity and profitability.

*Keywords: Cowpea, Finger millet, Foxtail millet, Intercropping, Proso millet, Paired row, Rice*

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is a staple food crop for billions of people worldwide and holds a prominent position among the food crops cultivated in India. In Kerala the predominant cropping system is rice- rice-fallow (John *et al.*, 2014). Crop diversification in the summer period is one of the ways to increase output and income (Varughese *et al.*, 2007). Intercropping is one of the promising methods of crop diversification in which two or more are grown together in the same field during the same growing season (Andrew and Kassam, 1976).

Growing rice using a paired row planting system with residual moisture can indeed provide additional income. However, the wider spacing between paired rows often leads to increased weed growth, necessitating labor-intensive intercultural operations for weed

management. An effective strategy to address this issue is to introduce intercrops between paired rows, which can help reduce weed infestation and enhance income per unit area. Studies, such as Shah *et al.* (2021), have shown that the addition of intercrops can significantly curb weed growth, creating a more efficient and profitable system for farmers.

Millets are increasingly recognized for their low water requirements and ability to thrive in arid and semi-arid conditions, which makes them suitable for intercropping in residual moisture settings, such as between rows of rice during summer. Their resilience against drought and high water efficiency enhances land productivity, while reducing the need for irrigation (Maitra *et al.*, 2021). This intercropping can also mitigate weed growth and improve soil health. In addition, leguminous crops like cowpea, with their nitrogen-fixing abilities, further support soil fertility and yield in mixed cropping systems, presenting an economically and ecologically promising approach (Bates, 2022).

## 2. MATERIALS AND METHODS

The experiment was conducted at Integrated Farming System Research Station (IFSRS), Karamana during summer season of 2023-2024. The experiment consisted of 10 treatments T<sub>1</sub>: Paired row of Rice + Grain Cowpea, T<sub>2</sub>: Paired row of Rice + Finger millet, T<sub>3</sub>: Paired row of Rice + Foxtail millet, T<sub>4</sub>: Paired row of Rice + Proso millet, T<sub>5</sub>: Sole crop of Paired row Rice, T<sub>6</sub>: Sole crop of Grain Cowpea, T<sub>7</sub>: Sole crop of Finger millet, T<sub>8</sub>: Sole crop of Foxtail millet, T<sub>9</sub>: Sole crop of Proso millet, T<sub>10</sub>: Fallow. The experiment was conducted in Randomized Complete Block Design with 3 replications. The varieties used for the experiment are Uma, Bhagyalakshmi, Co 15, Co (Te)7 and TNAU 202 for rice, grain cowpea, finger millet, foxtail millet and proso millet, respectively. The spacing followed for the crops were Rice – 35 cm-(15 cm x 15 cm)- 10 cm (35 cm between the pairs, 15 cm within the pair, 10 cm between plants), Grain cowpea – 30cm x15 cm, Finger millet -25 cm x 15 cm, Foxtail millet- 22 cm x 10 cm, Proso millet- 25 cm x 10 cm. The average temperature during the cropping period were 25.50° C (minimum) and 32.21°C (maximum), maximum relative humidity was 90.20% and minimum was 72.58% during the period with an average rainfall of 22.22 mm during the cropping period. The fertiliser recommendation used for the crops are given in the table below (Table 1).

## 3. RESULTS

### 3.1 Growth attributes

Intercropping with grain cowpea, finger millet, foxtail millet, and proso millet did not significantly affect the rice plant height (Table 2).

The analyzed data indicated that the plant height of intercrops were significantly influenced by the intercropping treatments. The sole crop of grain cowpea (T<sub>6</sub>) recorded the highest plant height of 75.29 cm at harvest. The sole crop of finger millet (T<sub>7</sub>) recorded the highest mean plant height, measuring 116.32 cm at harvest. The sole crop of foxtail millet recorded the highest plant height, measuring 133.73 cm at harvest. The analyzed data showed that the plant height of proso millet was highest in the sole crop of proso millet (T<sub>9</sub>), it recorded plant heights of 107.10 cm at harvest (Table 3).

The analysed data showed that the number of rice tillers was significantly influenced by intercropping treatments. At harvest, the highest number of tillers was observed in the sole crop of rice (432.3), which was on par with rice + grain cowpea (T<sub>1</sub>), rice + foxtail millet (T<sub>3</sub>) and rice + proso millet (T<sub>4</sub>) (Table 2).

Intercropping treatments had significant effect on number of tillers per m<sup>2</sup>. The highest number of tillers per m<sup>2</sup> (84.67) was recorded in the sole crop of finger millet (T<sub>7</sub>) at harvest compared to rice + finger millet (T<sub>2</sub>). The number of tillers per m<sup>2</sup> was highest (184.83) in sole foxtail millet (T<sub>8</sub>). A higher number of tillers per m<sup>2</sup> was in the sole crop of proso millet (T<sub>9</sub>) at harvest (98.80) (Table 3).

The results indicated that intercropping treatments had significant effect on the leaf area index (LAI) of rice. At harvest, the highest LAI (3.23) was observed in the sole crop of rice (T<sub>5</sub>), on par with T<sub>1</sub> and T<sub>3</sub>. The lowest LAI was recorded in T<sub>2</sub>, which was statistically similar to T<sub>4</sub> (Table 2).

The analyzed data indicated that intercropping treatments significantly influenced the LAI of intercrops. The highest LAI (3.70) was recorded in the sole crop of grain cowpea (T<sub>6</sub>). The highest LAI (1.48) values were recorded in the sole crop of finger millet at harvest. The highest LAI values were recorded in the sole crop of foxtail millet (T<sub>8</sub>) at harvest (0.97). The results indicated that the sole crop of proso millet recorded the highest LAI value of 1.09 at harvest (Table 3).

The analyzed data revealed that the number of branches per plant of grain cow pea was not influenced by intercropping treatments (Table 3).

Table 2. Growth parameters of rice at harvest stage

Treatments	At Harvest		
	Plant height (cm)	Number of tillers per m <sup>2</sup>	LAI
T <sub>1</sub> : Rice + Grain Cowpea	113.67	428.7	3.03
T <sub>2</sub> : Rice + Finger millet	106.33	382	2.10
T <sub>3</sub> : Rice + Foxtail millet	110.33	425.3	2.63
T <sub>4</sub> : Rice + Proso millet	112.33	427.3	2.28
T <sub>5</sub> : Sole crop of Rice	115.43	432.3	3.23
SEm (±)	3.425	3.1	0.21
CD (0.05)	NS	9.93	0.682

Table 3. Effect of intercropping on growth attributes of intercrops

Treatments	At harvest			
	Plant height (cm)	Number of branches per plant	LAI	Number of tillers per m <sup>2</sup>
T <sub>1</sub> : Rice + Grain Cowpea	1.92	4.52	1.92	
T <sub>6</sub> : Sole crop of Grain Cowpea	3.70	5.98	3.70	
T value	-7.74	-2.22	-7.74	
P value	0.001	0.091	0.001	

T <sub>2</sub> : Rice + Finger millet	110.36	-	0.96	33.00
T <sub>7</sub> : Sole crop of Finger millet	116.32	-	1.48	84.67
T value	-3.11	-	-3.18	-5.62
P value	0.036	-	0.033	0.005
T <sub>3</sub> : Rice + Foxtail millet	116.75	-	0.25	69.33
T <sub>8</sub> : Sole crop of Foxtail millet	133.73	-	0.97	184.83
T value	-4.67	-	-8.13	-11.66
P value	0.010	-	0.001	0.000
T <sub>4</sub> : Rice + Proso millet	98.37	-	0.687	44.47
T <sub>9</sub> : Sole crop of Proso millet	107.10	-	1.09	98.80
T value	-5.34	-	-8.17	-32.22
P value	0.006	-	0.001	0.000

### 3.2 Yield attributes

Intercropping had significant effects on number of productive tillers per m<sup>2</sup> of rice. The highest number of productive tillers per m<sup>2</sup> (409.7) was observed in the sole crop of rice (T<sub>5</sub>), which was statistically on par with T<sub>4</sub> (paired row of rice + proso millet), T<sub>3</sub> (paired row of rice + foxtail millet), and T<sub>1</sub> (paired row of rice + grain cowpea). In contrast, the lowest number of productive tillers (357) at harvest was recorded in T<sub>2</sub> (paired row of rice + finger millet) (Table 4).

The results indicated that the highest number of productive tillers per plant (3.03) was recorded in the sole crop of finger millet (T<sub>7</sub>), while the lowest (1.87) was observed in the intercropping treatment (T<sub>2</sub>). Similarly, the sole crop of foxtail millet (T<sub>8</sub>) exhibited a significantly higher number of productive tillers per plant (1.57) compared to intercropping. For proso millet, the sole crop (T<sub>9</sub>) recorded a higher number of productive tillers per plant (2.26) than the intercropped treatments (Table 5).

Intercropping had significant effect on number of pods per plant of grain cowpea. The analyzed data revealed that the sole crop of grain cowpea (T<sub>6</sub>) recorded the highest number of pods per plant (15.87) as compared to intercropping (Table 5).

A thorough analysis of the data indicated that the grain yield of rice was significantly affected by the treatments. The highest yield (5.85 t ha<sup>-1</sup>) was observed in the sole crop of rice (T<sub>5</sub>), which was superior to all other treatments. The yields recorded in T<sub>1</sub> (paired row of rice + grain cowpea) and T<sub>4</sub> (paired row of rice + proso millet) were statistically at par and significantly higher than T<sub>3</sub> (paired row of rice + foxtail millet). The lowest yield (4.97 t ha<sup>-1</sup>) was recorded in T<sub>2</sub> (paired row of rice + finger millet) (Table 4).

Results indicated that intercropping had a significant effect on grain yield of the intercrops. The grain yield was highest (3375 kg ha<sup>-1</sup>) in sole crop of grain cowpea (T<sub>6</sub>) compared to rice + grain cowpea (T<sub>1</sub>). The highest grain yield of finger millet (2223 kg ha<sup>-1</sup>) was recorded in its sole crop (T<sub>7</sub>), while the intercropping treatment of rice + finger millet (T<sub>2</sub>) yielded 666 kg ha<sup>-1</sup>. Similarly, the highest grain yield of foxtail millet (2352 kg ha<sup>-1</sup>) was observed in its sole crop (T<sub>8</sub>), whereas the intercropping treatment of rice + foxtail millet (T<sub>3</sub>) produced 642 kg ha<sup>-1</sup>. For proso millet, the sole crop (T<sub>9</sub>) recorded the highest grain yield

(949 kg ha<sup>-1</sup>), while the intercropping treatment of rice + proso millet (T<sub>4</sub>) yielded 269 kg ha<sup>-1</sup> (Table 5).

The analyzed data revealed that the straw yield of rice was significantly affected by the treatments. The highest straw yield (7.00 t ha<sup>-1</sup>) was recorded in the sole crop of paired row rice (T<sub>5</sub>), which was significantly superior to all intercropping treatments. Among the intercropping treatments, paired row rice + grain cowpea (T<sub>1</sub>) produced a straw yield of 6.61 t ha<sup>-1</sup>, followed by T<sub>4</sub> (paired row rice + proso millet) and T<sub>3</sub> (paired row rice + foxtail millet). The lowest straw yield (5.48 t ha<sup>-1</sup>) was observed in the paired row of rice + finger millet (T<sub>2</sub>) (Table 4).

Results on the haulm yield revealed that the highest haulm yield (4526 kg ha<sup>-1</sup>) was obtained in sole crop of grain cowpea (T<sub>6</sub>). The highest stover yield of finger millet (4927 kg ha<sup>-1</sup>) was recorded in its sole crop (T<sub>7</sub>), while the intercropping treatment of rice + finger millet (T<sub>2</sub>) produced 1666 kg ha<sup>-1</sup>. Similarly, the sole crop of foxtail millet (T<sub>8</sub>) recorded the highest stover yield (4067 kg ha<sup>-1</sup>), whereas the intercropping treatment of rice + foxtail millet (T<sub>3</sub>) yielded 1587 kg ha<sup>-1</sup>. For proso millet, the highest stover yield (2415 kg ha<sup>-1</sup>) was observed in its sole crop (T<sub>9</sub>), while the intercropping treatment of rice + proso millet (T<sub>4</sub>) yielded 904 kg ha<sup>-1</sup> (Table 5).

Table 4. Effect of intercropping on yield attributes of rice

Treatments	Productive tillers per m <sup>2</sup>	Grain yield (t ha <sup>-1</sup> )	Straw yield (t ha <sup>-1</sup> )	Harvest index
T <sub>1</sub> : Rice + Grain Cowpea	397.3	5.31	6.61	0.45
T <sub>2</sub> : Rice + Finger millet	357	4.97	5.48	0.48
T <sub>3</sub> : Rice + Foxtail millet	398.7	5.07	6.17	0.45
T <sub>4</sub> : Rice + Proso millet	401.7	5.25	6.3	0.46
T <sub>5</sub> : Sole crop of Rice	409.7	5.85	7.00	0.46
SEm (±)	4.2	0.02	0.04	0.00
CD (0.05)	13.72	0.066	0.14	0.006

Table 5. Effect of intercropping on yield attributes of intercrops

Treatments	At harvest				
	Number of pods per plant	Number of productive tillers per plant	Grain yield (kg ha <sup>-1</sup> )	Stover yield or Haulm yield (kg ha <sup>-1</sup> )	Harvest index
T <sub>1</sub> : Rice + Grain Cowpea	14.07	-	1122	2678	0.30
T <sub>6</sub> : Sole crop of Grain Cowpea	15.87	-	3375	4526	0.43
T value	-3.61	-	-16.78	-27.57	-15.06

P value	0.023	-	0.000	0.000	0.000
T <sub>2</sub> : Rice + Finger millet	-	1.87	666	1666	0.29
T <sub>7</sub> : Sole crop of Finger millet	-	3.03	2223	4927	0.31
T value	-	-3.26	-7.00	-6.18	-1.607
P value	-	0.031	0.002	0.003	0.183
T <sub>3</sub> : Rice + Foxtail millet	-	1.15	642	1587	0.29
T <sub>8</sub> : Sole crop of Foxtail millet	-	1.57	2352	4067	0.37
T value	-	-6.200	-23.83	-10.37	-3.35
P value	-	0.003	0.000	0.000	0.029
T <sub>4</sub> : Rice + Proso millet	-	1.98	269	904	0.23
T <sub>9</sub> : Sole crop of Proso millet	-	2.26	949	2415	0.28
T value	-	-11.05	-18.87	-48.17	-5.40
P value	-	0.000	0.000	0.000	0.006

### 3.3 Economics

The economic analysis revealed that the paired row of rice + grain cowpea (T<sub>1</sub>) generated the highest net income (₹129,686 ha<sup>-1</sup>), followed by the sole crop of grain cowpea (T<sub>6</sub>), which recorded net returns of ₹119,537 ha<sup>-1</sup>. In contrast, the lowest net returns were observed in the sole crop of proso millet (T<sub>9</sub>), with a negative value of ₹4,830 ha<sup>-1</sup>. The benefit-cost ratio (BCR) was highest (1.82) in the paired row of rice + grain cowpea (T<sub>1</sub>) intercropping system, followed by the sole crop of grain cowpea (T<sub>6</sub>) and the sole crop of paired row rice (T<sub>5</sub>), both recording a BCR of 1.79 (Table 6).

The monetary advantage index (MAI) was highest for the paired row of rice + grain cowpea (T<sub>1</sub>), recording ₹55,820 ha<sup>-1</sup>. This was followed by the paired row of rice + proso millet (T<sub>4</sub>) with ₹31,662 ha<sup>-1</sup>, paired row of rice + finger millet (T<sub>2</sub>) with ₹27,866 ha<sup>-1</sup>, and paired row of rice + foxtail millet (T<sub>3</sub>) with ₹27,073 ha<sup>-1</sup>, respectively (Table 6).

Table 6. Effect of intercropping on economics of the system

Treatments	Net income (₹ ha <sup>-1</sup> )	BCR	MAI (₹ ha <sup>-1</sup> )
T <sub>1</sub> : Paired row of Rice + Grain Cowpea	129686	1.82	55820
T <sub>2</sub> : Paired row of Rice + Finger millet	63097	1.42	27866
T <sub>3</sub> : Paired row of Rice + Foxtail millet	68246	1.45	27073
T <sub>4</sub> : Paired row of Rice + Proso millet	54122	1.36	31662
T <sub>5</sub> : Sole crop of Paired row Rice	93117	1.79	-

T <sub>6</sub> : Sole crop of Grain Cowpea	119537	1.79	-
T <sub>7</sub> : Sole crop of Finger millet	52686	1.73	-
T <sub>8</sub> : Sole crop of Foxtail millet	54181	1.75	-
T <sub>9</sub> : Sole crop of Proso millet	-4830	0.92	-
T <sub>10</sub> : Fallow	-	-	-

#### 4. DISCUSSION

The results revealed that the highest number of tillers (432.3 m<sup>2</sup>) was recorded in the sole crop of rice (T<sub>5</sub>). However, paired row planting with intercrops such as grain cowpea (T<sub>1</sub>), foxtail millet (T<sub>3</sub>), and proso millet (T<sub>4</sub>) produced tiller counts statistically on par with sole rice (Table 3). This suggested that paired row planting effectively accommodates intercrops while sustaining rice productivity. The reduced tiller count in T<sub>2</sub> (rice + finger millet) reflects higher competition between rice and finger millet in this treatment for nutrient and space.

Paired row planting in sole rice (T<sub>5</sub>) achieved the highest LAI (3.23), indicating optimal canopy development. Among intercropping treatments, rice + grain cowpea (T<sub>1</sub>) and rice + foxtail millet (T<sub>3</sub>) recorded comparable LAI values, reflecting effective utilization of sunlight and nutrients in these combinations. Intercropping rice with finger millet (T<sub>2</sub>) resulted in the lowest LAI (2.10), possibly due to increased competition for resources. For intercrops, sole cropping systems recorded higher LAI values than intercropped treatments, as seen in grain cowpea (3.70) and proso millet (1.09). This indicated that paired row planting supports intercrop growth while sole cropping remains more conducive to maximum canopy expansion.

Sole rice (T<sub>5</sub>) had the highest number of productive tillers (409.7 m<sup>2</sup>), followed closely by T<sub>4</sub> (rice + proso millet), T<sub>3</sub> (rice + foxtail millet), and T<sub>1</sub> (rice + grain cowpea). This indicates that paired row planting supports productive tiller formation, especially when paired with suitable intercrops. The lowest productive tillers in T<sub>2</sub> (357 m<sup>2</sup>) again highlight the competitive nature of finger millet in intercropping systems. For intercrops, sole cropping significantly outperformed intercropping, with finger millet and proso millet recording higher productive tiller counts in their respective sole treatments. Paired row planting offers distinct advantages for incorporating intercrops in rice-based systems. The wider spacing between rows reduces competition for light and enables efficient utilization of residual moisture. Additionally, the intercropping system suppresses weed growth, reducing the labour and cost associated with manual weeding. Rajani *et al.* (2022) recorded that sole foxtail millet had significantly higher grain weight per 0.5 m row length (99.40 g) in intercropping system with niger in different row ratios. According to Aravind *et al.* (2023), the grain yield of foxtail millet was significantly affected by intercropping. Sole cropping produced the highest grain yield (2208 kg ha<sup>-1</sup>), followed by intercropping with vegetable cowpea, green gram, and coriander in two different ratios. Saikumar *et al.* (2023) reported that aerobic rice

intercropped with black gram produced the highest number of spikelets per panicle (77.53). This result was on par with the sole crop of aerobic rice, black gram + aerobic rice with 100% RDN, black gram + aerobic rice with 50% RDN, and cowpea + aerobic rice with 100% RDN.

The highest grain yield for rice (5.85 t ha<sup>-1</sup>) was observed in the sole crop treatment (T<sub>5</sub>), reflecting the absence of competition. Paired row planting with grain cowpea (T<sub>1</sub>) and proso millet (T<sub>4</sub>) resulted in grain yields statistically superior to other intercropping treatments. This highlights the potential of these combinations to sustain rice productivity while diversifying farm income. Paired row planting of rice, combined with suitable intercrops such as grain cowpea and proso millet, demonstrates the potential to sustain rice productivity while diversifying cropping systems. This approach offers a balanced strategy to optimize resource use, suppress weeds, and enhance overall farm profitability in summer fallow conditions. Further refinement in crop combinations and management practices can enhance its viability and adoption.

Among intercrops, sole cropping yielded the highest outputs, with grain cowpea (3375 kg ha<sup>-1</sup>) and finger millet (2223 kg ha<sup>-1</sup>) outperforming their intercropped counterparts. This trend underscores the resource competition inherent in intercropping systems. Gibi (2023) conducted an experiment on red gram-cowpea intercropping and inferred that the number of pods per plant produced by bush cowpea was higher in sole crop of bush cowpea at (13.33) compared to intercropping treatments.

The straw yield of rice was highest in the sole crop (7.00 t ha<sup>-1</sup>), with paired row planting of rice + grain cowpea (T<sub>1</sub>) yielding 6.61 t ha<sup>-1</sup>. For intercrops, stover yields were significantly higher in sole cropping systems, with finger millet (4927 kg ha<sup>-1</sup>) and foxtail millet (4067 kg ha<sup>-1</sup>) leading.

Despite slightly lower yields compared to sole rice, intercropping system of rice + grain cowpea (T<sub>1</sub>) maintain productivity while enhancing ecological and economic benefits. These systems make efficient use of available land and resources, particularly in water-scarce summer fallows.

The economic analysis revealed that the paired row of rice + grain cowpea (T<sub>1</sub>) generated the highest net income (₹129,686 ha<sup>-1</sup>), followed by the sole crop of grain cowpea (T<sub>6</sub>), which recorded net returns of ₹119,537 ha<sup>-1</sup>. Erythrina *et al.* (2022) reported that intercropping resulted in a significantly higher gross margin compared to monocropping. Saikumar et al (2023) reported that the highest gross returns (Rs.180065 ha<sup>-1</sup>), net returns (Rs.103858 ha<sup>-1</sup>) and BCR (2.36) were recorded by cowpea + aerobic rice with 50% RDN.

## 5. CONCLUSION

The study highlights the benefits and limitations of intercropping in paired row rice systems during summer fallows. Sole cropping of rice achieved the highest grain and straw yields showed its potential for maximizing rice productivity. However, intercropping treatments, particularly rice + grain cowpea (T<sub>1</sub>), demonstrated comparable results in terms of productive tillers and overall yield, along with the highest economic returns. These findings suggest that paired row planting with suitable intercrops, such as grain cowpea, can balance productivity

with profitability while improving resource use and reducing weed pressure. This approach holds promise for sustainable cropping systems, especially in water-limited conditions, offering farmers a viable strategy to enhance income and maintain soil health.

## DISCLAIMER

I hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. ChatGPT
3. Grammarly
4. Quillbot

### Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## REFERENCES

1. John J, Rajasekharan P, Rajasree G, Bindu P. (2014). Cropping systems in Kerala. Thiruvananthapuram, Kerala: State Planning Board, 46.
2. Varughese K, Jacob J, Rani B, Vijayan M. (2007). Scope of crop diversification in paddy fields. In: Papers on special session: Paddy Cultivation in Kerala. 19th Kerala Science Congress; 2007 Jan 29–31; Kannur. Kerala State Council for Science, Technology and Government of Kerala, 59-70.
3. Andrews DJ, Kassam AH. (1976). The importance of multiple cropping in increasing world food supplies. *Multiple Cropping*, 27, 1–10.
4. Shah TM, Tasawwar S, Bhat MA, Otterpohl R. (2021). Intercropping in rice farming under the system of rice intensification—An agroecological strategy for weed control, better yield, increased returns, and social–ecological sustainability. *Agronomy*, 11(5), 1010.
5. Maitra S, Hossain A, Brestic M, Skalicky M, Ondrisik P, Gitari H et al. (2021). Intercropping—A low input agricultural strategy for food and environmental security. *Agronomy*, 11(2), 343.

6. Bates J. (2022). The fits and starts of Indian rice domestication: How the movement of rice across northwest India impacted domestication pathways and agricultural stories. *Frontiers in Ecology and Evolution*, 10, 924977.
7. Rajani CS, Patil SB, Kalaghatagi SB. (2022). Evaluation of productivity and profitability of millet and oilseed intercropping systems in the northern dry zone of Karnataka. *Journal of Farm Sciences*, 35(2), 185–91.
8. Aravind K, Vadivel N, Kumar GS, Ravichandran V, Bharathi C. (2023). Performance of foxtail millet-based intercropping system for improving the productivity, sustainability, and economics in Western Zone of Tamil Nadu under irrigated conditions. *International Journal of Plant & Soil Science*, 35(19), 1824–9.
9. Saikumar G, Jinsy VS, Sumesh KV. (2023). Agro-economic evaluation of aerobic rice + legume intercropping system under varying levels of nitrogen. *Mysore Journal of Agricultural Sciences*, 57 (3), 146-152.
10. Mariam G. (2023). *Performance of red gram (Cajanus cajan (L.) Millsp.) intercropping systems in lowlands* (Post graduate dissertation, Department of Agronomy, College of Agriculture, velayani).
11. Erythrina E, Susilawati S, Slameto S, Resiani NMD, Arianti FD, Jumakir J et. al. (2022). Yield advantage and economic performance of rice–maize, rice–soybean, and maize–soybean intercropping in rainfed areas of Western Indonesia with a wet climate. *Agronomy*, 12(10), 2326