

Original Research Article

CRISPR/Cas9 driven targeted editing of *GRAIN NUMBER 1a* gene: sgRNA constructs design for yield enhancement in rice (*Oryza sativa* L.)

ABSTRACT

Yield is the ultimate trait determined by various quantitative trait loci (QTLs). *GRAIN NUMBER 1a* (*GN1a*), encodes for an enzyme Cytokinin oxidase/ dehydrogenase (CKX2), negatively influences the yield by degrading the phytohormone cytokinin. Targeted genome editing of *GN1a* gene was performed to down-regulate the expression of *GN1a* using CRISPR/Cas9 for the increase in grain number and yield with improved cytokinin content in the panicle meristem. CRISPR/Cas9, a precise means for targeted editing of a gene for the improvement of a particular trait in plants. The single guide RNAs (sgRNAs) were designed using plant specific CRISPR-P v2.0 software. Two efficient sgRNAs were chosen critically entrenched on their GC content, on-target values, location on the gene, off-target sites and their location, secondary structures etc., adjacent to the Protospacer Adjacent Motif (PAM) NGG. The binary vector pRGEB32, with *BsaI* restriction site driven by rice U3 promoter was employed for cloning of sgRNAs and Cas9 by rice ubiquitin promoter. The sgRNAs were tempered, phosphorylated and astringed with the binary vector pRGEB32, transformed into *E.coli* DH5 α initially, then mobilized into *A.tumefaciens*EHA105. The present study helps in the development of elite lines which will lead to enhancement of grain number and overall yield for the growing population and farmer's welfare.

Keywords: Yield, Cytokinin, *GN1a* gene, Genome editing, CRISPR/Cas9.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is considered to be one of the most significant and primary cereal food crops, that provide sustenance for almost half of the world's population. Due to the continuous population growth in Asia, Africa, and Latin America, there will be a significant increase in the demand for rice. It is predicted that in order to meet the food demands of a growing global population, rice production will need to enhance proportionately. New superior varieties with unique plant structures which can produce more grain yields are required to meet this growing demand. An example of this can be seen in the "green revolution," where lodging-resistant semi-dwarf varieties of cereals like wheat and rice have significantly increased grain yields (Peng *et al.*, 1999).

Yield is the most complex and important physiological trait governed by various external (abiotic, biotic) and internal factors (genetic, biochemical). The yield of rice in the year 2023-2024 is 4.2 T/ha which was decreased from the year 2022-2023 which was 4.3 T/ha (<https://ipad.fas.usda.gov/updated> on April 11, 2024). Reduction in the yield can be caused by physiological, environmental, morphological bottlenecks which have huge impact on the growth and development of plant.

Cytokinins (CK) have a unique role in the growth and development of various [plant](#) organs. Plants developed precisely control in their concentration by taking benefit of biodegradation at spatial and temporal levels. [Eleven](#) CKX genes have been distinguished so far in rice; but, the majority of their tissue specific activities persist unknown (Ashikari *et al.*, 2005; Zalabak *et al.*, 2016), and were grouped together as clades. The first clade is comprised of *OsCKX1* and *OsCKX2*. *OsCKX1* is expressed in the apex of axillary buds as well as at the base of the shoot. Conversely, *OsCKX2*, typically found at elevated degree in the leaf collar and flowering, was also observed to be significantly expressed in the lateral root anlage, shoot plinth, and leaf blade. The second clade consists of *OsCKX6*, *OsCKX7*, and *OsCKX10*, which were exhibited very low expression level across every single tissue.

Comment [a1]: Citration needed for supporting your statement

The third clade included *OsCKX4*, *OsCKX5*, and *OsCKX9*. *OsCKX4* is usually highly expressed in vegetative tissues but expression specifically seen in the roots. The *OsCKX9* was normally expressed at a low concentration in almost all tissues but unveiled heightened expression throughout axillary buds and leaf blade. *OsCKX5* is typically shown high expression across all the organs, but it showed specific expression patterns in the leaves and roots. The *OsCKX3* and *OsCKX8* were included in the fourth clade. Although *OsCKX3* was anticipated to be most highly expressed in the shoot and young panicles, it also exhibited significant expression in base of the shoot. The *OsCKX8*, a gene tends to have lower expression levels in all vegetative organs than *OsCKX3*, displayed distinctly higher expression near the shoot outgrowth, in the primordia of flag leaf, and efflorescence. The *OsCKX11* showed a distinct nucleotide sequence compared to the other *OsCKX* genes included in the separate i.e., fifth clade and was generally expressed at greater levels in almost every tissue, especially in the roots, base of the shoot, and early flowering (Rong *et al.*, 2022).

Comment [a2]: Citration needed for supporting your statement

Comment [a3]: More citration needed

Cytokinins play a prominent role in the ordinance of panicle architecture which determines grain number in rice (Azizi *et al.*, 2015; Yeh *et al.*, 2015). A QTL (Quantitative Trait Locus) that governs the grain yield in rice is *GRAIN NUMBER 1a* (*GN1a*) (Ashikari *et al.*, 2005). Later it was found as a gene that encodes an enzyme Cytokinin Oxidase/dehydrogenase 2 (*OsCKX2*), which degrades plant hormone cytokinin and maintain its homeostasis (Ashikari *et al.*, 2005). *CKX2/ GN1a* was the first CKX gene to be known. Decreasing the expression of *OsCKX2* or *Osckx2* mutations resulted in plants or varieties that produced more vegetative tillers, grain number per inflorescence, and heavier grains (Ashikari *et al.*, 2005; Yeh *et al.*, 2015). The disruption of *OsCKX2* function increased the grain number, promoted secondary panicle branches, and boosted total grain yield by enhancing cytokinin levels in the panicle tissue (Rashid *et al.*, 2024).

In drought conditions, the *Osckx2* mutant retained more water and exhibited better water-saving characteristics, along with a survival response to manage dehydration stress. Additionally, *Osckx2* preserved the integrity of chloroplast membranes and demonstrated a notable improvement in photosynthetic function with enhanced antioxidant protection mechanisms. The elevated expression of *OsCKX2* adversely affects the spikelet count per panicle and drought resistance, but does not have a noticeable effect on salinity tolerance. These reports [suggests](#) the potential of *Osckx2* mutant to develop climate-resilient high-yielding varieties (Rashid *et al.*, 2024).

Comment [a4]: Citration needed

Now-a-days genetic engineering techniques such as gene silencing methods (Anti-sense technology, RNAi technology), and gene knock-out techniques (ZFN's, TALEN's, CRISPR/Cas system) are being used as efficient and precise tools for the development of elite varieties. Among these, CRISPR/Cas system is most widely used because of its high target specificity, and silencing efficiency.

Jansen *et al.* (2002) defined CRISPR as tandem repeats adjoined with non-recurrence DNA segments that were initially identified as defense mechanism of prokaryotes for

bacteriophages (Ishino *et al.*, 1987). The components of CRISPR technology for the cleavage process are (i) a sgRNA, a synthetic oligo of 20 base pairs that align to the desired DNA and (ii) a Cas9 nuclease enzyme that cut three bases before the PAM (generally 5' NGG; Jinek *et al.*, 2012). It comprises of two domains, (a) RuvC-like domain (member of RNase H family) and (b) an HNH domain (member of HNH endonuclease family), each cutting one DNA strand.

2. MATERIALS AND METHODS

CRISPR/Cas9 system binary vector

The pRGEB32, binary vector of CRISPR/Cas9 system was procured from Addgene, obtained as Bacterial stab culture. The vector encapsulated in the culture was retrieved on the Luria-Bertani (LB) agar with antibiotic Kanamycin (50mg/L).

Comment [a5]: Why you use only one vector pRGEB32? isn't other vector suitable? What is addgene? Is it a website or database or application??

Bacterial strains

The *E.coli*, *A. tumefaciens* strains DH5 α and EHA105 respectively were used in the study. The cultures were streaked and restored on LB agar plates with Nalidixic acid (25mg/L) and Rifampicin (25mg/L) antibiotics respectively.

Retrieval of gene sequence data of *GRAIN NUMBER 1a* gene

Rice *GRAIN NUMBER 1a* gene sequence was downloaded from Rice Genome Annotation Project (<http://rice.uga.edu/>) and Rice Annotation Project Database (<https://rapdb.dna.affrc.go.jp/>) in FASTA format and saved for further analysis. The Locus ID of the genome sequence for possible protospacer targets was taken from Rice Annotation Project Database.

Designing of sgRNAs

The sgRNAs were designed using plant specific CRISPR-P v2.0 software (<http://crispr.hzau.edu.cn/>). Two sgRNAs were chosen dependent on the off-target sites and their location, on-target scores, GC content, location on the gene, secondary structure (<https://rna.urmc.rochester.edu/RNAstructure.html>) etc., of sgRNAs with the Protospacer Adjacent Motif (PAM) as NGG. The sgRNAs were synthesized as sense and antisense strands with suitable complementary restriction enzyme (*Bsal*) sites as mentioned in Table1.

Synthesis of sgRNAs

The sgRNAs were analyzed and synthesized by Integrated DNA Technologies (IDT) with *Bsal* restriction sites.

Designing of primers for the study

Primers for M13 sequence of pRGEB32 vector was designed manually using the assistance of OligoAnalyzer™ tool of IDT.

CRISPR/Cas9 binary vector construct

In view of construction of CRISPR/Cas9 genome targeted editing cassette in binary vector, the pRGEB32 vector was isolated with GeNei@Puresol™ plasmid isolation Kit and was linearized with restriction enzyme *Bsal*-HF-v2, procured from New England Biolabs (NEB), UK. The restriction digestion reaction set up followed as mentioned in Table2. The restriction digested pRGEB32 vector was examined on 1% agarose gel for linearization and then vector was cleaned up with PCR Purification Kit purchased from QIAGEN QIAquick® PCR Purification Kit. The quality and quantity was analyzed using Nanodrop® (IMPLEN NP80) Spectrophotometer.

Ligation

The complementary strands of sgRNAs (100µM each) were annealed and the 5'ends were phosphorylated using T₄ Polynucleotide Kinase (PNK) as mentioned in Table 3. The annealed phosphorylated sgRNAs were diluted in (1:200) ratio and were ligated into digested pRGEB32 vector and then cloned into *E. coli* DH5α competent cells as mentioned in Table 4 and 5.

Table 1: The sgRNAs for targeted gene editing of *GN1a* with *Bsal* sites

Sl. No	Gene and sgRNA	Strand	Sequence (5' - 3')	PAM
1	<i>GN1a</i> sgRNA (OsGN1a # G1)	Sense strand Antisense strand	<u>GGCA</u> GCACGACGCGCGCAGCAGCG <u>AAAC</u> CGCTGCTGCGCGCGTTCGTGC	CGG
2	<i>GN1a</i> sgRNA (OsGN1a # G2)	Top strand Bottom strand	<u>GGCA</u> CGAGTGGCCACACCCCGCG <u>AAAC</u> CGCGGGGTGTGGCCACTCG	CGG

Table 2: Restriction digestion reaction set up of pRGEB32 vector

Components	Volume (µL)	Reaction conditions
Autoclaved distilled water	*to make up to 50 µL	Incubate at 37°C for 1 hr Heat inactivation at 65°C for 10min
10X CutSmart buffer	5	
<i>Bsal</i> -HF-v2 enzyme	1	
pRGEB32 vector	*5 micro gram	
Total reaction volume	50	

Table 3: Annealing and phosphorylation of sgRNAs

Components	Volume (µL)	Reaction conditions
Autoclaved distilled water	6.5	Incubation at 37°C for 3 min Incubation at 95°C for 5 min, slowly cooling down to 25°C at Ramp rate of 0.1°C/sec Store at -20°C
10X PNK Buffer	1.0	
sgRNA oligo sense strand (100µM)	1.0	
sgRNA oligo antisense strand (100µM)	1.0	
T ₄ Polynucleotide Kinase enzyme (PNK)	0.5	
Total reaction volume	10	

Table 4: Dilution of sgRNAs

Components	Volume (µL)	Reaction conditions
Annealed phosphorylated sgRNA	1	Mix it thoroughly Store at 4°C
Autoclaved distilled water	199	
Total reaction volume	200	

Table 5: Ligation of annealed sgRNAs and linearized vector

Components	Volume (μL)	Reaction conditions
Autoclaved distilled water	0.0	Incubation at 25°C for 15 min
10X ligation buffer	1.0	
Linearized vector (50ng)	7.0	Store at 4°C for shorter time
T ₄ DNA Ligase	1.0	
Diluted annealed PO ₄ ⁻ sgRNA	1.0	
Total reaction volume	10	
		Transformation into <i>E. coli</i> (DH5α strain) competent cells




Fig 1: A. Schematic representation of pRGEB32 vector.
B. Cloning of sgRNAs between *Bsa*I restriction sites expressed under OsU3 promoter.




Fig 2: A. Gel picture of isolated pRGEB32 vector. L - 1kb ladder, 1 - pRGEB32 vector.
B. Gel picture of *Bsa*I restriction site profile of pRGEB32 vector. L₁ - 1kb ladder, 1 - undigested vector, 2 and 3 - digested vector, L₂ - 100bp ladder.

Transformation of *E. coli* with recombinant vector

The ligated product consists of pRGEB32 vector incorporated with sgRNA was transformed into *E. coli* (DH5 α strain). For this process, the competent cells were prepared by Calcium-Chloride (CaCl₂) method (Tang *et al.*, 1994; Chang *et al.*, 2017) and transformation was done through heat-shock method @ 42° C for 90 sec (Chang *et al.*, 2017). After transformation, the bacterial culture was plated on LB agar containing selection marker Kanamycin (50mg/L), with 'L' shaped spreader and incubated at 37° C for 16 h.

Confirmation of recombinant construct in *E. coli*

The positive clones were randomly selected, streaked onto fresh LB agar plates containing Nalidixic acid (25mg/L) and Kanamycin (50mg/L) and analyzed by colony PCR. The PCR was performed through KAPA *Taq* PCR kit (Merck, USA) by using M13 reverse primer as forward primer and sgRNA antisense oligo as reverse primer as mentioned in Table 6 and 7 and the PCR product was visualized on 1% agarose gel using gel documentation system. Bands of expected size ~450 bp were observed. The positive clones confirmed by colony PCR was used for plasmid isolation.

Table 6: Colony PCR Master Mix

Components	Volume (μ L)
Autoclaved distilled water	14.25
10X buffer A (Forward reaction)	2.0
25 mM MgCl ₂	0.5
10 mM dNTP's	1.0
5U/ μ LTaq	0.25
10 μ M primer F (M13 R)	1.0
10 μ M primer R (sgRNAs)	1.0
Total reaction volume	20
Touch the colony edge with pipette tip or sterile tooth pick	
Tap in the reaction mixture	

Table 7: Colony PCR program

Steps	Temperature (°C)	Time	Cycles
Initial denaturation	95	10 min	} 38 cycles
Denaturation	95	30 s	
Annealing	60 - 62	30 s	
Extension	72	20 s	
Final Extension	72	10 min	
Final Hold	4	∞	

Table 8: Plasmid PCR Master Mix

Components	Volume (μL)
Autoclaved distilled water	12.05
10X buffer A	2.2
25 mM MgCl_2	0.5
10 mM dNTP's	1.0
100% DMSO	2.0
5U/ μL Taq	0.25
10 μM primer F (M13 R)	1.0
10 μM primer R (sgRNA)	1.0
Recombinant vector	2.0
Total reaction volume	22

Table 9: Plasmid PCR program

Steps	Temperature ($^{\circ}\text{C}$)	Time	Cycles
Initial denaturation	95	10 min	} 38 cycles
Denaturation	95	30 s	
Annealing	60 and 62 (gradient)	30 s	
Extension	72	1.5 min	
Final Extension	72	10 min	
Final Hold	4	∞	

Re-confirmation of recombinant construct in *E. coli*

The plasmid was isolated by Alkaline-lysis method (Ehrt and Schnappinger, 2003) and re-confirmation of recombinant construct was done by plasmid PCR. The quantity was analyzed using Nanodrop® Spectrophotometer, performed gel electrophoresis using 1% agarose and visualized under gel documentation system. The recombinant plasmid was also confirmed by PCR with KAPA TaqPCR kit by using M13 reverse as forward primer and sgRNA antisense oligo as reverse primer as mentioned in Table 8 and 9 and the product of PCR was visualized on 1% agarose gel using gel documentation system. Expected band size ~450 bp were seen. Further, the recombinant vector was confirmed by Sanger sequencing by GeneSpec Pvt. Ltd (<https://www.genespec.com>). The recombinant vector isolated was further purified using PCR purification kit purchased from QIAGEN QIAquick®PCR purification kit.

Mobilization of positive clones to *A. tumefaciens* strain EHA105

A. tumefaciens strain EHA105 culture was prepared to be competent by CaCl_2 method. The confirmed recombinant vector cloned and isolated from *E. coli* was then introduced into EHA105 strain following freeze-thaw method @ 37°C for 5 min (Holsters *et al.*, 1978). After transformation, the bacterial culture was spreaded on LB agar plates containing the antibiotics Rifampicin (25 mg/L) and Kanamycin (50 mg/L).

Confirmation of recombinant vector in *A. tumefaciens* strain EHA105

The positive clones were randomly selected, streaked onto fresh LB agar plates containing Rifampicin (25mg/L) and Kanamycin (50mg/L) and analyzed by colony PCR. The PCR was performed through KAPA Taq PCR kit by using M13 reverse as forward primer and sgRNA antisense oligo as reverse primer and the product of PCR was visualized on 1% agarose gel

using gel documentation system. The expected band size ~450 bp were observed. The positive clones were confirmed by colony PCR was used for plasmid isolation. The protocol mentioned above were followed for colony PCR, plasmid isolation and plasmid PCR.

3. RESULTS AND DISCUSSION

CRISPR/Cas9 system binary vector

The size of CRISPR/Cas9 system binary vector pRGEB32 is 15.9kbp (Xie *et al.*, 2014) and consisted of *Bsal* enzyme restriction site for sgRNA cloning, bacterial and plant selection markers Kanamycin (Kan^R) and Hygromycin (Hyg^R) resistant genes respectively under CaMV35 promoter, presence of Cas9 gene driven by rice ubiquitin promoter and also sgRNA cloned under pol III type promoter of rice U3 snoRNA as shown in Fig1A.

Retrieval of gene sequence data of *GRAIN NUMBER 1a* gene

The locus ID for *GN1a* gene sequence was designated in RGAP database as LOC_Os01g10110 and the gene was identified on chromosome 1 (Reverse orientation) (<http://rice.uga.edu/>). The complete sequence of the gene was downloaded in FASTA format from The Rice Annotation Project Database (RAP-DB) (<https://rapdb.dna.affrc.go.jp/>). The gene sequence length was 5576 bp, with a coding sequence of 1698 bp, covering 565 amino acids. The potential function was given as Cytokinin degradation. The gene had four exons and three introns (<http://rice.uga.edu/>).

Designing of sgRNAs

The sgRNAs of 20bp length for the specific targeting of *GN1a* were chosen from CRISPR-P v2.0 mentioned in Table 1. The two sgRNAs grounded on their on-target scores, GC content, location and position on the gene, off-target sites and their location, secondary structure were selected. The sgRNAs were located on the coding sequence region (CDS region) of the gene. The restriction sites of the typeII restriction enzyme *Bsal* were added to the 5'end of both sense and antisense strand of sgRNA to empower the cloning into pRGEB32 vector as shown in Fig1B.

Confirmation of recombinant construct in *E. coli*

The restriction digested plasmid with *Bsal* was analyzed for linearization on 1% agarose gel as shown in Fig2B. The restriction digested product was later ligated with annealed, phosphorylated sgRNAs; and were used for transformation into *E.coli* DH5 α . After overnight incubation, bacterial colonies with constructs were observed on the LB agar plates containing Nalidixic acid (25mg/L), and selection marker Kanamycin (50mg/L). The positive clones were randomly selected, streaked onto fresh LB agar plates containing Nalidixic acid (25mg/L) and Kanamycin (50mg/L) and analyzed by colony PCR. Expected bands of size ~450 bp were obtained on 1% agarose gel as shown in Fig3 and 4A. The positive colonies from PCR were chosen, used for plasmid isolation, plasmid PCR (Fig3 and 4B) and Sanger sequencing performed using universal M13 reverse primer and the Sanger sequencing result data was analyzed and sgRNAs insertion was confirmed using sequence alignment editor BioEdit7.2 software. Colony number 4 of sgRNA1 (pRGEB32:OsGN1a#G1) and colony 1 of sgRNA2 (pRGEB32:OsGN1a#G2) displayed insertion of respective sgRNAs with in pRGEB32 vector as shown in Fig 5A and B.




Fig 3: A. Colony PCR outline of *E. coli* DH5 α strain with the insertion of sgRNA. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G1-1, 2 - pRGEB32:OsGN1a # G1-2, 3 - pRGEB32:OsGN1a # G1-3, 4 - pRGEB32:OsGN1a # G1-4, 5 - pRGEB32:OsGN1a # G1-5, 6 - pRGEB32:OsGN1a # G1-6, 7 - pRGEB32:OsGN1a # G1-7, 8 - pRGEB32:OsGN1a # G1-8.
B. Plasmid PCR outline of selected colony pRGEB32:OsGN1a # G1-4. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G1 @ 60°C annealing temperature, 2 - pRGEB32:OsGN1a # G1 @ 62°C.




Fig 4: A. Colony PCR outline of *E. coli* DH5 α strain with the insertion of sgRNA. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G2-1, 2 - pRGEB32:OsGN1a # G2-2, 3 - pRGEB32:OsGN1a # G2-3, 4 - pRGEB32:OsGN1a # G2-4, 5 - pRGEB32:OsGN1a # G2-5, 6 - pRGEB32:OsGN1a # G2-6, 7 - pRGEB32:OsGN1a # G2-7, 8 - pRGEB32:OsGN1a # G2-8.
B. Plasmid PCR outline of selected colony pRGEB32:OsGN1a # G2-1. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G2 @ 60°C annealing temperature, 2 - pRGEB32:OsGN1a # G2 @ 62°C.



Fig 5: Sequence alignment showing cloning of sgRNA into pRGE32 vector backbone. **A.** pRGE32:OsGN1a # G1 construct, **B.** pRGE32:OsGN1a # G2 construct. BioEdit7.2 software was used for the analysis.

Mobilization of positive clones to *A. tumefaciens* strain EHA105

The positive clones of CRISPR/Cas9 sgRNA construct of both (OsGN1a#G1 and OsGN1a#G2) for *GN1a* confirmed after sequence analysis were then introduced into *A. tumefaciens* strain EHA105 through freeze-thaw method. After incubation period of 48h at 28° C, ~ 25 and 6 colonies for both sgRNA constructs were observed respectively on LB agar plates containing Rifampicin (25mg/L), selection marker Kanamycin (50mg/L).

Confirmation of recombinant vector in *A. tumefaciens* strain EHA105

The freshly streaked positive clones were verified by colony PCR. Eight colonies for *GN1a* (pRGE32:OsGN1a#G1) and six colonies for *GN1a* (pRGE32:OsGN1a#G2) were taken for analysis of colony PCR, further colony 4 for *GN1a* (OsGN1a#G1) and colony 6 for *GN1a* (OsGN1a#G2) were selected for plasmid isolation and plasmid PCR. Expected band size ~450 bp were observed on 1% agarose gel as shown in Fig 6A and B: Fig 7A and B.




Fig 6: A. Colony PCR outline of *A. tumefaciens* EHA105 strain with the insertion of sgRNA. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G1-1, 2 - pRGEB32:OsGN1a # G1-2, 3 - pRGEB32:OsGN1a # G1-3, 4 - pRGEB32:OsGN1a # G1-4, 5 - pRGEB32:OsGN1a # G1-5, 6 - pRGEB32:OsGN1a # G1-6, 7 - pRGEB32:OsGN1a # G1-7, 8 - pRGEB32:OsGN1a # G1-8. **B.** Plasmid PCR outline of selected colony pRGEB32:OsGN1a # G1-4. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G1 @ 60°C annealing temperature, 2 - pRGEB32:OsGN1a # G1 @ 62°C.




Fig 7: A. Colony PCR outline of *A. tumefaciens* EHA105 strain with the insertion of sgRNA. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G2-1, 2 - pRGEB32:OsGN1a # G2-2, 3 - pRGEB32:OsGN1a # G2-3, 4 - pRGEB32:OsGN1a # G2-4, 5 - pRGEB32:OsGN1a # G2-5, 6 - pRGEB32:OsGN1a # G2-6. **B.** Plasmid PCR outline of selected colony pRGEB32:OsGN1a # G2-6. L - 100bp ladder, 1 - pRGEB32:OsGN1a # G2 @ 60°C annealing temperature, 2 - pRGEB32:OsGN1a # G2 @ 62°C.

Rice grain yield can be efficiently driven by three characters (i) Number of panicles (ii) Grain number per panicle (GNPP) (iii) Grain weight (Zhou *et al.*, 2018). The improvement of grain number per panicle will increase the overall grain yield. Significantly, the rice grain yield per unit area will be high (Chen *et al.*, 2017; Chen *et al.*, 2018; Zhou *et al.*, 2018). Previous studies shown that grain yield was governed by over-expressing or suppressing phytohormone related genes. There is growing evidence that plant hormones primarily influence the transcriptional and post-transcriptional regulation of rice genes associated with GNPP in order to mediate the determination of GNPP. *GN4-1* from Wuyunjing 8 rice cultivar

positively regulated GNPP by 17% with promoting CK accumulation in rice inflorescences (Zhou *et al.*, 2018). GNP1 could boost the activity of the inflorescence meristem by enhancing cytokinin signaling and inhibiting gibberellin signaling, which may lead to an increase in the number of secondary rachis branches and GNPP (Wu *et al.*, 2016). *LARGER PANICLE (LP)/ERECT PANICLE 3 (EP3)* interacts with SKP1-like protein, when *OsCKX2* expression is up-regulated and CK levels are lowered during rice inflorescence, which increased primary and secondary rachis branch production and grain yield (Li *et al.*, 2011).

DROUGHT AND SALT TOLERANCE (DST), a zinc-finger transcription factor has a negative effect on cytokinin content, which leads to a lowering in the number of rachis branches and the GNPP in rice. *DST*, that modulates the expression of the *OsCKX2*, which encodes cytokinin oxidase, through a single base insertion that results in a loss of its ability to activate transcription. This change subsequently enhances the rachis branches number, GNPP, and overall yield of grain (Li *et al.*, 2013; Guo *et al.*, 2020).

The inactivation of *LONELY GUY (LOG)* led to an early cessation of shoot apical meristem (SAM) function by modulating the levels and spatial arrangement of CK, which reduced both the count of rachis branches and GNPP (Kurakawa *et al.*, 2007). *GRAIN NUMBER 1a*, *GRAIN LENGTH AND AWN DEVELOPMENT1 (GAD1)* acts as a negative regulator of GNPP, as the *GAD1* protein lowers the concentration of cytokinins (CK) by promoting the expression of *DST* and *OsCKX2*. This reduction results in decreased GNPP in wild rice. In cultivated rice, a mutation that alters the codon of *GAD1*, which disrupts the preserved cysteine structure, leading to the loss of *GAD1* function. Consequently, this enhances GNPP, decreases grain length, and inhibits the development of awns (Jin *et al.*, 2016). *GRAIN NUMBER 1a (GN1a) / Cytokinin Oxidase2 (OsCKX2)* acts as antagonist to GNPP by decreasing the cytokinin levels. A decrease in *OsCKX2* expression leads to an increase in cytokinin levels within the inflorescence meristem, which promotes the development of additional rachis branches and GNPP, ultimately enhancing grain yield (Ashikari *et al.*, 2005) and resistance to lodging (Tu *et al.*, 2022).

Plant vector pRGEB32 is the most pre-dominant binary vector system exploited for site-directed genome editing using CRISPR/Cas9 system, the efficient tool reported for precise mutagenesis in most crop species (Xie *et al.*, 2014). The vector system has sgRNA cloning sites fringed with *Bsal* restriction sites, driven by pol III type promoter of rice U3 snoRNA. Cas9 gene from *Streptococcus pyogenes* attached with nuclear-localizing signal (NLS), encodes for Cas9 nuclease (ribonucleoprotein), that directs DNA-targeted cleavage, whose expression driven by rice ubiquitin promoter. Kanamycin (Kan^R) and Hygromycin (Hyg^R) resistant genes as bacterial and plant selection markers respectively influenced by CaMV35S promoter.

The CRISPR/Cas9 recombinant cassettes were constructed by ligating the phosphorylated, annealed spacer oligos in pRGEB32 vector and then cloned into *E. coli* DH5 α strain by heat-shock method @ 42°C for 90s (Chang *et al.*, 2017). The cloning of this cassette poses a challenge because of larger plasmid size (~ 15.9kb) affecting the uptake of plasmid, possibly resulting in lesser number of bacterial transformants. Despite this expected hurdles, we obtained more than 50 transformants, among which 8 were selected for further sub-culturing. The confirmation of positive clones was done by colony PCR, followed by plasmid isolation and plasmid PCR. PCR positive clones were analyzed by sequence alignment after sequencing.

The positive clones were then mobilized into *A. tumefaciens* EHA105 with freeze-thaw method @ 37°C for 5min (Holsters *et al.*, 1978). Approx 6 - 20 colonies were observed in OsGN1a#G1 and OsGN1a#G2. *Agrobacterium tumefaciens*-mediated transformation became an adaptable

practice in lab conditions (Chan *et al.*, 1993 ; Hiei *et al.*, 1994), for its ideal capability to deliver a desired DNA fragment from a plasmid into a host plant (Gelvin, 2010), enables efficient integration, un-rearranged, one-copy DNA, which end up with stable expression than many gene copies or scrambled inserts which will lead to unwanted mutation (Iglesias *et al.*, 1997).

Comment [a6]: I can not align this part with result and discussion segment please go through it.

4. CONCLUSION

Yield is considered to be the significantly important physiological trait governed by many QTL's, some of which are negatively influencing the yield. In this study, CRISPR/cas9 technique, a genome editing tool used for site-targeted editing of *GRAIN NUMBER 1a* gene, which negatively regulates the yield by degrading Cytokinin. Precisely efficient sgRNAs were designed using CRISPR-P v2.0 software, synthesized as sense and antisense strands with *BsaI* restriction sites by IDT, phosphorylated, annealed, ligated into pRGEB32 vector, then transformed to *E.coli* DH5 α and then transferred to *A. tumefaciens* EHA105.

FUTURE SCOPE

The present study could be helpful for meeting the food demand for growing population and also for the farmer's welfare by increasing the yield through targeted genome editing using CRISPR/cas9 system.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The authors hereby declared that the following AI generative tools were used during the writing and editing of the manuscript.

Details of the AI tools used are given below:

1. Grammarly
2. ChatGPT
3. Quillbot

REFERENCES

- Ashikari, M., Sakakibara, H., Lin, S., Yamamoto, T., Takashi, T., Nishimura, A., Angeles, E.R., Qian, Q., Kitano, H. and Matsuoka, M. (2005). Cytokinin oxidase regulates rice grain production. *Science*, 309, 741-745.
- Azizi, P., Rafii, M.Y., Maziah, M., Abdullah, S.N.A., Hanafi, M.M., Latif, M.A., Rashid, A.A. and Sahebi, M. (2015). Understanding the shoot apical meristem regulation: a study of the phytohormones, auxin and cytokinin, in rice. *Mechanisms of Development*, 135, 1-15.
- Chan, M. T., Chang, H. H., Ho, S. L., Tong W. F. and Yu S. M. (1993). *Agrobacterium*-mediated production of transgenic rice plants expressing a chimeric alpha-amylase promoter/beta-glucuronidase gene. *Plant Molecular Biology*, 22 (3), 491-506.
- Chang, A.Y., Chau, V., Landas, J.A. and Pang, Y. (2017). Preparation of calcium competent *Escherichia coli* and heat-shock transformation. *Journal of Entrepreneurship, Management and Innovation methods*, 1, 22-25.
- Chen, H., Tang, Y., Liu, J., Tan, L., Jiang, J., Wang, M., Zhu, Z., Sun, X. and Sun, C. (2017). Emergence of a novel chimeric gene underlying grain number in rice. *Genetics*, 205, 993-1002.

- Chen, L., Bian, J., Shi, S., Yu, J., Khanzada, H., Wassan, G.M., Zhu, C., Luo, X., Tong, S., Yang, X. and Peng, X. (2018). Genetic analysis for the grain number heterosis of a super-hybrid rice WFYT025 combination using RNA-Seq. *Rice*, 11, 1-13.
- Ehrt, S. and Schnappinger, D. (2003). Isolation of plasmids from *E. coli* by alkaline lysis. *E. coli Plasmid Vectors: Methods and Applications*, 75-78.
- Gelvin, S. B. (2010). Plant proteins involved in *Agrobacterium*-mediated genetic transformation. *Annual Review of Phytopathology*, 48, 45-68.
- Guo, T., Lu, Z.-Q., Shan, J.-X., Ye, W.-W., Dong, N.-Q. and Lin, H.-X. (2020). *ERECTA1* acts upstream of the *OsMKKK10-OsMKK4-OsMPK6* cascade to control spikelet number by regulating cytokinin metabolism in rice. *Plant Cell*, 32, 2763-2779.
- Hiei, Y., Ohta, S., Komari, T. and Kumashiro, T. (1994). Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant Journal*, 6 (2), 271-282.
- Holsters, M., De Waele, D., Depicker, A., Messens, E., Van Montagu, M. and Schell, J. (1978). Transfection and transformation of *Agrobacterium tumefaciens*. *Molecular Genetics and Genomics*, 163, 181-187.
- Iglesias, V. A., Moscone, E. A., Papp, I., Neuhuber, F., Michalowski, S., Phelan, T., Spiker, S., Matzke M. and Matzke, A. J. (1997). Molecular and cytogenetic analyses of stably and unstably expressed transgene loci in tobacco. *Plant Cell*, 9 (8), 1251-1264.
- Ishino, Y., Shinagawa, H., Makino, K., Amemura, M. and Nakata, A. (1987). Nucleotide sequence of the *iap* gene, responsible for alkaline phosphatase isozyme conversion in *Escherichia coli*, and identification of the gene product. *Journal of bacteriology*, 169(12), 5429-5433.
- Jansen, R., Embden, J.D.V., Gaastra, W. and Schouls, L.M. (2002). Identification of genes that are associated with DNA repeats in prokaryotes. *Molecular Microbiology*, 43 (6), 1565-1575.
- Jin, J., Hua, L., Zhu, Z., Tan, L., Zhao, X., Zhang, W., Liu, F., Fu, Y., Cai, H., Sun, X. and Gu, P. (2016). *GAD1* encodes a secreted peptide that regulates grain number, grain length, and awn development in rice domestication. *The Plant Cell*, 28 (10), 2453-2463.
- Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A. and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*, 337 (6096), 816-821.
- Kurakawa, T., Ueda, N., Maekawa, M., Kobayashi, K., Kojima, M., Nagato, Y., Sakakibara, H. and Kozuka, J. (2007). Direct control of shoot meristem activity by a cytokinin-activating enzyme. *Nature*, 445, 652-655.
- Li, M., Tang, D., Wang, K., Wu, X., Lu, L., Yu, H., Gu, M., Yan, C. and Cheng, Z. (2011). Mutations in the F-box gene *LARGER PANICLE* improve the panicle architecture and enhance the grain yield in rice. *Plant Biotechnology Journal*, 9, 1002-1013.

- Li, S., Zhao, B., Yuan, D., Duan, M., Qian, Q., Tang, L., Wang, B., Liu, X., Zhang, J., Wang, J. and Sun, J. (2013). Rice zinc finger protein DST enhances grain production through controlling *Gn1a/OsCKX2* expression. *Proceedings of the National Academy of Sciences*, 110 (8), 3167-3172.
- Peng, J., Richards, D.E., Hartley, N.M., Murphy, G.P., Devos, K.M., Flintham, J.E., Beales, J., Fish, L.J., Worland, A.J., Pelica, F. and Sudhakar, D. (1999). 'Green revolution' genes encode mutant gibberellin response modulators. *Nature*, 400 (6741), 256-261.
- Rashid, A., Achary, V.M.M., Abidin, M.Z., Karippadakam, S., Parmar, H., Panditi, V., Prakash, G., Bhatnagar-Mathur, P. and Reddy, M.K. (2024). Cytokinin oxidase2 - deficient mutants improve panicle and grain architecture through cytokinin accumulation and enhance drought tolerance in indica rice. *Plant Cell Reports*, 43 (8), 207.
- Rong, C., Liu, Y., Chang, Z., Liu, Z., Ding, Y. and Ding, C. (2022). Cytokinin oxidase/dehydrogenase family genes exhibit functional divergence and overlap in rice growth and development, especially in control of tillering. *Journal of Experimental Botany*, 73 (11), 3552-3568.
- Tang, X., Nakata, Y., Li, H.O., Zhang, M., Gao, H., Fujita, A., Sakatsume, O., Ohta, T. and Yokoyama, K. (1994). The optimization of preparations of competent cells for transformation of *E. coli*. *Nucleic Acids Research*, 22 (14), 2857.
- Tu, B., Tao, Z., Wang, S., Zhou, L., Zheng, L., Zhang, C., Li, X., Zhang, X., Yin, J., Zhu, X. and Yuan, H. (2022). Loss of *Gn1a/OsCKX2* confers heavy-panicle rice with excellent lodging resistance. *Journal of Integrated Plant Biology*, 64 (1), 23-38.
- Wu, Y., Wang, Y., Mi, X.-F., Shan, J.-X., Li, X.-M., Xu, J.-L. and Lin, H.-X. (2016). The QTL *GNP1* encodes GA20ox1, which increases grain number and yield by increasing cytokinin activity in rice panicle meristems. *Public Library of Sciences Genetics*, 12, e1006386.
- Xie, K., Minkenberg, B. and Yang, Y. (2014). Targeted gene mutation in rice using CRISPR-Cas9 system. *Bio-Protocol*, 4 (17), e1225.
- Yeh, S.Y., Chen, H.W., Ng, C.Y., Lin, C.Y., Tseng, T.H., Li, W.H. and Ku, M.S. (2015). Down-regulation of cytokinin oxidase 2 expression increases tiller number and improves rice yield. *Rice*, 8, 1-13.
- Zalabak, D., Johnova, P., Plihal, O., Senkova, K., Samajova, O., Jiskrova, E., Novak, O., Jackson, D., Mohanty, A. and Galuszka, P. (2016). Maize cytokinin dehydrogenase isozymes are localized predominantly to the vacuoles. *Plant Physiology and Biochemistry*, 104, 114-124.
- Zhou, y., tao, y., yuan, y., zhang, y., miao, j., zhang, r., yi, c., gong, z., yang, z. And liang, g. (2018). Characterisation of a novel quantitative trait locus, *gn4-1*, for grain number and yield in rice (*oryza sativa* l.). *Theoretical and applied genetics*, 131, 637-648.