

Tsetse fly (*Diptera, Muscidae*) Population Dispersal in Endemic Areas of Animal Trypanosomiasis in Mali

Commented [II1]: No need for order in title, it should be in background

ABSTRACT

Tsetse flies are harmful to humans and animals through their bites and infestations. They are dependent on vegetation, islands, forests, galleries, etc. Visiting resting places, hunting grounds, or breeding grounds is a means of contact between human, animal, and insect populations. It encourages host-vector encounters and thus pathogen exchanges among humans, wildlife, and the environment. Controlling these insects considerably reduces their nuisance and breaks the pathogen transmission chain. Knowledge of the diversity of tsetse species and the main determinants of their geographical distribution is necessary to effectively combat this scourge. This study was initiated to determine the distribution of tsetse flies and identify the main species currently present in Mali. The 4 regions endemic for trypanosomiasis in Mali (Kayes, Koulikoro, Sikasso and Ségou) were selected for capture. In each region, 10 captures sites were randomly selected. The captures occurred from March to December 2020. The traps were placed on the banks of watercourses, in galleries, and in the Savannah. They were installed for 48 h and renewed every 24 h. A total of 1353 tsetse flies were captured, including 270 in Kayes, 527 in Koulikoro, 554 in Sikasso, and 4 in Ségou. Two riparian species were identified, and no Savannah species were recorded. Tsetse flies were heterogeneously distributed. All sites in the Sikasso region recorded the presence of at least one tsetse fly (100%), while representativeness was 80, 40 and 30%, respectively, in the Koulikoro, Kayes, and Ségou regions. These results show that tsetse flies are still present in these regions, albeit with an uneven distribution.

Commented [II2]: Re-structure the sentence

Commented [II3]: at

Commented [II4]: pitched

Commented [II5]: harvested

Commented [II6]: Abstract should not be in table format. Please convert into plane text

Keywords: Distribution, vector control, distribution area, Mali.

1. INTRODUCTION

Insects are arthropods with the largest number of species (nearly 1.3 million described). They account for more than two-thirds of all living organisms and are widespread throughout the world. In Africa south of the Sahara and in tropical zones, insect biodiversity remains predominant due to environmental conditions that favor their multiplication and induce outbreaks (Duvallat *et al.*, (2012)). Tsetse flies, which are responsible for human and animal trypanosomiasis, are widespread in Africa. Nearly 10 million km² are infested by the tsetse fly (FAO, (2002)). These tsetse fly species are highly sensitive to climatic conditions (Itard *et al.*, (2003)). Seasonal variations in temperature and humidity constantly drive them to seek their ecological optimum; they modify the amplitude of their niche according to seasonal conditions. The sensitivity of tsetse flies to climatic variations (Itard J. & al., (2003)) and their singular demographic behaviors provide information that can be used to model their spatial distributions. Tsetse are dependent on vegetation, islands, forests, and galleries. These areas are generally used by rural populations for hunting, farming, fishing, or market gardening. Anthropogenic pressure contributes to the degradation of the riparian cordon, leading to the disappearance of pupal sites or the retreat of tsetse flies to a more favorable biotope Nash, T.A.M., (1948). In addition, frequenting tsetse resting places, hunting grounds, or breeding grounds is a means of contact between human, animal, and insect populations. Host-vector encounters promote pathogen exchange between humans, wildlife, and the environment (Bouyer J. & al.,

Formatted: Font: Italic

Formatted: Font: Italic

Commented [II7]: -Only surname should be included in citations
-remove all '&' and replace with 'et'
-italize 'et al.', in all citations please
-maintain uniform font throughout the manuscript. E.g Times new roman

Commented [II8]: Same as above!

(2009a). The harmful bite of tsetse flies and their role in the cyclical transmission of African Animal Trypanosomiasis (AAT) and African-Human African Trypanosomiasis (AAT & HAT) make them very formidable insects Epstein, P.R. & al., (1993). It is transmitted to humans and animals by a variety of species, both male and female (Aksoy .S. et al., 2003 ; Courtin, F. et al., (2005 ; Mbida Mbida & al. (2009) ; Bosson-Vanga & al., (2012)). According to the WHO, vector-borne diseases are among the main causes of morbidity and mortality in humans and animals, and trypanosomoses are no exception. Their impact on human health is considerable. In many Central African countries, HAT alone accounts for 87% of the total number of patients diagnosed throughout Africa (WHO, World Health Organization, 2006). Approximately 50 million people and 48 million heads of livestock are at risk from trypanosomosis (Epstein et al., 1993). In Mali, nearly 80% of the population lives from agro-silvo - pastoral activities OMS, (2006). Tsetse fly over more than 20,000 km² from north to south, i.e., 16% of the national territory. Djitéye and al. (1997) reported the existence of three main species. After several years of control in the river basins of the Bani and Niger systems, the density is still high at 4.37 tsetse/trap/day in the Niger River basin. *G. palpalis gambiensis* was the predominant species in the Niger and Bani river basins Traoré A. & al., (2019). Bass & al., (2014) have nevertheless observed a decrease in the range of tsetse flies, especially toward the center of the country (Ségou region) Salou E. & al ; (2012). Vector control is a means of considerably reducing nuisance and breaking the pathogen transmission chain. The Tsetse fly may be present all year round or only during certain seasons. Knowledge of the ecosystem and target species is essential for effective control of these insects. The aim of this study was to update our knowledge of the distribution of tsetse flies and to identify the species currently encountered to contribute to the adaptation of a more effective control strategy against these insects in Mali.

2. MATERIAL AND METHODS

2.1. Study area

The study was conducted from March to December 2020 in four administrative regions endemic to animal trypanosomiasis in Mali. These are Sikasso's regions, Koulikoro, Kayes, and Ségou



Fig. 1: Geolocation of capture sites

Commented [II9]: Same!
Is there any other published research by same author in the same year?!
Please remove letter 'a'

Commented [II10]: Correct it pls!

Commented [II11]: Correct them!

Commented [II12]: Be consistent: trypanosomiasis

Commented [II13]: Edit!
What is OMS?

Commented [II14]: ?

Formatted: Font: Italic

Commented [II15]: Same as above!

Commented [II16]: Add coordinates of the locations used to capture flies

Rainfall variation in the study area

Table 1 shows the recorded rainfall and number of rainy days in the 4 regions during the 2021–2022 rainy season. A large variability was found not only in the annual rainfall but also in the number of rainy days per year. The region of Sikasso received a greater amount of rain (1185 mm) and a greater number of rainy days (95) than the other regions. The lowest amount of rain (655 mm) was observed in the region of Kayes (Sahelian zone) with a significant number of rainy days (53 days).

Table 1. Annual rainfall and number of rainy days in the study area during the 2021–2022 rainy season

Region	Annual rainfall	Number of rainy days
Kayes	655	53
Koulikoro	990	74
Sikasso	1185	95

2.2. Tsetse fly capture

Ten sites were randomly selected from each of the above regions. Catches were made using biconical and monotonical traps set along watercourses, galleries, and savannahs. Each trap was georeferenced and labeled with the initial site, date, and capture point (or trap) number. The temperature and hygrometry were recorded. Captured tsetse flies were collected 24 h after setting the traps, identified morphologically using the identification key and a magnifying glass, and their number and sex were determined.

2.3. Data analysis

Data were saved on the Windows 2010 Excel software, which was also used to perform proportion calculations and produce variation curves and histograms. For all analyses, the maximum significance level was 0.05. Because of its low numbers, *G. tachinoides* was excluded in the analyses. For the analysis of abundance, the Pearson correlation coefficient was applied using the number and species of tsetse and meteorological parameters as determinants of their dispersal.

■ Relative abundance

The number of individuals of one species in relation to the total number of individuals of all species combined was calculated using the following formula:

$$Ar = Aa/N \times 100$$

where N is the total number of individuals.

■ Apparent density (DAP)

This corresponds to the number of tsetse fly captured per number of traps per day.

3. RESULTS AND DISCUSSION

3.1. Spatial distribution of the tsetse flies

The number of tsetse fly captured was higher in Sikasso than in the other three regions (figure 2a).

Commented [II17]: Re-write!

Formatted: Font: Italic

Formatted: Indent: Left: 0.5", No bullets or numbering

Commented [II18]: Reference authority needed here
What is 'Ar and Aa'?




Fig. 2: Variation in the number of tsetse fly captured by region (a) and population distribution by capture site (b)

In the Koulikoro and Kayes regions, the number of tsetse flies captured is to a lesser extent high, with great intersite variability in the Koulikoro region in Mali's Sudanian zone ($p = 0.009$). Soudougouba (Baguinéda commune) recorded the highest number of tsetse flies captured (138), compared with only 34 at Samanko in the Mandé commune. Among the sites surveyed in the Sikasso region, Zèkorodougou recorded 109 tsetse flies, compared with 10 in Djandougou and Tonokalakoura. The sex ratio was almost identical in the Koulikoro and Ségou sites, whereas it was largely in favor of females in the Sikasso and Kayes sites. The number of tsetse fly captured in the Ségou region is relatively low and stable. The distribution of the captured population shows that captures of less than 20 flyers per site are the most frequent (figure 2b). Estimated average densities were 0.03 in Ségou, 1.96 tsetse fly/trap/day in Kayes, 3.15 tsetse fly/trap/day in Sikasso, and 3.56 tsetse fly/trap/day in Koulikoro.

Commented [II19]: Is not part of the 4 mentioned locations

3.2. Relative tsetse fly abundance by region

Of the 1,255 tsetse fly captured, 44% were from the Sikasso region, versus 34% from the Koulikoro region (figure 3). Tsetse flies were relatively scarce in the Ségou region (0.32%). The sex ratio was almost identical in the Koulikoro and Ségou sites, whereas it was largely in favor of females in the Sikasso and Kayes sites ($p < 0.005$) (figure 3).

Formatted: Font: Italic




Fig. 3: Relative tsetse fly abundance according to region

3.3. Tsetse diversity according to the region

Two species were identified in the study sites (*Glossina palpalis gambiensis* and *Glossina tachinoides*). These two species were recorded in the Sikasso and Kayes regions, whereas *G. palpalis gambiensis* was the only species caught in the Ségou and Koulikoro regions (figure 4).




Fig. 4: Number of tsetse fly captured by species and region

Glossina palpalis gambiensis was the predominant species, accounting for over 99.5% of the total, compared with only 0.5% for *G. tachinoides*. The representativeness of glossing was estimated at 100, 80, 40 and 30%, respectively, in Sikasso's regions, Koulikoro, Kayes, and Ségou. Average densities recorded were 0.03 in Ségou, 1.96 tsetse flies/trap/day in Kayes, 3.15 tsetse flies/trap/day in Sikasso, and 3.56 tsetse flies/trap/day in Koulikoro. *Glossina palpalis gambiensis* was the predominant species, accounting for over 99.5%, compared with just 0.5% for *G. tachinoides*.

3.4. Correlation between flies density and meteorological parameters

Average temperatures ranged from 28 to 33°C, with the lowest temperatures recorded in the Sikasso region and an average relative humidity of 51%. It was higher in Kayes, with an average relative humidity of 59%. Rainfall was highest in Sikasso and lowest in Kayes. The data show that the correlations are not significant at the 5% level under our rearing conditions (Table 2). However, the apparent density (DAP) of the flies was slightly sensitive to variations in meteorological parameters, particularly rainfall, compared with that outside temperature, for which the correlation coefficients were negative for the three observed parameters.

Table 2. Correlation between apparent density of flies and meteorological parameters under rearing conditions

Meteorological parameters	Correlation coefficient	p-value
Rainfall (mm)	0.72	0.28
Relative humidity (%)	-0.36	0.64
Outside temperature (°C)	-0.67	0.33

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [II20]: Re-write pls!

Formatted: Font: Italic

Formatted: Font: Italic

The maximum number of catches was observed in the Sikasso region and the minimum was recorded in the Ségou region. This is because the Sikasso region has a biotope more favorable to tsetse fly proliferation than the Ségou region, which is at the northern limit of tsetse fly distribution in Mali. Since 2019, the rarity of tsetse flies in this part of the country has been reported (Traoré *et al.*, 2019). The variability in the number of tsetse captured between the four regions could be explained by the diversity of bioclimates, as shown by Traoré A. & al., (2019). Two tsetse species, all riparian, were captured (*G. palpalis gambiensis* and *G. tachinoides*). These results confirm those of Bass *et al.*, (2014), in the Sikasso and Koulikoro regions, but differ from those of Djitéye & al. (1997), who reported the existence of three tsetse species (*G. palpalis gambiensis*, *G. tachinoides*, *G. morsitans*) in the tsetse range of Mali. *G. palpalis gambiensis* was present in all four regions surveyed and was the predominant species, whereas *G. tachinoides* was only caught in the Sikasso (Madinna Diassa) and Kayes (Taliko) regions. In a study conducted in Burkina Faso Salou E. & al., (2012) reported a strong predominance of *G. tachinoides* over *G. p. gambiensis* in Folonzo (84% versus 16% of catches, respectively), whereas in Kartasso, *G. p. gambiensis* was the only species present. Dibakou et al. (2015) in the forest zone and in the village of Doussala identified six species of tsetse fly, including *G. tachinoides*, but *G. p. gambiensis* was completely absent. According to Dibakou & al. (2015). *G. palpalis gambiensis* and *G. tachinoides* frequently coexist and occur in gallery-type vegetation, but *G. tachinoides* remains the dominant species near permanent watering holes. Given that most of the watercourses surveyed, especially in the Kayes region, were temporary, it is obvious that the numbers of this species caught were lower. The rarity of this species (*G. tachinoides*) and the absence of *G. morsitans* could be due to their ecological requirements or anthropic effects. Some authors [Darchen, R., (1978) ; Solano, P., (2008) ; Mbida Mbida (2009) , Buxton PA. (1955) reported that the infestation of an environment by tsetse flies is conditioned by the simultaneous presence of suitable environmental factors (temperature between 15°C and 25°C, luminosity, relative humidity) and feeding vertebrate hosts. Solano, P., (2008), indicated that the very low presence of certain tsetse species such as *G. tachinoides* in a given biotope reflects their ecological requirements. It is clear that in our catchment areas, feeder hosts are scarce, and the climatic conditions were very unfavorable in Kayes and Ségou. The results show that tsetse flies were present in all four regions surveyed but were heterogeneously dispersed. All the sites surveyed in the Sikasso region had an apparent density greater than or equal to one tsetse fly/trap/day, giving a representativeness of 100%. In contrast, only 30% of the sites surveyed were positive in the Ségou region, with an average density of 0.03 tsetse fly/trap/day. In Koulikoro and Kayes, representativeness was 80% and 40%, with average densities of 1.96 and 3.56 tsetse fly/trap/day, respectively. The results of the Pan-African Animal Trypanosomiasis Control Campaign (PATTEC, 2009) showed that the Sikasso region was 100% infested, the Koulikoro region 76%, and the Kayes and Ségou regions 46% and 44%, respectively. A study of the distribution of tsetse flies in the Savannah region showed that they are present almost everywhere in the river system, as long as there is sufficient tree vegetation to create the microclimate they need. Tsetse flies have been reported in all three of Mali's river basins. We note the absence of tsetse *Glossina morsitans* and the rarity of *G. tachinoides*. The *morsitans* group is more xerophilous (optimum 50-60% relative humidity) and disperses widely during the rainy season in wooded savannahs and in the dry season in the most favorable microclimates of streamside vegetation. According to Djitéye & al. (1997), the *morsitans* group is located in southern Mali. In a range covering approximately 200,000 km² south of 14° 30' N and west of 4° 30' W, tsetse flies are found only in the Sahelian, Sudanian, and pre-Guinean zones. In Sahelian zones such as the Ségou region, the climate is arid or semi-arid, and tsetse flies are few or non-existent depending on the microclimate, as our results show. The Sudanian zone, a tsetse fly zone par excellence, offers a favorable biotope for tsetse proliferation and dynamics. In the homogeneous landscapes found in protected areas such as that of the Sikasso region, the forest gallery is preserved along the entire length of the river,

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Commented [II21]: Correct it as above!

Formatted: Font: Italic

providing tsetse with favorable living conditions (temperature, humidity, pupation sites), hence the existence of three species of tsetse in this part of the country. Under these conditions, the tsetse fly has a strong tendency to disperse, which explains their wide dispersal (100%) in this part of the country. The microclimate of the Koulikoro region (Mandé commune) borders the pre-Guinean climate. These conditions are very similar to those in the south of the country, with representativeness estimated at 80%. On the other hand, in the Kayes region, the favorable biotope is spatially fragmented and gradually degrades under the influence of climate, anthropogenic and agricultural effects, lower relative humidity, and higher temperatures. In this region, the flies are less distributed (40%) than in the previous two. (Bouyer J. & al., 2009a), showed that agricultural practices lead to the degradation of the riparian cordon, resulting in the disappearance of pupation sites, lower relative humidity, and higher temperatures. According to Soberón J. Peterson A. T. (2005), abiotic conditions such as altitude, climate or cover, biotic factors, the species' ability to disperse spatially, and its capacity to adapt are factors that influence the spatial distribution of a species. According to our results and Pearson's matrix analysis, the correlation between climatic variables (temperature, relative humidity, rainfall) and tsetse fly dispersal is not always positive. There is a negative correlation between relative abundance, temperature, and humidity. Diversity, rainfall, and relative humidity were positively correlated ($P < 0.05$). Traoré A. & al., (2023), showed that there was no direct correlation between the vegetation index and the number of tsetse fly captured. Variables such as hygrometry and temperature are the main determinants of variation in plain tsetse fly density, regardless of the site surveyed. The uneven distribution of tsetse could therefore be explained by the diversity and structure of the environment, which can create specific microhabitats that are more or less favorable to tsetse development. The absence of *Glossina morsitans* does not necessarily mean that the species is extinct. In addition to abiotic factors, the survey period could also be a determining factor in tsetse dispersal. At this time of year, tsetse flies are more focused on their hunting grounds, larval breeding sites, and the microclimate suitable for this purpose, as set out (Frezil and Carnevale, (1976); Laveissière and Hervouët, (1981); Gouteux and Kiénoú, (1982); Zinga & al., (2013) and (2024). Such conditions seem to be found in the southern part of Mali (Sikasso and Koulikoro regions) and would explain the high level of tsetse fly dispersal in these areas.

Commented [II22]: ???

Commented [II23]: Modify pls!

Commented [II24]: Change as above

Commented [II25]: ??

4. CONCLUSION

This study aimed to document the diversity of tsetse species present in the four regions and their geographical distribution. The abundance and distribution of the tsetse fly has not been uniform. The number of tsetse fly captured varied according to the biotopes surveyed in the four administrative regions. Two species were present, with a clear predominance of *G. palpalis gambiensis*. Despite numerous tsetse control campaigns, tsetse are still present in the four regions mentioned above, and their density remains high outside the Ségou region. A longitudinal survey coupled with tsetse trapping at several sites along the bioclimatic gradient is required to determine the actual or seasonal density of these species.

Formatted: Font: Italic

REFERENCES

- [1] Duvallet G. and Gentile L. (2012). Protection Personnelle Antivectorielle. IRD Editions, 25 p
- [2] Alain B. (2007). Parasitoses and mycoses in temperate and tropical regions. Edition Elsevier, pp 321
- [3] Itard J., Cuisance D. and Tacher G. (2003). Trypanosomes: History: Geographical Distribution. In Lefèvre PC, Blancou J., and Chermette R, dir. Main infectious and parasitic diseases of livestock: Europe and Hot Regions. Paris, France: Lavoisier Tec and Doc, pp. 1607- 1615,
- [4] Nash TAM (1948). Tsetse Ejes in british west africa : the gold coast. Her Majesty Stationary Office, London, UK, 28: 47
- [5] Bouyer J (2009). Tsetse flies, intelligent flies? Part 2. The dispersal of tsetse flies. Insects (153): 21-24.
- [6] Epstein PR, Ford TE, Colwell RR (1993). Marine ecosystems: emerging diseases as indicators of change. Lancet, 342: 1217-1219.

Commented [II26]: -Remove all square brackets
-Italicize journal name
-Bold volume numbers
-arrange from A-Z in ascending order
-do the same for all references

Formatted: Font: Italic

- [7] Aksoy S (2003). Control of tsetse flies and trypanosomes using molecular genetics. *Vet Parasitol.*, 115 : 125-145.
- [8] Courtin F, Dupont S, Zeze DG, Jamonneau V, Sané B, Coulibaly B, Cuny G, Solano P (2005). Human African Trypanosomiasis: Urban Transmission in the Bonon Focus (Côte d'Ivoire). *Tropical Medicine and International Health* 10(4): 340-346.
- [9] Mbida JA, Mimpfoundi R, Njiokou F, Manga L, Laveissière C (2009). Distribution and ecology of vectors of human African trypanosomiasis of the savannah type in degraded forest areas in southern Cameroon: the case of the Doumé outbreak. *Bulletin of the Exotic Pathological Society* 102: 101-105.
- [10] Bosson-Vanga AH, Acapovi-Yao G, Kaba D, Dofini F, Coulibaly B, N'dri Louis, Koné M (2012). Infection of *Glossina palpalis palpalis* by trypanosomes along the Comoé River in the ABENGOUROU region (Ivory Coast). *J. Sci. Pharm. Biol* 3(1): 31-37.
- [11] (INSTAT, (2021). The National Institute of Statistics (INSTAT) collects, produces, analyses and disseminates information on the Malian economy and society. <https://www.instat-mali.org>
- [12] Djitéye A, Moloo SK., Foua BIK, Touré M, Boiré S, Bengaly S, Coulibaly E, Diarra M, Traoré D, Ouattara I and Coulibaly Z (1997). Update of data on the distribution of tsetse flies in Mali. *Revue Elev. Med. Vet. Pays Trop* 50 (2): 126-132.
- [13] Traoré A, Yaro AS, Bass B, Diarra B and Koné T (2019). Effect of deltamethrin and diminazene acetonate treatments on tsetse flies and cattle as a prelude to a tsetse eradication campaign in the Niger River Basin. *Am. J. innov. res. appl. sci.* 9 (4): 320-327
- [14] Bass B, Traoré D, Maiga BM, Bengaly S, Diakité B, Koné F (2014). Impact of climate change on the spatial distribution of tsetse flies in the Bougouni circle in Mali. *Malian Journal of Infectiology and Microbiology*.
- [15] Salou E, Rayaissé JB, Laveissière C, Sanon A & Solano P (2012). Interactions between tsetse species in Burkina Faso. *Parasite* 19: 217-225
- [16] Itard J (1968). Entomological survey in the savannah region (Republic of Togo). *Rev. Elev. Med. Vet. Trop. Countries* 21(3): 375.85
- [17] Darchen R (1978). Populations of *Agelena consociata* Denis, a social spider, in the Gabonese primary forest. Their distribution and density. *Annales de sciences naturelles, Zoologie* 14 (2): 19-26.
- [18] Solano P, Bouyer J, Itard J, Cuisance D 2008? (2010). The cyclical vectors of trypanosomes. *Infectious and parasitic diseases of livestock* (13): 155-183.
- [19] Mbida JA, Mimpfoundi R, Njiokou F, Manga L, Laveissière C (2009). Distribution and ecology of vectors of human African trypanosomiasis of the savannah type in degraded forest areas in southern Cameroon: the case of the Doumé outbreak. *Bulletin of the Exotic Pathological Society* (102): 101-105.
- [20] Buxton PA (1955). The natural history of tsetse flies. An account of the biology of the genus *Glossina* (Diptera). London, UK: London School of Hygiene and Tropical Medicine
- [21] (PATTEC, 2014 unpublished)
- [22] Soberón J, Peterson AT (2005). Interpretation of models of fundamental ecological niche and species distribution areas. *Biodiversity Informatics.* (2):1-10
- [23] Traoré A, Bass B, Yaro AS, Kéita YF, Ballo S, Ly B, and Sodio BA (2023). The influence of the vegetation index on the apparent density of tsetse flies in the communes of Mandé and Kita, Mali. *Innovative Space of Scientific Research Journals.* 66 (2): 279-288
- [24] Frezil JL, Carnevale P (1976). Use of dry ice for the capture of *Glossina fuscipes quanzensis* Pires, 1948, with the Chailier Laveissière trap. *Epidemiological consequences. ORSTOM Notebooks, Medical Entomology and Parasitology Series,* 14): 225 - 233.
- [25] Laveissière C, Hervouët JP (1981). Tsetse fly population and space occupation. Preliminary entomological survey in the Lobo region (Ivory Coast). *ORSTOM Notebooks, Medical Entomology and Parasitology Series,* 19: 247-260.
- [26] Gouteux JP, Kiéno JP (1982). Observations on tsetse flies in a forest focus of human trypanosomiasis in Côte d'Ivoire. 5. Population of some characteristic biotopes: Plantations, forests and forest galleries, in the rainy season. *ORSTOM Notebooks, Medical Entomology and Parasitology Series,* 20: 41-61.
- [27] Zinga Koumba CR, Bouyer J, Mavoungou JF, Acapovi-Yao GL, Kohagne Tongué L, Mbang Nguema OA, Obame Ondo KP, Mutambwe S (2013). Assessment of the diversity of hematophagous diptera in a clearing.
- [28] Zinga Koumba CR, Mbang Nguema OA, Kohagne TL, Acapovi Yao GL, Obame OKP, Mutambwe S, Mavoungou JF (2014). Contribution to the evaluation of the diversity of biological vectors of Human African Trypanosomiasis and their daily activity in the Ivindo National Park (North-East Gabon). *Journal of Applied Biosciences,* 80:7060-707.