

Original Research Article

Non alcoholic fatty liver disease examined by FIBROSCAN-AST (FAST) score and the risk of colorectal carcinoma- A case controlled study

ABSTRACT

Background: Non-alcoholic fatty liver disease (NAFLD) is an entity with its evident role in systemic inflammation and tumorigenesis. The association between colorectal cancer (CRC) and NAFLD identified by the FibroScan-AST (FAST) score allows the simultaneous assessment of steatosis and fibrosis in CRC patients. However, there is a lack of studies to date.

Objective: To assess the association between the risk factor of NAFLD identified by FAST scoring in colorectal cancer.

Methods: This was a case-control study where the case group involves patients with colorectal cancer before initiating chemotherapy and the elapsed time between the Vibration-controlled transient elastography exam by Fibroscan and blood collection is less than 6 months whereas the control group without any documented chronic liver disease or prior malignancies. The demographic and clinical details of the patients were recorded for each patient. The quantitative values of steatosis and fibrosis were measured according to controlled attenuation parameter (CAP) and liver stiffness measurement (LSM), both measured using FibroScan and FAST score calculated. Statistical analyses was performed using SPSS software of version 25.

Results: The mean age was 53.7 years in the case group and 50.8 years in the control group. There were 37 males and 16 females in the case group, whereas 33 males and 20 females in the control group. About 77.3 % of CRC and 71% of control population had steatosis, defined by CAP ≥ 238 dB/m ($p=0.23$). About 68% of the case group and 50% of the control group had fibrosis, defined by LSM ≥ 7.5 kPa ($p=0.02$). With rule-in cut off value for FAST score as ≥ 0.78 , 63.4% of the case group and 36.6% of the control group had NASH with a statistically significant of p-value of 0.045. Overall, it was found that 61.9% of the CRC patients shown positive for NAFLD whereas 30.1% of the patients shown negative for NAFLD. The odds ratio for association of NASH in CRC patients was 3.12, CI 1.27-7.58, $p=0.0015$.

Conclusion: It was noted that there was a relationship between CRC and NAFLD with 61.9% of the colorectal cancer patients shown positive for NAFLD detected by FAST scoring.

Keywords: Non alcoholic fatty liver disease, Colorectal carcinoma, FibroScan, FAST scoring, Correlation

1. INTRODUCTION

The entity of non-alcoholic fatty liver disease (NAFLD) comprises a spectrum from simple steatosis to non-alcoholic steatohepatitis (NASH), and it can progress to liver fibrosis, cirrhosis, and hepatocellular carcinoma. The global prevalence of NAFLD is estimated to be 25% [1], with a higher prevalence seen in the Middle East and South America and the lowest in Africa. The prevalence of NASH is estimated to be 1.5%–6.5% [1]. The prevalence of adult NAFLD in India has been reported between 6.7% and 55.1% [2]. Out of all cases with an asymptomatic elevation of liver enzymes, NAFLD may be responsible for almost one-third [3]. Furthermore, explant histology data from liver transplant centres suggest that two-third of the patients with 'cryptogenic' cirrhosis had NAFLD [4].

Colorectal cancer is the third most prevalent cancer and the second most common cause of cancer-related mortality worldwide, respectively [5]. Colorectal adenoma is 1.5 times higher in patients with NAFLD than in the general population [6,7]. This implies that the importance of NAFLD in colorectal tumorigenesis is increasing. A previously postulated mechanism of the relationship between NAFLD and colorectal tumorigenesis is that the pro-inflammatory status, presence of metabolic syndrome, increased insulin resistance, and decreased adiponectin in patients with NAFLD may be associated with an increased risk of colorectal adenoma [8]. The NAFLD, regarded as a chronic systemic inflammatory state characterized by increased inflammation. This chronic inflammatory condition or secretory molecules such as reactive oxygen species, interleukin (IL)-6, and tumor necrosis factor (TNF)- α could be associated with the occurrence of colorectal adenoma [6]. However, the exact mechanism underlying the close association between NAFLD and colorectal tumorigenesis remains unclear.

Liver biopsy is considered as the gold standard diagnostic tool for evaluating hepatic steatosis and fibrosis. However, it is an invasive technique and can cause complications, such as massive bleeding. Abdominal ultrasonography is the standard modality for evaluating NAFLD; however, it is a subjective procedure and thus has limitations due to inter-observer variation [9]. FibroScan is a novel modality for evaluating fatty liver disease and has the advantages of convenience, reproducibility, and non-invasive [10]. It measures a novel, non-invasive value called the controlled attenuation parameter (CAP) by acquiring a considerable part of the liver volume. The accuracy of CAP in the evaluation of hepatic steatosis has been examined in many studies, including the studies of NAFLD [11].

Recently, Newsome et al. derived the FibroScan-AST (FAST) score in a cohort of patients with NAFLD from England and validated it in multiple international cohorts. The FAST score clearly demonstrated an impressive AUROC of 0.74–0.95 for detecting patients with NASH, elevated NAFLD activity score (NAS ≥ 4) and significant fibrosis ($\geq F2$) on liver biopsy. By incorporating serum AST with FibroScan-derived parameters of LSM and CAP, the FAST score allows the simultaneous assessment of steatosis (CAP), inflammation (AST) and fibrosis (LSM) [12]. However, no studies on the relationship between NAFLD diagnosed by FibroScan-AST (FAST) score and colorectal cancer have been conducted to date. Thus, this study aimed to evaluate the relationship between incidence of colorectal cancer and NAFLD detected by FAST scoring, as a risk factor.

2. METHODOLOGY

This was a case-control study that was conducted in the Department of medical gastroenterology at the Government Super-specialty Hospital, Tirunelveli, Tamil Nadu from January 2023 to January 2024. The inclusion criteria were individuals over 18 years of age with biopsy proven colorectal cancer before initiating chemotherapy and the elapsed time

between the VCTE exam by Fibroscan and blood collection is less than 6 months. The exclusion criteria were the presence of liver metastasis, history of inflammatory bowel disease and history of chronic hepatitis B and C, liver cirrhosis of any cause, alcoholic liver disease, or autoimmune hepatitis. Control group included adults over 18 years of age attending the gastroenterology outpatient ward without any documented chronic liver disease or prior malignancies. The scientific and ethical committee of the institution approved the study and all the patients included in the study provided written informed consent.

The sample size was calculated in openepi.com using Fleiss/ Kelsay sample size formula for case control studies. Keeping the alpha error as 0.05 (confidence level = 95%), power as 80%, ratio of cases to control as 1 and with Odd ratio of NAFLD in Colorectal CA as 3.31 (from previous study), the total sample size calculated was 106 (53/group).

The demographic details such as age, sex, body-mass index (BMI), and presence of diabetes, hypertension, and hypercholesterolaemia were recorded for each patient. Smoking and drinking habits were documented. The quantitative values of steatosis and fibrosis were measured according to CAP and liver stiffness measurement (LSM), respectively, both measured using FibroScan 502 Touch devices equipped with M probe (Echosens, Paris, France). Patients were asked to fast for at least 3 hours before the examination. Patients were placed in the supine position with their right arm fully abducted, and measurements done by scanning the right liver lobe through an intercostal space. CAP is an average estimate of ultrasound attenuation at 3.5 MHz and is expressed in dB/m. LSM by Vibration-controlled transient elastography (VCTE) is an average estimate of stiffness (Young's modulus) at a shear wave frequency of 50 Hz and is expressed as kPa. Only examination results with at least ten valid individual measurements were deemed valid.

The severity of steatosis were graded as normal, mild, moderate, and severe according to the CAP score defined by the cut-off value of FibroScan; these were marked as normal, S1, S2, and S3, respectively. The cut-off values of the CAP score were taken as ≥ 238 dB/m, ≥ 260 dB/m and ≥ 290 dB/m for S1, S2 and S3, respectively. Under the value of S1 grade were described as normal. The grade of fibrosis was defined by the cut-off values of LSM by Fibroscan: the values are ≥ 7.5 kPa, ≥ 10 kPa and ≥ 14 kPa for F2, F3 and F4, respectively. The value of <7.5 kPa was described as F0-1.

All patients undergone routine blood tests including a complete haemogram, liver function test, lipid profile and viral markers. USG/CT abdomen reports were included in the data. The FAST score was calculated using the equation:

$$\frac{[e^{(-1.65+1.07 \times \ln(\text{LSM})+2.66 \times 10^{-8} \times \text{CAP}^3-63.3 \times \text{AST}^{-1})}]}{[1+e^{(-1.65+1.07 \times \ln(\text{LSM})+2.66 \times 10^{-8} \times \text{CAP}^3-63.3 \times \text{AST}^{-1})}]}$$

as carried out by Newsome et al. [12] The optimal rule-out (FAST: ≤ 0.55) and rule-in (FAST: ≥ 0.78) cut-offs were applied.

Statistical analysis

Continuous variables were expressed as the means with standard deviations or as percentages, as appropriate. Categorical variables were expressed as numbers or as percentages and were compared using student 't' test or Fishers exact test, as appropriate. The comparison of the association between CAP, LSM, AST and FAST score values between the case and control groups was done by Mann Whitney U test. All statistical analyses performed using the Statistical Package for Social Sciences (SPSS) software of version 25. Statistical significance was defined as a two-sided p value of <0.05 .

3. RESULTS

Basic characteristics and risk factors of study participants

The mean age was 53.7 years in the case group and 50.8 years in the control group. Also, the mean height and weight in the case group was 162.3 cms and 63.3 kg while it was 160.6 cms and 63.2 kg in the control group respectively. In addition, the mean BMI was 24.1 in the CRC group and 24.5 in the control group (Table 1). There were 37 males and 16 females in the case group, whereas 33 males and 20 females in the control group. The presence of comorbidities such as hypertension, diabetes mellitus, dyslipidemia and social history were described in Table 2.

Table 1: Comparison of baseline characteristics between the case and control groups

Variables	Control (n=53)		Case (n=53)		t value	p value
	Mean	SD	Mean	SD		
Age (years)	50.8	10.1	53.7	10.2	1.46	0.147
Height (cm)	160.6	4.7	162.3	5.5	1.72	0.087
Weight (Kg)	63.2	5.2	63.3	6.3	0.05	0.96
BMI (Kg/m ²)	24.5	2.2	24.1	1.8	1.36	0.176

Table 2: Comparison of various risk factors between case and control groups

Variables		Control		Case		Fisher's statistic	p value
		n	%	n	%		
Gender	Female (n=36)	20	55.6	16	44.4	0.673	0.539
	Male (n=70)	33	47.1	37	52.9		
Diabetes	No (n=67)	31	46.3	36	53.7	1.01	0.421
	Yes (n=39)	22	56.4	17	43.6		
Hypertension	No (n=71)	37	52.1	34	47.9	0.384	0.681
	Yes (n=35)	16	45.7	19	54.3		
Dyslipidemia	No (n=61)	33	54.1	28	45.9	0.965	0.432
	Yes (n=45)	20	44.4	25	55.6		
Smoking	No (n=67)	32	47.8	35	52.2	0.365	0.687
	Yes (n=39)	21	53.8	18	46.2		
Alcoholism	No (n=62)	30	48.4	32	51.6	0.155	0.844
	Yes (n=44)	23	52.3	21	47.7		

Comparison of CAP, LSM, AST and FAST score values between the case and control groups

The median CAP value was 283 dB/m in the case group and 290 dB/m in the control group (Figure 1). Further, the median LSM values were 13 kPa and 8 kPa for the case and control groups, respectively which was statistically significant (P=0.002) (Figure 2). The median AST was higher in the case group of 76 IU/L and 60 IU/L in the control group which was also statistically significant (p=0.013) (Figure 3). There were increased median FAST score in the CRC group of 0.78 and for the control group of 0.52 which was statistically significant of P value of 0.004.

Table 3: Comparison of CAP, LSM, AST and FAST score values between the case and control groups

Parameters	Control (n=53)	Case (n=53)	Mann	p value
------------	----------------	-------------	------	---------

	Median	IQR	Median	IQR	Whitney U test	
CAP (dB/m)	290	217 - 332	283	241 - 315	1273.5	0.408
LSM (kPa)	8	5 - 11	13	7 - 49	1895.5	0.002*
AST (IU/L)	60	46 - 93	76	62 - 92	1796.5	0.013*
FAST score	0.52	0.32 - 0.78	0.78	0.5-0.9	1861.5	0.004*

*P<0.05, significant




Figure 1. Comparison of median CAP value between case and control group




Figure 2: Comparison of median LSM value between case and control group




Figure 3: Comparison of median AST values between case and control group

Severity of hepatic steatosis and fibrosis between case and control groups

About 77.3 % of CRC (41/53) and 71% of control population (38/53) had steatosis, defined by CAP ≥ 238 dB/m ($p=0.23$). About 68% of the case group (36/53) and 50% of the control group (27/53) had fibrosis, defined by LSM ≥ 7.5 kPa ($p=0.02$) (Table 4).

Table 4: Comparison of hepatic steatosis and fibrosis between case and control groups

Parameter		Control		Case		Fisher's statistic	df	p value
		n	%	n	%			
Hepatic steatosis	Normal (n=28)	15	53.6	12	44.4	4.3	3	0.23
	S1 (n=9)	2	22.2	8	77.8			
	S2 (n=19)	9	47.4	10	52.6			
	S3 (n=50)	27	54	23	46			
Hepatic fibrosis	F0-F1 (n=43)	26	60.5	17	39.5	9.42	3	0.02*
	F2 (n=19)	12	63.2	7	36.8			
	F3 (n=7)	4	57.1	3	42.9			
	F4 (n=37)	11	29.7	26	70.3			

*P<0.05, significant

Association of NAFLD in color cancer

With rule-in cut off value for FAST score as ≥ 0.78 , 63.4% of the case group and 36.6% of the control group had NASH with a statistically significant of p-value of 0.045. Also, 52.2% of the patients in the case group had intermediate NASH while 47.8% of patients in the control group had intermediate NASH (Table 5). Overall, it was found that 61.9% of the CRC patients shown positive for NAFLD whereas 30.1% of the patients shown negative for NAFLD (Figure 4). The odds ratio for association of NASH in CRC patients was 3.12, CI 1.27-7.58, $p=0.0015$ (Table 6).

Table 5: Comparison of NAFLD occurrence (based on FAST score category) between case and control groups

Fast score	Control	Case	Fisher's	p value
------------	---------	------	----------	---------

category	n	%	n	%	statistic	
NAFLD negative	27	64.3	15	35.7	6.42	0.045*
Indeterminate	11	47.8	12	52.2		
NAFLD positive	15	36.6	26	63.4		

*P<0.05, significant




Figure 4: Association of NAFLD with colon cancer observed

Table 6: Calculation of odd ratio for association of NAFLD in colon cancer patients

2X2 table	NAFLD Positive		NAFLD negative		Odds ratio	95% Confidence interval	P value
	n	%	n	%			
Colon cancer present (n=41)	26	61.9	15	30.1	3.12	1.3 to 7.5	0.0015*
Colon cancer absent (n=42)	15	35.7	27	64.3			

*P<0.05, significant

4. DISCUSSION

Non-alcoholic fatty liver disease (NAFLD) has become the most common chronic liver disease worldwide. Non-alcoholic steatohepatitis (NASH) is the more active form of NAFLD and is characterised by the presence of hepatic steatosis, inflammation, and hepatocyte ballooning. NASH is emerging as one of the leading causes of cirrhosis, cirrhotic complications, hepatocellular carcinoma (HCC) and liver-related death [13]. Studies have shown that, apart from HCC, NAFLD is also associated with other types of gastrointestinal cancers, including colorectal cancer (CRC) [14]. Around 18%–24% and 9%–20% of advanced colon neoplasm occurs in NAFLD men and women respectively compared to 9%–5% and 3%–8% in non-NAFLD men and women respectively [7].

In this study, the mean age in the case group and the control group was 53.7 years and 50.8 years which highlighted that disease prevalence across the middle age groups. Further, there were majority of male patients in both groups which showed the male predominance in

this study. According to FibroScan, the population was first classified as having steatosis. The steatosis group consisted of people with S1 or more steatosis, whereas the normal group consisted of subjects with normal values. Age, sex, height, weight, smoking, comorbidities, smoking and alcohol intake did not significantly differ across groups. This was inline with Kim KW et al., however, they found significant difference noted in BMI and alcohol intake among CRC patients with NAFLD [16]. In addition, there were statistically significant higher median AST levels in the case group which correlates with the existence of NAFLD.

Moreover, one of the major constraints in understanding NAFLD is the absence of a disease specific biomarker. By incorporating serum AST with fibroscan- derived parameters of LSM and CAP, the FAST score allows the simultaneous assessment of steatosis, inflammation and fibrosis. In our study, with rule-in FAST score as ≥ 0.78 , 63.4% of the colorectal cancer group and 36.6% of the control group had NASH with a statistically significant p-value of 0.045. It suggests that steatohepatitis may be associated with colorectal tumorigenesis. It is evident from the previous studies that fat accumulation in hepatocytes can lead on to an inflammatory state in the liver [15]. Hepatic steatosis can further accelerate or induce disarray of the gastrointestinal tract via pro-inflammatory cytokines that promote tumorigenesis in the colon [16].

Fibroscan also has the advantages of providing more information than does liver biopsy by acquiring images of a larger liver volume and being non-invasive, requiring only a probe to touch the skin. It helps in early detection of the disease and identifying patients with NASH who are at high risk of progressive disease who are likely to be benefited by the emerging therapies in NASH [17]. Fibroscan and FAST scoring system can be utilised as a good screening test for NAFLD in the general population and thereby, identify patients in their early disease course. This may help to actively intervene before they develop complications of the disease, including various malignancies.

Our study points towards the importance of NAFLD diagnosis in its earlier stages and to consider the utilisation of Fibroscan and its derivative Scoring systems in the screening programs. However, our study had some limitations. First, this was a single-centre study, and thus, the results may have limited generalizability. Secondly, the study population was not large enough and we could not evaluate other confounding factors such as abdominal visceral adiposity and show more clear statistical values in various aspects of colorectal neoplasia. Third, there was very few data about the stages of fibrosis and its statistical significance in the tumorigenesis process. Finally, we choose only one of the reference values for Fibroscan that have been previously reported in the validity study. There are various reference values of CAP and LSM for assessing hepatic steatosis and fibrosis. It means that the cut off values could vary due to many factors such as countries, ethnics, statistical methods, underlying clinical characteristics and so on. Further large scale prospective studies are required to conform our results and the effect of managing hepatic steatosis in reducing colorectal tumorigenesis.

5. CONCLUSION

In this study, it was found that 61.9% of the colorectal cancer patients shown positive for NAFLD detected by FAST scoring. Therefore, clinicians should be aware of the relationship between colorectal cancer and NAFLD assessed using FAST scoring and its clinical implications.

REFERENCES

1. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease-Meta-analytic assessment of prevalence, incidence, and outcomes. *Hepatology*. 2016;64:73–84.
2. Shalimar, Elhence A, Bansal B, Gupta H, Anand A, Singh TP, Goel A. Prevalence of Non-alcoholic Fatty Liver Disease in India: A Systematic Review and Meta-analysis. *J Clin Exp Hepatol*. 2022;12(3):818-829.
3. Madan K, Batra Y, Panda SK, et al. Role of polymerase chain reaction and liver biopsy in the evaluation of patients with asymptomatic transaminitis: implications in diagnostic approach. *J Gastroenterol Hepatol*. 2004;19:1291–1299.
4. Nayak NC, Jain D, Vasdev N, Gulwani H, Saigal S, Soin A. Etiologic types of end-stage chronic liver disease in adults: analysis of prevalence and their temporal changes from a study on native liver explants. *Eur J Gastroenterol Hepatol*. 2012;24:1199–1208.
5. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394–424.
6. Hwang ST, Cho YK, Park JH, Kim HJ, Park DI, Sohn CI, et al. Relationship of non-alcoholic fatty liver disease to colorectal adenomatous polyps. *J Gastroenterol Hepatol*. 2010;25(3):562–7.
7. Wong VW, Wong GL, Tsang SW, Fan T, Chu WC, Woo J, et al. High prevalence of colorectal neoplasm in patients with non-alcoholic steatohepatitis. *Gut*. 2011; 60(6): 829–36.
8. Divella R, Mazzocca A, Daniele A, Sabba C, Paradiso A. Obesity, Nonalcoholic Fatty Liver Disease and Adipocytokines Network in Promotion of Cancer. *Int J Biol Sci*. 2019;15(3):610–6.
9. Lee DH. Imaging evaluation of non-alcoholic fatty liver disease: focused on quantification. *Clin Mol Hepatol*. 2017;23(4):290–301.
10. Singh S, Muir AJ, Dieterich DT, Falck-Ytter YT. American Gastroenterological Association Institute Technical Review on the Role of Elastography in Chronic Liver Diseases. *Gastroenterology*. 2017;152(6):1544–77.
11. Karlas T, Petroff D, Garnov N, Bohm S, Tenckhoff H, Wittekind C, et al. Non-invasive assessment of hepatic steatosis in patients with NAFLD using controlled attenuation parameter and 1H-MR spectroscopy. *PLoS One*. 2014;9(3):e91987.
12. Newsome PN, Sasso M, Deeks JJ, et al. FibroScan-AST (FAST) score for the non-invasive identification of patients with non-alcoholic steatohepatitis with significant activity and fibrosis: a prospective derivation and global validation study. *Lancet Gastroenterol Hepatol*. 2020;5:362–373.
13. Huang DQ, Singal AG, Kono Y, et al. Changing global epidemiology of liver cancer from 2010 to 2019: NASH is the fastest growing cause of liver cancer. *Cell Metab*. 2022; 34: 969-977.
14. Muhidin SO, Magan AA, Osman KA, et al. The relationship between nonalcoholic fatty liver disease and colorectal cancer: the future challenges and outcomes of the metabolic syndrome. *J Obes*. 2012; 2012:637538.
15. Liu Q, Bengmark S, Qu S. The role of hepatic fat accumulation in pathogenesis of non-alcoholic fatty liver disease (NAFLD). *Lipids Health Dis*. 2010;9:42.

16. Kim KW, Kang HW, Yoo H, Jun Y, Lee HJ, Im JP, et al. Association between severe hepatic steatosis examined by Fibroscan and the risk of high-risk colorectal neoplasia. *PLoS One*. 2022;17(12):e0279242.
17. Singh S, Muir AJ, Dieterich DT, Falck-Ytter YT. American Gastroenterological Association Institute Technical Review on the Role of Elastography in Chronic Liver Diseases. *Gastroenterology*. 2017;152(6):1544–77.

UNDER PEER REVIEW