

Original Research Article

Coat-protein based characterization of yellow mosaic virus infecting blackgram in northern Karnataka

ABSTRACT

Aims: Coat-protein based characterization of yellow mosaic virus infecting blackgram.
Place and Duration of Study: The study was carried out during *Kharif* 2023 at College of Agriculture, Vijayapur.
Methodology: The whole genomic DNA was isolated from the leaf tissues of both healthy blackgram plants and plants infected with the yellow mosaic virus by using a modified CTAB approach. Specific primers for yellow mosaic viruses were tried to amplify coat protein area of blackgram YMV.
Results: Sequence analysis revealed that the CP gene of the begomovirus under study shared 99.8 per cent similarity with MYMV Shivamogga isolate (OM106035.1) at the nucleotide level. When comparing the deduced amino acid sequences of individual proteins from YMV infecting blackgram at Vijayapur with those of other begomoviruses, the highest identity was found with the MYMV Shivamogga isolate (OM106035.1), showing 84.59 per cent similarity. A phylogenetic tree constructed based on the full-length coat protein gene sequence of MYMV, along with 37 other *Geminivirus* sequences, formed two major clusters in which MYMV-Vijayapur isolate appeared in Cluster I.
Conclusion: The results of the current study, based on similarities in coat protein sequence at both the nucleotide and amino acid levels, as well as phylogenetic analysis, confirmed the prevalence of MYMV strain rather than MYMIV, HgYMV and DoYMV in blackgram in northern parts of Karnataka.

Keywords: Coat protein, Geminivirus, Blackgram, Yellow mosaic virus

1. INTRODUCTION

Blackgram (*Vigna mungo* L.), commonly known as urdbean in India, belongs to the *Leguminosae* family. It is a significant short-duration, self-pollinating legume. Renowned for its rich nutritional profile, blackgram contains proteins (25-26%), carbohydrates (60%), fats (1.5%), minerals, amino acids and vitamins. Consequently, it serves as an economical protein source for vegetarians. It also aids in the process of atmospheric nitrogen fixation into the soil. This highly valued pulse crop is cultivated across various regions of India (Archana et al., 2018).

The optimal climate for blackgram cultivation ranges between 27-30°C, with moderate rainfall and loamy soil that has high water-holding capacity. Its maturity typically occurs within 90-100 days (Swaminathan et al., 2023). This crop is grown during both *Kharif* and summer seasons, particularly in central Indian conditions. Blackgram has been found to be beneficial in reducing elevated cholesterol levels. It has gained prominence in Indian cuisine, especially in the preparation of dishes such as Dal, Idli, Vada, Dosa and Papad (Marabi et al., 2017).

Blackgram originates from the Indo-Burma region and stands as one of the crucial leguminous crops extensively cultivated in major tropical and subtropical nations worldwide (Biswas et al., 2008). In India, it ranks as the third most significant pulse crop following chickpea and pigeon pea. Blackgram demonstrates drought tolerance, allowing for biannual cultivation and integration into India's crop rotation scheme (Salam et al., 2009).

India holds the title of the world's largest producer of blackgram, with cultivation spanning approximately 40.02 lakh hectares, resulting in a production of 26.31 lakh tonnes and a productivity rate of 657 kg/ha. The primary blackgram cultivating states include Andhra Pradesh, Bihar, Karnataka, Maharashtra, Madhya Pradesh, Orissa, Rajasthan, Tamil Nadu, Uttar Pradesh and West Bengal. Andhra Pradesh leads the chart as the top blackgram producing state, with cultivation across approximately 3.45 lakh hectares, with a production of 4.31 lakh tonnes and achieving a productivity of 1249 kg/ha. In Karnataka, it is cultivated across roughly 0.71 lakh hectares, with a production of 0.33 lakh tonnes and productivity rate of 466 kg/ha. Notable blackgram cultivating districts in Karnataka include Kalburgi, Bidar, Mysuru, Belagavi, Dharwad, Bagalkot and Vijayapur (Anon., 2023).

The low yield of blackgram is attributed to various factors, primarily due to attacks by numerous insects (such as pod borers and sucking pests) and diseases (including *Alternaria* leaf spot, *Cercospora* leaf spot, Powdery mildew and viral diseases), as well as other abiotic factors from seedling to maturity stages. Among these constraints, Yellow Mosaic Disease (YMD) caused by Mungbean Yellow Mosaic Virus (MYMV) stands out as the most serious disease and a significant bottleneck for blackgram cultivation and production (Biswas et al., 2012).

In the realm of pulses, yellow mosaic disease is attributed to four distinct begomoviruses based on viral nucleotide sequence identity. These include Mungbean Yellow Mosaic Virus (MYMV), Mungbean Yellow Mosaic India Virus (MYMIV), Horsegram Yellow Mosaic Virus (HgYMV) and Dolichos Yellow Mosaic Virus (DoYMV) (Mishra et al., 2020). YMV is a single-stranded DNA virus belonging to the genus *Begomovirus* within the family *Geminiviridae* (Malathi and John, 2008). In India, Nariani (1960) initially documented MYMV in mungbean fields at the Indian Agricultural Research Institute (IARI) in New Delhi during the 1950s.

YMV's belong to the genus *Begomovirus*, are transmitted by the whitefly (*Bemisia tabaci*), and possess bipartite genomes. DNA-A encodes proteins essential for encapsidation, replication and transcription regulation, while DNA-B encodes proteins responsible for intra and intercellular movement. The DNA-A and DNA-B components of begomoviruses share a highly conserved intergenic common region (CR), which contains a stem-loop structure. Within this loop lies the invariant nononucleotide motif (TAATATTAC), serving as the origin of viral strand replication (Bhaskara et al., 2015).

The disease is transmitted by the hemipteran whitefly, *Bemisia tabaci*, in a circulative and persistent manner. This vector can acquire and transmit the virus within a minimum acquisition and inoculation access period of 10-15 minutes each. The optimal acquisition access period (AAP) ranges between 4-6 hours, while the inoculation access period (IAP) is typically 4 hours. Mungbean Yellow Mosaic Virus (MYMV) is highly destructive and primarily spreads through whiteflies. This virus is not transmitted through seeds, soil or sap (Nair and Nene, 1973).

The initial symptoms of the disease manifest as small yellow flecks in the veinlets of young emerging leaves. Subsequently, developing leaves display more prominent and irregular yellow and green patches, alternating with each other. There is not much impact on leaf size. However, in susceptible genotypes, affected plants produce only a few pods that are smaller

in size and deformed (Verma et al., 1992). The degree of yield loss largely depends on the age of the plant at the time of infection and the severity of the disease (Naimuddin et al., 2011). The YMD causes 85-100 per cent yield loss in the plants that are infected at the seedling stage (Nene, 1972).

The goal of the current study was to identify and characterize the coat protein gene of the blackgram yellow mosaic virus at the molecular level. Since, not much work has been done on the molecular characterization of the virus and to determine whether a single strain or mixed strains of YMV are prevalent in northern Karnataka, the present study was initiated.

2. MATERIAL AND METHODS

The present investigation on blackgram yellow mosaic virus was carried at College of Agriculture, Vijayapur, University of Agricultural Sciences, Dharwad, Karnataka, India during *Kharif*, 2023.

Extraction of total DNA using modified Cetyl Trimethyl Ammonium Bromide (CTAB) method

Blackgram plants exhibiting severe yellow mosaic symptoms were collected from the field (Fig. 1). Total genomic DNA was extracted from leaf tissues of both healthy plants and those infected with yellow mosaic virus disease, using the modified CTAB method developed by Rouhibakshet *al.* (2008). The quality and quantity of the DNA were assessed through agarose gel electrophoresis and spectrophotometry.

Amplification of the coat protein gene of blackgram yellow mosaic virus

To amplify the coat protein of the Blackgram yellow mosaic virus, specific primers documented in the literature for yellow mosaic viruses were used to target the coat protein region, approximately 1,000 bp in size (Naimuddin and Akram, 2010). Primers specific to MYMIV (NM1/AV1-F: GTATTTGCAKCAWGTTC AAGA and NM2/AV1-R: AGGDGTCATTAGCTTAGC), MYMV (MYMV-CPF: ATGGG(T/G)TCCGTTGTATGCTTG and MYMV-CPR: GCGTCATTAGCATAGGCAAT) and HgYMV (HgYMV-CPF: ATGCTTGCAATTAAGTACTTGCA and HgYMV-CPR: TAGGCGTCATTAGCATAGGCA) were employed to amplify the coat protein gene of blackgram yellow mosaic virus. PCR was performed using these primers, with samples taken along with a positive control (CTAB-extracted MYMV DNA) and a negative control (healthy DNA). After preparing the PCR mixture, the DNA template was added, the tubes were briefly spun, and placed into the wells of a thermal cycler (Eppendorf thermocycler) for amplification.

PCR was performed in Thermocycler programmed for one step of initial denaturation at 94°C for 2 min and 35 cycles of denaturation at 94°C for 1 min, annealing at 55°C for 2 min for primers NM1/AV1-F/ NM2/AV1-R, MYMV-CP-F/MYMV-CP-R, HgYMV CP-F/HgYMV-CP-R and extension at 72°C for 3 min, followed by one step of final extension at 72°C for 10 min. PCR mixture reaction mixture volume of 25 µl that contained PCR buffer - 12.5 µl; SDW - 4 µl; forward and reverse primers (20 pmole/µl) - 2 µl each; Taq Polymerase (3 U/µl) - 0.5 µl; DNA - 2 µl; DNTPs - 2 µl. After completion of the reaction, the products were kept at 4°C prior to gel analysis. PCR products were subjected to electrophoresis in 1% agarose at 60 V for 1-2 hr in Electrophoresis system.

Sequence analysis of coat protein gene and phylogenetic tree construction

The 1000 bp amplicon amplified from YMV infected blackgram samples using MYMV specific primers were sent to direct sequencing. The sequence obtained was searched in NCBI- BLAST to know the sequence similarity with other begomoviruses. Sequence identity matrix tool, Bio Edit software were used to assemble and analyze the data. Multiple alignments were made and sequence was obtained by ClustraWprogramme. The phylogenetic tree was constructed based on the available sequences using MEGA 11 Software (Valli et al., 2022).

3. RESULTS AND DISCUSSION

The modified C-TAB method incorporated several key chemicals, including 100 mM (0.1 M) Tris-HCl (pH 8.0), 20 mM (0.02 M) EDTA (pH 8.0), 1 M NaCl, CTAB (Cetyl Trimethyl Ammonium Bromide) (2%), SDS (1%), PVP 40 (2%), sodium sulfate (0.65%), β -mercaptoethanol (0.2%) and 1.4-2.0 M sodium chloride. These components were crucial for isolating high quality DNA. This modified C-TAB method was utilized to isolate viral DNA from both infected and healthy blackgram plants (Rouhibakhshet *et al.*, 2008). High quality DNA was successfully obtained from YMV infected blackgram leaves using this technique.

The DNA extracted through the modified C-TAB method was then subjected to Polymerase Chain Reaction (PCR). The PCR effectively amplified the CP-gene from YMV infected blackgram leaf samples. Amplification of the coat protein gene of YMV in blackgram resulted in a DNA fragment of approximately 1000 bp (Fig. 2), while no PCR product was obtained from DNA extracted from healthy samples or water controls.

Sequencing of PCR products containing the CP-gene of YMV using specific primers was performed at Eurofins Genomics India Pvt. Ltd., Bengaluru. The nucleotide sequence of MYMV-Vijayapur isolate was deposited in NCBI database and the accession number obtained was PP470806.1. The results of a BLAST search (<http://www.ncbi.nih.gov/BLAST>) were used to identify sequence homology. The nucleotide sequence of the coat protein gene of the yellow mosaic virus infecting blackgram was compared with selected begomovirus sequences obtained from the NCBI database (Table 1). The sequence data revealed that the CP gene of the begomovirus under study shared 99.80 per cent similarity with MYMV Shivamogga isolate (OM106035.1) at the nucleotide level. When the deduced amino acid sequence of individual proteins of YMV infecting the blackgram at Vijayapur was compared with those of other begomoviruses, the highest identity was found with the MYMV Shivamogga isolate (OM106035.1), showing 84.59 per cent similarity. Based on this sequence comparison, the test isolate was confirmed as mungbean yellow mosaic virus. The nucleotide sequence identity and deduced amino acid sequence identity of MYMV with other begomoviruses ranged from 92.62-99.8 per cent and 70.03-84.59 per cent, respectively.

A phylogenetic tree was constructed based on the full-length coat protein gene sequence of MYMV of blackgram and 37 other *Geminivirus* sequences downloaded from the NCBI GenBank. The phylogenetic neighbor-joining trees and evolutionary analysis were conducted using MEGA 11 with 1,000 bootstrap values and it formed two major clusters. The MYMV-Vijayapur isolate appeared in Cluster I and showed similarity with MYMV-Shivamogga isolate (Fig. 3). Pairwise per cent identities of coat protein nucleotide sequences of MYMV and MYMIV isolates was done by using Sequence Demarcation Tool (SDT) and the analysis result showed that highest identity was with MYMV-Shivamogga isolate (Fig. 4). Based on these sequence comparisons, the Vijayapur test isolate was confirmed as mungbean yellow mosaic virus.

These results are consistent with those of several other researchers. Archana et al. (2018) demonstrated that MYMV infecting blackgram from Mandya (Karnataka) was a strain of the MYMV group, with 95 per cent similarity, rather than MYMIV and HgYMV.

Phylogenetic analysis showed that the blackgram yellow mosaic virus clustered with MYMV isolates, having a nucleotide sequence identity of 94.4 to 98.7 per cent with MYMV and 79 to 80.7 per cent with MYMIV. Amino acid sequence comparisons showed high homology with MYMV isolates. This indicated that the virus was a variant of the Mungbean yellow mosaic virus (MYMV), and not of Mungbean yellow mosaic India virus (MYMIV) (Prema and Rangaswamy, 2018).

4. CONCLUSION

The results of the current study, based on similarities in coat protein sequence at both the nucleotide and amino acid levels, as well as phylogenetic analysis, confirmed the prevalence of MYMV strain rather than MYMIV, HgYMV and DoYMV in blackgram in northern parts of Karnataka.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

- Anonymous.(2023). Area, production, productivity of blackgram in India, Ministry of Agriculture and Farmer welfare, Government of India. (ON2930) www.indiastat.com.
- Archana. S., Venkatesh, P, A., Nagaraju, N.,& Manjunatha, N.(2018). Management of yellow mosaic disease (YMD) of blackgram (*Vigna mungo* L.) in Southern dry zone of Karnataka. Journal of Entomology and Zoology Studies, 6(3), 860-863.
- Bhaskara, R. B. V., Obaiah, S., Prasanthi, L., Sivaprasad, Y., Sujitha, A.,& Giridhara, K. T.(2015).Mungbean yellow mosaic India virus is associated with yellow mosaic disease of blackgram (*Vigna mungo* L.) in Andhra Pradesh, India. Archives of Phytopathology and Plant Protection, 48(4), 345-353.
- Biswas, K. K., Malathi, V. G.,&Varma, A.(2008). Diagnosis of symptom less yellow mosaic begomovirus infection in pigeonpea by using cloned Mungbean yellow mosaic India virus as probe. Journal of Plant Biochemistry and Biotechnology,17(1), 9-14.

- Biswas, K. K., Tarafdar, A., & Biswas, K. (2012). Viral diseases and its mixed infection in mungbean and urdbean: Major biotic constraints in production of food pulses in India. (Ed. Asha S, Sharma B K and Manisha S), Modern trends in microbial biodiversity of natural ecosystem, New Delhi, pp. 301-317.
- Malathi, V. G., & John, P. (2008). Geminiviruses infecting legumes. Characterization, diagnosis and management of plant viruses, 3, 97-123.
- Marabi, R. S., Chaukikar, K., Das, S. B., & Bhowmick, A. K. (2017). Population dynamics of whitefly, *Bemisia tabaci* and incidence of Mungbean yellow mosaic India virus (MYMIV) on blackgram. International Journal of Bio-resource and Stress Management, 8(6), 846-852.
- Mishra, G. P., Dikshit, H. K., Ramesh, S. V., Tripathi, K., Kumar, R. R., Ask, i M., Singh, A., Roy, A., Kumari, N., Dasgupta, U., Kumar, A., Praveen, S., & Nair, R. M. (2020). Yellow mosaic disease (YMD) of mungbean (*Vigna radiata* (L.) Wilczek): Current status and management opportunities, Frontiers in Plant Science, 11, 918.
- Naimuddin & Akram, M. (2010). Detection of mixed infection of begomoviruses in cowpea and their molecular characterization based on CP gene sequences. Journal of Food Legumes, 23, 191-195.
- Naimuddin. K., Mohammad, A., & Gupta, S. (2011). Identification of mungbean yellow mosaic India virus infecting *Vigna mungo* var. *silvestris* L. Phytopathologia Mediterranea, 50 (1), 94-100.
- Nair, N. G., & Nene, Y. L. (1973). Studies on the yellow mosaic of urd bean (*Phaseolus mungo* (L.) caused by mungbean yellow mosaic virus. 1. Transmission studies, Indian Journal of Farm Sciences, 1(1), 62-70.
- Nariani, T. K. (1960). Yellow mosaic of mung (*Phaseolus aureus*). Indian Phytopathology, 13, 24-29.
- Nene, Y. L. (1972). A survey of the viral diseases of pulse crops in Uttar Pradesh. First Annual Report. F.G-In-358, Uttar Pradesh Agricultural University, pp. 1-25.
- Prema, G. U., & Rangaswamy, K. T. (2018). Molecular characterization of coat protein gene of blackgram yellow mosaic virus (BGYMV) from Karnataka, India. International Journal of Current Microbiology and Applied Sciences, 7(7), 2225-2235.

Rouhibakhsh, A., Priya, J., Periasamy, M., Haq, Q. M. I., & Malathi, V. G. (2008). An improved DNA isolation method and PCR protocol for efficient detection of multicomponents of begomovirus in legumes. *Journal of Virological Methods*, 147(1), 37-42.

Salam, S. A., Patil, M. S., & Byadgi, A. S. (2009). IDM of mungbean yellow mosaic disease. *Annals of Plant Protection Sciences*, 17, 157-160.


Swaminathan, C., Surya, R., Subramanian, E., & Arunachalam, P. (2023). Challenges in pulses productivity and agronomic opportunities for enhancing growth and yield in blackgram [*Vigna mungo* (L.) Hepper]: A review. *Legume Research*, 46(1), 1-9.

Valli, N. S., Karthikeyan, G., Harish, S., Sudha, M., & Sharmila, D. (2022). Coat protein characterization of mungbean yellow mosaic virus infecting blackgram (*Vigna mungo* L. Hepper) in Tamil Nadu. *International Journal of Plant and Soil Science*, 34(22), 638-644.

Verma, A., Dhar, A. K., & Mandal, B. (1992). MYMV transmission and control in India, Mungbean yellow mosaic disease, *Proceedings of Shanhua, Tainan AVRDC Taiwan*, pp.8-27.



Fig. 1: Blackgram plants showing typical yellow mosaic disease symptoms



Lane:

M- 1Kb Marker (NEB 1 kb DNA ladder)

Lane 1 - Healthy blackgram leaf DNA

Lane 2, 3, 4 - Specific PCR product of 1000 bp from YMV infected blackgram leaf sample

Lane 5 - Water control

Fig. 2: Amplification of coat protein gene of YMV infecting blackgram using MYMV-CP-F/MYMV-CP-R primer pair

UNDER PEER REVIEW






Fig. 3: Phylogenetic relationship of CP gene of Vijayapur strain of YMV infecting blackgram with other reference strain

MN698293.1_HgYMVBelgaum
 AJ132575.1_MYMV_Tamil_Nadu
 PP738873.1_MYMVCoimbatore
 MK409376.1_MYMVBangalore
 MH885653.1_MYMVBengaluru
 PP470806.1_MYMV_Vijayapur
 OM106035.1_MYMVShivamogga
 OM106038.1_MYMV_Chikkamagaluru
 OM106036.1_MYMVShivamogga
 OM106037.1_MYMVGadag
 DQ389144.1_MYMVGujarat
 OQ123521.1_MYMVDharwad
 MN698275.1_MYMVHyderabad
 MN698273.1_MYMVHyderabad
 JX244175.1_MYMNVietnam
 JX244176.1_MYMNVietnam
 JX244172.1_MYMNVietnam
 AB017341.1_MYMVThailand
 AY271892.1_MYMVCambodia
 OQ732573.1_MYMVTirupati
 MW814715.1_MYMVBagalkot
 KY824799.1_MYMVChamarajanagar
 MN602422.1_MYMVDhaward
 MN602427.1_MYMVBelgaum
 DQ389150.1_MYMVMaharashtra
 AF314530.1_MYMVMaharashtra
 DQ865201.1_MYMVTamil_Nadu
 MW736048.1_MYMNVamakkal
 MW736047.1_MYMNVamakkal
 MW736053.1_MYMVCoimbatore
 MW736043.1_MYMVBangalore
 KC911718.1_MYMVCoimbatore
 ON421034.1_MYMVKanpur
 ON421030.1_MYMVKanpur
 FM242701.1_MYMVPakistan
 MN885479.1_MYMVPakistan
 MN922309.1_MYMVPakistan
 MW736054.1_MYMVMamban



MN698293.1_HgYMVBelgaum
 AJ132575.1_MYMV_Tamil_Nadu
 PP738873.1_MYMVCoimbatore
 MK409376.1_MYMVBangalore
 MH885653.1_MYMVBengaluru
 PP470806.1_MYMV_Vijayapur
 OM106035.1_MYMVShivamogga
 OM106038.1_MYMV_Chikkamagaluru
 OM106036.1_MYMVShivamogga
 OM106037.1_MYMVGadag
 DQ389144.1_MYMVGujarat
 OQ123521.1_MYMVDharwad
 MN698275.1_MYMVHyderabad
 MN698273.1_MYMVHyderabad
 JX244175.1_MYMNVietnam
 JX244176.1_MYMNVietnam
 JX244172.1_MYMNVietnam
 AB017341.1_MYMVThailand
 AY271892.1_MYMVCambodia
 OQ732573.1_MYMVTirupati
 MW814715.1_MYMVBagalkot
 KY824799.1_MYMVChamarajanagar
 MN602422.1_MYMVDhaward
 MN602427.1_MYMVBelgaum
 DQ389150.1_MYMVMaharashtra
 AF314530.1_MYMVMaharashtra
 DQ865201.1_MYMVTamil_Nadu
 MW736048.1_MYMNVamakkal
 MW736047.1_MYMNVamakkal
 MW736053.1_MYMVCoimbatore
 MW736043.1_MYMVBangalore
 KC911718.1_MYMVCoimbatore
 ON421034.1_MYMVKanpur
 ON421030.1_MYMVKanpur
 FM242701.1_MYMVPakistan
 MN885479.1_MYMVPakistan
 MN922309.1_MYMVPakistan
 MW736054.1_MYMVMamban

Fig. 4: Pairwise per cent identities of coat protein nucleotide sequences of MYMV and MYMIV isolates along with MYMV Vijayapur isolate

Table 1: Nucleotide and amino acid sequence identities of coat protein gene of YMV infecting blackgram with other *Geminiviruses*

| Sl. No. | Accession number | Virus species | Geographical origin | Nucleotide sequence identity (%) | Amino acid sequence identity (%) | Host crop |
|---------|------------------|------------------------------------|----------------------|----------------------------------|----------------------------------|----------------------------|
| 1 | OM106035.1 | Mungbean yellow mosaic virus | MYMV-Shivamogga | 99.80 | 84.59 | <i>Vigna radiata</i> |
| 2 | OM106038.1 | Mungbean yellow mosaic virus | MYMV- Chikkamagaluru | 99.20 | 83.08 | <i>Vigna radiata</i> |
| 3 | OM106037.1 | Mungbean yellow mosaic virus | MYMV-Gadag | 99.20 | 83.08 | <i>Vigna radiata</i> |
| 4 | OM106036.1 | Mungbean yellow mosaic virus | MYMV-Shivamogga | 99.10 | 82.78 | <i>Vigna radiata</i> |
| 5 | MN602422.1 | Mungbean yellow mosaic virus | MYMV-Dhaward | 98.90 | 82.87 | <i>Vigna radiata</i> |
| 6 | OQ123521.1 | Mungbean yellow mosaic virus | MYMV-Dharwad | 98.80 | 82.87 | <i>Vigna radiata</i> |
| 7 | MN698275.1 | Mungbean yellow mosaic virus | MYMV-Hyderabad | 98.60 | 81.87 | <i>Vigna radiata</i> |
| 8 | MN698273.1 | Mungbean yellow mosaic virus | MYMV-Hyderabad | 98.60 | 81.87 | <i>Vigna radiata</i> |
| 9 | OQ732573.1 | Mungbean yellow mosaic virus | MYMV-Tirupati | 98.50 | 81.96 | <i>Vigna radiata</i> |
| 10 | AF314530.1 | Mungbean yellow mosaic virus | MYMV-Maharashtra | 98.30 | 80.66 | <i>Glycine max</i> |
| 11 | JX244175.1 | Mungbean yellow mosaic virus | MYMV-Vietnam | 98.20 | 80.97 | <i>Vigna radiata</i> |
| 12 | MW814715.1 | Mungbean yellow mosaic virus | MYMV-Bagalkot | 98.10 | 80.36 | <i>Vigna radiata</i> |
| 13 | JX244176.1 | Mungbean yellow mosaic virus | MYMV-Vietnam | 98.10 | 81.27 | <i>Vigna radiata</i> |
| 14 | JX244172.1 | Mungbean yellow mosaic virus | MYMV-Vietnam | 98.10 | 81.57 | <i>Vigna radiata</i> |
| 15 | DQ865201.1 | Mungbean yellow mosaic virus | MYMV-Tamil Nadu | 98.20 | 81.27 | <i>Vigna aconitifolia</i> |
| 16 | MN602427.1 | Mungbean yellow mosaic virus | MYMV-Belgaum | 98.00 | 81.27 | <i>Vigna radiata</i> |
| 17 | MW736053.1 | Mungbean yellow mosaic virus | MYMV-Coimbatore | 98.00 | 80.73 | <i>Vigna radiata</i> |
| 18 | AJ132575.1 | Mungbean yellow mosaic virus | MYMV-Tamil Nadu | 98.00 | 80.97 | <i>Vigna radiata</i> |
| 19 | DQ389144.1 | Mungbean yellow mosaic virus | MYMV-Gujarat | 99.58 | 83.44 | <i>Glycine max</i> |
| 20 | MK409376.1 | Mungbean yellow mosaic virus | MYMV-Bangalore | 97.90 | 80.66 | <i>Vigna mungo</i> |
| 21 | MW736048.1 | Mungbean yellow mosaic virus | MYMV-Namakkal | 97.90 | 81.57 | <i>Vigna radiata</i> |
| 22 | MW736047.1 | Mungbean yellow mosaic virus | MYMV-Namakkal | 98.00 | 81.27 | <i>Vigna radiata</i> |
| 23 | MW736043.1 | Mungbean yellow mosaic virus | MYMV-Bangalore | 97.80 | 80.36 | <i>Phaseolus vulgaris</i> |
| 24 | AY271892.1 | Mungbean yellow mosaic virus | MYMV-Cambodia | 97.70 | 78.62 | <i>Vigna radiata</i> |
| 25 | AB017341.1 | Mungbean yellow mosaic virus | MYMV-Thailand | 97.41 | 74.20 | <i>Vigna radiata</i> |
| 26 | KC911718.1 | Mungbean yellow mosaic virus | MYMV-Coimbatore | 95.92 | 72.78 | <i>Vigna mungo</i> |
| 27 | PP738873.1 | Mungbean yellow mosaic virus | MYMV-Coimbatore | 97.85 | 79.15 | <i>Vigna radiata</i> |
| 28 | FM242701.1 | Mungbean yellow mosaic virus | MYMV-Pakistan | 94.91 | 75.23 | <i>Rhynchosia capitata</i> |
| 29 | MN885479.1 | Mungbean yellow mosaic virus | MYMV-Pakistan | 94.11 | 74.32 | <i>Glycine max</i> |
| 30 | MN922309.1 | Mungbean yellow mosaic virus | MYMV-Pakistan | 94.19 | 76.23 | <i>Vigna aconitifolia</i> |
| 31 | MN698293.1 | Horsegram yellow mosaic virus | HgYMV-Belgaum | 93.11 | 74.02 | <i>Vigna aconitifolia</i> |
| 32 | KY824799.1 | Mungbean yellow mosaic virus | MYMV-Chamarajanagar | 96.02 | 73.79 | <i>Vigna mungo</i> |
| 33 | MW736054.1 | Mungbean yellow mosaic virus | MYMV-Vamban | 92.62 | 70.03 | <i>Vigna radiata</i> |
| 34 | ON421034.1 | Mungbean yellow mosaic virus | MYMV-Kanpur | 97.93 | 81.78 | <i>Phaseolus vulgaris</i> |
| 35 | ON421030.1 | Mungbean yellow mosaic virus | MYMV-Kanpur | 97.93 | 81.78 | <i>Phaseolus vulgaris</i> |
| 36 | DQ389150.1 | Mungbean yellow mosaic India virus | MYMIV-Maharashtra | 99.05 | 81.90 | <i>Glycine max</i> |

| | | | | | | |
|----|------------|------------------------------------|-----------------|-------|-------|----------------------|
| 37 | MH885653.1 | Mungbean yellow mosaic India virus | MYMIV-Bengaluru | 97.26 | 77.06 | <i>Vigna radiata</i> |
|----|------------|------------------------------------|-----------------|-------|-------|----------------------|

UNDER PEER REVIEW