

Original Research Article

“Effect of phytoremediated sewage water irrigation, organic manures & biofertilizers on growth and yield of okra variety *Kashi chaman*”

Abstract

Okra belongs to family Malvaceae, it is a good source of nutrition as well as for farmer economy. This experiment was based on two years field work. During the year 2021-22 and 2022-23 it was aimed to study the Effect of phytoremediated sewage water irrigation, organic manures & biofertilizers on growth and yield of okra var. *Kashi chaman* in Prayagraj agro climatic condition. The experiment was designed as Randomized block design consist of 16 treatments and 3 replicates of each. Among which the treatment T₁₅ with the combination of Azotobacter @4kg/ha + PSB(*Pseudomonas*) @2kg/ha + Vermicompost + treated sewage water, emerged as most effective in promoting highest Number of branches (4.61) number of leaves (28.52) leaf area LAI (124.12 cm²) Root length per plant (15.15 cm) maximum fruit yield (464.72 g). Whereas maximum plant height (127.11 cm) was recorded in T₉ (Vermicompost+ Azotobacter @4kg/ha and TSW). So it can be concluded that this treatment combination can enhance plant growth and yield attributes of okra variety *Kashi chaman* along with sustaining nutrient levels in the soil.

Keywords: *Okra, Bio fertilizers, Organic manure, Growth, Yield.*

Introduction

Okra, scientifically known as *Abelmoschus esculentus* L., is a member of the Malvaceae family. It is referred to by various names such as Lady's finger, okra, bhindi, okura, quingombo, bamia, gombo, and lailongma, depending on the region where it is grown (Jain *et al.*, 2012). Okra is said to be originated from Africa and is grown in various countries across Africa, Asia, Southern Europe, and America. It thrives in tropical, subtropical, and warm temperate climates (Durazzo *et al.*, 2019). Okra is a shrub that is grown in tropical and subtropical regions worldwide. It is a commonly cultivated plant, both in gardens and on farms. This particular crop is extensively grown for consumption and is recognized worldwide for its delicious taste (Dhaliwal, 2010). The young, unripe pods of okra are commonly eaten as vegetables. Additionally, the extract from these pods is often used as a thickening agent in various soup and sauce recipes to increase their thickness (Kumar *et al.*, 2013). Polysaccharides found in okra have been utilized in sweetened frozen foods like ice creams and bakery products. These compounds offer various health benefits and contribute to extending the shelf life of these food items (Archana *et al.*, 2015).

In order to achieve success in utilizing treated wastewater for crop production, several factors must be carefully considered. These include selecting the most suitable crops, choosing appropriate irrigation methods, and implementing effective strategies to maximize crop yields and quality while also preserving soil productivity and protecting the environment. According to Elnasikh and Satti (2017) organic manures play a crucial role in improving soil structure and fertility, while biofertilizers have the ability to increase nutrient availability and stimulate beneficial microbial activity (Youssef and Farag, 2021). Bio-fertilizers, on the other hand, are microorganisms with agricultural value. They have the ability to convert non-stable components into usable forms through biological processes (Kumar *et al.*, 2011). According to Singh *et al.* (2014), a significant portion of conventional N-based fertilizers, up to 50%, is lost to the soil and environment. These activities have the potential to have a substantial impact on the environment, leading to increased levels of greenhouse gases, soil acidification, depletion of non-renewable resources, and the release of nitrates into groundwater and surface water.

Materials and methods

Geographical location of the experimental site:

In Prayagraj, Uttar Pradesh 211007, India, the investigation was carried out in the research area of the Department of Biological Sciences at Sam Higginbottom University of Agriculture, Technology and Sciences. The experimental field is located in close proximity to the Yamuna River, on the left side of the Allahabad-Rewa Road, and is roughly 8 kilometers from the metropolis of Allahabad. The coordinates of the experimental site are 81.51°E longitude and 25.57°N latitude.

Soil characteristics of the experimental site:

The soil in the experimental field, which is located in the central Gangetic alluvium region, has a neutral pH level and is characterized by its depth. Soil samples were collected prior to sowing by using an auger at a depth of 15 cm. The composite samples were utilized for the purpose of conducting chemical and mechanical analysis. The soil had a sandy loam texture and contained low levels of organic carbon. It had medium levels of available nitrogen, phosphorus, and potassium (Table no 1).

Crop and Experimental Details: The study was designed using a randomized block design with fifteen treatments and three replications, and it ran from January to June (2021–22 & 2022–23). Crop selected was okra (*Abelmoschus esculentus* L. Moench) family Malvaceae, Variety Kashi chaman. The plants were strategically spaced at a distance of Plant to Plant and row to row 60 cm × 45 cm. Total 432 plants and experimental area of 57.6 m². The treatment combinations consisted of alternatives such as the Vermicompost- @2.5 t/ ha (per plot 600g), Azotobacter- @4 kg/ ha (per plant: 1.20g), PSB- @2 kg/ ha (per plant: 0.50g), Quantity of sewage water – 168 litre per irrigation taken under study as shown in Table 2. The treated sewage water which was used in the experiment was firstly treated with *Ecchornia crassipes* for 21 days after that applied in the form of irrigation.

Sampling and parameters: studied Growth attributes are observed under various parameters like – plant height (cm), number of branches per plant, number of leaves per plant, root length per Plant (cm), number of nodes at first flowering, leaf area, leaf area index. Similarly yield attributes are - days to first fruit harvest, number of pods per plant, pod length (cm), pod

diameter (mm), average fresh pod weight (g), total fruit yield per plant (g).

Statistical analysis: The experiment was organized in a Randomized block design manner with three replicates of each treatment during two years 2021-2022 and 2022-2023. Statistical analysis was done to determine the interdependencies between different variables the collected data were submitted to two way analysis of variance (ANOVA) approach were compared at the 0.05 level. An open source agricultural data analysis tool i.e. OPSTAT was used for statistical analysis. The data incorporated in the tables are based on the pooled data of both the years.

Table 1. Physical and chemical properties of experimental soil:

Sl. No.	Particulars	Values (2021-22)	Values (2022-23)	Method employed
I	Physical Properties			
	Sand(%)	60.41	60.38	Bouyoucos Hydrometer method (Piper, 1966)
	Clay(%)	22.89	22.93	
	Silt (%)	16.35	16.36	
	Texture	Sandy loam	Sandy loam	USDA Triangle, Soil Survey, 1975
II	Chemical properties			
	pH	7.2	7.3	Glass Electrode, pH meter, (Jackson, 1967)
	Electrical conductivity (dS/m)	0.26	0.28	Conductivity bridge (Piper, 1966)
	Organic carbon (%)	0.48	0.49	Wet method, (Walkley, 1947 and Black 1965)
III	Available Nutrients			
	Available Nitrogen (kg/ha)	171.48	173.54	Modified kjeldhal's method (Jackson, 1967)
	Available Phosphorus (kg/ha)	13.6	15.41	Olsen method (Muhretal., 1963)
	Available Potassium (kg/ha)	215.4	218.34	Flame photometer method (Muhretal., 1963)

Table no 2. Details of treatment combinations:

S.No.	Notation	Treatment combinations
1	T0	Normal water(control)
2	T1	Treatedsewagewater
3	T2	Vermicompost+Normal water
4	T3	Vermicompost+Treatedsewagewater
5	T4	Azotobacter@4kg/ha+Normal water
6	T5	Azotobacter@4kg/ha+treatedsewagewater
7	T6	PSB(Pseudomonas)@2kg/ha +Normal water
8	T7	PSB(Pseudomonas)@2kg/ha +treatedsewage water
9	T8	Vermicompost+Azotobacter@4kg/ha+Normal water
10	T9	Vermicompost+Azotobacter@4kg/ha+treatedsewage water
11	T10	Vermicompost+PSB(Pseudomonas)@2kg/ha+Normal water
12	T11	Vermicompost+PSB(Pseudomonas)@2kg/ha+treatedsewageWater
13	T12	Azotobacter@4kg/ha+PSB(Pseudomonas)@2kg/ha+TapWater
14	T13	Azotobacter@4kg/ha+PSB(Pseudomonas)@2kg/ha+ treated sewagewater
15	T14	Azotobacter@4kg/ha+PSB(Pseudomonas)@2kg/ha+Vermicompost+Normal water
16	T15	Azotobacter@4kg/ha+PSB(Pseudomonas)@2kg/ha +Vermicompost+treatedsewagewater

Results and discussion:-

Growth attributes of okra

According to observation recorded it was resulted that the treatment T15 with the combination of Azotobacter @4kg/ha + PSB(Pseudomonas) @2kg/ha +Vermicompost + treated sewage water shows the maximum growth attributes based on pooled data that includes number of branches per plant (4.61) number of leaves per plant (28.52) leaf area (124.12) leaf area index (1.11) Number of nodes at first flowering (6.55) Root length per plant (15.15cm)whereas maximum plant height (127.11 cm) was recorded in T9.Over all treatments during both the years of experiment as shown in Table 3-4.Whereas lowest growth attributes was recorded in T₀ [Control (no treatment)].

The application of plant nutrients through organic sources likes vermicompost, farm yard manure and bio fertilizers remain the alternative choice of the growers for maintaining its sustainable production (**Gayathri and Reddy, 2013**). Vermicompost may have enhanced the microbial activity in the soil, leading to benefits such as improved oxygen availability, optimal soil

temperature, increased soil porosity and water infiltration, enhanced nutrient content, and ultimately, increased plant height (**Arora et al., 2011**).

Bio- inoculants such as Azotobacter and PSB have been found to enhance the bio-availability of Nitrogen and Phosphorous in the soil through fixation by converting non-usable form into usable form, as demonstrated in studies by **Zaidi et al. (2009)** and **Hafez et al. (2016)**. Pseudomonas can help to overcome the negative effects of contaminated waters, and other environmental and biological stresses, as described by **Gardezi et al. (2019)**.

In addition, the consistent supply of water and nutrients through treated sewage water irrigation guarantees that the plants have the necessary resources to maintain robust growth and branch development all year round (**Khan, 2018**).

Yield attributes of okra

A similar trend has been observed on various yield attributes. In the continuation the treatment T₁₅ recorded on Daystofirstfruit harvest (39.36) Number of pods per plant (22.86) Pod length (14.32cm) Pod diameter (13.51mm) Average fresh pod weight (20.50g) Total fruit yield per plant (464.72 g) as depicted in Table 5.

In addition, vermicompost promotes mutually beneficial interactions between plants and beneficial soil microorganisms, such as mycorrhizal fungi. These microorganisms play a vital role in improving nutrient absorption and transportation within the plant. These interactions contribute to enhanced nutrient absorption and distribution towards reproductive organs, leading to a notable increase in the length of okra pods (**Rehman et al., 2023**).

Azotobacter, a bacterium that fixes nitrogen, has the ability to improve nitrogen availability. This, in turn, promotes the growth of plants and their reproductive processes, leading to an increase in flower and fruits (pods) production. In addition, PSB has the ability to solubilize insoluble phosphorus, allowing plants to access this crucial nutrient for seed development and maturation (**Iftikhar et al., 2024**).

According to **Hashem and Qi (2021)**, treated sewage water can provide vital macro and micronutrients that support both vegetative and reproductive growth. Vermicompost enhances the soil by adding organic matter, resulting in improved soil structure, increased water retention, and enhanced nutrient availability. These benefits have a positive effect on the development of flowers and pods, as demonstrated by **Rehman et al. (2023)**.

Conclusion

In view of two years of experimental investigation it was found that phytoremediation of sewage water with *Ecchornia crassipes* improves the physico- chemical characteristics of raw sewage water and found suitable for okra production in sandy loam soil.

Growth parameters including Number of branches, number of leaves, leaf area LAI was found maximum in T15 [Azotobacter @4kg/ha + PSB(*Pseudomonas*) @2kg/ha + Vermicompost + treated sewage water] whereas maximum plant height was recorded in T9 (Vermicompost+ Azotobacter @4kg/ha and TSW) and T15 treatment combination resulted in maximum fruit yield (464.72g) and other yield parameters. So, overall it can be concluded that the T15 (Azotobacter @4kg/ha + PSB(*Pseudomonas*) @2kg/ha + Vermicompost + treated sewage water) had found most effective in terms of growth and yield of okra among all the treatments.

Reference:-

1. Archana, G., Babu, P. A. S., Sudharsan, K., Sabina, K., Raja, R. P., Sivarajan, M. and Sukumar, M. (2015). Evaluation of Fat Uptake of Polysaccharide Coatings on Deep-Fat Fried Potato Chips by Confocal Laser Scanning Microscopy.; *Int.J.Food Prop.*; 19:1583– 1592.
2. Arora, V.K., Singh, C.B., Sidhu, A.S. and Thind, S.S. (2011). Irrigation, tillage and mulching effects on Soybean yield and water productivity in relation to soil texture. *Agric Water Manag.* 98(4):563–568.
3. Dhaliwal, M. S. (2010). *Okra (Abelmoschus esculentus) L. (Moench)* Kalyani Publishers; New Delhi, India.
4. Durazzo, A., Lucarini, M., Novellino, E., Souto, E. B., Daliu, P. and Santini, A. (2019). *Abelmoschus esculentus*(L.): Bioactive components' beneficial properties—Focused on antidiabetic role—For sustainable health applications; *Molecules*, 24:38.
5. Elnasikh, M. H. and Satti, A.A. (2017). Potentiality of Organic Manures in Supporting Sustainable Agriculture in Sudan; *Environment and Natural Resources International Journal*, 2(1): 01- 26.
6. Gardezi, A. K., Berber, S. R. M., Castañeda, G. C., Flores-Magdaleno, H., & Aguilar, G. H. (2019). Sewage Water Effects on Okra (*Abelmoschus esculentus* (L) *Moench*) Growth Affected by Organic Matter, and Avm of *Pseudomonas fluorescens*. *World J Agri & Soil Sci.* 1 (5): WJASS. MS. ID, 524.
7. Gayathri, K. and Syam Sundar Reddy, P. 2013. Effect of integrated nutrient management growth and yield of okra (*Abelmoschus esculentus* (L). *Moench*) cv. Arka Anamika. *Veg. Sci.* 40(2): 246-248.

8. Hafez, S. K., El-Beik, A. K., El-Tantawy, A. M. and ElKasas, A. I. (2016). Effect of Organic and Bio-Fertilizer sources on Summer Squash under El-Arish conditions. *Sinai Journal of Applied Sciences*, 5(2): 155-172.
9. Hashem, M. S. and Qi, X. (2021). Treated Wastewater Irrigation—A Review; *Water*, 13(11), 1527.
10. Iftikhar, A., Aijaz, N., Farooq, R. and Aslam, S. (2023). Beneficial Role of Phosphate Solubilizing Bacteria (PSB) In Enhancing Soil Fertility Through A Variety Of Actions On Plants Growth And Ecological Perspective: An Updated Review; *Journal of Xi'an Shiyou University, Natural Science Edition*, ISSN: 1673-064X
11. Jain, N., Jain, R., Jain, V. and Jain, S. (2012). A review on: *Abelmoschus esculentus*. *Pharmacia*; 1: 84–89.
12. Khan, N. (2018). Natural Ecological Remediation and Reuse of Sewage Water in Agriculture and Its Effects on Plant Health; In book: *Natural Ecological Remediation and Reuse of Sewage Water in Agriculture and Its Effects on Plant Health* (pp.1-16).
13. Kumar, S., Girwani, G., Reddy, S. and Bhagwan, A. (2011). Studies on nutrient management in custard apple" banalagar; *Acta horticulturae*, 890: 381-385.
14. Kumar, A., Kumar, P. and Nadendla, R. (2013). A review on: *Abelmoschus esculentus* (Okra); *Int. Res. J. Pharm. Appl. Sci.*, 3: 129–132.
15. Rehman, S., DECastro, F., Aprile, A., Benedetti, M. and Fanizzi, F.P. (2023). Vermicompost: Enhancing Plant Growth and Combating Abiotic and Biotic Stress; *Agronomy*, 13(4): 1134.
16. Singh, A., Maji, S. and Kumar, S. (2014). Effect of biofertilizers on yield and biomolecules of anti-cancerous vegetable broccoli; *International Journal of Bio-resource and Stress Management*, 5(2): 262-268.
17. Zaidi, A., Khan, M., Ahemad, M. and Oves, M. (2009). Plant growth promotion by phosphate solubilizing bacteria. *Acta microbiologica et immunologica Hungarica*, 56(3): 263-284.
18. Piper, C. S. (1966). *Soil and Plant Analysis*. Academic Press, New York, pp. 367.
19. USDA (1975). *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys*. Soil Survey Staff, Coord., Soil Conservation Service. Agriculture Handbook 436, US Department of Agriculture, Washington DC, 754 p.

20. Walkley, A. and Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science*, 37(1): 29-38.
21. Youssef, M. A. and Farag, M. I. H. (2021). Co-application of Organic Manure and Bio-fertilizer to Improve Soil Fertility and Production of Quinoa and Proceeding Jew's Mallow Crops; *Journal of Soil Science and Plant Nutrition*, 21(1): 1-5.
22. Muhr, G. R., Dutia, N. P., Sankar, S. H., Dever, R. F., Leley, V. K. and Dohalur, R. L. (1963). *Methods of analysis soil testing in India*. United States agency for international development mission to India, New Delhi, pp. 37-39.
23. Jackson, M. L. (1967). *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi,

Table 3. Effect of Phyto remediated sewage water irrigation, organic manures, and biofertilizers on growth attributes of okra var. Kashi chaman.

TreatmentSymbol	Plantheight(cm)			Numberofbranchesperplant			Numberofleavesperplant		
	20 DAS	40 DAS	60 DAS	20 DAS	40 DAS	60 DAS	20 DAS	40 DAS	60 DAS
T0	10.12	35.19	72.98	0.81	2.51	3.53	6.33	11.68	19.73
T1	10.87	37.85	79.63	0.92	2.65	3.76	6.63	12.17	20.34
T2	15.63	44.40	109.11	1.03	2.96	4.28	7.91	14.57	24.10
T3	17.88	48.36	114.76	1.54	3.13	4.32	8.35	15.12	26.30
T4	11.72	40.56	85.28	1.07	2.71	3.85	6.95	12.85	21.18
T5	12.84	42.70	88.57	1.19	2.78	3.91	7.19	13.14	21.83
T6	12.30	40.59	85.22	1.11	2.75	3.89	6.95	12.85	21.18
T7	13.39	42.32	90.94	1.15	2.78	3.94	7.19	13.32	22.23
T8	16.48	53.26	107.95	1.44	3.05	4.21	8.03	14.84	25.18
T9	20.12	58.78	127.11	1.79	3.32	4.58	8.68	15.70	27.20
T10	17.03	53.24	110.37	1.62	3.22	4.21	8.15	14.84	25.17
T11	19.27	56.00	120.40	1.66	3.24	4.44	8.68	15.70	27.20
T12	14.24	46.40	99.15	1.34	2.86	4.06	7.58	13.94	23.19
T13	14.78	50.52	99.48	1.30	2.88	4.06	7.58	13.93	23.18
T14	18.73	56.01	120.21	1.69	3.25	4.44	9.02	15.26	28.13
T15	20.06	58.75	127.05	1.83	3.35	4.61	9.14	15.44	28.52
F-test	S	S	S	S	S	S	S	S	S
S.E.(m)(±)	0.17	0.38	0.86	0.02	0.01	0.01	0.05	0.07	0.19
C.D.(P=0.05)	0.47	1.09	2.44	0.05	0.03	0.04	0.14	0.21	0.58

Table 4. Effect of Phyto remediated sewage water irrigation, organic manures, and biofertilizers on growth attributes of okra var. Kashi chaman.

Treatment symbol	Leaf area			Leaf area index			Number of nodes at first flowering	Rootlengthperplant(cm)
	20 days	40	60 days	20 days	40	60 days		
T0	15.69	31.20	56.88	0.13	0.26	0.47	6.01	12.59
T1	16.55	33.49	62.80	0.14	0.28	0.52	6.04	12.73
T2	23.62	47.12	91.91	0.20	0.39	0.77	6.28	13.95
T3	26.83	53.40	106.60	0.22	0.45	0.89	6.42	14.44
T4	17.72	35.79	68.73	0.15	0.30	0.57	6.11	13.02
T5	19.45	39.16	74.40	0.16	0.33	0.62	6.16	13.17
T6	18.58	37.47	71.56	0.15	0.31	0.60	6.13	13.02
T7	20.40	40.84	77.23	0.17	0.34	0.64	6.16	13.27
T8	24.80	49.42	97.84	0.21	0.41	0.82	6.35	14.13
T9	28.02	55.69	112.53	0.24	0.46	1.02	6.47	14.73
T10	25.66	51.10	100.67	0.21	0.43	0.84	6.35	14.13
T11	28.88	57.38	115.36	0.24	0.48	0.96	6.47	14.73
T12	21.59	43.14	83.16	0.18	0.36	0.69	6.23	13.68
T13	22.45	44.82	85.99	0.19	0.37	0.72	6.23	13.68
T14	30.05	59.67	121.29	0.25	0.50	1.04	6.53	15.04

T15	30.91	61.36	124.12	0.26	0.51	1.11	6.55	15.15
F-test	S	S	S	S	S	S	S	S
S.E.(m)(±)	0.24	0.47	0.94	0.01	0.02	0.03	0.01	0.04
C.D.(P=0.05)	0.69	1.33	2.65	0.03	0.05	0.10	0.03	0.12

UNDER PEER REVIEW

Table 5. Effect of Phyto remediated sewage water irrigation, organic manures, and biofertilizers on yield attributes of okra var. Kashi chaman.

Treatment Symbol	Daystofirst fruit harvest	Numberof podsperplant	Podlength (cm)	Poddiameter(mm)	Average freshpodweight(g)	Total fruit yield per plant (g)
T0	46.61	16.66	11.20	11.11	13.98	245.70
T1	46.15	17.83	11.48	11.34	14.94	266.38
T2	41.45	19.73	12.89	12.35	15.05	296.94
T3	42.66	19.17	13.56	12.89	14.17	271.57
T4	45.30	18.45	11.85	11.57	15.49	285.79
T5	44.59	19.24	12.11	11.73	15.81	304.18
T6	45.31	18.77	11.85	11.65	17.10	320.88
T7	44.60	17.65	12.05	11.81	17.54	309.58
T8	41.88	21.25	13.22	12.58	18.11	384.73
T9	39.01	21.69	14.43	13.12	19.31	418.63
T10	41.89	19.64	12.53	12.66	18.18	356.87
T11	40.20	21.52	13.22	13.20	18.64	401.03
T12	43.57	18.82	13.90	12.04	19.60	388.37
T13	43.58	19.27	12.53	12.12	19.46	391.14
T14	40.02	22.23	13.91	13.43	20.37	441.52
T15	39.36	22.86	14.32	13.51	20.50	464.72
F-test	S	S	S	S	S	S
S.E.(m)(±)	0.18	0.18	0.05	0.04	0.01	0.30
C.D.(P=0.05)	0.54	0.54	0.14	0.11	0.03	0.89

UNDER PEER REVIEW