

Original Research Article

Evaluation of the use of organic and mineral fertilisers on amaranth productivity in the locality of Kombé, Republic of Congo.

ABSTRACT

The use and high cost of chemical fertilisers, as well as the dangers associated with their misuse, are leading growers to look for an alternative, sustainable form of fertilisation. A survey was carried out in the Kombé area of Brazzaville to assess the effectiveness and use of organic and mineral fertilisers on amaranth growth and yield. During the survey, 50 market gardeners growing amaranth were randomly selected and interviewed using a semi-structured questionnaire. The results obtained from our survey were analysed statistically, revealing significant differences between the different types of fertiliser used at Agricongo de Kombé ($p < 0.05$). The variance analysis revealed five (5) heterogeneous groups (a, b, c, d, cd), the most significant being poultry manure (group d) and NPK (group cd). The results of the survey revealed that amaranth (32.31%) was the dominant previous crop, poultry manure (34%) was the main and least expensive fertiliser, so it is capable of improving the growth and yield of amaranth, compared with other manures. Poultry droppings could be recommended for amaranth production.

Keywords: Amaranth, organic fertilizer, mineral fertilizer, NPK, Agricongo de Kombé

1. INTRODUCTION

Market gardening is characterised by the cultivation of leafy vegetables, fruit, roots and tubers for food use. Among the great diversity of vegetables in Africa are leafy vegetables of which amaranth is the most widely grown and the most important. Amaranth is one of the world's oldest food crops, having been cultivated in Mexico for 6,000 years [Srivastava \(2011\)](#). Some species of amaranth are leafy vegetables that are essentially needed to maintain health, especially in preventing human diseases, as they are a good source of vitamins, mineral nutrients and antioxidants [Prasad et al. \(2014\)](#) and [Wouyou et al. \(2017\)](#).

However, the yield of amaranth production can be improved by the use of chemical fertilisers, but the use of chemical fertilisers can lead to high toxicity, environmental pollution and possible side-effects on human health when they end up in foodstuffs. In addition, most vegetable growers in tropical Africa are smallholders who cannot afford the high cost of inorganic fertilisers [Makinde et al. \(2008\)](#). This is why new alternatives have been explored to reduce the use of chemical fertilisers. One solution is to use organic and mineral fertilisers to feed the soil and improve amaranth production by providing the necessary nutrients. Organic fertiliser sources with a C/N ratio below 20 contain a high concentration of nutrients [Chaves et al. \(2007\)](#); [Tognetti et al. \(2008\)](#). Mineral fertilisers provide nutrients that can be directly

assimilated by plants. They are easy to use and act quickly on plants Graines Hubert (2024). These fertilisers are inexpensive rather than chemical fertilisers, because they are easy to mass-produce.

Several studies have focused on the chemical characterisation of organic and mineral fertilisers and their contribution to plant nutrition in vegetable crops Laroungaet al. (2020) and Ballot et al. (2016). All these studies confirm the importance of nitrogen in the growth and production of market garden crops. Although organic and mineral fertilisers are used on market gardening sites in the Republic of Congo in general and in the city of Brazzaville in particular, very little work has been done on guiding market gardeners in their choice of fertilisers for amaranth production.

The aim of the study is to assess the effectiveness and use of organic and mineral fertilisers on amaranth growth and yield in Brazzaville, in order to recommend the choice of fertiliser types for optimum amaranth growth and development in the Kombé area.

2. MATERIAL AND METHODS

2.1 Study area

The choice of the Agricongo centre in Kombé as the survey site was based on the level of vegetable production and the fact that it is one of the major vegetable production areas in Brazzaville Massengo et al. (2004).

The Agricongo center in Kombé was set up on 16 January 1986 on the initiative of the Congolese government and with the support of the Agri-sud international group, after the failure of the pre-cooperative groups set up in earlier years (70-80). It is located seventeen kilometres (17 km) south of the city of Brazzaville, in district 8 Madibou. The centre lies at latitude 4°19'34 South, longitude 15°9'49 East, at an altitude of 295 metres (Kombé agrometeorological station).

2.2 Plant materials and Fertilising

The plant material used by market gardeners consists of *Amaranthus hybridus* seeds.

Fertilising material include organic and mineral fertilizers.

Table 1. List of fertilizers used

Type of fertiliser	Commercial name
Organics	Straw (dried <i>Echinochloastagnina</i>)
	Poultry manure
	Pig manure Wheat waste
Minerals	NPK
	Urea

2.3 Survey

Aware of the need to master techniques for using fertilisers on amaranth production, we conducted a survey of growers from August to October 2022. The aim was to assess the type of fertiliser used for amaranth production in the Kombé area. Fifty (50) market gardeners were interviewed as a sample using a pre-established questionnaire consisting of individual forms (Table 2).

Table 2. Individual survey forms

Date.....			
Locality.....			
Proprietor			
AgricongoKombéSite			
1. Characterisation of the operation			
1.1 Variety (s)			
1.2 Age of culture	<input type="text"/>	month(date of plantation:	
1.3 Cultural profil plat	<input type="text"/>	butteridges	<input type="text"/>
1.4 Plantation density	<input type="text"/>	strong	<input type="text"/>
1.5 Siol Type light heavy	<input type="text"/>	other	<input type="text"/>
1.6 cultural system	<input type="text"/>	pure	<input type="text"/>
1.7 Cultivation Precedent (4 years ago)	2022.....	2021.....	
	2020.....	2019.....	
1.8 Agricultural input	<input type="text"/>	fertilizer	<input type="text"/>
		herbicide	<input type="text"/>
			<input type="text"/>
nature.....		
and dosage.....		

2.4 Statistical analysis

The data collected in the field were analysed with the aid of computer-assisted statistical tools, using SPSS (Statistical Package For Social Science) version 22.0 and R version 4.1.3. The statistical methods used are linked to the sampling methods and experimental set-up adopted. The normality of the residuals and the homogeneity of the variances were verified. These methods included: two-factor ANOVA and linear correlations. Means were separated using the Student Newman and Keuls test and the Kruskal-Wallis test at the 5% probability threshold.

3. RESULTS

A field survey was carried out in the locality of Kombé in Brazzaville to evaluate the different types of fertilizer used on amaranth production.

Statistical analysis reveals significant differences between the different speculations of the Kombé center ($p < 0.05$). The analysis of variance revealed six (6) homogeneous groups (a, b, c, bc, ab, d), the most significant of which are amaranth (group d) and lettuce (group c). Table 3 represents the average percentage of crop precedents (PC) obtained after a survey of the respondents. There is a high occurrence of amaranth (32.31%) in the site compared to other speculations. It is followed by lettuce (23.08%) and spinach (13.84%), spring onion (7.69%), black nightshade (6.15%), baselle (4.62%), endive (4.62%), tomato (3.08%), eggplant (1.54%), cabbage (1.54%) and okra leaf (1.54%).

Table 3. Average percentage of market garden crops per farmer in the KombéAgricongo.

Speculation	N	Percentage (%)
Amaranth	21	32,31d
Aubergine	1	1,54a
Baselle	3	4,62a
Cabbage	1	1,54a
Chives	5	7,69b
Endive	3	4,62a
Spinach	9	13,85bc
Okra leaf	1	1,54a
Lettuce	15	23,08c
Black nightshade	4	6,15ab
Tomato	2	3,08a

N= number of patients per culture. Figures preceded by different are statistically different at the threshold of $P < 0.05$ according to the Student Newmann and Keul test.

Figure 1 shows the percentage use of fertilisers obtained from the survey. The results show that the most commonly used fertiliser is poultry manure, with the highest percentage (34%). In contrast, wheat waste has the lowest percentage of use. The percentage of other

fertilisers used shows that: NPK (25.5%) > pig manure (12.8%) > straw (10.6%) > urea (8.6%) (Figure 2).

The results obtained from the statistical analysis reveal significant differences between the different types of fertiliser used at Agricongo de Kombé ($p < 0.05$). The analysis of variance revealed five (5) heterogeneous groups (a, b, c, d, cd), the most significant being poultry manure (group d) and NPK (group cd).




Fig.1. Fertiliser use of organic and mineral in the Agricongo centre

The straw was made from the dried waste of *Echinochloa stagnina*.

Méthode de Student Newman et Keul : Les différences sont statistiquement différentes au seuil de $P < 0,05$.

4. DISCUSSION

The average percentage of previous crops (PC) obtained after surveying the respondents showed that amaranth was the dominant previous crop before the crops planted. This is because amaranth produces higher yields than other seed plants. It is a short-cycle plant. It requires less money and time to grow because it grows in poor soil, is resistant to disease and pests, resists water shortages and can be cultivated all year round. Its seeds are readily available on the market and from market gardeners. On local markets, it is the most common leafy vegetable to be found on stalls, because of its low price and the fact that the quantity sold per bunch is sufficient. Amaranth is a widely grown vegetable on market gardens. In fact, amaranth is a widely grown vegetable on market gardening farms. These results are quite similar to the results of work carried out by Marandet (2016) in market gardening in the city of Brazzaville: the dominant leafy vegetables include amaranth (*Amaranthus hybridus* L.), baselle (*Basella alba* L.), black nightshade (*Solanum nigrum* L.), bitter nightshade (*Solanum aethiopicum*), etc. According to the results of a study by Mialoundama (2021) on strategies for managing bioaggressors in urban market gardening in Brazzaville (Congo), the most widely grown vegetables are tomatoes, spring onions, amaranth and chillies.

The results of our survey revealed that poultry manure is the main fertiliser used at Agricongo's Kombé site. Organic matter feeds and shelters a large number of useful soil

organisms, such as earthworms and micro-organisms, which work continuously to improve soil fertility and structure Ouédraogo et al. (2008), hence the attachment of the market gardeners of Kombé to the use of poultry manure, they understood that organic manure improves the physical properties of the soil and feeds the plant by providing the nutrients it needs. Their preference for poultry manure is due to the fact that it is affordable and more accessible than other manures. Poultry manure is a very economical fertiliser, because market gardeners harvest it from poultry farms for the most part, and spend only on the purchase of the shavings. Onana (2006), for example, shows that a 35-40 kg bag of poultry manure is sold for 2200 CFA francs in the villages, compared with 17500 CFA francs for a 50 kg bag of compound mineral fertiliser on the Yaoundé market. At Agricongo in Kombé, market gardeners use poultry manure in nurseries and open fields to reduce the use of chemical fertilisers and pesticides. These results are similar to those obtained by Ruf et al. (2016), who conducted a survey on the use of poultry manure in the Duékoué region of Côte d'Ivoire. According to their results, poultry manure is widely used by cocoa farmers. In the Duékoué region as a whole, 90% of farmers use poultry manure. Another reason why market gardeners in Kombé prefer to use poultry manure is to improve the productivity of amaranth. This corroborates the observations made by Tchaniley et al. (2020), who showed that treatment T2 with poultry manure increased lettuce yield with a product surplus of 134.00%. The same results were obtained by Biauou et al. in (2017) after using compost enriched with poultry droppings for carrot production. In addition, the results of our survey show poor results on the use of pig manure. This is because pork is sold at high prices and is difficult to access. Ognika et al. (2016) pointed out that pork production in the city of Brazzaville and its outskirts is not yet able to satisfy domestic demand and guarantee the sustainability of the activity.

According to data collected in 2011 on fertiliser consumption and nutrient balances, nitrogen fertilisers accounted for the vast majority (77.4%) of nutrients consumed. Consumption is estimated at 59 kg per hectare in the European Union (EU), ranging from 19 kg per hectare in Portugal to 125 kg per hectare in the Netherlands (Statistics Explained, 2015). These data differ from our results. Our results collaborate with the experimental results of Kahu et al. (2019) have shown that poultry manure had higher growth performance on the two varieties of Amaranth when compared NPK compound fertilizer. The poor results on the use of NPK by market gardeners in Kombé is due to the low productivity of amaranth. Ognalaga et al. (2015) noted the poor growth of amaranth plants on plots fertilised with NPK. This means that the mineral elements alone are not sufficient for good amaranth growth.

5. CONCLUSION

The aim of this study was to assess the effectiveness and use of organic and mineral fertilisers on the growth and yield of amaranth in the Kombé area of Brazzaville. The results of the survey revealed that amaranth (32.31%) was the dominant previous crop, poultry manure (34%) was the main fertiliser and was able to improve amaranth growth and yield compared with other manures. As a result, organic fertilisers such as poultry manure could well be substituted, even partially, for synthetic chemical fertilisers.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. The authors declare that NO generative artificial intelligence technologies such as large language models (ChatGPT, COPILOT, etc.) and text-image generators were used during the writing or editing of this manuscript.

2.

3.

REFERENCES

1. Srivastava. (2011). Nutritional quality of some cultivated and wild species of *Amaranthus L.* International Journal of Pharmaceutical Sciences and Research, 2(12),3152.
2. Prasad SM., Parihar P., Singh VP. (2014). Effect of Salt Stress on Nutritional Value ofVegetables. Biochem. Pharmacol , 3(2),1-2.
3. Wouyou, A.D., Ahissou, E.A., Gandonou, C.B., Assogba, K.F., Hounghèmè, A., Gbaguidi F.A., Ahissou, H., Lagnika, L., Zanklan, S.A. and Lutts, S. (2017). Salinity increased vitamins concentration in *Amaranthus cruentus* leaves, African Journal of Biotechnology, 16(44), 2106-2111.
4. Makinde EA., Ayeni LS., Ojeniyi SO., Odedina JN. (2008). Effect of organic, organomineral and NPK fertilizer on nutritional quality of *Amaranthus* in Lagos, Nigeria. Nigerian Journal of Soil Science, 2, 91-96.
5. Chaves B, De Neve S, Piulats LM, Boeckx P, Van Cleemput O, Hofman G. (2007). Manipulating Nitrogen Release from Nitrogen-Rich Crop Residues using Organic

Wastes under Field Conditions. Soil Science Society of America Journal, 71(4), 1435-0661.

6. Tognetti, C., Mazzarino MJ., Laos, F. (2008). Compost of municipal organic waste: effects of different management practices on degradability and nutrient release capacity. *Soil Biology Biotechnology*, 49, 2290-2296.
7. Graines Hubert (2024). Why use a fertiliser and which one to choose? Site web :https://www.graines-hubert.com/module/stblog/39_pourquoi-utiliser-un-engrais-et-lequel-choisir-.html
8. Larounga, C., Kwasi D. A., Kodjo, A. D. K., (2020). Effect of combining organic and mineral fertilisers (NPK 15-15-15 and urea) on lettuce (*Lactuca sativa* L.) yield in southern Togo. *Journal of Applied Biosciences*, 151, 15540 – 15549.
9. Ballot, C. S. A., Semballa, S., Atakpama, W., Yangakola, T. M., Bo-Mbi, A. D., Blavet, D., Zinga, I., Wala, K., Batawila, K., & Akpagana, K. (2016). Effect Of Mineral Manures On The Yield And Organoleptic Quality Of Manioc (*Manihot Esculenta Crantz*) In The Savannah Zone In The Centre-South Of The Central African Republic. *European Scientific Journal*, *ESJ*, 12(6), 185.
10. Massengo, D., and Ngoyi, J.-I. (2004). Training in urban market gardening in Brazzaville, the action of Agricongo, Grain de sel n° 29. https://www.inter-reseaux.org/wp-content/uploads/6.2_initiatives_maraichage_urbain.pdf
11. Marandet, M. (2016). Sustainable agriculture in Congo: the case of market garden production on the right bank of the Djoué (Brazzaville region). Master's thesis. Agronomic sciences. Brazzaville: University of Marien Ngouabi - National School of Agronomy and Forestry.
12. Mialoundama, B.G. (2021). Strategies for managing pests and diseases in urban market gardening in Brazzaville (Congo). *African Journal of Environment and Agriculture*, 02, 30-39.
13. Ouédraogo, A., Yombi, L., Doumbia, S., Eyhorn, F., DISCHL, R. (2008). Organic and fair-trade cotton production guide (reference manual for West Africa). Helvetas. 49 p. http://www.slire.net/download/1825/guide-coton-bio-afrique_helvetas_08_1_.pdf
14. Onana Onana, L.G. (2006). Fertilization practices and soil characteristics in the peri-urban market gardening zone of Yaoundé: case of the Nkolondom lowlands. Final dissertation, University of Dschang (Cameroon), 73 pp.

15. Ruf F., Kiendré J., (2016). Innovation «Chicken droppings» in cocoa farms. Inter-Réseaux, développement rural, CIRAD. https://www.inter-reseaux.org/wp-content/uploads/l_innovation_dans_les_cacaoyeres_fevrier_2016-3.pdf
16. Biaou, O. D. B. A. Saidou, F-X. Bachabi, G. E. Padonou et I. Balogoun. (2017). Effect of organic manure on ferralitic soil and carrot (*Daucus carota* L.) production in southern Benin. *Int. J. Biol. Chem. Sci.* 11(5), 2315-2326.
17. Ognika, A., Missoko, R., Mopoundza, P., Akouango, P. (2016). Farm dynamics and characteristics of pig producers in and around Brazzaville (Republic of Congo). *Journal of Animal & Plant Sciences*, 29(1), 4426-4536.
18. Statistics Explained (2015). Fertilizer consumption statistics and nutrient balances. Eurostat, <http://ec.europa.eu>.
19. Kahu, J. C., C. C. Umeh, and A. E. Achadu. 2019. "Effects of Different Types of Organic Fertilizers on Growth Performance of *Amaranthus Caudatus* (Samaru Local Variety) and *Amaranthus Cruentus* (NH84/452)". *Asian Journal of Advances in Agricultural Research*, 9 (2), 1-12.
20. Ognalaga, M., Boussiengui-Boussiengui, G., Oyanadigui Odjogui P. (2015). Contribution à la restauration de la fertilité des sols du périmètre maraîcher de l'IGAD DJAMITI (Franceville) par l'apport raisonné des amendements organiques et minéral. *Journal of Animal & Plant Sciences*, (24)3, 3843-3853.