

Effect of Organic Manures on Growth and Yield attributes of Citronella Java (*Cymbopogon winterianus*) under Moringa (*Moringa oleifera*) based agroforestry and open-field systems

Abstract

~~A field investigation was conducted to evaluate the effects of farmyard manure (FYM), vermicompost, and neem cake on the growth and yield of Citronella Java (*Cymbopogon winterianus*) under moringa-based agroforestry and open-field systems during 2022 and 2023. Field investigation was carried out to study the effect of FYM, Vermicompost and Neem cake on the growth and yield of Citronella Java (*Cymbopogon winterianus*) cultivated under moringa based agroforestry and open field systems during 2022 and 2023. The experiment was laid out in a randomized block design with four levels of each component.~~ Growth parameters like plant height, number of leaves per plant and number of tillers per plant and yield attributing parameters like fresh weight (kg ha^{-1}), Dry Weight (kg ha^{-1}) and total herbage yield (t ha^{-1}). The findings highlight the superior performance of the treatment combining FYM, vermicompost, and neem cake (T13), particularly in shaded conditions provided by moringa trees, which created a favourable microclimate. This study demonstrates the significant role of integrated organic amendments in enhancing citronella productivity while fostering environmental sustainability. The findings offer valuable insights for farmers and policymakers, advocating for the adoption of sustainable practices in aromatic crop production through agroforestry systems. ~~These results provide valuable insights for farmers and policymakers, promoting sustainable aromatic crop production through agroforestry systems.~~

Formatted: Font color: Red, Strikethrough

Keywords: ~~*Citronella Java, Moringa, Agro forestry and Organic manures*~~

Agroforestry, Citronella Java, Moringa, and Organic manures

Formatted: Strikethrough

Introduction

Sustainable agriculture seeks to balance productivity with environmental stewardship by promoting practices that enhance soil health, biodiversity, and economic viability. Agroforestry, the integration of trees with crops or livestock, is increasingly recognized as a sustainable farming system due to its ability to improve soil fertility, optimize resource use, and provide environmental benefits such as carbon sequestration and microclimate regulation (Nair, 2011). In such systems, medicinal and aromatic plants like Citronella Java (*Cymbopogon winterianus*), known for its high-value essential oil used in the cosmetic, pharmaceutical, and insect-repellent industries, offer significant economic potential (Singh *et al.*, 2015).

The integration of organic amendments like farmyard manure (FYM), vermicompost, and neem cake in agroforestry systems further enhances soil health by providing essential nutrients, improving soil structure, and stimulating beneficial microbial activity. In combination with the resource benefits provided by tree species, such as *Moringa oleifera*, these organic inputs can create a synergistic environment that supports optimal growth conditions for crops like citronella (Singh *et al.*, 2015). The adoption of organic manures, including farmyard manure (FYM), vermicompost, and neem cake, is a key strategy for enhancing soil fertility in sustainable cropping systems. Organic amendments supply essential nutrients, improve soil organic matter, and stimulate beneficial soil microbial activity, contributing to better plant growth and yield (Manna *et al.*, 2005). These benefits are particularly relevant in agroforestry systems, where the interplay between tree species and crops affects nutrient availability, light conditions, and soil moisture dynamics. In systems combining *Moringa oleifera*, a fast growing and nutrient rich tree species, with citronella, the shade and nutrient contribution from moringa may complement the use of organic manures, further enhancing the growth and yield of citronella (Kaushal *et al.*, 2014).

Addmoringa details also :Moringa oleifera Lam is an important tree whose pods, leaves, flowers, barks and roots have been advocated for traditional and medicinal uses for thousands of years and native to India (Yadav et al., 2021, Yadav et al., 2022, Yadav et al., 2023, Yadav et al., 2024). Despite the recognized benefits of organic inputs and agroforestry systems, the specific effects of FYM, vermicompost, and neem cake on citronella's growth and yield under both shaded (moringa-based) and open-field conditions remain underexplored. Understanding these interactions is critical to optimizing the productivity and sustainability of citronella cultivation, particularly for resource-limited farmers seeking to adopt eco-friendly practices.

Although previous studies have highlighted the benefits of organic inputs and agroforestry systems in other crops, there is a lack of detailed research on how specific organic amendments, such as FYM, vermicompost, and neem cake, influence the growth and yield of citronella in shaded versus open-field conditions. This knowledge gap is particularly important for optimizing citronella cultivation, as farmers increasingly seek sustainable practices to improve productivity and minimize environmental impact. This study aims to evaluate the effects of FYM, vermicompost, and neem cake on the growth and yield attributes of citronella under moringa-based agroforestry and open-field systems. The findings will provide valuable insights into sustainable aromatic crop production, highlighting the potential of integrating organic manures with agroforestry practices to achieve higher yields while maintaining environmental sustainability.

Materials and Methods

The study was conducted during the Kharif season of 2022-2023 at the central research farm of the Department of Agroforestry and Silviculture, College of Forestry, Sam Higginbottom University of Agriculture, Technology, and Science. The study was conducted during the Karif season of 2022 to 2023. The research took place at a central research farm belonging to the Department of Agro forestry and Silviculture at College of Forestry, which is affiliated with Sam Higginbottom University of Agriculture, Technology, and Science. The experimental site was in the Prayagraj district, positioned at a geographical latitude and longitude of 250 40' N and 810 85' E, with an altitude of 92 meters above sea level. This site was situated near the Yamuna Riverbank in the east and the indalpur village in the north.

Comment [DY1]: Add reference:

Yadav, L. P. and Gangadhara, K. 2021. Cultivation of Drumstick for Socio-Economic Upliftment. In: Current Horticulture: Improvement, Production, Plant Health Mangement and Value-Addition Volume II, Singh, B., Singh, A.K., Tomar, B.S., Ranjan, J.K. and Dutt, S. (eds.) Brillion Publishing, New Delhi, ISBN: 978-93-90757-79-4, e-ISBN: 978-93-90757-91-6 Pp. 201-206

Yadav, L. P.; Gangadhara, K. and Apparao, V. V. (2022). Evaluation of drumstick variety Thar Harsha under rainfed semi-arid conditions for growth, yield and quality along with antioxidant potentiality and nutrient content. *S. Afr. J. Bot.* 148: 112-122.
<https://doi.org/10.1016/j.sajb.2022.04.005>

Yadav, L. P., Gangadhara, K., Apparao, V.V., Singh, S. and Saroj, P. L., 2018. Drumstick: a multi-purpose plant. *Indian Horticulture* 63(5), 89-94.

Yadav, L. P.; Gangadhara, K.; Apparao, V.V.; Singh, A. K. (2023). Antioxidants and nutritional counters of drumstick (*Moringa oleifera*) germplasm under rainfed semi-arid region. *Indian Journal of Agricultural Sciences* 93 (10): 1073–1079.
<https://doi.org/10.56093/ijas.v93i10.131824>

Yadav, L. P.; Gangadhara, K.; Apparao, V.V.; Singh, A. K.; Rane, J.; Kaushik, P.; Sekhawat, N.; Malhotra, S. K.; Ramniwas; Rai, A. K.; Yadav, S. L. and Berwal, M. K. (2024). Nutritional, antioxidants and protein profiling of leaves of *Moringa oleifera* germplasm. *S. Afr. J. Bot.* 165: 443-454.
<https://doi.org/10.1016/j.sajb.2024.01.012>

The present study examined the effects of FYM, vermicompost, and neem cake on the growth and yield attributes of citronella, using a Randomized Block Design (RBD) to account for spatial variations and assess the independent and combined influences of these factors on citronella yield.~~The present study investigated the effect of FYM, Vermicompost and neem cake on growth and yield attributes of citronella crop. A Randomized Block Design (RBD) was utilized to address potential spatial variations within the research site. This design enables the concurrent examination of the independent and combined influences of various factors on the dependent variable's citronella yield attributes.~~ Farmyard Manure (FYM) application was implemented 12 t ha^{-1} at distinct rates: 25%, 33.3%, 50%, 75%, and 100%. Vermicompost (VC) were applied 6 t ha^{-1} at different rates: 25%, 33.3%, 50%, 75%, and 100%. Neem-cake (NC) were incorporated 4 t ha^{-1} at different rates: 25%, 33.3%, 50%, 75%, and 100%. With following treatment combinations T_0 -Control, T_1 - 50%FYM+50%Neem cake, T_2 - 50%FYM+50% Vermicompost, T_3 - 50%NC+50% Vermicompost, T_4 - 75%FYM+25% Vermicompost, T_5 - 75%FYM+25%NC, T_6 - 75% NC+25% Vermicompost, T_7 - 75%NC+25% FYM, T_8 - 75%VC+25%NC, T_9 - 75%VC+25% FYM, T_{10} - 100%FYM, T_{11} - 100% Vermicompost, T_{12} - 100%Neem-cake, T_{13} - 33.3%FYM+33.3% VC+33.3% NC. Citronella, a perennial grass, is propagated vegetatively using slips, as it does not produce viable seeds. Mature clumps, typically yielding about 50 slips after a year, are carefully divided into slips with 1-3 tillers each. Before planting, fibrous roots and leaves are clipped off the slips. Planting was conducted in July using slips sourced from CIMAP, Lucknow, Uttar Pradesh. The field was ploughed 2-3 times and divided into 2×2 meter plots with irrigation channels beside each plot. Organic manures—FYM, vermicompost, and neem cake—were evenly applied across the field. Slips were planted vertically at $60 \text{ cm} \times 60 \text{ cm}$ spacing with a $90 \text{ cm} \times 90 \text{ cm}$ gap between rows, at a depth of 10 cm. Proper drainage was ensured to avoid waterlogging, which can stunt growth, and planting on slopes was preferred. If rain did not occur within 24 hours of planting, irrigation was provided immediately.

The growth attributes of the citronella were quantified at various time intervals following planting, specifically at 30, 60, 90, 120, 150 and 180 DAS (Days after sowing), encompassing parameters such as plant height, no. of leaves per plant and no. of tiller per plant and yield attributes of the citronella were observed such as fresh weight of plant, dry weight of plant and total herbage yield during the consecutive years of 2022 and 2023.

The experimental data were analysed using the Randomized Block Design (RBD) to assess the effects of organic manures (FYM, vermicompost, and neem cake) and cultivation conditions (moringa-based agroforestry and open-field systems) on the growth and yield

attributes of citronella (*Cymbopogon winterianus*). Analysis of variance (ANOVA) was performed to determine the significance of differences among treatments, with the main factors being Cultivation condition (shaded vs. open field), Organic manure type, and Interaction between cultivation condition and manure type. The treatment means were compared using the least significant difference (LSD) test at a 5% level of significance. Statistical analyses were performed using SPSS. This approach ensured robust evaluation of treatment effects and their interactions on the measured parameters.

Results and Discussion:

Plant height

The effect of citronella treatments on plant height across different growth stages (30, 60, 90, 120, 150, and 180 DAS) under both open and shaded conditions is presented in Tables 1 and 2. ~~The detailed of results were showed in table 1 and 2 on effect of citronella treatments on plant height was observed across different growth stages (30, 60, 90, 120, 150, and 180 DAS) under both open and shaded conditions.~~ Treatment T13 consistently resulted in the highest plant height at all stages, highlighting its superior efficacy in promoting growth. At 30 DAS, T13 showed a marked improvement over the control (T0), indicating its ability to stimulate early-stage growth. By 60 and 90 DAS, the advantage of T13 became more pronounced, particularly under shaded conditions, which generally supported greater growth compared to open environments. Treatments such as T11 and T12 also demonstrated notable results, though they fell short of T13's performance.

At 120 DAS, the growth pattern remained consistent, with T13 continuing to outperform, while other treatments like T5, T6, and T9 exhibited competitive growth. By 150 DAS, plant heights across treatments stabilized, but T13 maintained its leading position, reflecting its sustained effectiveness in later growth stages. Finally, at 180 DAS, the height gains began to plateau, but T13's dominance persisted under both open and shaded conditions, significantly outperforming the control and other treatments.

Overall, the application of organic manure treatments positively influenced plant height, with T13 emerging as the most effective treatment throughout the study period. The shaded conditions further enhanced the growth, suggesting that citronella responds well to partial shade, which likely optimizes its physiological processes for height gain. These results underscore the importance of treatment selection and environmental conditions in maximizing citronella growth.

Comment [DY2]: Citation in Discussion part is missing in this section; add references with citation

Number of leaves/Plant

The effect of citronella treatments on the number of leaves per plant was assessed across various growth stages (30, 60, 90, 120, 150, and 180 DAS) under open and shaded conditions. Treatment T13 consistently recorded the highest leaf count across all stages, indicating its superior efficacy in enhancing vegetative growth. At 30 DAS, T13 significantly outperformed the control (T0), reflecting its ability to promote early leaf initiation and development. By 60 and 90 DAS, the trend became more pronounced, with shaded conditions generally resulting in higher leaf numbers compared to open conditions, suggesting enhanced adaptability and growth potential under reduced light intensity. Treatments T11 and T12 also exhibited competitive performance during this period, albeit at slightly lower levels than T13.

At 120 DAS, T13 showed superior performance, with significant increases in leaf production under both shaded and open conditions. Treatments T5, T6, and T9 also exhibited improved leaf counts, contributing to vegetative biomass. By 150 DAS, the number of leaves had substantially increased across all treatments, with T13 maintaining its lead. Shaded conditions continued to promote higher leaf production, highlighting citronella's preference for partial shade. At 180 DAS, leaf production peaked, with T13 achieving the highest leaf count in both environments, significantly outperforming the control and other treatments.

~~At 120 DAS, T13 continued to exhibit superior performance, with notable increases in leaf production under both conditions. Other treatments, such as T5, T6, and T9, also demonstrated enhanced leaf counts, contributing to the vegetative biomass. By 150 DAS, the number of leaves had substantially increased across treatments, with T13 maintaining its dominance. Shaded conditions continued to support higher leaf production compared to open conditions, emphasizing the plant's preference for partial shade. At 180 DAS, leaf production reached its peak, with T13 achieving the maximum leaf count in both environments, significantly outperforming the control and all other treatments.~~

The statistical analysis confirmed that the differences among treatments were significant (F-test: Significant) at all stages, demonstrating the efficacy of the applied organic manures. The consistent performance of T13 across all growth stages highlights its potential for maximizing foliage production, particularly under shaded conditions. These findings underscore the critical role of treatment selection and environmental optimization in enhancing the vegetative performance of citronella.

Number of Tiller/Plant

The number of tillers per plant in citronella was assessed at different growth stages (30, 60, 90, 120, 150, and 180 DAS) under open and shaded conditions. Treatment T13 consistently recorded the highest tiller count, demonstrating its superior ability to enhance tillering throughout the growth period. At 30 DAS, T13 significantly outperformed the control (T0), reflecting its effectiveness in promoting early tiller initiation. By 60 and 90 DAS, the tiller count showed a marked increase across all treatments, with shaded conditions generally producing higher tiller numbers compared to open conditions. Other treatments, such as T11 and T12, also exhibited notable performance, albeit less pronounced than T13.

At 120 DAS, T13 continued to lead in tiller production under both open and shaded environments, while treatments such as T5, T6, and T9 also contributed substantially to tiller growth. By 150 DAS, the trend of increasing tiller numbers remained consistent, with T13 maintaining its dominance, particularly in shaded conditions. At 180 DAS, tiller production reached its peak, with T13 achieving the maximum tiller count under both conditions, significantly surpassing the control (T0) and other treatments.

Statistical analysis indicated that the differences among treatments were significant (F-test: significant) at all growth stages, confirming the efficacy of the treatments in enhancing tillering. The superior performance of T13 across all stages underscores its potential for maximizing tiller production, especially in shaded environments, which consistently supported better tillering. These results highlight the importance of organic manure treatments and environmental optimization in promoting vegetative propagation in citronella.
~~The statistical analysis indicated that the differences among treatments were significant (F test: Significant) at all growth stages, confirming the efficacy of the treatments in enhancing tillering. The superior performance of T13 across all stages underscores its potential for maximizing tiller production, especially in shaded environments, which were consistently favourable for tillering. These results highlight the importance of organic manure treatments and environmental optimization in promoting vegetative propagation in citronella.~~

Fresh Weight

The effect of different organic manures on the fresh weight of citronella java plants at harvest under open and shaded conditions in a moringa-based agroforestry system was significant.

Across the treatments, T13 consistently recorded the highest fresh weight, indicating its superior ability to enhance biomass accumulation.

Under shaded conditions, the fresh weight was generally higher compared to open conditions, reflecting the favourable microenvironment created by the moringa canopy. Specifically, T13 achieved the maximum fresh weight in both conditions, demonstrating its effectiveness in optimizing plant growth in agroforestry settings. Treatments T11 and T12 also showed competitive results, although they were outperformed by T13.

The statistical analysis (F-test: Significant) confirmed that differences among treatments were substantial. These results suggest that the integration of organic manures, particularly T13, within moringa-based agroforestry systems can significantly improve the biomass productivity of citronella java, making it a promising strategy for sustainable agricultural practices [add citation](#).

Dry Weight

The application of different organic manures significantly influenced the dry weight of citronella java plants at harvest under open and shaded conditions within a moringa-based agroforestry system. Among the treatments, T13 consistently demonstrated the highest dry weight accumulation, highlighting its superior efficacy in enhancing the physiological and biomass attributes of citronella.

Dry weight under shaded conditions was generally higher than in open environments, likely due to the moderated microclimatic conditions provided by the moringa canopy, which optimized plant growth processes. Treatments such as T11 and T12 also recorded competitive dry weights but were consistently outperformed by T13. The control (T0) exhibited the lowest dry weight under both conditions, underscoring the necessity of organic manure application for improved biomass production.

Statistical analysis indicated that the differences among treatments were significant (F-test: Significant) at harvest. These findings demonstrate the potential of integrating high-performance organic manures, particularly T13, in moringa-based agroforestry systems to maximize the dry matter yield of citronella java. This underscores the importance of selecting appropriate amendments and leveraging agroforestry practices for sustainable and productive biomass management [add citation](#).

Total Herbage Yield

The application of different organic manures significantly impacted the total herbage yield of citronella java under open and shaded conditions in a moringa-based agroforestry system. Among all treatments, T13 consistently recorded the highest herbage yield, highlighting its superior efficacy in enhancing overall biomass production. Shaded conditions generally resulted in higher yields compared to open conditions, likely due to the favourable microclimatic effects of the moringa canopy, which improved moisture retention and moderated temperature fluctuations.

The statistical analysis (F-test: Significant) revealed substantial differences among treatments, confirming the effectiveness of organic manure applications. T13 demonstrated the most pronounced effect on herbage yield under both conditions, followed by T11 and T12, which showed competitive performance. The control (T0) consistently recorded the lowest yield, emphasizing the importance of organic amendments for optimal biomass accumulation.

These results underscore the potential of T₁₃ as a high-performance organic input for enhancing herbage yield in citronella java, particularly when cultivated within agroforestry systems. The integration of organic manures with moringa-based shading practices offers a sustainable strategy to maximize biomass productivity, optimize resource use, and improve overall system efficiency [add citation](#).

Conclusion

[The study highlights the significant benefits of integrating organic manures with agroforestry practices to boost citronella growth and yield. The treatment combining FYM, vermicompost, and neem cake \(T13\) consistently delivered superior results, yielding higher plant biomass and herbage compared to other treatments. The moringa-based agroforestry system further enhanced productivity by providing shaded conditions, demonstrating the synergy between organic inputs and agroforestry. This research offers valuable insights for resource-limited farmers, showing how eco-friendly practices can optimize agricultural output while improving long-term soil and environmental health. By showcasing the economic and ecological advantages of sustainable citronella cultivation, the study provides important guidance for advancing sustainable agricultural practices in regions where aromatic crops and agroforestry systems are viable.](#)

The study establishes the value of integrating organic manures with agroforestry practices to enhance citronella growth and yield. The treatment combining FYM, vermicompost, and neem cake (T13) consistently shown the superior results, achieving higher plant biomass and herbage yield compared to other treatments. The moringa-based agroforestry system provided shaded conditions that further amplified productivity, underscoring the synergy between organic inputs and agroforestry. This research provides actionable guidance for resource-limited farmers, illustrating how eco-friendly practices can optimize agricultural outputs and contribute to long-term soil and environmental health. By demonstrating the economic and ecological benefits of sustainable citronella cultivation, the study has significant implications for advancing sustainable agricultural practices in regions where both aromatic crops and agroforestry systems are viable.

References

- Nair, P. K. R. (2011). Agroforestry systems and environmental quality: Introduction. *Journal of Environmental Quality*, 40(3), 784-790.
- Singh, B., & Prasad, R. (2015). Role of aromatic plants in sustainable agriculture. *Agricultural Research*, 4(4), 358-367.
- Manna, M. C., et al. (2005). Long-term effect of fertilizers and manure on soil fertility and sustainability. *Journal of Sustainable Agriculture*, 26(2), 5-24.
- Kaushal, R., et al. (2014). Nutrient cycling in *Moringa oleifera* based agroforestry systems. *Agroforestry Systems*, 88(2), 201-213.
- Bhattacharyya, R., et al. (2015). Effects of organic manures on soil fertility and crop productivity under sustainable agricultural systems. *Agriculture, Ecosystems & Environment*, 213, 322-332.
- Kumar, S., et al. (2014). Integration of medicinal and aromatic plants in agroforestry systems: A sustainable approach. *Industrial Crops and Products*, 62, 25-32.
- Arancon, N.Q., Edwards, C.A., & Atiyeh, R.M. (2004). Effects of vermicomposts on growth and marketable fruits of field-grown tomatoes, peppers, and strawberries. *Pedobiologia*, 47(5-6), 731-735.
- Rao, P.S.C., & Reddy, G.B. (2009). Use of neem cake for improving nutrient use efficiency and managing soil health. *Journal of Sustainable Agriculture*, 33(1), 73-92.
- Nath, A.J., & Das, A.K. (2011). Managing agroforestry systems for biodiversity conservation and sustainable production: A case study of citronella-based systems. *Agroforestry Systems*, 81(2), 123-134.

Gomez, K.A., & Gomez, A.A. (1984). *Statistical Procedures for Agricultural Research*. John Wiley & Sons.

Add reference:

Yadav, L. P. and Gangadhara, K. 2021. Cultivation of Drumstick for Socio-Economic Upliftment. In: Current Horticulture: Improvement, Production, Plant Health Mangement and Value-Addition Volume II, Singh, B., Singh, A.K., Tomar, B.S., Ranjan, J.K. and Dutt, S. (eds.) Brillion Publishing, New Delhi, ISBN: 978-93-90757-79-4, e-ISBN: 978-93-90757-91-6 Pp. 201-206

Yadav, L. P.; Gangadhara, K. and Apparao, V. V. (2022). Evaluation of drumstick variety Thar Harsha under rainfed semi-arid conditions for growth, yield and quality along with antioxidant potentiality and nutrient content. *S. Afr. J. Bot.* 148: 112-122. <https://doi.org/10.1016/j.sajb.2022.04.005>

Yadav, L. P., Gangadhara, K., Apparao, V.V., Singh, S. and Saroj, P. L., 2018. Drumstick: a multi-purpose plant. *Indian Horticulture* 63(5), 89-94.

Yadav, L. P.; Gangadhara, K.; Apparao, V.V.; Singh, A. K. (2023). Antioxidants and nutritional counters of drumstick (*Moringa oleifera*) germplasm under rainfed semi-arid region. *Indian Journal of Agricultural Sciences* 93 (10): 1073–1079. <https://doi.org/10.56093/ijas.v93i10.131824>

Yadav, L. P.; Gangadhara, K.; Apparao, V.V.; Singh, A. K.; Rane, J.; Kaushik, P.; Sekhawat, N.; Malhotra, S. K.; Ramniwas; Rai, A. K.; Yadav, S. L. and Berwal, M. K. (2024). Nutritional, antioxidants and protein profiling of leaves of *Moringa oleifera* germplasm. *S. Afr. J. Bot.* 165: 443-454. <https://doi.org/10.1016/j.sajb.2024.01.012>

Table.1Effect of different organic manures on plant height (cm) at 30, 60, 90, 120, 150 and 180 DAS of *citronella java* under open

Treatments	30 DAS		60 DAS		90 DAS		120 DAS		150 DAS		180 DAS	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
T₀	33.43	35.43	41.23	40.21	50.12	49.25	75.21	72.23	100.21	100.25	115.21	111.35
T₁	35.23	40.12	46.21	42.23	66.22	61.23	86.23	82.23	115.12	115.23	126.12	126.32
T₂	38.26	41.21	46.28	42.23	67.15	61.22	88.12	82.32	120.21	120.12	126.23	126.42
T₃	40.35	41.25	46.32	42.35	67.32	62.32	87.21	82.35	121.12	125.32	127.21	126.35
T₄	41.25	41.24	47.22	42.26	67.25	62.25	88.23	85.23	120.22	121.32	127.12	127.35
T₅	41.32	42.15	47.26	43.53	67.35	63.45	88.32	85.35	121.23	125.23	127.24	130.12
T₆	42.23	41.26	47.25	42.35	67.28	63.52	88.23	86.45	123.21	125.32	127.25	131.23
T₇	42.23	41.32	47.32	42.53	67.35	63.45	89.25	86.35	123.25	124.32	128.12	132.23
T₈	43.25	40.23	47.28	41.36	67.29	65.26	89.28	85.54	123.24	125.42	128.21	134.25
T₉	43.21	42.32	48.21	43.34	68.25	65.55	89.23	87.65	123.25	126.32	128.23	134.32
T₁₀	43.32	40.21	48.12	41.26	68.34	65.23	90.01	86.35	124.25	128.35	128.25	134.35
T₁₁	43.54	41.32	48.12	42.31	69.21	65.32	89.26	86.23	124.26	128.32	128.27	134.35
T₁₂	43.26	40.25	48.21	41.34	68.26	64.25	89.21	86.35	124.21	127.45	128.28	134.36
T₁₃	43.62	44.25	55.21	43.25	70.21	65.35	90.25	88.23	125.12	130.23	128.31	140.23
F-test	NS	NS	S	S	S	S	S	S	S	S	S	S

condition

Table.3Effect of different organic manures on No. of leaves/plant at 30, 60, 90, 120, 150 and 180 DAS of *citronella java* under open

Treatments	30 DAS		60 DAS		90 DAS		120 DAS		150 DAS		180 DAS	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
T₀	4.42	5.21	6.62	7.25	7.42	9.23	8.21	12.23	9.26	16.15	10.22	20.12
T₁	5.52	6.23	7.45	8.42	8.46	9.21	9.23	12.21	10.02	17.12	11.23	20.32

condition

UNDER PEER REVIEW

T ₂	6.32	6.32	7.12	8.23	8.54	9.12	9.26	12.24	10.35	17.24	11.26	22.31
T ₃	6.62	7.25	8.12	8.23	9.21	9.35	10.22	12.32	11.52	17.25	12.26	22.16
Treatments	6.43	6.32	7.25	8.32	8.42	9.34	9.12	12.35	10.23	15.23	11.24	18.24
T ₄	6.43	6.32	7.25	8.32	8.42	9.34	9.12	12.35	10.23	15.23	11.24	18.24
T ₅	6.43	6.32	7.25	8.32	8.42	9.34	9.12	12.35	10.23	15.23	11.24	18.24
T ₆	7.45	8.23	8.28	10.25	9.45	11.54	10.28	13.45	11.33	17.37	18.64	24.56
T ₇	7.44	7.45	8.38	8.48	9.46	10.63	10.63	13.43	11.34	17.45	15.25	20.32
T ₈	7.45	7.54	8.45	8.26	9.32	9.45	10.42	15.21	11.62	18.24	12.24	25.62
T ₉	8.54	8.32	9.26	10.21	10.21	11.12	11.21	15.24	12.63	18.45	13.22	24.65
T ₁₀	7.43	8.35	8.24	10.12	9.28	11.23	10.21	15.45	12.42	18.35	13.25	24.51
T ₁₁	7.48	8.65	8.54	10.13	9.31	11.32	10.32	15.46	12.52	18.65	13.24	26.84
T ₁₂	8.42	8.25	8.65	10.32	9.45	11.25	10.26	15.54	12.52	18.35	13.26	25.65
T ₁₃	8.42	8.65	9.12	10.25	10.32	11.34	11.32	15.64	13.32	18.67	13.85	28.45
F-test	S	S	S	S	S	S	S	S	S	S	S	S

Table.4 Effect of different organic manures on No. of leaves/plant at 30, 60, 90, 120, 150 and 180 DAS of *citronella java* under shade condition in moringa based agro-forestry system.

T ₂	6.42	7.54	7.58	9.10	8.48	10.66	9.64	14.45	10.45	14.35	15.46	22.32
T ₃	7.77	8.84	8.45	9.45	9.75	11.15	10.11	14.65	12.54	15.65	18.45	22.15
T ₄	7.45	8.45	8.75	9.46	9.49	11.62	12.72	14.85	13.15	16.26	19.45	25.45
Treatments	30 DAS	60 DAS	90 DAS	120 DAS	150 DAS	180 DAS						
T ₅	8.22	8.23	8.32	8.23	8.23	8.23	8.23	8.23	8.23	8.23	8.23	8.23
F ₀	8.44	7.33	6.33	6.38	6.75	7.31	9.24	13.62	18.43	18.64	21.34	24.35
F ₁	8.44	6.63	6.33	6.38	6.48	8.51	8.43	13.23	16.32	18.46	19.12	23.43
T ₈	8.78	8.45	9.47	9.62	10.46	13.64	13.32	17.26	16.45	15.65	22.34	25.56
T ₉	9.75	9.14	10.49	10.84	10.50	12.45	12.34	15.16	16.45	14.35	23.42	24.59
T ₁₀	8.78	8.45	9.15	9.65	11.45	12.15	14.64	15.66	18.15	16.64	24.42	24.15
T ₁₁	7.87	8.65	8.78	9.85	10.65	12.45	10.54	15.43	18.45	16.53	23.44	26.85
T ₁₂	8.45	8.45	9.85	9.84	9.78	13.15	14.24	18.16	18.45	16.65	24.44	25.59
T ₁₃	9.45	9.55	10.75	10.85	10.46	13.65	15.31	18.32	18.15	16.66	25.12	28.95
F-test	S	S	S	S	S	S	S	S	S	S	S	S

Table.5 Effect of different organic manures on No. of tiller/plant at 30, 60, 90, 120, 150 and 180 DAS of *citronella java* under open condition

T ₂	5.53	7.25	7.45	8.52	8.26	9.23	9.54	11.26	10.42	15.26	11.45	21.15
T ₃	5.65	7.26	7.54	8.65	8.32	9.26	9.35	12.25	10.54	15.25	11.65	21.25
Treatments	5.39 DAS	7.45	7.62 DAS	8.35	8.87 DAS	10.12	9.33 DAS	12.35	10.65 DAS	16.21	11.25 DAS	21.26
T ₅	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
T ₆	5.45	6.12	7.45	7.84	9.57	10.56	10.13	14.56	15.45	18.54	24.44	20.55
T ₆	6.52	7.15	8.71	8.45	9.56	10.26	10.35	12.65	11.45	16.24	12.54	22.45
T ₇	5.58	7.24	7.23	8.62	8.52	10.28	9.62	12.63	10.53	16.41	11.52	22.35
T ₈	6.26	8.24	8.62	9.21	9.35	10.56	10.26	12.62	11.34	15.42	12.45	23.45
T ₉	6.35	8.45	8.58	9.12	9.65	11.21	10.24	13.21	11.43	15.24	12.43	23.41
T ₁₀	6.45	8.42	8.65	9.67	9.34	11.26	10.23	12.23	11.65	15.41	12.34	24.45
T ₁₁	6.75	8.64	8.63	9.32	9.43	11.45	10.52	12.41	11.45	15.53	12.42	23.54
T ₁₂	6.36	8.35	8.54	9.81	9.75	11.65	10.85	12.43	11.65	15.60	12.75	24.21
T ₁₃	7.24	8.84	9.53	9.92	10.21	12.12	11.25	13.23	12.42	16.21	13.68	25.21
F-test	S	S	S	S	S	S	S	S	S	S	S	S

Table.6 Effect of different organic manures on No. of tiller/plant at 30, 60, 90, 120, 150 and 180 DAS of *citronella java* under shade condition in moringa based agro-forestry system.

Table.7Effect of different organic manures on Fresh weight of plant (kg plant⁻¹) at harvest of *citronella java* under open and shade condition in moringa based agro-forestry system.

Treatments	Open			Shade		
	2022	2023	Pooled	2022	2023	Pooled
T₀	9.10	12.56	11.40	10.23	14.45	11.78
T₁	10.12	13.26	12.75	12.23	17.45	13.79
T₂	10.22	15.56	14.91	14.25	17.45	13.84
T₃	11.21	14.56	15.21	15.85	17.64	14.43
T₄	12.32	15.47	15.02	14.56	18.74	15.53
T₅	13.22	16.56	15.61	14.65	18.85	16.04
T₆	12.52	15.56	14.54	13.52	17.84	15.18
T₇	14.22	18.46	17.36	16.25	21.45	17.84
T₈	10.22	16.26	15.76	15.26	18.75	14.49
T₉	12.23	17.84	16.00	14.15	19.85	16.04
T₁₀	12.26	18.48	17.37	16.25	21.45	16.86
T₁₁	12.65	19.45	18.86	18.26	21.54	17.10
T₁₂	12.23	17.45	16.85	16.25	19.54	15.89
T₁₃	15.56	22.45	20.66	18.86	24.45	20.01
F-test	S	S	S	S	S	S
C.D. (P=0.005)	0.74001	0.907	0.49822	0.46	0.8031	0.46131
SE(m)	0.24583	0.3013	0.1655	0.15	0.26678	0.15324
SE(d)	0.34765	0.4261	0.23406	0.21	0.37729	0.21672
C.V.	3.54653	3.12334	1.80578	1.78	2.40088	1.69861

Table.8Effect of different organic manures on dry weight of plant (kg plant⁻¹) at harvest of *citronella java* under open and shade condition in moringa based agro-

Treatments	Open			Shade		
	2022	2023	Pooled	2022	2023	Pooled
T₀	3.45	3.47	3.46	3.41	3.81	3.63
T₁	4.65	4.65	4.65	4.54	4.86	4.70
T₂	4.65	4.58	4.62	3.56	4.68	4.12
T₃	3.45	4.65	4.05	4.56	4.70	4.62
T₄	4.95	4.85	4.90	4.58	4.98	4.77

forestry system.

T ₅	4.85	4.75	4.80	4.51	4.67	4.57
T ₆	4.95	4.84	4.90	5.45	4.61	5.02
T ₇	5.45	4.95	5.20	4.65	4.73	4.70
Treatments	4.56	Open	4.69	3.45	Shade	3.99
	4.82			4.59		
	2022	2023	Pooled	2022	2023	Pooled
T ₉	4.86	4.41	4.64	4.56	4.34	4.46
T ₀	10.58	16.12	13.35	14.52	17.32	15.92
T ₁₀	5.21	4.62	4.92	5.01	4.60	4.82
T ₁₁	14.87	17.12	16.00	16.51	18.45	17.48
T ₁	4.87	4.25	4.56	4.65	5.03	4.80
T ₂	14.59	16.62	15.61	16.53	18.65	17.59
T ₁₂	4.95	4.75	4.85	4.54	4.87	4.70
T ₃	14.95	17.65	16.30	16.25	18.15	17.20
T ₁₃	5.89	5.56	5.73	5.50	5.73	5.58
T ₄	14.78	17.48	16.13	16.56	18.65	17.61
F-test	S	S	S	S	S	S
	14.55	16.65	15.60	16.35	18.56	17.46
C.D. (P=0.005)	0.22701	0.24036	0.17002	0.19869	0.20192	0.14888
	14.35	16.75	15.55	16.26	18.65	17.46
SE(m)	0.07541	0.07985	0.05648	0.066	0.06708	0.04946
	14.75	16.45	15.60	16.34	18.62	17.48
SE(d)	0.10665	0.11292	0.07987	0.09334	0.09486	0.06994
C.V.	2.73987	2.97188	2.07675	2.5417	2.45663	1.86079

Table.9 Effect of different organic manures on total herbage yield (t ha⁻¹) at harvest of *citronella java* under open and shade condition in moringa based agro-forestry system.

T₈	14.14	16.65	15.40	16.44	18.35	17.40
T₉	14.51	16.15	15.33	16.56	18.45	17.51
T₁₀	14.25	16.35	15.30	16.54	18.65	17.60
T₁₁	14.65	16.45	15.55	16.64	18.26	17.45
T₁₂	14.35	16.56	15.46	16.57	18.36	17.47
T₁₃	15.26	18.35	16.81	16.78	19.62	18.20
F-test	S	S	S	S	S	S
C.D. (P=0.005)	0.45678	0.79938	0.44315	0.55373	0.81278	0.53521
SE(m)	0.15174	0.26555	0.14721	0.18395	0.27	0.17779
SE(d)	0.16969	0.15876	0.0853	0.14892	0.16687	0.11237
C.V.	0.3612	0.71933	1.09257	0.67474	0.75607	1.36421

UNDER PEER REVIEW