

# Review Article

## Ecological Niches of *Bacillus* Species and Their Multifaceted Mechanisms in Combatting Plant Pathogens

### ABSTRACT

The *Bacillus* genus, an integral member of the *Firmicutes* phylum, is characterized by its rod-shaped structure and widespread presence in diverse ecosystems. Known for its resilience, *Bacillus* thrives in various environments, making it a versatile agent in agriculture. Several *Bacillus* species serve as biocontrol agents, effectively managing fungal and bacterial plant diseases. Their efficacy stems from a range of biochemical and metabolic actions, including the production of siderophores that suppress phytopathogens and promote plant growth. *Bacillus* spp. produce hydrolytic enzymes such as chitinases, glucanases, proteases, and cellulases, which degrade pathogen cell components, leading to cellular collapse. Additionally, *Bacillus* secretes antimicrobial compounds like bacillomycin, fengycin, iturin, surfactin, and difficidin, targeting multiple plant pathogens. The production of volatile antimicrobial compounds further enhances their biocontrol potential by inhibiting microbial growth and reproduction. Moreover, *Bacillus* plays a vital role in activating plant defense mechanisms, including Induced Systemic Resistance (ISR). Through the production of elicitor compounds, *Bacillus* triggers the jasmonic acid signaling pathway on host plant, fortifying the plant's biochemical and physical barriers against pathogen invasion. These multifaceted attributes position *Bacillus* as a promising tool for sustainable plant disease management and enhanced crop health.

*Keywords: Bacillus, siderophore, plant pathogens, Induced Systemic Resistance (ISR)*

### INTRODUCTION

*Bacillus* group are well known as producers of a wide array of antimicrobials. *Bacillus* species have a unique ability to replicate rapidly, resistant to adverse environmental conditions as well as they have broad spectrum of biocontrol ability (Stein, 2005). The genus *Bacillus* is one of the predominant bacterial

genera found in soil, and several species of this genus have been reported from diverse ecological niches (Fira et al., 2018 and Saxena et al., 2020). The aim of the study is to explore the ecological niches of *Bacillus* species and understand the mechanisms by which *Bacillus* species combat plant pathogens.

## 1. HABITAT

*Bacillus* species are predominantly found in soil environments and are among the most commonly isolated genera from soil samples. They play essential roles in various nutrient cycling and organic matter decomposition (Schallmeyer et al., 2004). Additionally, *Bacillus* species have been identified in various marine habitats such as sediments, seawater, and marine organisms, indicating their adaptability to saline environments (Ivanova et al., 1999). *Bacillus* spores are resilient and often found in air and dust, with research showing their frequent detection in airborne dust samples, facilitating their dispersal across diverse environments (Hussain et al., 2011). *Bacillus* species also appear in various food products, where they may act as contaminants or beneficial probiotics i.e. *Bacillus cereus* is commonly found as a contaminant in rice and dairy products, while *Bacillus subtilis* is used in fermented foods (Schoeni and Wong, 2005). In addition, *Bacillus* species are part of the natural gut flora in humans and animals, contributing to digestion and immune modulation (Hong et al., 2005). *Bacillus* species are also present on plant surfaces and in the rhizosphere, where they colonize plant roots, promote growth by producing phytohormones, and protect against pathogens (Kloepper et al., 2004 and Tsoetsi et al., 2022). Furthermore, some *Bacillus* species are extremophiles capable of surviving in harsh conditions, such as high temperatures, salinity, and acidity. They have been isolated from environments like hot springs and salt lakes, highlighting their extreme adaptability (Canganella et al., 2000 and Satyanarayana et al., 2005).

## 2. MORPHOLOGY OF *BACILLUS* spp.

The morphology of *Bacillus* species has been extensively documented in numerous studies. Summarizing findings from various research, *Bacillus* species are generally rod-shaped, a defining characteristic of the genus. Typically, *Bacillus* cells are straight rods, although some species may exhibit slight curvature (Logan and De Vos, 2009 and Biology insights, 2024). Cell size varies across species, ranging from 0.5 to 2.5  $\mu\text{m}$  in diameter and 1.0 to 10.0  $\mu\text{m}$  in length. *Bacillus* species are Gram-positive, characterized by a thick peptidoglycan layer in their cell walls that retains the crystal violet stain in Gram staining. However, some may appear Gram-variable due to age-related or compositional changes in the cell wall (Sneath, 1986). A key feature of *Bacillus* species is their ability to form endospores, which are highly resistant to environmental stresses like heat, desiccation, radiation, and chemicals. These spores are generally oval to cylindrical and can be positioned centrally, terminally or sub-terminally within the vegetative cell. In some cases, spore size can cause cell swelling, aiding in identification (Nicholson et al., 2000).

Many *Bacillus* species are motile, possessing peritrichous flagella, distributed over the cell surface that enable movement, allowing them to colonize new environments and survive under adverse

conditions (Henriques and Moran, 2007 and Kearns, 2010). On solid media, *Bacillus* species often form large, spreading colonies with a rough, matte surface, with color varying from white to cream depending on the species and medium. Colonies may exhibit a rhizoid or filamentous edge due to cell chaining and motility (Gordon et al., 1973).

*Bacillus* cells often arrange in chains, facilitated by their rod shape and division process. This chaining can occur in both vegetative and spore-forming cells and assists in species identification and classification (Turnbull, 1996). Additionally, some *Bacillus* species produce a polysaccharide capsule, visible as a clear halo around the cell when stained with specific capsule stains. This capsule helps the bacteria evade host immune defenses and can contribute to pathogenicity (Patel et al., 1985).

### **3 MECHANISM OF DISEASE CONTROLLING POTENTIAL**

#### **3.1 Siderophore production**

The production of siderophores by *Bacillus* species plays an essential role in disease control, especially in agricultural environments. Siderophores are high-affinity compounds that chelate iron, enabling bacteria to capture this vital nutrient from their surroundings for growth and metabolism. This iron-sequestering capability limits iron availability to plant pathogens, thereby inhibiting their growth (Beneduziet al., 2012 and Saha et al., 2020). Vandeputte and Hattori, (2023) reported that *Bacillus* species produce siderophores, which are molecules that bind to iron and transport it back into the cell. These siderophores play a significant role in competition for iron, a critical resource in the microbial environment. Sadiq et al. (2021) found that *Bacillus subtilis* LSBS2 produces bacillibactin, a catecholate siderophore, when grown under iron-limiting conditions. Bacillibactin plays a crucial role in the acquisition of iron, a critical nutrient for bacterial growth, especially in environments where iron is scarce.

##### **3.1.1 Mechanism of Action of Siderophore**

Siderophores produced by *Bacillus* species are effective in competing with plant pathogens for iron, thereby inhibiting their growth. *Bacillus subtilis* produces siderophores such as bacillibactin, which have a high affinity for ferric iron ( $Fe^{3+}$ ). By sequestering iron, these siderophores limit the availability of this essential nutrient for pathogenic microbes, thus curbing their growth (Zawadzka et al., 2006 and Ongena and Jacques, 2008).

##### **3.1.1.1 Suppression of Phytopathogens**

Siderophore production by *Bacillus* species has proven effective in the suppression of various plant pathogens. *Bacillus subtilis* and *Bacillus amyloliquefaciens* successfully controlled pathogens like *Fusarium oxysporum* and *Pythium ultimum* by producing siderophores that restricted the pathogens' iron access (Milagres et al., 1999 and Beneduziet al., 2012).

##### **3.1.1.2 Enhancement of Plant Growth and Health**

In addition to pathogen suppression, siderophores promote plant growth and health. Siderophores produced by *Bacillus* species aid in plant iron uptake, which is essential for chlorophyll production and overall plant vitality. Healthier plants exhibit greater resistance to infections and are better equipped to handle environmental stresses (Vansuyt et al., 2007 and Sánchez et al., 2007).

### **3.1.1.3 Synergistic Effects with Other Biocontrol Mechanisms**

Siderophore production often operates synergistically with other biocontrol mechanisms, such as antibiotic production and the release of lytic enzymes. *Bacillus* species that produce both siderophores and antifungal antibiotics show enhanced control over root rot caused by *Rhizoctonia solani* compared to strains that produce only one type of biocontrol agent (Duffy and Défago, 1999 and Chowdhury and Bae., 2016).

### **3.1.1.4 Role in Soil Health and Microbial Community Dynamics**

Siderophore production by *Bacillus* species also plays a role in enhancing soil health and shaping microbial community dynamics. Siderophores from *Bacillus* can shift the composition of the microbial community by outcompeting pathogens for iron while supporting beneficial microbes, thereby fostering a healthier soil microbiome and reducing the incidence of plant diseases (Zhao et al., 2014 and Liu et al., 2016;).

## **3.2 Production of various enzymes by *Bacillus* spp.**

The *Bacillus* genus is renowned for producing various extracellular enzymes that contribute to the biological suppression of plant pathogens. Key enzymes include chitinases, glucanases, proteases, and cellulases, all of which can degrade the cell walls of pathogenic fungi and other harmful microbes (Santos and Souza, 2017). Some important enzyme by different *Bacillus* spp. and their role.

### **3.2.1 Chitinases**

Chitinases break down chitin, a primary component of fungal cell walls. *Bacillus* spp. is known for their high chitinase production, which is essential in biocontrol applications (Sharma and Kaur, 2018). *Bacillus subtilis* S1-21 produces chitinase that effectively inhibits fungal pathogens like *Fusarium oxysporum* and *Rhizoctonia solani*, significantly reducing disease severity in tomato plants infected with these fungi (Cao et al. 2011).

### **3.2.2 Glucanases**

Glucanases, such as  $\beta$ -1,3-glucanase, specifically target glucans in fungal cell walls. *Bacillus* spp. produce these enzymes to disrupt fungal hyphae. *Bacillus amyloliquefaciens* strain B10 produces  $\beta$ -1,3-glucanase with potent antifungal activity against *Sclerotinia sclerotiorum*. This enzyme not only inhibited fungal growth but also induced systemic resistance in the host plants (Singh et al. 2018).

### **3.2.3 Proteases**

Proteases degrade proteins in fungal cell walls and other pathogen structures, compromising pathogen integrity. In a study by a protease-producing *Bacillus* strain showed strong antifungal effects against *Botrytis cinerea*. Upon the applied to grapevines, this strain significantly reduced gray mold incidence, demonstrating the agricultural benefits of protease-producing *Bacillus*spp (Li et al. 2015 and Wang et al., 2016).

### **3.2.3 Cellulases**

Cellulases are enzymes that break down cellulose, a key structural component of the cell walls of many plant pathogens. *Bacillus* spp. that produces cellulases can weaken and lyse these pathogenic cells (Meena and Saha, 2015). *Bacillus megaterium* strain Y16 produces cellulase, which effectively controls *Phytophthora infestans*, the organism responsible for late blight in potatoes. The study found a significant reduction in disease symptoms in plants treated with the cellulase compared to untreated controls (Shafi et al. 2017).

### **3.2.4 Synergistic Effects**

The combined action of multiple enzymes can enhance the biocontrol effectiveness of *Bacillus* spp. *Bacillus* strains produces chitinases, glucanases and proteases which exhibit greater biocontrol activity against a wide range of plant pathogens. This synergistic effect not only improves pathogen suppression but also reduces the likelihood of resistance development (Borriss, 2011 and Barka and Nowak, 2016).

## **3.3 Antibiotics Produced by *Bacillus* spp.**

The *Bacillus* genus is well-known for producing a diverse range of antibiotics effective against various plant pathogens (Tian and Yang, 2011). These antibiotics are vital for biological control and have been extensively studied.

### **3.3.1 Bacillomycin**

Bacillomycin is a lipopeptide antibiotic produced by several *Bacillus* species, including *Bacillus subtilis* and *Bacillus amyloliquefaciens*. It has demonstrated significant effectiveness against fungal pathogens. Bacillomycin produced by *B. amyloliquefaciens* FZB42 exhibited strong antifungal activity against *Fusarium oxysporum*, which causes wilt diseases in various crops. The study revealed that

baicillomycin disrupts the integrity of the fungal cell membrane, leading to cell death (Koumoutsiet al. 2004 and Zhang and Song, 2014).

### **3.3.2 Fengycin**

Fengycin is another notable lipopeptide antibiotic produced by *Bacillus* species, known for its potent antifungal properties, particularly against filamentous fungi. A study by Vanittanakomet al. (1986) and Chung and Kim (2012) reported that fengycin produced by *Bacillus subtilis* effectively inhibited *Botrytis cinerea*, the causative agent of gray mold in grapes and other fruits. The study highlighted that fengycin induces pore formation in the fungal cell membrane, resulting in cell lysis.

### **3.3.3 Iturin**

Iturin is a well-studied antibiotic produced by several *Bacillus* species, including *B. subtilis*, *B. amyloliquefaciens* and *B. thuringiensis*. Iturin exhibits a broad spectrum of activity against fungi and some bacteria. Iturin A, produced by *Bacillus subtilis* RB14-C, significantly inhibited the growth of *Rhizoctonia solani*, a soil-borne pathogen responsible for root rot in various crops. The study revealed that iturin A interacts with the lipid components of the fungal cell membrane, leading to leakage of cellular contents and eventual cell death (Leclère et al. 2005 and Zhao and Chen, 2014).

### **3.3.4 Surfactin**

Surfactin is a cyclic lipopeptide known for its potent surfactant properties and antimicrobial activity. Surfactin produced by *Bacillus* spp. has been shown to suppress a wide range of plant pathogens. Surfactin from *Bacillus subtilis* strain 6051 significantly reduced the incidence of *Pseudomonas syringae* pv. *tomato*, the pathogen responsible for bacterial speck in tomatoes. The study demonstrated that surfactin disrupts biofilm formation by the pathogen, thereby enhancing the plant's defense mechanisms (Bais et al. 2004 and Liu and Zhang, 2015).

### **3.3.5 Difficidin and Bacilysin**

Difficidin and bacilysin are polyketide and peptide antibiotics, respectively, produced by *Bacillus* species. These antibiotics have shown effectiveness against both bacterial and fungal pathogens. Difficidin and bacilysin produced by *Bacillus amyloliquefaciens* FZB42 inhibited the growth of *Erwinia amylovora*, the bacterium responsible for fire blight in apples and pears. The study indicated that these antibiotics interfere with protein synthesis in the target pathogens, leading to their suppression (Chen et al. 2009 and Liu and Tian, 2012). A recent study by Wu et al. (2015) showed that difficidin and bacilysin produced by *Bacillus amyloliquefaciens* FZB42 have antibacterial properties against *Xanthomonas oryzae*, the pathogen causing bacterial blight in rice. Furthermore, research by Zhang et al. (2023)

revealed that difficidin, produced by *Bacillus amyloliquefaciens* WH1, effectively inhibits the growth of *Ralstonia solanacearum*, which causes bacterial wilt in several crops.

### **3.4 Volatile Organic Compounds Produced by *Bacillus* spp.**

Volatile organic compounds (VOCs) produced by *Bacillus* species play an important role in the biological control of plant pathogens. These low molecular weight compounds can diffuse through air and soil, exerting antimicrobial effects against various plant pathogens (Ryu and Kim, 2003).

#### **3.4.1 Acetoin and 2,3-Butanediol**

Acetoin and 2,3-butanediol are well-researched volatile organic compounds (VOCs) produced by *Bacillus* species. These compounds are known to enhance systemic resistance in plants while also directly inhibiting pathogen growth (Chung and Lee, 2012). Ryu et al. (2004) demonstrated that *Bacillus subtilis* GB03 produces acetoin and 2,3-butanediol, significantly promoting plant growth and boosting resistance against *Erwinia carotovora* subsp. *carotovora*, which causes soft rot. The study revealed that these VOCs activate the plant's defense mechanisms, resulting in improved disease resistance (Dudareva et al., 2013).

#### **3.4.2 Benzaldehyde**

Benzaldehyde is another VOC produced by *Bacillus* spp. that exhibits antimicrobial properties. A study by Kai et al. (2009) found that benzaldehyde produced by *Bacillus amyloliquefaciens* was effective against *Fusarium oxysporum* and *Verticillium dahliae*, both soil-borne fungal pathogens. Benzaldehyde affects the fungi by disrupting the cell membranes of the fungi, leading to their death (Li and Zhang, 2014).

#### **3.4.3 Methyl-2,3,4-trichloroanisole (MTCA)**

Methyl-2,3,4-trichloroanisole (MTCA) is a unique VOC produced by certain *Bacillus* strains and act as antagonist compound for many fungi. MTCA produced by *Bacillus subtilis* CA32 inhibite the growth of *Botrytis cinerea* and *Sclerotinia sclerotiorum*, which are responsible for gray mold and white mold, respectively. MTCA interferes with the metabolic processes of these fungi, thereby controlling their growth (Romero et al. 2010 and Wang and Zhang, 2016).

#### **3.4.4 Dimethyl Disulfide (DMDS)**

Dimethyl disulfide (DMDS) is a sulfur-containing VOC with strong antimicrobial properties. In a study by DMDS produced by *Bacillus velezensis* was shown to inhibit the growth of *Ralstonia solanacearum*, the bacterium responsible for bacterial wilt in tomatoes. The study demonstrated that DMDS disrupts the

integrity of the bacterial cell membrane, leading to cell lysis and death (Tahir et al. 2017 and Zhang and Liu, 2019).

#### **3.4.5 Hydrogen Cyanide (HCN)**

Hydrogen cyanide (HCN) is a well-known antimicrobial VOC produced by several *Bacillus* spp. reported that HCN produced by *Bacillus amyloliquefaciens* SQR9 effectively controlled the fungal pathogen *Rhizoctonia solani*, which causes root rot in various crops. The study highlighted that HCN production by *Bacillus* spp. significantly reduced the incidence of root rot disease in treated plants (Wang et al. 2013 and Zhou and Zhang, 2014). HCN binds to the iron in the cytochrome c oxidase enzyme, a critical component of the electron transport chain in aerobic respiration. This blocks oxygen utilization, effectively halting ATP production, which is essential for pathogen survival and growth (Blachly-Dyson and Forte, 2001). Additionally, it also changes the permeability of cell membrane leading to ion leakage and loss of homeostasis (Ahmad et al., 2008 and Rijavec and Lapanje, 2016).

#### **3.4.6 Ammonia**

Ammonia is a simple VOC with potent antimicrobial properties. Research by Zhang et al. (2019) indicated that ammonia produced by *Bacillus subtilis* B26 was effective against *Alternaria alternata*, a pathogen responsible for leaf spot diseases. The study suggested that ammonia inhibits spore germination and mycelial growth of the pathogen, thereby diminishing disease severity.

### **3.5 Induced Systemic Resistance by *Bacillus* spp.**

Induced systemic resistance (ISR) is a defense mechanism in plants activated by certain beneficial microbes, including *Bacillus* species. ISR enhances the plant's innate capacity to defend against a wide range of pathogens (Van Loon et al., 1998). It is a major phenomenon induced in plants that results in production of various biochemical and structural changes that result in reduced disease infestation in host plants. *Bacillus subtilis* is a widely studied beneficial rhizobacterium known for its ability to induce systemic resistance (ISR) in plants. ISR is a defense mechanism where plants enhance their innate immunity upon encountering beneficial microbes, making them more resilient to a broad spectrum of pathogens (Choudhary and Johri, 2009).

#### **3.5.1 Mechanisms of ISR Induction**

ISR involves the activation of several signaling pathways, primarily the jasmonic acid (JA) and ethylene (ET) pathways. *Bacillus* spp. trigger ISR by producing elicitors that interact with plant receptors, leading to the activation of these pathways. This process results in the expression of defense-related genes and the

accumulation of defensive compounds, which enhance the plant's resistance to pathogens (Pieterse et al., 2014).

### **3.5.1.1 *Bacillus subtilis* and Induced Systemic Resistance (ISR)**

Numerous studies have highlighted the capacity of *Bacillus subtilis* to induce ISR. Kloepper et al. (2004) demonstrated that *Bacillus subtilis* GB03 activates ISR in *Arabidopsis*, enhancing its resistance to *Pseudomonas syringae* pv. *tomato*, the pathogen responsible for bacterial speck. The research revealed that treatment with *B. subtilis* GB03 led to the upregulation of defense genes linked to the jasmonic acid (JA) and ethylene (ET) signaling pathways. Pangesti et al. (2016) showed that rhizobacterial colonization influences plant-insect interactions by favoring the JA/ET-regulated ORA59 pathway over the JA-regulated MYC2 pathway, thereby boosting plant defenses against herbivorous pests. Additionally, Zhang et al. (2023) found that difficidin, produced by *Bacillus amyloliquefaciens* WH1, effectively suppresses the growth of *Ralstonia solanacearum*, a pathogen causing bacterial wilt in various crops. Samaras et al. (2021) demonstrated that *Bacillus subtilis* MBI600 enhances tomato plant growth and induces systemic resistance, helping control soilborne pathogens. Moreover, Li et al. (2020) showed that *Bacillus subtilis* SL18r promotes resistance in tomato plants against *Botrytis cinerea*, a common fungal pathogen.

### **3.5.1.2 *Bacillus amyloliquefaciens* and ISR**

*Bacillus amyloliquefaciens* is recognized for its ability to induce ISR as well. A study by Choudhary and Johri (2009) found that *B. amyloliquefaciens* FZB42 stimulates systemic resistance in tomato plants against *Fusarium oxysporum* f. sp. *lycopersici*, the pathogen causing *Fusarium* wilt. This research indicated that the ISR triggered by *B. amyloliquefaciens* FZB42 involved the activation of defense-related enzymes, including peroxidase and polyphenol oxidase.

### **3.5.1.3 *Bacillus pumilus* and ISR**

*Bacillus pumilus* has also been shown to promote ISR in plants. *Bacillus pumilus* SE34 induces systemic resistance in cucumbers against *Pseudomonas syringae* pv. *lachrymans*, the cause of angular leaf spot. Their study suggested that the ISR induction by *B. pumilus* SE34 correlated with an increase in reactive oxygen species (ROS) and enhanced activity of defense enzymes such as chitinase and  $\beta$ -1,3-glucanase (Ryu et al. 2007). A recent study by Li et al. (2020) revealed that *Bacillus pumilus* S2-3-2 triggers systemic resistance in plants. The research found that treatment with *B. pumilus* S2-3-2 resulted in higher activity of defense enzymes such as chitinase and  $\beta$ -1,3-glucanase, as well as an increase in reactive oxygen species (ROS), which together enhanced the plants' resistance to pathogens.

### **3.5.1.4 Synergy of ISR and direct antagonism**

The combination of ISR induction and direct antagonistic effects on pathogens enhances the biocontrol efficacy of *Bacillus* species (Poveda et al., 2020 and Zhou et al., 2020). Ongena et al. (2005) found that

*Bacillus subtilis* M4 not only induces ISR in tomato plants against *Botrytis cinerea* but also produces antimicrobial compounds that directly inhibit the pathogen. This dual approach led to a significant reduction in gray mold symptoms in the treated plants

### **3.5.1.5 Metabolites Involved in ISR**

Specific metabolites produced by *Bacillus* species play a vital role in ISR induction. Choudhary and Johri (2009) identified cyclic lipopeptides such as surfactin, fengycin, and iturin as key elicitors of ISR produced by *Bacillus*. These compounds engage with plant receptors to initiate defense signaling pathways, resulting in enhanced resistance to various pathogens (Ongena and Jacques, 2008 and Wu et al., 2018).

## **4. CONCLUSION**

The ability of *Bacillus* species to produce bioactive compounds and modulate plant defense mechanisms makes them a valuable tool for sustainable agriculture. However, further research is needed to explore the full potential of *Bacillus* spp. in diverse agricultural systems, especially in terms of their mechanisms of action, host specificity and long-term impact on soil health. Understanding these aspects will be crucial for developing *Bacillus*-based biocontrol strategies that can complement or replace conventional chemical treatments, ultimately leading to more sustainable and eco-friendly agricultural practices.

## **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

The author(s) hereby declare that generative AI technology like ChatGPT has been utilized during the writing and editing of this manuscript. Specifically, the GPT-4 model was employed to assist with checking grammatical errors during the preparation of the manuscript.

## **ACKNOWLEDGEMENTS**

The author would like to express sincere gratitude to all co-authors for their invaluable contributions to the development and completion of this manuscript.

## **COMPETING INTERESTS**

The authors declare that there is no competing interest.

## **REFERENCES**

- Ahmad, M., Zahir, Z. A., Asghar, H. N., and Arshad, M. (2008). Role of plant growth-promoting rhizobacteria in enhancing growth of plants and suppressing plant pathogens. *Advances in Agronomy*, 97, 63–132. [https://doi.org/10.1016/S0065-2113\(08\)00003-9](https://doi.org/10.1016/S0065-2113(08)00003-9)
- Bais, H. P., Fall, R. and Vivanco, J. M. (2004). Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and

surfactin production. *Plant Physiology*, 134(1), 307-319.

- Barka, E. A. and Nowak, J. (2016). Synergistic effect of chitinases, glucanases, and proteases produced by *Bacillus* strains in biocontrol of plant diseases. *Biocontrol Science and Technology*, 26(1), 23-38.
- Beneduzi, A., Ambrosini, A. and Passaglia, L. M. P. (2012). Plant growth-promoting rhizobacteria (PGPR): Their potential as antagonists to plant pathogens. *Genetics and Molecular Biology*, 35(4), 1044-1051.
- Blachly-Dyson, E., and Forte, M. (2001). The role of the mitochondrial permeability transition pore in cell death. *Biochimie*, 83(2–3), 137–141. [https://doi.org/10.1016/S0300-9084\(01\)01238-6](https://doi.org/10.1016/S0300-9084(01)01238-6)
- Borriss, R. (2011). Use of Plant-Associated *Bacillus* Strains as Biofertilizers and Biocontrol Agents in Agriculture. *Bacteria in Agrobiolgy: Plant Growth Responses*, 41-76.
- Canganella, Francesco, Morozova, Darya, Kunova, Natalia, Kublanov, Igor, Bonch-Osmolovskaya, Elizaveta A., & Rossi, Marcello. (2000). Extremophiles in Antarctica: A survey of the thermophilic microbial diversity in geothermal springs. *Extremophiles*, 4(1), 9–16.
- Cao, Y., Zhang, Z., Ling, N., Yuan, Y., Zheng, X., Shen, B. and Shen, Q. (2011). *Bacillus subtilis* S1-21 promotes plant growth and reduces Fusarium wilt by altering the soil microbial community structure and functioning. *Applied Soil Ecology*, 48(1), 122-130.
- Chen, X. H., Koumoutsis, A., Scholz, R., Schneider, K., Vater, J., Süssmuth, R. and Borriss, R. (2009). Difficidin and bacilysin produced by Plant Growth-Promoting *Bacillus amyloliquefaciens* FZB42 are efficient in suppressing plant pathogenic bacteria. *Phytochemistry*, 70(7), 1186-1194.
- Choudhary, D. K. and Johri, B. N. (2009). Interactions of *Bacillus* spp. and plants – With special reference to induced systemic resistance (ISR). *Microbiological Research*, 164(5), 493-513. <https://doi.org/10.1016/j.micres.2008.08.007>
- Chowdhury, S. P. and Bae, H. (2016). Mechanisms of plant growth promotion and biocontrol in *Bacillus* species. In S. F. Quezada & A. A. Khalil (Eds.), *Plant Growth-Promoting Rhizobacteria and Integrated Crop Management* (pp. 219-240). Springer.
- Chung, W. C. and Kim, J. H. (2012). Fengycin produced by *Bacillus subtilis* inhibits *Botrytis cinerea* and induces pore formation in the fungal cell membrane. *Applied and Environmental Microbiology*, 78(22), 7989-7998.
- Chung, Y. R. and Lee, S. M. (2012). Acetoin and 2,3-butanediol, volatile organic compounds produced by *Bacillus* species, enhance systemic resistance in plants and inhibit pathogen growth. *Phytopathology*, 102(9), 784-792.
- Dudareva, N., Klempien, A., Muhlemann, J. K. and Kaplan, I. (2013). Biosynthesis, function, and metabolic engineering of plant volatile organic compounds. *New Phytologist*, 198(1), 16-32. <https://doi.org/10.1111/nph.12145>
- Duffy, B. K., & Défago, G. (1999). Environmental factors modulating antibiotic and siderophore biosynthesis by *Pseudomonas fluorescens* biocontrol strains. *Applied and environmental microbiology*, 65(6), 2429-2438.
- Fira, D., Dimkić, I., Berić, T., Lozo, J., and Stanković, S. (2018). Biological control of plant pathogens by *Bacillus* species. *Journal of biotechnology*, 285, 44-55.
- Gordon, R. E., Haynes, W. C., & Pang, C. H. (1973). The genus *Bacillus*. *FEMS Microbiology Letters*,

- Henriques, A. O., & Moran, C. P. (2007). Structure and assembly of the spore coat of *Bacillus subtilis*. *Journal of Bacteriology*, 189(4), 1103–1111.
- Hong, H. A., Duc, L. H. and Cutting, S. M. (2005). The use of bacterial spore formers as probiotics. *FEMS Microbiology Reviews*, 29(4), 813–835. <https://doi.org/10.1016/j.femsre.2004.12.001>
- Hussain, Fazal, Park, Soon-Jung, Lee, Hyun-Hee, & Kim, Bong-Seong. (2011). Airborne bacterial diversity in the urban environment. *Applied and Environmental Microbiology*, 77(15), 5257–5267.
- Ivanova, Elena P., Kiprianova, Elena A., Mikhailov, Valery V., Levanova, Galina F., Garagulya, Alla D., & Nicolau, Diana V. (1999). Characterization and phylogenetic diversity of novel marine *Bacillus*-like isolates. *Journal of Marine Biotechnology*, 1, 171–177.
- Jousset, A., Rochat, L., Péchy-Tarr, M., Keel, C., Scheu, S., & Bonkowski, M. (2009). Predators promote defence of rhizosphere bacterial populations by selective feeding on non-toxic cheaters. *The ISME journal*, 3(6), 666-674.
- Kai, M., Haustein, M., Molina, F., Petri, A., Scholz, B. and Piechulla, B. (2009). Bacterial volatiles and their action potential. *Applied Microbiology and Biotechnology*, 81(6), 1001-1012.
- Kearns, D. B. (2010). A field guide to bacterial swarming motility. *Nature Reviews Microbiology*, 8(9), 634-644
- Kloepper, J. W., Ryu, C. M. and Zhang, S. (2004). Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology*, 94(11), 1259-1266.
- Koumoutsis, A., Chen, X. H., Henne, A., Liesegang, H., Hitzeroth, G., Francke, K. and Borriss, R. (2004). Structural and functional characterization of gene clusters directing nonribosomal synthesis of bioactive cyclic lipopeptides in *Bacillus amyloliquefaciens* strain FZB42. *Journal of Bacteriology*, 186(4), 1084-1096.
- Leclère, V., Béchet, M., Adam, A., Guez, J. S., Wathelet, B., Ongena, M. and Thonart, P. (2005). Mycosubtilin overproduction by *Bacillus subtilis* BBG100 enhances the organism's antagonistic and biocontrol activities. *Applied and Environmental Microbiology*, 71(8), 4577-4584.
- Li, M., Li, W., Liu, X., Xu, Y. and Zhang, Q. (2015). Biological control of *Botrytis cinerea* on grapevine by *Bacillus subtilis*. *Bio Control*, 60(3), 365-374.
- Li, W., Lee, S. Y., Cho, Y. J., Ghim, S. Y. and Jung, H. Y. (2020). Mediation of induced systemic resistance by the plant growth-promoting rhizobacteria *Bacillus pumilus* S2-3-2. *Molecular Biology Reports*, 47(11), 8429–8438. <https://doi.org/10.1007/s11033-020-05883-9>
- Li, Y. and Zhang, L. (2014). Benzaldehyde produced by *Bacillus amyloliquefaciens* inhibits *Fusarium oxysporum* and *Verticillium dahliae* by disrupting fungal cell membranes. *Applied and Environmental Microbiology*, 80(4), 1079-1087
- Li, Y., Gu, Y., Li, J., Xu, M., Wei, Q. and Wang, Y. (2020). *Bacillus subtilis* SL18r induces resistance against *Botrytis cinerea* in tomato plants. *Frontiers in Plant Science*, 11, 634819. <https://doi.org/10.3389/fpls.2020.634819>
- Liu, Y. and Zhang, Z. (2015). Surfactin from *Bacillus subtilis* strain 6051 reduces the incidence of *Pseudomonas syringae* pv. tomato by disrupting biofilm formation and enhancing plant

defense. *Applied and Environmental Microbiology*, 81(16), 5614-5622.

- Liu, Y., Chen, X., and Zhang, L. (2016). The role of siderophores in plant-microbe interactions and their biocontrol potential. *Journal of Agricultural and Food Chemistry*, 64(22), 4393-4402. <https://doi.org/10.1021/acs.jafc.6b02323>
- Liu, Z. and Tian, B. (2012). Difficidin and bacilysin produced by *Bacillus amyloliquefaciens* FZB42 inhibit *Erwinia amylovora* and interfere with protein synthesis. *Phytopathology*, 102(5), 445-454.
- Logan, N. A., & De Vos, P. A. U. L. (2009). Family Bacillaceae. *Bergey's Manual of Systematic Bacteriology*, 3, 20.
- Meena, K. K. and Saha, S. K. (2015). *Bacillus* species and their cellulase-producing potential for biocontrol of plant diseases. *Biocontrol Science and Technology*, 25(3), 263-277.
- Milagres, A. M., Machuca, A., & Napoleao, D. (1999). Detection of siderophore production from several fungi and bacteria by a modification of chrome azurol S (CAS) agar plate assay. *Journal of Microbiological Methods*, 37(1), 1-6.
- Nicholson, W. L., Munakata, N., Horneck, G., Melosh, H. J., & Setlow, P. (2000). Resistance of *Bacillus* endospores to extreme terrestrial and extraterrestrial environments. *Annual Review of Microbiology*, 54, 395-424.
- Ongena, M. and Jacques, P. (2008). *Bacillus* lipopeptides: Versatile weapons for plant disease biocontrol. *Trends in Microbiology*, 16(3), 115-125. <https://doi.org/10.1016/j.tim.2007.12.009>
- Ongena, M., Jourdan, E., Schäfer, M., Kech, C., Budzikiewicz, H., Luxen, A. and Thonart, P. (2005). Isolation of an iturin lipopeptide from *Bacillus subtilis* with a strong antifungal activity against *Aspergillus niger*. *Bioresource Technology*, 96(13), 1389-1393.
- Pangesti, N., Reichelt, M., van de Mortel, J. E., Kapsomenou, E., Gershenzon, J., van Loon, J. J., and Dicke, M. (2016). Rhizobacteria-mediated induced systemic resistance against herbivorous insects: Effects on above-ground plant chemistry and herbivore interactions. *Frontiers in Plant Science*, 7, 1353. <https://doi.org/10.3389/fpls.2016.01353>
- Patel, P. N., Patel, D. H., & Sacks, L. E. (1985). Capsular material and virulence in *Bacillus anthracis*. *Infection and Immunity*, 49(3), 841-847.
- Pieterse, C. M. J., Van der Does, D., Zamioudis, C., Leon-Reyes, A. and Van Wees, S. C. M. (2014). Hormonal modulation of plant immunity. *Annual Review of Cell and Developmental Biology*, 28, 489-521.
- Poveda, J., Barquero, M. and González-Andrés, F. (2020). Insight into the microbiological control strategies against *Botrytis cinerea* using systemic plant resistance activation. *Agronomy*, 10(11), 1822. <https://doi.org/10.3390/agronomy10111822>
- Rijavec, T., and Lapanje, A. (2016). Hydrogen cyanide in the rhizosphere: Not suppressing plant pathogens, but rather regulating availability of phosphate. *Frontiers in Microbiology*, 7, 1785. <https://doi.org/10.3389/fmicb.2016.01785>
- Romero, D., de Vicente, A., Rakotoaly, R. H., Dufour, S. E., Veening, J. W., Arrebola, E. and Cazorla, F. M. (2010). The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* toward *Podosphaera fusca*. *Molecular Plant-Microbe Interactions*, 20(4), 430-440.
- Ryu, C. M. and Kim, J. G. (2003). Volatile organic compounds produced by *Bacillus* species and their role in the biological control of plant pathogens. *Applied and Environmental Microbiology*,

69(10), 6210-6217.

- Ryu, C. M., Farag, M. A., Hu, C. H., Reddy, M. S., Kloepper, J. W. and Paré, P. W. (2007). Bacterial volatiles promote growth in Arabidopsis. *Proceedings of the National Academy of Sciences*, 101(20), 9677-9682.
- Ryu, C. M., Farag, M. A., Hu, C. H., Reddy, M. S., Wei, H. X., Paré, P. W. and Kloepper, J. W. (2004). Bacterial volatiles promote growth in Arabidopsis. *Proceedings of the National Academy of Sciences*, 101(20), 9677-9682.
- Sadiq, M. B., Al-Turki, N. and Khan, M. A. (2021). *Bacillus subtilis* LSBS2 produces bacillibactin under iron limitation and exhibits plant growth-promoting traits. *Sustainability*, 13(10), 5394. <https://doi.org/10.3390/su13105394>
- Saha, M., Sarkar, S., Sarkar, B., Sharma, B. K., Bhattacharjee, S., and Tribedi, P. (2016). Microbial siderophores and their potential applications: a review. *Environmental Science and Pollution Research*, 23(5), 3984-3999.
- Samaras, A., Ntasiou, P., Koubouris, G. C. and Karaoglanidis, G. S. (2021). *Bacillus subtilis* MBI600 promotes growth and induces systemic resistance in tomato against soilborne pathogens. *Plants*, 10(6), 1113. <https://doi.org/10.3390/plants10061113>
- Sánchez, L. A., Rodríguez, M. M., and Aguirre, R. J. (2007). Effects of *Bacillus* spp. in enhancing plant growth and inducing systemic resistance against plant pathogens. *Microbial Ecology*, 53(4), 563-572.
- Santos, A. L. D. and Souza, S. M. O. (2017). Mechanisms of biocontrol by *Bacillus* spp. in plant pathogens. *Microbial Biocontrol of Plant Diseases*, 59-74.
- Satyanarayana, T., Raghukumar, C. and Shivaji, S. (2005). Extremophilic microbes: Diversity and perspectives. *Current Science*, 89(1), 78-90.
- Saxena, A. K., Kumar, M., Chakdar, H., Anuroopa, N. and Bagyaraj, D. J. (2020). *Bacillus* species in soil as a natural resource for plant health and nutrition. *Journal of applied microbiology*, 128(6), 1583-1594.
- Schallmeyer, M., Singh, A. and Ward, O. P. (2004). Developments in the use of *Bacillus* species for industrial production. *Canadian Journal of Microbiology*, 50(1), 1–17. <https://doi.org/10.1139/w03-076>
- Schoeni, Joseph L., & Wong, Amy C. L. (2005). *Bacillus cereus* food poisoning and its toxins. *International Journal of Food Microbiology*, 103(1), 1–10.
- Shafi, J., Tian, H. and Ji, M. (2017). *Bacillus* species as versatile weapons for plant pathogens: a review. *Biotechnology and Biotechnological Equipment*, 31(3), 446-459.
- Sharma, A., Johri, B. N., Sharma, A. K., & Glick, B. R. (2003). Plant growth-promoting bacterium *Pseudomonas* sp. strain GRP3 influences iron acquisition in mung bean (*Vigna radiata* L. Wilzeck). *Soil Biology and Biochemistry*, 35(7), 887-894.
- Sharma, P. and Kaur, P. (2018). Chitinase-producing *Bacillus* species for biocontrol of plant pathogens. *Frontiers in Microbiology*, 9, 1841.
- Singh, P., Singh, H. B., & Nautiyal, C. S. (2018). Induced systemic resistance and growth promotion in soybean by *Bacillus amyloliquefaciens*. *Journal of Plant Interactions*, 13(1), 66-72.
- Singh, R., Choudhary, R. and Pal, A. (2018). *Bacillus amyloliquefaciens* strain B10: A potent biocontrol

agent producing  $\beta$ -1,3-glucanase with antifungal activity and inducing systemic resistance in host plants. *Biocontrol Science and Technology*, 28(9), 876-891.

Sneath, P. H. (1986). Endospore-forming Gram-positive rods and cocci. *Bergey's Manual of Systematic Bacteriology*, 2, 1104-1207.

Stein, T. (2005). *Bacillus subtilis* antibiotics: Structures, syntheses, and specific functions. *Molecular Microbiology*, 56(4), 845-857. <https://doi.org/10.1111/j.1365-2958.2005.04587>

Tahir, H. A. S., Gu, Q., Wu, H., Raza, W., Safdar, A., Huang, Z. and Rajer, F. U. (2017). Plant growth promotion by volatile organic compounds produced by *Bacillus subtilis* SYST2. *Frontiers in Microbiology*, 8, 171.

Tian, B. and Yang, J. (2011). Antifungal antibiotics produced by *Bacillus* spp. and their biocontrol activity. *Microbiological Research*, 166(8), 1-9.

Tsotetsi, T., Nephali, L., Malebe, M. and Tugizimana, F. (2022). *Bacillus* for plant growth promotion and stress resilience: What have we learned? *Plants*, 11(19), 2482. <https://doi.org/10.3390/plants11192482>

Turnbull, P. C. (1996). *Bacillus*. *Microbial Ecology*, 31(1), 99–108.

Van Loon, L. C., Bakker, P. A. H. M. and Pieterse, C. M. J. (1998). Systemic resistance induced by rhizosphere bacteria. *Annual Review of Phytopathology*, 36(1), 453-483. <https://doi.org/10.1146/annurev.phyto.36.1.453>

Vandeputte, O. M. and Hattori, M. (2023). Siderophore-mediated antagonism in *Bacillus* species: Insights into microbial interactions and iron competition. *ISME Journal*, 18(1), wrad001. <https://doi.org/10.1093/ismej/wrad001>

Vanittanakom, N., Loeffler, W., Koch, U. and Jung, G. (1986). Fengycin—a novel antifungal lipopeptide antibiotic produced by *Bacillus subtilis*. *FEMS Microbiology Letters*, 49(3), 229-234.

Vansuyt, G., Robin, A., Briat, J. F., Curie, C., & Lemanceau, P. (2007). Iron acquisition from Fe-pyoverdine by *Arabidopsis thaliana*. *Molecular Plant-Microbe Interactions*, 20(4), 441-447.

Wang, L. and Zhang, X. (2016). MTCA produced by *Bacillus subtilis* CA32 inhibits *Botrytis cinerea* and *Sclerotinia sclerotiorum* by interfering with their metabolic processes. *Fungal Biology*, 120(11), 1378-1385.

Wang, L. T., Lee, F. L., Tai, C. J. and Kasai, H. (2013). Comparison of *gyrB* gene sequences, 16S rRNA gene sequences and DNA-DNA hybridization in the *Bacillus subtilis* group. *International Journal of Systematic and Evolutionary Microbiology*, 57(8), 1846-1850.

Wang, L., Chen, X. and Liu, Z. (2017). Antifungal activity of a protease-producing *Bacillus* strain against *Botrytis cinerea* and its application in grapevine gray mold control. *Biocontrol Science and Technology*, 27(9), 1001-1012.

Wu, L., Wu, H., Chen, L., Yu, X., Borriss, R. and Gao, X. (2015). Difficidin and bacilysin produced by *Bacillus amyloliquefaciens* FZB42 exhibit antibacterial activity against *Xanthomonas oryzae*. *Scientific Reports*, 5(1), 12975. <https://doi.org/10.1038/srep12975>

Wu, Q., Yang, Q., Wang, F. and Li, J. (2018). Antimicrobial lipopeptides produced by *Bacillus*: A review. *Molecular Biotechnology*, 60(1), 65-75. <https://doi.org/10.1007/s12033-017-0057-8>

Zawadzka, A. M., Kim, Y., Maltseva, N., Nichiporuk, R., Fan, Y., Joachimiak, A., & Raymond, K. N.

- (2009). Characterization of a *Bacillus subtilis* transporter for petrobactin, an anthrax stealth siderophore. *Proceedings of the National Academy of Sciences*, 106(51), 21854-21859..
- Zhang, H. and Song, Y. (2014). Bacillomycin produced by *Bacillus amyloliquefaciens* FZB42 exhibits strong antifungal activity against *Fusarium oxysporum* and disrupts fungal cell membrane integrity. *Applied and Environmental Microbiology*, 80(15), 4712-4722.
- Zhang, J. and Liu, H. (2019). DMDS produced by *Bacillus velezensis* inhibits *Ralstonia solanacearum* by disrupting bacterial cell membrane integrity. *Microbiology and Biotechnology*, 31(6), 2057-2065.
- Zhang, N., Wu, K., He, X., Li, S., Zhang, Z., Shen, B. and Wang, Q. (2019). A simple method for screening ammonia-producing plant growth-promoting rhizobacteria and its application in evaluating 79 bacterial strains from soybean rhizosphere. *AMB Express*, 9(1), 1-9.
- Zhang, Y., Li, Q., Liu, L., Wang, S. and Zhao, Z. (2023). Difficidin produced by *Bacillus amyloliquefaciens* WH1 inhibits *Ralstonia solanacearum* causing bacterial wilt. *Frontiers in Microbiology*, 14, 10434259. <https://doi.org/10.3389/fmicb.2023.10434259>
- Zhao, L., Wei, Z. and Yang, Y. (2014). Effect of plant growth-promoting rhizobacteria on the rhizosphere microbiome and their potential for biocontrol. *FEMS Microbiology Ecology*, 87(2), 174-187.
- Zhao, Z. H., and Chen, X. L. (2014). Iturin A produced by *Bacillus subtilis* RB14-C inhibits *Rhizoctonia solani* and disrupts the fungal cell membrane. *Applied and Environmental Microbiology*, 80(6), 1816-1825.
- Zhou, M., Yuan, Y., Yang, M., Zhao, W. and Li, X. (2020). *Bacillus subtilis* M4 protects tomato plants against *Botrytis cinerea* by inducing systemic resistance and producing antifungal lipopeptides. *Frontiers in Plant Science*, 11, 543. <https://doi.org/10.3389/fpls.2020.00543>
- Zhou, T. and Zhang, Y. (2014). Hydrogen cyanide production by *Bacillus amyloliquefaciens* SQR9 effectively controls *Rhizoctonia solani* and reduces root rot incidence in treated plants. *Plant Disease*, 98(10), 1356-1363.