

Standardization of seed to solution ratio and hydropriming duration for seed priming of fenugreek (*Trigonella foenum-graecum* L.)

ABSTRACT

A laboratory experiment was carried out to standardize the seed to solution ratio and hydropriming duration for seed priming in fenugreek in the laboratory of department of seed science & technology, College of agriculture, UAS, Raichur during *Rabi* 2023-24. The experiment was laid out in two factorial Completely Randomized Design. The first factor consisted of seed to solution ratios viz., 1:1, 1:1.5, 1:2, 1:2.5 and 1:3. The second factor was hydro-priming durations viz., 8, 12, 16, 20 and 24 hours. The results revealed that priming in 1:2.5 seed to solution for 12 hours has recorded higher seed quality parameters viz., seed germination (97.25 %), seedling length (21.50 cm), seedling dry weight (11.24 mg) seedling vigour index-I and II (2091 & 1093). While, seed priming in 1:1 seed to solution for 8 hours recorded lower seed quality parameters (89.50 %, 15.99 cm, 8.30 mg, 1431 and 743, respectively). Based on the results, it can be concluded that for hydropriming in fenugreek 1:2.5 seed to solution ratio for 12 hours duration to get good quality seedlings.

Key words: Germination, hydro priming, seed to solution ratio, seedling vigour index.

INTRODUCTION

Fenugreek (*Trigonella foenum-graecum* L.) belongs to the Fabaceae family. It is a short duration crop and mainly grown as a leafy vegetable as well as a seed spice. It is one of the oldest medicinal plants with an exceptional medicinal and nutritional profile. Fenugreek seeds are frequently used in human medicine to manage hypercholesterolemia, cancer and diabetes as they possess hypoglycaemic, antilipidemic, anticarcinogenic and cholagogic properties (Murlidhar and Goswami, 2012). Its fresh tender leaves are also consumed as vegetable which are rich in iron, calcium, vitamins and essential amino acids like lysine, leucine and phenylalanine (Mungofa *et al.*, 2022). Seeds contain a substantial amount of fiber, phospholipids, glycolipids, oleic acid, linolenic acid, linoleic acid, choline, vitamins A, B1, B2, C, nicotinic acid, niacin, and other elements (Awulachew, 2022).

Vegetables are considered as essential building blocks of any diet, provides good source of vitamins, minerals, high dietary fiber and complex carbohydrates with low fat content. But per capita consumption of fruits and vegetables in India is very low (160 to 184 g/person/day) against world health organization standards of 400 g/person/day. So, we are in the stage of increasing the production and productivity of vegetable crops. Fenugreek is a valuable plant for its culinary and medicinal uses, but its growth is often hindered by poor seed quality including low germination rates and seed-borne diseases. The success of pre-sowing seed management techniques depends on the duration of priming and the seed to solution ratio. However, these factors vary with crops depending on their chemical composition. The efficacy

of treatments depends on the level of moisture content to which the seeds are hydrated just before radicle emergence. This study was aimed to standardize the appropriate seed to solution ratio and hydropriming duration to optimize fenugreek seed performance. To provide higher quality seeds, many researchers have developed new technologies called seed quality enhancement techniques. In the last two decades, seed priming, an effective seed invigoration method, has become a common seed treatment to increase the rate and uniformity of emergence and crop establishment in most vegetable and flower crops especially in advanced countries. The priming is influenced by several factors. Among them the most important factors are selection of priming agent, its concentration, the method of priming and duration of priming. As a first step studies were initiated to standardize the method of priming by evolving seed to solution ratio to be adopted for priming and the duration of priming using water, the universal priming agent suitable for all kinds of seed. Keeping these in view, the present experiment was conducted in fenugreek to standardize the seed to solution ratio and hydropriming duration for seed priming and its influence on seed quality characters.

MATERIAL AND METHODS

The laboratory experiment was conducted at Department of Seed Science and Technology, University of Agricultural Sciences, Raichur, located at 16° 15' North latitude and 77° 20' East longitude with an altitude of 389 m above mean sea level at Raichur, Karnataka during 2023-24. The experiment was laid out in Factorial completely randomised design with four replications. The first factor comprised of five seed to solution ratios *viz.*, 1:1 (S₁), 1:1.5 (S₂), 1:2 (S₃), 1:2.5 (S₄) and 1:3 (S₅). The second factor consisted five hydropriming durations *viz.*, 8 (D₁), 12 (D₂), 16 (D₃), 20 (D₄) and 24 hours (D₅). The cleaned and graded fenugreek seeds were hydroprimed with different seed to solution ratios (weight by volume) as per the treatments mentioned above, after soaking the seeds were taken out from water as per the different durations and they were uniformly spread on a blotting paper and kept for air drying under shade to re-dry to their original moisture level and then those seeds were further used for assessing various seed quality parameters. The observations were recorded on seed germination (%), seedling length (cm), seedling dry weight (mg), seedling vigour index-I and II.

Seed germination (%): The standard germination test was conducted in four replications of 100 seeds in between paper method and rolled towels were incubated in the walk-in seed germination chamber maintained with 25±2°C temperature and 90±5 % relative humidity. The

final count was taken on 14th day. The number of normal seedlings from each replication were counted and the mean germination was expressed in percentage (ISTA, 2024).

Seedling length (cm): From the germination test, ten normal seedlings were randomly selected from each treatment. The seedling length was measured from the tip of shoot to tip of root of the seedlings. The mean seedling length was expressed in centimeter (ISTA, 2024).

Seedling dry weight (mg): The ten normal seedlings which were used for seedling length measurement were placed in butter paper and dried in hot air oven maintained at 70 °C temperature for 24 hours. Later the seedlings were removed from hot air oven and allowed to cool in a desiccator for 20 minutes and weighed using electronic balance. The average weight was calculated and expressed in milligrams (ISTA, 2024).

Seedling vigour index-I and II: The seedling vigour index was calculated as per the formula given by Abdul -Baki and Anderson (1973).

Seedling vigour index - I = Seed germination (%) × Seedling length (cm)

Seedling vigour index - II = Seed germination (%) × Seedling dry weight (mg)

RESULTS AND DISCUSSION

The results showed that different seed to solution ratio and hydropriming durations had shown significance difference for seed germination (%), seedling length (cm), seedling dry weight (mg) were presented in table 1. And seedling vigour index-I and II were presented in table 2..

The seed germination is an important seed quality parameter. The significantly highest seed germination was recorded in 1:2.5 seed to solution ratio (96.55 %) compared to 1:1 seed to solution ratio (91.95 %). Among hydropriming durations 12 hours had recorded significantly highest seed germination (95.20 %) than 24 hours (93.35 %). In case of interactions, priming in 1:2.5 seed to solution ratio for 12 hours had recorded significantly highest seed germination (97.25 %) compared to 1:1 seed to solution ratio for 8 hours duration (89.50 %). The probable reason for enhancement of percentage and uniformity of germination of the hydroprimed seed might have completed pre-germination process, repair and synthesis of nucleic acids (DNA and mRNA), proteins (Ibrahim, 2019) and initiation of biochemical reactions and enzyme

activation. Wattanakulpakin *et al.* (2012) observed an increased germination percentage in maize due to hydropriming for 6 hours. The availability of optimum moisture at an appropriate duration to enhance germination and growth by hydrolyzing complex molecules into simpler sugars, which are readily utilized in the synthesis of auxins and proteins. The auxins produced help to soften cell walls, facilitating growth, while the proteins are readily used in the production of new tissues. Soaking in the equal volume of water for brinjal and chilli seeds for 12 hours significantly resulted in higher seed germination than untreated seeds which was observed by Ponnuswamy and Vijayalakshmi, (2011). Similar findings were reported by Georghiou *et al.* (1982) in tomato and Anbarasan and Srimathi (2015) in Redgram.

The 1:2.5 seed to solution ratio had significantly recorded longest seedling length (20.26 cm) compared to 1:1 seed to solution (17.02 cm). Among hydropriming durations, hydropriming for 12 hours had recorded significantly longest seedling length (18.97 cm) than 24 hours (17.73 cm). Among interactions, hydropriming in 1:2.5 seed to solution ratio for 12 hours duration (S₄D₂) had recorded significantly longest seedling length (21.50 cm) compared to hydropriming in 1:1 seed to solution ratio for 8 hours duration (S₁D₁) which had recorded shortest (15.99 cm) seedling length (Table 1). This may be due to better physiological activities which is associated with the washing away of the inhibitor ABA. The increased porosity of the seed coat enhanced the water and gas exchange, which in turn accelerated the germination process. This improvement in resource uptake supports more vigorous metabolic activities as a result, seedlings exhibit greater seedling length. These results are in conformity with Mohsen *et al.* (2013) in chickpea, Kamithi *et al.* (2016) in chickpea and Tanwar *et al.* (2023) in wheat.

The significantly highest seedling dry weight was recorded in 1:2.5 seed to solution (10.93 mg) compared to 1:1 seed to solution (8.96 mg). Among hydropriming durations, 12 hours had recorded significantly highest seedling dry weight (10.18 mg) than 24 hours (9.46 mg). Among interactions, 1:2.5 seed to solution with 12 hours had recorded significantly highest seedling dry weight (11.24 mg) compared to 1:1 seed to solution with 8 hours (8.30 mg) which had recorded lowest (Table 1). Higher accumulation of dry matter in hydroprimed seeds in appropriate seed to solution ratio for appropriate duration. Complete imbibition of seeds occurs when seeds absorb water fully for an effective duration leading to several physiological changes. As a result, the seed coat swells and becomes thinner, making it easier for the seed to leach out of chemicals that are responsible for seed dormancy, such as abscisic acid or other inhibitors. A reduction in the inhibitory effect, facilitates the seeds to proceed with

germination and vigorous seedling growth which resulted in the accumulation of more seedling dry weight. Similar findings of higher seedling dry weight was observed by Anbarasan and Srimathi (2015) in pulses.

There was significant difference for the seedling vigour indices among treatments. The significantly highest seedling vigour index-I (1956) and II (1056) were recorded in 1:2.5 seed to solution compared to 1:1 seed to solution (1565 and 824). Among hydropriming durations 12 hours had recorded significantly highest seedling vigour index-I (1808) and II (970) than 24 hours (1657 and 885). In case of interactions, 1:2.5 seed to solution with 12 hours priming had recorded significantly highest seedling vigour index-I (2091) and II (1093) compared to 1:1 seed to solution with 8 hours priming (1431 and 743) which had recorded lowest (Table 2). Comparing the different seed to solution ratios and durations of hydropriming, better shoot length, root length and seed germination were observed in seeds hydroprimed in 1:2.5 seed to solution ratio for 12 hours which finally resulted in higher seedling vigour indices and better performance of seedlings. The seedling vigour index increased mainly due to reduction in imbibition lag time for priming treatments (Bradford, 1986).

CONCLUSION

From the study, it can be concluded that fenugreek seeds hydroprimed in 1:2.5 seed to solution ratio for 12 hours to get better quality seedlings.

Table 1. Influence of seed to solution ratio and hydropriming duration on seed germination, seedling length and seedling dry weight of fenugreek

Seed germination (%)							Seedling length (cm)						Seedling dry weight (mg)					
Treatment	D ₁	D ₂	D ₃	D ₄	D ₅	Mean (S)	D ₁	D ₂	D ₃	D ₄	D ₅	Mean (S)	D ₁	D ₂	D ₃	D ₄	D ₅	Mean (S)
S ₁	89.50	93.00	92.75	92.50	92.00	91.95	15.99	17.49	17.36	17.24	17.02	17.02	8.30	9.29	9.24	9.06	8.89	8.96
S ₂	94.00	94.25	94.25	93.75	93.25	93.90	18.00	18.16	18.07	17.88	17.72	17.97	9.73	9.85	9.77	9.47	9.42	9.65
S ₃	95.75	96.00	95.75	95.00	94.75	95.45	18.77	19.02	18.88	18.51	18.39	18.71	10.26	10.37	10.30	10.06	9.93	10.18
S ₄	96.75	97.25	96.50	96.25	96.00	96.55	20.90	21.50	20.12	19.57	19.19	20.26	11.13	11.24	10.95	10.78	10.59	10.93
S ₅	95.25	95.50	94.50	91.50	90.75	93.50	18.59	18.69	18.25	16.54	16.33	17.68	10.09	10.15	9.88	8.63	8.50	9.45
Mean (D)	94.25	95.20	94.75	93.80	93.35	94.27	18.45	18.97	18.54	17.95	17.73	18.33	9.90	10.18	10.03	9.60	9.46	9.83
Factors	S.E.m. ±		CD @ 1 %			S.E.m. ±		CD @ 1 %			S.E.m. ±		CD @ 1 %					
S	0.23		0.85			0.11		0.41			0.09		0.32					
D	0.23		0.85			0.11		0.41			0.09		0.32					
S × D	0.51		1.90			0.25		0.92			0.19		0.72					

Legend:

Factor I: Seed to solution ratio (S):

S₁ = 1:1 S₂ = 1:1.5 S₃ = 1:2 S₄ = 1:2.5 S₅ = 1:3

Factor II: Duration of hydropriming (D):

D₁ = 8 hours D₂ = 12 hours D₃ = 16 hours D₄ = 20 hours D₅ = 24 hours

Table 2. Influence of seed to solution ratio and hydropriming duration on seedling vigour indices of fenugreek

Seedling vigour index-I							Seedling vigour index-II					
Treatment	D ₁	D ₂	D ₃	D ₄	D ₅	Mean (S)	D ₁	D ₂	D ₃	D ₄	D ₅	Mean (S)
S ₁	1431	1626	1610	1595	1566	1565	743	864	857	838	817	824
S ₂	1692	1712	1704	1676	1652	1687	915	929	921	888	879	906
S ₃	1798	1826	1808	1758	1742	1786	983	995	987	956	940	972
S ₄	2022	2091	1942	1883	1842	1956	1076	1093	1056	1037	1016	1056
S ₅	1770	1784	1725	1514	1482	1655	961	970	934	790	772	885
Mean (D)	1743	1808	1758	1685	1657	1730	935	970	951	902	885	929
Factors	S.Em. ±			CD @ 1 %			S.Em. ±			CD @ 1 %		
S	10.88			40.68			8.68			32.43		
D	10.88			40.68			8.68			32.43		
S × D	24.34			90.96			19.40			72.52		

Legend:

Factor I: Seed to solution ratio (S):

S₁ = 1:1 S₂ = 1:1.5 S₃ = 1:2 S₄ = 1:2.5 S₅ = 1:3

Factor II: Duration of hydropriming (D):

D₁ = 8 hours D₂ = 12 hours D₃ = 16 hours D₄ = 20 hours D₅ = 24 hours

REFERENCES

1. Abdul-Baki AA and Anderson JD. Vigour determination of soybean seeds by multiple criteria. *Crop Sci.* 1973; 13(1): 630-633.
2. Anbarasan R and Srimathi P. Standardization of hydro priming technique for seed invigoration in pulses. *Trends in Biosci.* 2015; 8(12): 3068-3073.
3. Awulachew, M. T., 2022, Health benefits and improvements of Fenugreek (*Trigonella foenum-graecum* L.) crop. *J. Agric. Res. Adv.* 4(3): 33-44.
4. Bradford, K. J., 1986, Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *Horti. Sci.* 21(5):1105-1112.
5. Fischer RA. Statistical methods for research workers, biological monographs and manuals. 1950; No. V. (11th ed.).
6. Georghiou K, Thanos CA, Tafas TP and Mitrakos K. Tomato seed germination osmotic pretreatment and far-red inhibition. *J. Exp. Bot.* 1982; 3: 1068-1073.
7. Ibrahim EAA. Fundamental processes involved in seed priming. *Priming and Pretreatment of Seeds and Seedlings: Implication in Plant Stress Tolerance and Enhancing Productivity in Crop Plants.* 2019; 1: 63-115.
8. ISTA. International Rules for Seed Testing. *Seed Sci. Technol.* 2024; 27: 25-30.
9. Kamithi KD, Wachira F and Kibe A. Effects of different priming methods and priming durations on enzyme activities in germinating chickpea (*Cicer arietinum* L.). *Am. J. Appl. Sci.* 2016; 1(1): 1-9.
10. Krishnakumary K, Venkatasubramaniam A and Mini C. Optimization of hydropriming techniques for cowpea seed invigoration. *Seed Res.* 2008; 36(1): 33-36.
11. Malarkodi K, Jeyavelan M and Ananthi M. Standardization of seed to solution ratio and soaking duration for priming of brinjal seeds. *Indian J. Agric. Res.* 2023; 57(2): 266-269.

12. Mohsen J, Adel B, Reza AS and Naser S. Impact of pre-sowing seed treatments and fertilizers on growth and yield of chickpea (*Cicer arietinum* L.) under rainfed conditions. *Nat. Monten.* 2013; 12(1): 217-229.
13. Mungofa N, Sibanyoni JJ, Mashau ME and Beswa D. Prospective role of indigenous leafy vegetables as functional food ingredients. *Molecule.* 2022; 27(22): 1-23.
14. Murlidhar MM and Goswami TK. A review on the functional properties, nutritional content, medicinal utilization and potential application of fenugreek. *J. Food Process. Tech.* 2012; 3(9): 181-184.
15. Panse VG and Sukhatme PV. Statistical methods for agricultural workers, ICAR, New Delhi. 1954; 167-174.
16. Ponnuswamy, AS and Vijayalakshmi V. Standardization of soaking duration and volume of solution for fortification of Tomato, Brinjal and Chilli Seeds. *Madras Agric. J.* 2011; 98: 370-371.
17. Tanwar H, Mor VS, Sharma S, Khan M, Bhuker A, Singh V, Yadav J, Sangwan S, Singh J, Yashveer S and Singh K. Optimization of 'on farm' hydropriming conditions in wheat: Soaking time and water volume have interactive effects on seed performance. *Plos one.* 2023; 18(1): 1-14.
18. Wattanakulpakin P, Photchanachai S, Miyagawa S and Ratanakhanokchai K. Loss of maize seed vigour as affected by biochemical changes during hydropriming. *Crop Sci.* 2012; 52(6): 2783-2793.