

## Effective Management of Pearl Millet Blast (*Pyricularia grisea*) Through Combined Chemical and Biological Tactics

### ABSTRACT

Pearl millet is a staple cereal grown in India. It encounters number of diseases which attack the crop during its growth, cause low yield and economic loss to the peasant and finally to the nation as a whole. The blast also referred as leaf spot caused by *Pyricularia grisea* has emerged as a serious disease affecting both forage and grain production in pearl millet. In view of this a field experiment was conducted over three consecutive *kharif* seasons (2021, 2022 and 2023) at the Pearl Millet Research Station, JAU, Jamnagar, to assess to evaluate the efficacy of different fungicide and bio agents in for reducing the pearl millet blast disease as well as identify the most effective fungicide/bio agents and application rates for minimizing blast intensity. On the basis of field and based on the pooled data,

Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 and 35 DAS found minimum blast disease intensity (22.96%), Highest grain yield (2472 kg ha<sup>-1</sup>) and fodder yield (46.80 q ha<sup>-1</sup>) over treatment and sprays of *Pseudomonas fluorescens*, 10 g L<sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS (26.75%), found statistically at par, with blast intensity (26.75%), grain yield (2377 kg ha<sup>-1</sup>) and fodder yield 44.28 q ha<sup>-1</sup>).

**Keywords:** Pearl millet Blast, *Kharif*, Tebuconazole + Trifloxystrobin, *Pseudomonas fluorescens*, disease intensity, Yield

### Introduction

During 2023-24, pearl millet area in India was 7.36 million ha with an average production of 10.67 million tons and 1449 kg ha<sup>-1</sup> productivity (DA&FW, 2024). The major pearl millet growing states are Rajasthan, Maharashtra, Uttar Pradesh, Gujarat and Haryana contributing to 90% of total production in the country. Rajasthan contributes nearly 45% followed by Uttar Pradesh (19%), Haryana (9%), Gujarat (9%), Maharashtra (6%) and Tamil Nadu (2%). Most of pearl millet in India is grown in rainy (*kharif*) season (June/July-September/October). Pearl millet is also cultivated during summer season (February-May) in parts of Gujarat, Rajasthan and Uttar Pradesh; and during the post-rainy (*rabi*) season (November-February) at a small scale in Maharashtra and Gujarat. In Gujarat it is grown in 26 out of 33 districts covering an area of 2.03 lakh ha in *kharif* with an average production 3.04 lakh tonnes and average yield 1787 kg ha<sup>-1</sup> (DA&FW, 2024). In 2023, Hon'ble prime minister of India rebranded millets as "Shree Anna" for their climate resilience and nutritional superiority and declared ICAR-IIMR, Hyderabad as "Global Centre of Excellence for Millets". In order to mainstream and exploit nutritionally superiority of millets and promote their cultivation, Govt. of India declared Year 2018 as the "Year of Millets" and after declaration of FAO Committee on Agriculture (COAG) forum in 2021, Year 2023 was celebrated as "International Year of Millets" (Anon., 2024). Among the diseases of pearl millet, blast caused by *Pyricularia grisea* (Cooke) Sacc. [Teleomorph: *Magnaporthe grisea*

(Herbert) Barr], a disease of minor importance in past years, has gained status of major constraint to pearl millet production in India (Lukose et al., 2007). The blast disease caused by *Magnaporthe grisea* has emerged as one of the major production constraints during the last decade in pearl millet cultivation, causing severe loss in grain yield and quality (Singh et al. 2021). Bajra blast also referred as leaf spot caused by *Pyricularia grisea* (Cooke) Sacc. [Teleomorph: *Magnaporthe grisea* (Herbert) Barr.] has emerged as a serious disease affecting both forage and grain production in pearl millet (Kaurav et al., 2018), resulting economic loss. Recently intensity of blast increased at alarming rate in commercial hybrids cultivation (Thakur et al., 2009) In view of these, chemical control is taken to manage this disease. *Magnaporthe grisea* is externally seed borne and also survives as chlamydoconidia or as free saprophytic mycelium in the soil/leaf debris which serves as a source of primary inoculum (Singh and Pavgi, 1977).

## MATERIALS AND METHODS

Three-year field experiments were conducted during *kharif* 2021, *kharif* 2022 and *kharif* 2023 at Pearl Millet Research Station, JAU, Jamnagar to find out the bio efficacy of different fungicide and bio agents against the minimized blast disease intensity at natural condition.

Experiment conducted with randomized block design (RBD), each having four replications. The plot size was 4.2 m × 2.4 m and distance between row to row and plant to plant was 60 cm and 10 cm, respectively. Four row were maintained in each treatment (plot) during all experimental season. Total six chemicals and bio agents (Table 1) including control was used as treatment for management of pearl millet blast disease intensity.

Seed treatments were given initially at the time of sowing. Foliar application of different fungicides and bio agents was carried out management of pearl millet blast. The first spray was given at 20 DAS and second spray given at 35 DAS.

For observation, ten plants were selected randomly and labeled from each plot for scoring the disease intensity. These labeled plants were observed for disease intensity from upper, middle and lower leaves using disease rating scale of 0-9. Observations on disease intensity was recorded at 30, 45 and 60 DAS.

**Table 1: Treatments details:**

Tr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	a. i g ha <sup>-1</sup>	Quantity of formulation kg or l ha <sup>-1</sup>
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> kg seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	-	0.015 kg
		1 × 10 <sup>8</sup> cfu g <sup>-1</sup>	100 g	-	5.0 kg
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> kg seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	-	0.015 kg
		1 × 10 <sup>8</sup> cfu g <sup>-1</sup>	100 g	-	5.0 kg
3.	Sprays of <i>P.s fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	1 × 10 <sup>8</sup> cfu g <sup>-1</sup>	100 g	-	5.0 kg
		0.04	5.33 g	200	0.267 kg
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	200	0.267 kg
		1 × 10 <sup>8</sup>	100 g	-	5.0 kg

		cfu g <sup>-1</sup>			
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	200	0.267 kg
6.	Control	-	-	-	-

Per cent disease intensity (PDI) will be calculated by using the following formula (Wheeler, 1969).

$$\text{Disease intensity (\%)} = \frac{\text{Sum of total rating}}{\text{Total number of leaves observed}} \times \frac{\text{Maximum disease rating}}{\text{Maximum disease rating}} \times 100$$

### Blast disease rating scale (0-9)

Scale	Description	Scale	Description
0	: No lesions	5	: Typical blast lesions infecting 2-10% of the leaf area
1	: Small brown specks of pinhead size without sporulating center	6	: Blast lesions infecting 11-25% leaf area
2	: Small roundish to slightly elongated, necrotic grey spots, about 1-2 mm in diameter with a distinct brown margin, lesions are mostly found on the lower leaves	7	: Blast lesions infecting 26-50% leaf area
3	: Lesion type is the same as in scale 2, but significant number lesions are on the upper leaves	8	: Blast lesions infecting 51-75% leaf area
4	: Typical sporulating blast lesions, 3 mm or longer, infecting less than 2% of the leaf area	9	: More than 75% leaf area affected

### Observations recorded:

1. Seedling germination (%)
2. Per cent blast disease intensity at 30, 45 and 60 DAS.
3. Grain yield kg ha<sup>-1</sup> and fodder yield q ha<sup>-1</sup>.

Grain and fodder yield was recorded from net plot area at harvest and data obtained was analyzed statistically.

### RESULTS AND INTERPRETATION

A field experiments was conducted with different six treatments including bio agents and chemical during *kharif* 2021, 2022 and 2023. The three year pooled result of all parameters presented in Table 2 to 7. All the treatment found effective to suppress blast disease intensity significantly.

The three-year pooled data analysis (Table 2) shows that none of the treatments had a statistically significant impact on seed germination (Fig. 1). This result suggests that the treatments applied did not effectively enhance germination rates. Consequently, no treatment was superior, highlighting the need for further investigation to identify potential improvements or alternative approaches to influence seed germination effectively.

Looking to results of blast disease, three year pooled observation on 30 DAS (Table 3, Fig. 2) stated that the treatment spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 and 35 DAS (9.17%) found significantly superior over treatment and which was at par with remaining all treatment without control, viz., sprays of *P. fluorescens*, 10 g L<sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS (10.98%), spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and *B. subtilis*, 10 g L<sup>-1</sup> at 35 DAS (11.63%), seed treatment with chitosan, 3.75 g kg<sup>-1</sup> seed + two sprays of *B. subtilis*, 10 g

L<sup>-1</sup> at 20 & 35 DAS (12.18%) and seed treatment with chitosan, 3.75 g kg<sup>-1</sup> seed + two sprays of *P. fluorescens*, 10 g L<sup>-1</sup> at 20 & 35 DAS (12.42%).

The pooled data (Table 4 and Fig. 3) at 45 DAS shows that the treatment involving two sprays of Tebuconazole 50 + Trifloxystrobin 25 WG at 0.04% concentration, applied at 20 and 35 DAS, resulted in a blast intensity of 12.85%, which was significantly superior to other treatments. Among the remaining treatments, only the spray of *P. fluorescens* at 10 g L<sup>-1</sup> at 20 DAS followed by Tebuconazole 50 + Trifloxystrobin 25 WG at 0.04% at 35 DAS (15.48%) was statistically at par with the superior treatment. The control registered highest blast intensity at 25.48%,

The treatment Tebuconazole 50 + Trifloxystrobin 25 WG treatment, particularly when applied at both 20 and 35 DAS, consistently resulted in significantly lower blast intensity compared to other approaches. This suggests it as a highly effective choice for disease control, outperforming other treatments and the untreated control.

More or less similar trend was observed in results of blast intensity on 60 DAS (Table 5 and Fig. 4) pooled results same as 45 DAS data, treatment spray Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 and 35 DAS (22.96%) found superior over treatment and sprays of *P. fluorescens*, 10 g L<sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS (26.75%), found statistically at par. Maximum blast intensity (47.72%) recorded in control.

Above result supported the previous worked done by Chaudhari *et al.* (2024) showed that Azoxystrobin 11% + Tebuconazole 18.30% SC, 0.05% recorded the lowest blast intensity at 30.20%, which was statistically at par with Tebuconazole 50% + Trifloxystrobin 25% WG, 0.05% (31.65%). Sharma *et al.* (2018) reported that, the disease can be effectively managed in pearl millet with two to three sprays of propiconazole or tebuconazole + trifloxystrobin at 15 day intervals with the first spray at 20–25 days after sowing. Patro *et al.* (2020) mentioned that initial spray of *P. fluorescens* and Trifloxystrobin + Tebuconazole as second spray was found superior in managing the blast disease.

#### **Grain and fodder yield:**

The three-year pooled results (Table 6 and Fig. 5) for grain yield indicate that the highest yield was achieved with the treatment involving two sprays of Tebuconazole 50 + Trifloxystrobin 25 WG at 0.04%, applied at 20 and 35 DAS, resulting in 2472 kg ha<sup>-1</sup>. This yield was statistically on par with the treatment of *P. fluorescens* spray at 10 g L<sup>-1</sup> at 20 DAS followed by Tebuconazole 50 + Trifloxystrobin 25 WG at 0.04% at 35 DAS, which produced a yield of 2377 kg ha<sup>-1</sup>. These findings suggest that both treatment regimens effectively enhance grain yield, with Tebuconazole 50 + Trifloxystrobin 25 WG showing a slightly higher yield. Field experiment results of Sharma *et al.*, (2018) revealed that three sprays of Tebuconazole + Trifloxystrobin or propiconazole was superior in reducing blast incidence with higher yields in pearl millet. Pramesh *et al.*, (2016) reported that rice blast was effectively controlled with Tebuconazole + Trifloxystrobin and resulted in higher yield. Chaudhari *et al.* (2024) recorded that highest grain yield (2135 kg ha<sup>-1</sup>) and fodder yield (44.38 q ha<sup>-1</sup>) recorded in treatment Azoxystrobin 11% + Tebuconazole 18.30% SC, 0.05% which was at par with Tebuconazole 50% + Trifloxystrobin 25% WG, 0.05% (2054 kg ha<sup>-1</sup>).

Three year pooled result (Table 7 and Fig. 6) for fodder yield indicated that the highest fodder yield also found in treatment spray Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 and 35 DAS (46.80 q ha<sup>-1</sup>) and which was at par with treatment sprays of *P. fluorescens*, 10 g L<sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS (44.28 q ha<sup>-1</sup>), spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and *B. subtilis*, 10 g L<sup>-1</sup> at 35 DAS (43.26 q ha<sup>-1</sup>) and seed treatment with chitosan, 3.75 g kg<sup>-1</sup> seed + two sprays of *B. subtilis*, 10 g L<sup>-1</sup> at 20 & 35 DAS (42.84 q ha<sup>-1</sup>). Minimum grain yield (1755 kg ha<sup>-1</sup>) and fodder yield (36.41 q ha<sup>-1</sup>) recorded in control. The result supported by Patro (2020) on foliar application of *P. fluorescens* at 20 DAS and Trifloxystrob in +

Tebuconazole at 35 DAS was found effective with least disease intensity of blast (14.1%), highest grain (26.0 q ha<sup>-1</sup>) and fodder yield (60.9 q ha<sup>-1</sup>).

#### **Economics:**

Looking to the economics of different bio agents and fungicidal treatments (Table 8), the highest additional income ₹17129 ha<sup>-1</sup>, highest net realization of ₹13393 ha<sup>-1</sup> and maximum ICBR 1: 4.58 was obtained in the treatment, spray of *P. fluorescens* (100 g per 10 litre of water) at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% (5.33 00 g per 10 litre of water) at 35 DAS.

#### **Conclusion:**

It can be concluded from the above results that the spraying of *P. fluorescens* (100 00 g per 10 litre of water) at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% (5.33 00 g per 10 litre of water) at 35 DAS or spray Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 and 35 DAS (5.33 00 g per 10 litre of water) in pearl millet against blast disease were found effective to minimize blast intensity, higher grain and fodder yield and additional income also.

#### **Disclaimer (Artificial intelligence):**

There has been NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### **competing interests:**

Authors have declared that no competing interests exist.


#### List of abbreviations

@	:	At a rate of
a.i.	:	Active Ingredient
DAS	:	Days After Sowing
DMRT	:	Duncan's Multiple Range Test
lb	:	Pound
ICBR	:	Incremental Cost Venefit Ratio
JAU	:	Junagadh Agricultural University
kg ha <sup>-1</sup>	:	Kilogram Per Hectare
PDI	:	Per cent Disease Intensity
q ha <sup>-1</sup>	:	Quintal Per Hectare
WG	:	Water dispersible Granules

**Table 2: Effect of chemical and bio-agents treatments on seedling emergence (%)**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Seedling emergence (%)			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	44.81 <sup>ab</sup>	57.93 <sup>ab</sup>	82.42 <sup>a</sup>	61.72 <sup>a</sup>
		-	100 g				
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	44.81 <sup>ab</sup>	56.93 <sup>ab</sup>	77.25 <sup>a</sup>	59.66 <sup>a</sup>
		-	100 g				
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	43.68 <sup>b</sup>	57.05 <sup>ab</sup>	79.41 <sup>a</sup>	60.04 <sup>a</sup>
		0.04	5.33 g				
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	46.13 <sup>a</sup>	60.48 <sup>a</sup>	80.24 <sup>a</sup>	62.28 <sup>a</sup>
		-	100 g				
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	44.43 <sup>ab</sup>	56.13 <sup>b</sup>	83.58 <sup>a</sup>	61.38 <sup>a</sup>
6.	Control	-	-	43.02 <sup>b</sup>	57.74 <sup>ab</sup>	80.59 <sup>a</sup>	60.45 <sup>a</sup>
	<b>S. Em. ±</b>			<b>0.66</b>	<b>1.09</b>	<b>1.98</b>	<b>0.78</b>
	<b>C. D. at 5%</b>			<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>
	<b>C. V. %</b>			<b>2.96</b>	<b>3.79</b>	<b>4.91</b>	<b>4.46</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>0.55</b>
	<b>C. D. at 5%</b>						<b>1.58</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>1.36</b>
	<b>C. D. at 5%</b>						<b>NS</b>

Data were transformed (angular transformed) before analysis. Treatment means with letters(s) in common are at par as per DNMR at 5% level of significance.




**Table 3: Effect of chemical and bio-agents treatments on per cent disease intensity at 30 DAS**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Per cent disease intensity (30 DAS)			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	22.02 <sup>b</sup> (14.06)	23.42 <sup>b</sup> (15.80)	16.46 <sup>a</sup> (8.06)	20.64 <sup>b</sup> (12.42)
		-	100 g				
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	21.15 <sup>bc</sup> (13.02)	23.66 <sup>b</sup> (16.10)	16.46 <sup>a</sup> (8.06)	20.42 <sup>b</sup> (12.18)
		-	100 g				
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	21.38 <sup>b</sup> (13.33)	21.15 <sup>c</sup> (13.01)	15.53 <sup>a</sup> (7.22)	19.35 <sup>b</sup> (10.98)
		0.04	5.33 g				
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	22.66 <sup>b</sup> (15.00)	20.68 <sup>cd</sup> (12.48)	16.48 <sup>a</sup> (8.06)	19.94 <sup>b</sup> (11.63)
		-	100 g				
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	18.68 <sup>c</sup> (10.28)	18.67 <sup>d</sup> (10.25)	15.53 <sup>a</sup> (7.22)	17.63 <sup>b</sup> (9.17)
6.	Control	-	-	27.91 <sup>a</sup> (21.94)	27.53 <sup>a</sup> (21.37)	16.46 <sup>a</sup> (8.06)	23.97 <sup>a</sup> (16.50)
	<b>S. Em. ±</b>			<b>0.75</b>	<b>0.61</b>	<b>0.67</b>	<b>1.00</b>
	<b>C. D. at 5%</b>			<b>2.2506</b>	<b>1.829</b>	<b>NS</b>	<b>3.16</b>
	<b>C. V. %</b>			<b>6.70</b>	<b>5.39</b>	<b>8.25</b>	<b>6.65</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>0.28</b>
	<b>C. D. at 5%</b>						<b>0.79</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>0.68</b>
	<b>C. D. at 5%</b>						<b>1.93</b>

Figures in parenthesis are retransformed arc sine values.

Data were transformed (angular transformed) before analysis. Treatment means with letters(s) in common are at par as per DNMRT at 5% level of significance.




**Fig. 2.** Effect of chemical and bio-agents treatments on per cent disease intensity at 30 DAS.

**Table 4: Effect of chemical and bio-agents treatments on per cent disease intensity at 45 DAS**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Per cent disease intensity (45 DAS)			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	27.66 <sup>bc</sup>	26.14 <sup>bc</sup>	21.65 <sup>b</sup>	25.15 <sup>bc</sup>
		-	100 g	(21.55)	(19.41)	(13.61)	(18.06)
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	28.68 <sup>b</sup>	27.91 <sup>b</sup>	21.86 <sup>b</sup>	26.15 <sup>b</sup>
		-	100 g	(23.03)	(21.91)	(13.87)	(19.42)
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	25.73 <sup>cd</sup>	24.08 <sup>cd</sup>	19.70 <sup>c</sup>	23.17 <sup>cd</sup>
		0.04	5.33 g	(18.84)	(16.64)	(11.36)	(15.48)
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	30.15 <sup>b</sup>	27.89 <sup>b</sup>	20.92 <sup>bc</sup>	26.32 <sup>b</sup>
		-	100 g	(25.23)	(21.89)	(12.75)	(19.66)
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	23.00 <sup>d</sup>	21.86 <sup>d</sup>	18.15 <sup>d</sup>	21.00 <sup>d</sup>
				(15.27)	(13.87)	(9.70)	(12.85)
6.	Control	-	-	35.07 <sup>a</sup>	31.79 <sup>a</sup>	24.08 <sup>a</sup>	30.31 <sup>a</sup>
				(33.02)	(27.75)	(16.64)	(25.48)
	<b>S. Em. ±</b>			<b>0.86</b>	<b>0.84</b>	<b>0.44</b>	<b>0.70</b>
	<b>C. D. at 5%</b>			<b>2.58</b>	<b>2.52</b>	<b>1.34</b>	<b>2.20</b>
	<b>C. V. %</b>			<b>6.04</b>	<b>6.28</b>	<b>4.22</b>	<b>5.81</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>0.30</b>
	<b>C. D. at 5%</b>						<b>0.56</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>0.74</b>
	<b>C. D. at 5%</b>						<b>NS</b>

Figures in parenthesis are retransformed arc sine values. Data were transformed (angular transformed) before analysis. Treatment means with letters(s) in common are at par as per DNMRT at 5% level of significance.




**Fig. 3.** Effect of chemical and bio-agents treatments on per cent disease intensity at 45 DAS.

**Table 5: Effect of chemical and bio-agents treatments on per cent disease intensity at 60 DAS**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Per cent disease intensity (60 DAS)			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	39.23 <sup>c</sup>	41.48 <sup>b</sup>	25.92 <sup>bc</sup>	35.54 <sup>b</sup>
		-	100 g	(40.00)	(43.88)	(19.11)	(33.80)
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	40.52 <sup>c</sup>	41.15 <sup>b</sup>	26.16 <sup>bc</sup>	35.94 <sup>b</sup>
		-	100 g	(42.22)	(43.30)	(19.43)	(34.46)
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	36.07 <sup>d</sup>	33.71 <sup>c</sup>	23.65 <sup>cd</sup>	31.14 <sup>c</sup>
		0.04	5.33 g	(34.67)	(30.80)	(16.09)	(26.75)
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	43.57 <sup>b</sup>	40.30 <sup>b</sup>	27.14 <sup>b</sup>	37.00 <sup>b</sup>
		-	100 g	(47.50)	(41.83)	(20.81)	(36.22)
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	33.03 <sup>c</sup>	31.06 <sup>c</sup>	21.81 <sup>d</sup>	28.63 <sup>c</sup>
				(29.71)	(26.62)	(13.81)	(22.96)
6.	Control	-	-	49.48 <sup>a</sup>	49.64 <sup>a</sup>	31.95 <sup>a</sup>	43.69 <sup>a</sup>
				(57.79)	(58.07)	(28.01)	(47.72)
	<b>S. Em. ±</b>			<b>0.71</b>	<b>1.36</b>	<b>0.90</b>	<b>1.08</b>
	<b>C. D. at 5%</b>			<b>2.14</b>	<b>4.10</b>	<b>2.70</b>	<b>3.40</b>
	<b>C. V. %</b>			<b>3.53</b>	<b>6.88</b>	<b>6.86</b>	<b>5.81</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>0.42</b>
	<b>C. D. at 5%</b>						<b>1.19</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>0.02</b>
	<b>C. D. at 5%</b>						<b>NS</b>

Figures in parenthesis are retransformed arc sine values. Data were transformed (angular transformed) before analysis. Treatment means with letters(s) in common are at par as per DNMRT at 5% level of significance




**Fig. 4.** Effect of chemical and bio-agents treatments on per cent disease intensity at 60 DAS.

**Table 6 : Effect of chemical and bio-agents treatments on grain yield (kg ha<sup>-1</sup>)**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Grain yield (kg ha <sup>-1</sup> )			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	1750 <sup>b</sup>	2035 <sup>ab</sup>	2375 <sup>bc</sup>	2053 <sup>bc</sup>
		-	100 g				
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	1595 <sup>b</sup>	1912 <sup>bc</sup>	2348 <sup>bc</sup>	1952 <sup>cd</sup>
		-	100 g				
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	2193 <sup>a</sup>	2206 <sup>ab</sup>	2732 <sup>a</sup>	2377 <sup>a</sup>
		0.04	5.33 g				
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	1866 <sup>b</sup>	2111 <sup>ab</sup>	2511 <sup>ab</sup>	2163 <sup>b</sup>
		-	100 g				
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	2361 <sup>a</sup>	2309 <sup>a</sup>	2746 <sup>a</sup>	2472 <sup>a</sup>
6.	Control	-	-	1547 <sup>b</sup>	1627 <sup>c</sup>	2092 <sup>c</sup>	1755 <sup>d</sup>
	<b>S. Em. ±</b>			<b>98.46</b>	<b>103.91</b>	<b>112.38</b>	<b>60.66</b>
	<b>C. D. at 5%</b>			<b>296.72</b>	<b>313.16</b>	<b>338.67</b>	<b>172.93</b>
	<b>C. V. %</b>			<b>10.44</b>	<b>10.22</b>	<b>9.11</b>	<b>9.87</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>43.90</b>
	<b>C. D. at 5%</b>						<b>122.28</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>105.07</b>
	<b>C. D. at 5%</b>						<b>NS</b>

Treatment means with letters(s) in common are at par as per DNMRT at 5% level of significance.




**Fig. 5, Effect of chemical and bio-agents treatments on grain yield (kg ha<sup>-1</sup>),**

**Table 7: Effect of chemical and bio-agents treatments on fodder yield (q ha<sup>-1</sup>)**

Sr. No.	Treatment	Con. (a. i.)	Quantity in g or ml in 10 liter of water or 1 kg seed	Fodder yield (q ha <sup>-1</sup> )			
				2021	2022	2023	Pooled
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	40.61 <sup>a</sup>	50.34 <sup>abc</sup>	37.58 <sup>a</sup>	42.84 <sup>ab</sup>
		-	100 g				
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	-	3.75 g kg <sup>-1</sup>	37.08 <sup>a</sup>	47.47 <sup>bc</sup>	36.22 <sup>ab</sup>	40.26 <sup>bc</sup>
		-	100 g				
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	-	100 g	40.11 <sup>a</sup>	54.9 <sup>ab</sup>	37.84 <sup>a</sup>	44.28 <sup>ab</sup>
		0.04	5.33 g				
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	0.04	5.33 g	42.71 <sup>a</sup>	48.62 <sup>abc</sup>	38.44 <sup>a</sup>	43.26 <sup>ab</sup>
		-	100 g				
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	0.04	5.33 g	42.62 <sup>a</sup>	57.5 <sup>a</sup>	40.29 <sup>a</sup>	46.80 <sup>a</sup>
6.	Control	-	-	35.84 <sup>a</sup>	42.37 <sup>c</sup>	31.02 <sup>b</sup>	36.41 <sup>c</sup>
	<b>S. Em. ±</b>			<b>2.52</b>	<b>2.88</b>	<b>1.71</b>	<b>1.40</b>
	<b>C. D. at 5%</b>			<b>NS</b>	<b>8.69</b>	<b>5.15</b>	<b>3.98</b>
	<b>C. V. %</b>			<b>12.65</b>	<b>11.49</b>	<b>9.26</b>	<b>11.44</b>
	<b>Y</b>						
	<b>S. Em. ±</b>						<b>0.99</b>
	<b>C. D. at 5%</b>						<b>2.82</b>
	<b>Y×T</b>						
	<b>S. Em. ±</b>						<b>2.42</b>
	<b>C. D. at 5%</b>						<b>NS</b>

Treatment means with letters(s) in common are at par as per DNMRT at 5% level of significance.



**Fig. 6.** Effect of chemical and bio-agents treatments on fodder yield (q ha<sup>-1</sup>).

**Table 8: Economics of various treatments for the management pearl millet blast**

Tr. No.	Treatment	Yield (kg ha <sup>-1</sup> ) Pooled		Yield increase over control (kg ha <sup>-1</sup> )		Income (₹)		Additional income (₹)	Cost of treatment (fungicides/bio agents, labour charge, etc.) (₹ ha <sup>-1</sup> )	Net realization (₹)	ICBR
		Grain	Fodder	Grain	Fodder	Grain*	Fodder**				
1	2	3	4	5	6	7	8	9	10	11 (9-10)	12 (9/10)
1.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	2053	4284	298	643	7455	1286	8741	2117	6625	1:4.13
2.	Seed treatment with chitosan, 3.75 g kg <sup>-1</sup> seed + two sprays of <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 20 & 35 DAS	1952	4026	197	385	4913	770	5683	2117	3566	1:2.68
3.	Sprays of <i>P. fluorescens</i> , 10 g L <sup>-1</sup> at 20 DAS and Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 35 DAS	2377	4428	622	787	15555	1574	17129	3736	13393	1:4.58
4.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 DAS and <i>B. subtilis</i> , 10 g L <sup>-1</sup> at 35 DAS	2163	4326	408	685	10192	1370	11562	3736	7826	1:3.09
5.	Spray of Tebuconazole 50 + Trifloxystrobin 25 WG, 0.04% at 20 & 35 DAS	2472	4680	717	1039	17924	2078	20002	3136	16866	1:6.38
6.	Control	1755	3641	0	0	0	0	0	0	0	0

\* Price of bajra grain: ₹25 kg<sup>-1</sup>, \*\* Price of bajra fodder: ₹2 kg<sup>-1</sup>

## REFERENCES

1. Ajay K, Pandya RK, Bhagyashree S. Performance of botanicals and fungicides against blast of pearl millet (*Pennisetum glaucum*). *ANN. Plant Soil Res.* 2018; 20(3):258–262.
2. Anonymous. AICRP on Pearl Millet 2023-24. 2024. pp. 1-2.
3. Chaudhari, R. J., K. D. Mungra, G. M. Parmar, S. K. Parmar, and J. S. Sorathiya. 2024. “Mitigating Pearl Millet Blast (*Pyricularia grisea*): Effective Fungicidal Treatments”. *Advances in Research*, 25(5):223-35. <https://doi.org/10.9734/air/2024/v25i51156>.
4. DA&FW. Area, production and yield of bajra in Gujarat 2023-24. 2024. (<https://dag.gujarat.gov.in/estimate.html>)
5. Kaurav Ajay, Pandya R. K. And Bhagyashree Singh (2018). Performance of botanicals and fungicides against blast of pearl millet (*Pennisetum glaucum*). *Annals of Plant and Soil Research*. 2018;20(3): 258–262.
6. Lukose, C.M., Kadvani, D.L., Dangaria, C.J. Efficacy of fungicides in controlling blast disease of pearl millet. *Indian Phytopathol.* 2007;60: 68–71.
7. Patro, T. S. S. K., Georgia, K.E., Raj Kumar, S., Anuradha, N., Sandhya Rani, Y. and Triveni, U. Management of pearl millet blast through fungicides and biocontrol agents. *International Journal of Chemical Studies* 2020; 8(4): 1357-1359.
8. Pramesh D, Maruti, Muniraju K M, Mallikarjun K, Guruprasad G S, Mahantashivayogayya K, Reddy B G M, Gowdar S B and Chethana B S. Bio-efficacy of a Combination Fungicide against Blast and Sheath Blight Diseases of Paddy. *J. Exp. Agric. Int.* 2016;14(4): 1-8.
9. Sharma R., Gate, V.L. and Madhavan, S. Evaluation of fungicides for the management of pearl millet [*Pennisetum glaucum* (L.)] blast caused by *Magnaporthe grisea*. *Crop Protection*. 2018; 112:209-213.
10. Singh S, Sharma R, Chandra Nayaka S, Tara Satyavathi C, Raj C. Understanding pearl millet blast caused by *Magnaporthe grisea* and strategies for its management. *Blast disease of cereal crops: evolution and adaptation in context of climate change*. 2021:151-72.
11. Singh, S.D. and Pavgi, M.S. Perpetuation of *Pyricularia penniseti* causing brown leaf spot of bajara. *Indian Phytopathol.* 1977;30: 242–244.
12. Thakur, R.P., Sharma, R., Rai, K.N., Gupta, S.K., & Rao, V.P. Screening techniques and resistance sources for foliar blast in pearl millet. *Journal of SAT Agril. Res.*, 2009;7:1-15.
13. Wheeler, B. E. J. An Introduction of Plant Disease, John Wiley and Sons Ltd., London. 1969; pp. 301.

UNDER PEER REVIEW