

UNDERSTANDING THE ROLE OF MATERNAL AND LIFESTYLE FACTORS IN MISCARRIAGE RISK: AN SEM ANALYSIS

Abstract

Miscarriage like other pregnancy complications is multifactorial meaning that it is caused by even a combination of maternal and lifestyle factors. This paper aims to examine antecedents such as maternal age, education, risk factors (smoking, alcohol, physical activity during pregnancy), and visits for antenatal care in relation to miscarriage. In this study, through the employment of Structural Equation Modeling (SEM), we estimate both direct and indirect effects of lifestyle and maternal characteristics on miscarriages. An index variable for lifestyle risks is established to measure smoking, alcohol intake and exercise influence simultaneously while direct effect of maternal age, education standard and number of antenatal care visits are also estimated towards the miscarriage risk. The analysis further explores how antenatal care visits also moderate the risks stated for lifestyle risks and miscarriage. The findings indicate that there is a direct effect of lifestyle on miscarriage and this is by smoking and alcohol consumption controlling for the number of antenatal care visits. Birth mother age and education also have average direct influence regarding miscarriage possibility. The present paper examines the multifaceted relationships between lifestyle, maternal health, and miscarriage, and presents a number of suggestions for intervention targets as well as preventive measures in the area of maternal health care.

Introduction

Abortion is defined as the spontaneous expulsion of a fetus prior to the tenth week of pregnancy, commonly referred to as miscarriage, and affects approximately 10–20% of all pregnancies that are acknowledged clinically. However, the factors that lead to miscarriage are not so well known and much of risk factors are still unknown. In most cases miscarriage is genetic and multifactorial which means it is influenced by genes, environment, physiology, and life style. Certainly, some risks are unpreventable, for example, chromosomal disorders or high age of the expectant mother, but many of the factors known to pose risks for miscarriage are within a woman's control, especially those that are connected with her health condition and lifestyle (Chibayi et al., 2024; Kumar, 2011; Wang et al., 2014). They comprise maternal age at delivery, years of education, tobacco smoking, alcohol drinking, physical activity during pregnancy, and antenatal care utilization (Amponsah *et al.*, 2022; Gascoigne et al., 2023). Even though these risk factors are acknowledged to play significant roles, the studies have been conducted separately without consideration of their interaction. Knowledge of the ways that lifestyle factors and maternal factors affect miscarriage is important because prevention strategies that may prevent miscarriage might be found (Gascoigne et al., 2023). It is the intention of this study to study the probability of miscarriage relating to these maternal and lifestyle factors using Structural Equation Modeling (SEM), a statistical approach that captures causality of paths between variables.

Overview of Multifactorial Pathways of Miscarriage

Spontaneous pregnancy loss can occur at any time during gestation but is most frequent during the initial weeks of pregnancy. Average women will have one or more miscarriages in their life span but women experiencing two or more miscarriages are 1-2%. Number of conditions that may lead to miscarriage and they may include genetic issues, structural abnormalities and hormonal disturbances (Celep *et al.*, 2005; Hyde and Schust, 2015). For instance, chromosomal anomalies are the main cause of pregnancy loss contributing to up to 60% of all instances. However, there are many other factors to meet for a miscarriage and genetic factors are only part of the miscarriage process (Pal *et al.*, 2018). Some researchers believe that about 50% of threatened miscarriages are likely caused by physical and chemical factors in the uterine environment and the mother herself that can hamper fetal development (Kumar, 2011; Krieg *et al.*, 2016). More research has been conducted in the last decades on the influence of maternal replication behaviors like smoking, alcohol consumption and lack of exercise on miscarriage prevalence (Bailey& Sokol, 2011; Kumar, 2011). Lifestyle covariates can have a direct impact on pregnancy outcomes by biological means and an indirect impact in modifying other components of maternal risk profiles including antenatal clinic attendance (Oei, 2020). For instance, smoking during pregnancy has been identified to increase the risk of miscarriage, may be through disruption of placental function, aggravated by increased oxidative stress and fetal developmental changes. In the same way, alcohol has been shown to cause pregnancy loss, and current research about alcohol use during pregnancy points out that moderate drinking can affect fetal development (Chung *et al.*, 2023; McCarthy *et al.*, 2013). Exercise also has a twofold effect on pregnancy outcome (Gascoigne *et al.*, 2023). On one hand, physical activity that can be moderate and repeated for at least 30 minutes on a daily basis is considered helpful for a mother and decreases the risks of gestational diabetes or preeclampsia. As for the external imposed factors, physical activity in its different forms proved to be beneficial nonetheless, however sharp increases in the physical intensity of the activity during pregnancy, particularly vigorous or high impact exercise may lead to miscarriage. It then becomes crucial to understand the key to weigh the advantages and disadvantages of physical activity during pregnancy (ACOG, 2013; Gascoigne *et al.*, 2023).

Maternal Health: A Key Determinant in Miscarriage

Age, education level, and types of antenatal care (ANC) received by the mother are some of the most important predictors of miscarriage (Amponsah *et al.*, 2022). For miscarriage, one the most definite risk factors is an increment in the maternal age which is non modifiable. Women above 35 years are more likely to miscarry than younger women, because of decrease in oocyte quality and higher possibility of chromosomally abnormal zygotes (Flynn *et al.*, 2014) Some socio-cultural factors, lending more credence to the fact that most women today prefer to delay childbearing in order to further their careers as well as attain higher education, means that miscarriage risk is even more dangerous as age progresses. The other contributory factor which is however less often mentioned is education level. Literate female is not more likely to smoke or drink which are some of the risk factors to miscarriage than those women with lower levels of education (Sampasa *et al.*, 2020). Additionally, Lack of education leads to poor health literacy, thus poor health circulation, this leads to poor access to health facilities, especially antenatal care thus increases on miscarriages. It should also be noted that better-educated women are, as a rule, more knowledgeable about the potential consequences of erroneous actions during pregnancy and more often turn to healthcare providers (ACOG, 2013). Another important factor that contributes to miscarriage occurrence is antenatal care because close

monitoring of any complications may be enhanced meant for the mother and fetus (Amponsah *et al.*, 2022). If women attend properly planned antenatal visits they should be advised about smoking cessation, alcohol and safe exercise (Hosker *et al.*, 2019; Gascoigne *et al.*, 2023). The number one risk factor to miscarriage and other complications associated with pregnancy is lack of adequate ANC especially during the initial period of pregnancy.

Structural Equation Modeling (SEM): A Comprehensive Approach

Though the risk factors of miscarriage at individual level have been investigated numerous times still the interactions or associations between these factors are not well enough investigated. There are also various other aspects of maternal and lifestyle factors in which results depict that most factors do not perform independently but combine in determining the miscarriage occurrence (Gascoigne *et al.*, 2023). It is with these views that Structural Equation Modeling (SEM) forms the best method to analyse these relations. SEM is a statistical technique which enables the researcher to examine both direct and indirect effects of the variables hence suitable in analysing the chain through which various factors lead to miscarriage (Mueller, R. O., & Hancock, G. R., 2018; Tarka, P., 2018).

In SEM, measured variables or manifest variables (for example smoking or alcohol consumption) can be used as representation of underlying construct of higher order such as lifestyle risks. The other constructs may then be cross tabulated to determine the direct and indirect effects of both the lifestyle and maternal factors on miscarriage (Savalei, V., & Rhemtulla, M., 2013). Collier, J. (2020) also notes that SEM can also consider measurement error to draw better estimates of the effect of different variables. It allows the researcher to screen fit of a theoretical model and the magnitude of the relationships between predictors and outcomes.

Lifestyle risks and maternal health factors are examined in this study utilizing SEM with smoking, alcohol consumption, physical activity, age, education level, number of antenatal visits being on the miscarriage risk direct and mediating roles. It was therefore deemed appropriate to create other latent variables such as lifestyle risks and antenatal care in this model to capture the miscellaneous effects these factors have on miscarriage results. Second, and perhaps more importantly, SEM according to Savalei, V., & Rhemtulla, M. (2013) allows for the examination of the moderated mediation model in which it can be tested whether antenatal care moderates the effect of the proposed lifestyle risks to miscarriage risk, a topic that has not been investigated in prior studies.

Pathways to Miscarriage: Conceptual Model

In the present research, the conceptual model is used to test the direct and mediating relationships between maternal and lifestyle miscarriage risk factors. Specifically, the model includes the following key relationships:

- 1. Direct effects of maternal age and education level:** Socio demographic variables are specifically maternal age and education level are postulated to have direct impact on miscarriage risk (Chung *et al.*, 2023). Higher maternal age is thought to raise the risk of miscarriage, and lower education levels are also thought to be linked with the risk due to higher rates of risky behaviours and lower health service utilization (Fishman SH & Min S., 2018).

2. **Direct effects of antenatal care:** It is postulated that improved antenatal clinic attendance decreases the rates of miscarriage because problems that may lead to miscarriage are promptly identified and they are dealt with accordingly (Amponsah et al.,2022). After examining these correlations separately, the effect of direct antenatal care on miscarriage, as well as how antenatal care mediates risk factors and miscarriage are described using the proposed model.

3. **Indirect effects of lifestyle risks:** Cigarette smoking, alcohol intake and physical inactivity are hypothesized to have only an indirect influence on miscarriage via a mediatorial role of the number of antenatal care visits made (Chung *et al.*, 2023; Gascoigne *et al.*, 2023) For instance, while unhealthy women behaviours may include missing regular antenatal visits, they are likely to result in increased miscarriage rates. Second, patterns in lifestyle risks are modeled to have a direct impact on miscarriage and an indirect impact through antenatal care.

4. **Latent construct for lifestyle risks:** Our data analysis fits an unspecified number of indicators for lifestyle risks under the assumption that it can be regressed on smoking, alcohol, and physical activity. This makes it possible to obtain a better assessment of the extent to which these behaviors are associated with an increased risk of miscarriage.

Materials and Methods

Study Design

In this study cross-sectional data was mimicked for exploring the impact of various maternal and lifestyle factors on risk of miscarriage using Structural Equation Modeling (SEM). The simulation was performed to assess the patterns of interaction between manifest and hidden variables and to assess possible links between these factors and miscarriage. In particular, by using simulation data we could examine most of the hypothetical interactions in the safest possible way and could test the SEM models, based on how well they performed under different conditions using the approach described in Normande et al. (2024) and Kananura *et al.*, (2017) This approach is frequently used in the assessment of theories or validation of model specifications when realistic samples maybe difficult to come by.

Simulated Data Generation

The data used in the study were artificial and those utilized in calculating the probability of miscarriage were based on research hypothesis and theoretical interaction between maternal features and lifestyle and miscarriage. The data were derived from assuming reasonable underlying distributions of the variables and their relationships as described in Rhemtulla *et al.* (2012).

Key Assumptions for the Simulated Data:

- Maternal Age (MA): As if it were a real continuum, normally distributed with a mean age of 30 years and an age deviation of 5 years.
- Education Level (EL): Operationalised as an ordinal variable; (1) Less than a primary education level, (2) Secondary school, or (3) College or university level. These categories were given probabilities taken from an assumed distribution of education levels within the total population.
- Lifestyle Risks (LS): Two of these constructs are smoking frequency and alcohol intake, while physical activity level is its measure, which are themselves latent variables indicated by three

manifest variables: smoking, alcohol consumption and physical activity. These factors were modeled as binary variables, where:

o Smoking: This variable is a binary measure with score 1, if a subject smoked regularly, otherwise score 0, if they did not smoke.

o Alcohol Consumption: 1 (where they consume it regularly) or 0 (where they do not consume at all).

o Physical Activity: 1 if they perform regular physical activity or 0 if they do not perform regular activity at all.

Lifestyle risk is the other construct that has been modeled into a latent variable using factor loading with the usual assumption that factor loading is positively related

. • Antenatal Care Visits (ACV): Imitated as a quantitative variable that ranged from 0 to 15 visits. The distribution of the numbers of visits was taken to be normal with an expected value of 8 visits and standard deviation of 3.

• Miscarriage (M): The outcome variable was imputed as a dummy variable (1 = miscarriage; 0 = live birth) with reference to maternal age, education levels, and risk factors, as well as antenatal clinic visits. The risk of miscarriage was believed to rise with higher Lifestyle Risk Score, older maternal age, and fewer visits to health care providers during pregnancy.

Simulation Process

The data were simulated in accordance to Rhemtulla et al. (2012) using the following steps:

- (a) Generate Maternal Age (MA): Employed as a rational variable with a normal probability distribution with mean value of 30years and a standard deviation of 5 years.
- (b) Generate Education Level (EL): Indicated here as ordinal with probability values of each category of the education level (Agresti, A., 2012). For example:
 - No formal education or primary education: 25%
 - Secondary education: 45%
 - Higher education: 30%
- (c) Generate Lifestyle Risks (LS): o Simulated as a latent variable with three indicators: tobacco use, drinking routines and exercise or movement routines. Both smoking and alcohol consumption were coded as 0 and 1, probability of which were 0.3 and 0.6 respectively for physical activity. The endogenous construct was specified to have the same factor loadings with the three measured variables.
- (d) Generate Antenatal Care Visits (ACV): It is assumed as being an ordinal variable with values simulated from a normal distribution with a mean of 8 visits and a standard deviation of 3 visits.
- (e) **Generate Miscarriage (M):** Miscarriage risk was simulated as a binary variable, where the probability of miscarriage was modeled using the following logistic regression equation derived as follows:

Let Miscarriage risk be denoted with y such that:

$$y = \begin{cases} 1 & \text{patient experienced miscarriage risk} \\ 0 & \text{did not experienced miscarriage risk} \end{cases}$$

Therefore, the model is given as

$$\log\left(\frac{p(y = 1)}{1 - p(y = 1)}\right) = \beta_0 + \beta_1\text{MA} + \beta_2\text{EL} + \beta_3\text{LS} + \beta_4\text{ACV}.$$

where

- (i) $p(y = 1)$ is the probability of miscarriage (dependent variable)
- (ii) β_0 is the intercept which represents the baseline **log-odds** of miscarriage when all predictor variables are set to their reference categories or zero (for continuous variables).
- (iii) $\beta_1, \beta_2, \beta_3, \beta_4$ are the independent variable's corresponding coefficients.

Regression coefficients (β) in (iii) were estimated based on theoretical assumptions as described in Agresti, A. (2012), Bollen, K. A. (2014) and Gelman, A. (2007): In the same equation, whereas positive association with maternal age is shown by $\beta_1 = 0.05$. $\beta_2 = -0.1$ (negative correlation with education level), Generally: $\beta_3 = 0.3$ (indicating a positive relationship with lifestyle risks). $\beta_4 = 0.02$ (indicating a negative relationship with antenatal care visits). Having obtained the outcome of the logistic regression equation, a binary variable for miscarriage was created from the probability score for each simulated participant, assigned either 0 or 1.

Data Analysis

The simulated dataset underwent structural equation modeling analysis as described in Malaju (2023; Xia & Yang, 2019) to determine maternal and lifestyle factors' direct and mediated effects on miscarriage risk. SEM enables the estimation of both manifest and unobserved constructs and their association; it is used more frequently in research with intricate structures of variables and processes.

1. **Model Specification:** The hypothesized model was defined in the following way:
 - Latent Variable: Lifestyle risks (LS) were estimated as a form of unobserved construct with three manifest variables: smoking, alcohol intake, and exercise.
 - Direct Effects: The relationships between MA, EL, LS, and M were defined in terms of the direct effects, where MA, EL, and LS were specified as causing M.
 - Indirect Effects: The mediated path analysis involved the idea that LS are associated with miscarriage through a causal mediator; antenatal care visits.
2. **Model Estimation:** SEM model was analysed under ML estimation procedure as described in Li *et al.* (2016) and Rhemtulla *et al.* (2012). All of the parameters of the model, including regression coefficients and their standard errors, factor loadings, variances and covariances, were freely estimated.
3. **Model Fit Assessment:** To assess the model fit the following indices have been used as described in the paper of Xia, Y., & Yang, Y. (2019):
 - **Chi-square test:** This, in return, imply that the Chi-square value is insignificant, mean that the model fit the data appropriately.
 - **Root Mean Square Error of Approximation (RMSEA):** Mathematically, values below 0.05 SSD depict a good fit while those values in the range of 0.05-

0.08 are acceptable. o Comparative Fit Index (CFI): Closeness of fit is a vital test and, specifically, a CFI of above 0.90 is considered desirable.

- **Tucker-Lewis Index (TLI):** Any TLI value more than 0.90 indicates the satisfaction of this requirement of model fit.
4. **Indirect Effects:** To estimate the indirect effects of lifestyle risks on miscarriage through ANC visits, products of coefficients along the indirect paths were obtained.
 5. **Sensitivity Analysis:** Thus, to examine the sensitivity of the model, various aspects such as the association between lifestyle risks and miscarriage, were adjusted in order to assess the goodness of the fit under these circumstances.
 6. **Data Visualization:** Covariance-based path diagrams were used to depict the SEM model and with estimates of the between variables. These diagrams were useful in communicating the hypothesized and estimated effect of both maternal and lifestyle factors on miscarriage risk.

Ethical Considerations

Pseudo-data was used in this study but adherent to all ethical standards towards the use of data and modeling. The simulation was carried out with the purpose to analyze theoretical relations and to check SEM's performance. When it comes to the source, the authors do not involve real data hence no ethical issues such as using data involving matters such as miscarriages.

Limitations

Some drawbacks of this simulated research are as follows. These results depend on assumed relationships and potentially misrepresent actual data. Another limitation is that data and hence model assumptions pertain to distribution of the lifestyle behaviors and relation between the variables may not be similar across populations. Furthermore, given that the data are not actual data collected from real people, there is no problem such as socially desirable bias or measurement errors involved. But there is always a possibility of refining the model and these studies using real life data could support these findings.

Results and Discussion

Results

Model Fit and Evaluation

Based on the simulated data, the SEM model was estimated, and several fit indices were examined. The results showed good model fit based on common fit indices:

- **Chi-Square Test:** The Chi-square statistic for the model was not significant ($p= 0.441$) meaning that observed data does not differ significantly from the model-implied data- the model fit is acceptable.
- **Root Mean Square Error of Approximation (RMSEA):** The chi-square test of fitness yielded a value of 200.30 which compares well with the suggested sequence value of 344.48, and the RMSEA value was 0.003, which is below the threshold of 0.05 hence the model had a good fit. For RMSEA, we obtained a 90% confidence interval of 0.000 to 0.046, therefore being also in line with the necessity of the model.

- Comparative Fit Index (CFI): Hence the total value of CFI was 1.000; that is, it can be noted that values closer to 1.0 present a better fit to the data.
- Tucker-Lewis Index (TLI): The TLI value of 1.024 confirms the acceptable model of fit. The calculated value of TLI was higher than 1 showing that the proposed model is better compared to a baseline model. With these results, it can be concluded that the proposed model fits the simulated data well and provides reassurance of the estimates of the relationships between the variables.

Parameter Estimates

The analysis identifies a numeral of maternal and lifestyle characteristics associated with miscarriage risk based on the estimated parameters in the SEM analysis. Below are the key findings:

1. Maternal Age (MA): Estimate: -0.004 ($p = 0.221$)
 Interpretation: First, we could not establish a relationship between miscarriage risk and maternal age. However, the attained coefficient was negative and also insignificant implying that maternal age was not so influential in determining miscarriage rates as seen in this simulated model. This result may be contrary to clinical research about the detrimental effect of increased maternal age on miscarriage, further study is needed using actual data.
2. Education Level (EL): Estimate: -0.001 ($p = 0.977$)
 Interpretation: Unfortunately, there was no correlation between education level and miscarriage risk. Thus, we can conclude that according to the assembled dummy data, there is no dependence between miscarriage results and maternal education. This result could be explained by the fact that, in the current simulation of the 'education level' variable, it was categorized into only three ordered groups.
3. Lifestyle Risks (LS): Estimate: 0.226 ($p = 0.000$)
 Interpretation: Smoking, alcohol consumption, and physical activity, established lifestyle risks, were confirmed to be related to miscarriage risk. This implies that risks associated with elevated lifestyle increase the chances of miscarriage as presented by the positive coefficient above. This accords with existing studies associating miscarriages with specific lifestyle factors such as smoking and drinking alcohol. But, it is important to remember, in the real world, lifestyle choices do not exist in isolation and may combine with others such as genetic predisposition or coexisting diseases.
4. Antenatal Care Visits (ACV): Estimate: -0.027 ($p = 0.533$)
 Interpretation: The extent of antenatal care visit miscarriage risk was not affected in this synthetic model. This may be because, in the previous analysis, antenatal care visits were measured as a continuous variable instead of using more complex approaches that control for factors like the quality of care and timing of visits. The non-statistical outcome is most definitely worthy of further examination, especially given that antenatal care is a factor that helps decrease pregnancy complications.

Indirect Effects

The indirect effect of lifestyle risks on miscarriage through the number of antenatal care visits was also modeled. We also found lifestyle risks were independent of the number of antenatal care visits, as indicated by the insignificant value for the path coefficient

between the two variables ($p = 0.931$). Therefore, the above analysis of total and direct mediation of antenatal care visits showed that the impact of lifestyle risks on miscarriage was actually direct rather than to do with the number of antenatal care visits.

Discussion

The finding of this study offers us preliminary evidence regarding the associations between the studied maternal and lifestyle factors and miscarriage risk. Significant direct effects of lifestyle risks, especially smoking and alcohol consumption, on miscarriage risk were indicated by the SEM model, which was in line with the previous work. Nonetheless, maternal age, education level, and number of ANC visits were not found to have any impacts in this simulation. Possible reasons for insignificant results in the study that relate to maternal age, education level, and number of antenatal care visits are explained below. First, the simulation relied on simplified assumptions of relationships between various factors; it is possible that these or even other relationships between the factors in this context may not hold as they are in real life. For example, maternal age may have a curvilinear, adverse relation to miscarriage, or this relation may depend on health status, prior pregnancy experience, and genetic factors. Furthermore, education level was assessed as an ordinal measure comprising three extended categories, which might not classify education level with sufficient accuracy to identify a given impact, like ANC visits which are also crucial in reducing pregnancy complications may not have been well captured in this study. The actual data from real life would possibly suggest that the timing of the ANC visits, the complete quality, and the frequency of ANC visits are the critical determinants of miscarriage, and these were not effectively evaluated in the simulation. Although there was no substantial relationship between the visits for antenatal care and the risk of miscarriage, it may be because of the weaknesses inherent in the basic model. On the other hand, the strong impact of analyzed lifestyle risks on the risk of miscarriage underlines the necessity of effective lifestyle modification in pregnancy. Indeed, smoking and alcohol are established causes of miscarriage and this study equally confirms these factors. Lifestyle risks were significantly associated with miscarriage, after controlling for antenatal care visits thus supporting the idea that strategies to modify the identified lifestyle risks could have a significant potential to reduce the risks of miscarriage. Smoking and alcohol cessation media brief interventions during pregnancy could therefore be useful in preventing miscarriages.

Limitation

The main limitations of this study are as follows: First, because this was an experimental simulation, results might be generalized cautiously in that the observed relationships among variables of interest may not necessarily capture the relationships in actual population(s). Some assumptions made in the simulation may not hold in real data for instance the assumption that maternal age is Normally distributed and the fact that education level is categorized simply as low middle and high. Further, some possible confounding factors are not included in the model, these are diseases that pregnant women could have had in the past any genetic disorder that runs in families, or a woman's social status. In addition, the simulation also had no interaction terms

whereby main effects can be conditioned of each other making it difficult for researchers to determine the effect of one on the other between the maternal characteristics and the lifestyle factors on the risk of miscarriage. For example, the relationship between maternal age and miscarriage weighting may be moderated by lifestyle or by the number of antenatal visits. Finally, only a few factors were studied in the present study; thus, it may be possible that there are other factors, not captured within the above model, for example, environmental conditions, stress levels, or, perhaps the use of fertility treatments. The following additional factors can come into research works with real-world data to help enhance the understanding of the risk of miscarriage;

Conclusion and Recommendation

Therefore, the author of this study stresses the importance of lifestyle risks in miscarriage and their need to develop measures designed for the modification of lifestyle factors for the enhancement of pregnancy results. Future work will have to extend the presented model and will have to use it to validate these results with actual data. Future research should come up and test the findings from this simulation study to see whether the results hold up when actual data is used to compare the interrelation between maternal characteristics and lifestyle and miscarriage risk. This would include wider sampling, and using finer measures of maternal characteristics, lifestyle behaviors, and antenatal care to test the model. Thus, it may be also interesting to analyze the mechanisms of interactions between lifestyle factors and other individual characteristics, for example, health status, and miscarriage risk.

REFERENCE

- Agresti, A. (2012). *Categorical data analysis* (Vol. 792). John Wiley & Sons.
- American College of Obstetricians and Gynecologists. (2013). ACOG Committee opinion no. 548: weight gain during pregnancy. *Obstetrics and gynecology*, 121(1), 210-212.
- Amponsah-Tabi S, Dassah ET, Asubonteng GO, Ankobea F, Annan JJK, Senu E, Opoku S, Opoku E, Opore-Addo HS. An assessment of the quality of antenatal care and pregnancy outcomes in a tertiary hospital in Ghana. *PLoS One*. 2022 Oct 12;17(10):e0275933. doi: 10.1371/journal.pone.0275933. PMID: 36223426; PMCID: PMC9555636.
- Bailey, B. A., & Sokol, R. J. (2011). Prenatal alcohol exposure and miscarriage, stillbirth, preterm delivery, and sudden infant death syndrome. *Alcohol Research & Health*, 34(1), 86.
- Bollen, K. A. (2014). *Structural equations with latent variables*. John Wiley & Sons.
- Celep F, Karagüzel A, Ozeren M, Bozkaya H. The frequency of chromosomal abnormalities in patients with reproductive failure. *Eur J Obstet Gynecol Reprod Biol*. 2006;127:106–9. doi: 10.1016/j.ejogrb.2005.12.019.
- Chibayi, J. O., Ranyimbo, A. O., & Akang'o, A. A. (2024). Modeling the Risk Factors of Miscarriage Using Survival Analysis Techniques. *Asian Journal of Probability and Statistics*, 26(9), 160-184.

- Chung, Y., Melo, P., Pickering, O., Dhillon-Smith, R., Coomarasamy, A., & Devall, A. (2023). The association between dietary patterns and risk of miscarriage: a systematic review and meta-analysis. *Fertility and Sterility*, *120*(2), 333-357.
- Collier, J. (2020). *Applied structural equation modeling using AMOS: Basic to advanced techniques*. Routledge.
- Fishman SH, Min S. Maternal Age and Offspring's Educational Attainment. *J Marriage Fam*. 2018 Aug;*80*(4):853-870. doi: 10.1111/jomf.12490. Epub 2018 May 22. PMID: 30555181; PMCID: PMC6289533.
- Flynn H, Yan J, Saravelos SH, Li TC. Comparison of reproductive outcome, including the pattern of loss, between couples with chromosomal abnormalities and those with unexplained repeated miscarriages. *J Obstet Gynaecol Res*. 2014;*40*:109–16. doi: 10.1111/jog.12133.
- Gascoigne EL, Webster CM, Honart AW, Wang P, Smith-Ryan A, Manuck TA. Physical activity and pregnancy outcomes: an expert review. *Am J Obstet Gynecol MFM*. 2023 Jan;*5*(1):100758. doi: 10.1016/j.ajogmf.2022.100758. Epub 2022 Sep 26. PMID: 36174931; PMCID: PMC9772147.
- Gelman, A. (2007). *Data analysis using regression and multilevel/hierarchical models*. Cambridge university press.
- Hosker, D. K., Elkins, R. M., & Potter, M. P. (2019). Promoting mental health and wellness in youth through physical activity, nutrition, and sleep. *Child and Adolescent Psychiatric Clinics*, *28*(2), 171-193.
- Hyde KJ, Schust DJ. Genetic considerations in recurrent pregnancy loss. *Cold Spring Harb Perspect Med*. 2015;*5*:a023119. doi: 10.1101/cshperspect.a023119.
- Kananura, R. M., Wamala, R., Ekirapa-Kiracho, E., Tetui, M., Kiwanuka, S. N., Waiswa, P., & Atuhaire, L. K. (2017). A structural equation analysis on the relationship between maternal health services utilization and newborn health outcomes: a cross-sectional study in Eastern Uganda. *BMC pregnancy and childbirth*, *17*, 1-12.
- Krieg, S. A., Shahine, L. K., & Lathi, R. B. (2016). Environmental exposure to endocrine-disrupting chemicals and miscarriage. *Fertility and sterility*, *106*(4), 941-947.
- Kumar, S. (2011). Occupational, environmental and lifestyle factors associated with spontaneous abortion. *Reproductive sciences*, *18*(10), 915-930.
- Li, C. H. (2016). The performance of ML, DWLS, and ULS estimation with robust corrections in structural equation models with ordinal variables. *Psychological methods*, *21*(3), 369.
- Malaju, M. T. (2023). A structural equation modelling of the direct and indirect factors associated with functional status over time as measured by WHODAS-32 items among postpartum women in Northwest Ethiopia. *Archives of Public Health*, *81*(1), 41.

- McCarthy, F. P., O'keeffe, L. M., Khashan, A. S., North, R. A., Poston, L., McCowan, L. M., ... & Kenny, L. C. (2013). Association between maternal alcohol consumption in early pregnancy and pregnancy outcomes. *Obstetrics & Gynecology*, *122*(4), 830-837.
- Mueller, R. O., & Hancock, G. R. (2018). Structural equation modeling. In *The reviewer's guide to quantitative methods in the social sciences* (pp. 445-456). Routledge.
- Normande MEMEM, da Silva LC, de Menezes RCE, Florêncio TMMT, Clemente APG. Association of parental level of education and child factors on length-for-age indicator among socially vulnerable children aged 6-24 months from a Brazilian state using structural equation modelling. *Br J Nutr*. 2024 Jul 28;132(2):192-199. doi: 10.1017/S0007114524001119. Epub 2024 May 28. PMID: 38804182.
- Oei, J. L. (2020). Alcohol use in pregnancy and its impact on the mother and child. *Addiction*, *115*(11), 2148-2163.
- Pal AK, Ambulkar PS, Waghmare JE, Wankhede V, Shende MR, Tarnekar AM. Chromosomal Aberrations in Couples with Pregnancy Loss: A Retrospective Study. *J Hum Reprod Sci*. 2018 Jul-Sep;11(3):247-253. doi: 10.4103/jhrs.JHRS_124_17. PMID: 30568354; PMCID: PMC6262666.
- Rhemtulla, M., Brosseau-Liard, P. É., & Savalei, V. (2012). When can categorical variables be treated as continuous? A comparison of robust continuous and categorical SEM estimation methods under suboptimal conditions. *Psychological methods*, *17*(3), 354.
- Sampasa-Kanyinga, H., Colman, I., Goldfield, G. S., Janssen, I., Wang, J., Podinic, I., ... & Chaput, J. P. (2020). Combinations of physical activity, sedentary time, and sleep duration and their associations with depressive symptoms and other mental health problems in children and adolescents: a systematic review. *International Journal of Behavioral Nutrition and Physical Activity*, *17*, 1-16.
- Savalei, V., & Rhemtulla, M. (2013). The performance of robust test statistics with categorical data. *British Journal of Mathematical and Statistical Psychology*, *66*(2), 201-223.
- Tarka, P. (2018). An overview of structural equation modeling: its beginnings, historical development, usefulness and controversies in the social sciences. *Quality & quantity*, *52*, 313-354.
- Wang, B. T., Chong, T. P., Boyar, F. Z., Kopita, K. A., Ross, L. P., El-Naggar, M. M., ... & Anguiano, A. L. (2014). Abnormalities in spontaneous abortions detected by G-banding and chromosomal microarray analysis (CMA) at a national reference laboratory. *Molecular cytogenetics*, *7*, 1-6.
- Xia, Y., & Yang, Y. (2019). RMSEA, CFI, and TLI in structural equation modeling with ordered categorical data: The story they tell depends on the estimation methods. *Behavior research methods*, *51*, 409-428.