

**Profile of Lipid and Antioxidant Activity Lipids Extracted from *Pangasius Pangasius*: Complementary Analysis**

---

**ABSTRACT**

Extraction of lipids from *Pangasius Pangasius* employed a unique simple route, such as the solvent extraction method, which is better useful for extract lipids from fish, cost effective and comparatively easier than others process where ethanol is a diluent and chloroform acts as a solvent. The prime objective of this manuscript is to determine the chemical composition and antioxidant activity of lipids from *Pangasius Pangasius*. The chemical composition determination by Gas Chromatography Mass Spectrometry (GC-MS). The most abundant lipid components were Dodecanoic acid 1,2,3 Propanetriyl ester 86.67 % found in GC-MS which have strong bactericidal properties. It is utilized as an algal and plant metabolite and possesses antimicrobial qualities. For determination of the antioxidant activity, the Ultraviolet-Visible Spectrometer (UV) was introduced. The degree of hydrolysis, DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity the antioxidant activity can be estimated. The control value of the UV is 1.297 a. u. and the absorption value by sample minimum value is 1.168 a. u. and the maximum value is 1.208 a. u. The inhibition rate maximum of 0.0995 % and a minimum of 0.0686 % by the human body. The absorption of UV light by the sample is near the control

value which indicates lipids contain a little percentage of antioxidant activity. The sample contains a small percentage of 0.33 of L (+) ascorbic acid 2,6 di-hexa-decanoate which is an antioxidant agent that reduces free radicals in the human body. This outcome highlighted its possibility as an origin of bioactive chemicals and expanded our knowledge of its nutritive and therapeutic properties.

**Keywords:** Antioxidant activity, Chloroform, Ethanol, *Pangasius Pangasius*, Solvent extraction.

## 1. Introduction

Fish is an invaluable source of vital vitamins essential for the proper functioning of the human body [1]. Fish liver oil, in particular, is an extraordinary reservoir of vitamins A and D where Vitamin A maintains healthy skin and promotes bone development and vitamin D plays a pivotal role strength of teeth and bones [1]. It is noteworthy that oily fish, in particular, constitute an excellent source of vitamin D [1]. A large number of peptides produced from hydrolyzed dietary proteins have antioxidative properties [2]. Antioxidative action has been observed for fish protein hydrolysates, such as skin gelatin hydrolysate from Alaska Pollack [2], yellowfin sole [3] and phytochemicals [4]. Hydrolyzed fish protein may also be a promising source of anticancer peptides [5], angiotensin I-converting enzyme (ACE) inhibitors [6], an anti-anemia drug [7] and a component of microbial growth media [8] considering preliminary evidence. On the other hand, not much is known about the antioxidative properties of collagen peptides derived from *pangasius pangasius* or striped catfish [9]. The *pangasius hypophthalmus* fish is a popular food source because it is rich in nutrients, including fat, protein, vitamins and minerals, all of which have a substantial positive impact on human health [9]. *Pangasius* boasts minerals, vitamins A, B, and E, and these antioxidants efficiently reduce oxidative stress [9]. Dietary sources of *pangasius hypophthalmus* fish include vitamins and minerals that function as antioxidants, thereby mitigating the effects of oxidative stress [10,23]. *Pangasius hypophthalmus* oil's omega-

3 content improves zinc transfer to the cell membrane which assists in regulating insulin, slows down its breakdown, and improves insulin sensitivity [10]. In food preparation items, gelatin can be utilized as a stabilizer and emulsifier. Ethical and theological arguments have shown that fish gelatin is a more efficient substitute for animal gelatin [11]. Usually, freshwater fish are utilized as a source of fresh gelatin [11]. However, numerous investigations have demonstrated that after the edible portions of some freshwater fish are removed, a significant amount of by-product remains [12]. High gelatin yields are therefore anticipated from fish such as tilapia [12], Nile perch [13] and striped catfish (*Pangasius sutchi fowler*) [14]. Gelatin extraction and the characterization of several freshwater fish as substitute sources of gelatin revealed that the gelatin derived from *pangasius pangasius* catfish had the best rheological characteristics when compared to the other fish [14,23]. Its characteristics include a total amino acid content, a melting point of 32.0 °C, a gelling temperature of 12.0 °C, a melting strength of 273.58 g and a viscosity of 36.5 cP [14,26]. The lipids extracted from *Pangasius pangasius* and *Pangasius sutchi* have undergone nutrient content analysis. The fishes' specific gravity, refractive index and viscosity coefficient were recorded at  $0.97 \pm 0.01$  and  $0.94 \pm 0.02$  (at 30.0 °C),  $4.88 \pm 0.25$  and  $5.49 \pm 0.30$  (at 30.0 °C), and  $448.96 \pm 2.5$  and  $421.76 \pm 2.1$ , respectively. Chemical characterization of the lipids in both fish species has yielded significant results [14,23,26]. Furthermore, *P. pangasius* and *P. sutchi* have demonstrated carbohydrate, protein, cholesterol and lipid contents of  $1.89 \pm 0.35$  % and  $0.75 \pm 0.11$  %,  $35.0 \pm 2.1$  % and  $38.37 \pm 2.5$  %,  $17.46 \pm 1.15$  % and  $13.79 \pm 0.37$  % and  $10.01 \pm 1.11$  % and  $6.16 \pm 0.54$  %, respectively. Adequate levels of humidity, solids and cinder percentages were discovered. Furthermore, *P. pangasius's* mineral levels such as calcium, phosphorus and zinc were inferior, while iron levels were greater than those of *P. sutchi* [15,25]. The fatty acid compositions of *P. pangasius* and *P. sutchi* consisted amount of lauric acid 13.36 and 4.26 %, palmitic acid 26.15 and 29.32 %, oleic acid 46.07 and 59.16 % and stearic acid 14.40 and 7.24 % correspondingly [15]. Additionally, consuming

pangasius fish provides a rich source of essential proteins which are crucial for combating protein insufficiency. In contrast to other fish, pangasius is economically viable and offers a higher concentration of protein and beneficial cholesterol [16]. In Bangladesh, two distinct varieties of the revered Pangas fish can be discovered- the first being the cultivated Pangas and the latter being the natural Pangas that thrive in the rivers [25]. While the former is raised in household-installed tanks and ponds, the latter is preferred by the majority of the population owing to its delectable taste [16]. However, as a result of its scarce availability, the natural river pangasius catfish comes with a hefty price tag.

Freshwater catfish or *Sperata aor* are highly preferred by consumers and make a substantial contribution to the production of fisheries in tropical rivers [16]. Nutritional profiles of pangasius catfish, both wild (*Pangasius pangasius*) and farmed (*Pangasius hypophthalmus*) were gathered from several river and culture pond sources in Bangladesh [17]. The indigenous Pangas exhibited an elevated protein content of  $26.06 \pm 1.27$  % and fat content of  $14.79 \pm 2.47$  %, in contrast to the hybrid Pangas which demonstrated a protein content of  $23.18 \pm 2.11$  % and a fat content of  $11.11 \pm 1.75$  % [18]. It has been stated that the muscles of farmed pangasius are exported to more than 80.0 nations worldwide, including the US, Germany and the Netherlands where frozen fillets devoid of skin and bone are much sought after [19]. The fillets exhibited elevated moisture levels ranging from 80.0 % to 85.0 % coupled with diminished protein levels ranging from 12.60 % to 15.60 % and lipid levels hovering between 1.10 % and 3.00 % [20,26]. A study was designed to ascertain the proximate composition of fatty acids, amino acids and mineral contents of frozen fillets from Pangasius catfish (*Pangasius hypophthalmus*) imported to Poland, Germany and Ukraine to establish the nutritional composition of these fish [21]. One of Vietnam's primary seafood products is Pangasius and it was studied more than 1.3 million tonnes cultivated in 2018 [22]. However, due to its superior taste compared to farmed pangasius and the

wild pangasius (*Pangasius pangasius*) species of riverine catfish is in high demand locally [24]. Due to high consumption with high cost and for this point of the study chemical composition and antioxidant activity of lipids from *Pangasius pangasius* profiling is the prime object of this manuscript.

## **2.0 Materials and Methods**

### **2.1 Materials**

The DPPH (2,2-diphenyl-1-picrylhydrazyl), ethanol and chloroform were procured from Sigma-Aldrich, Germany and DI water was collected from the ACCE Lab plant. The *Pangasius pangasius* was also collected from the local market Sheikhpara, Jhenaidah -7300, Bangladesh.

### **2.2 Methodology Selection for this study**

There are some previous methodologies by which researchers extract lipids from *Pangasius pangasius*.

#### **a. First method:**

To estimate total lipids, the Pangus fish (*P. pangasius*) was caught. The Bligh and Dyer method called for the use of a 2:1 v/v mixture of methanol and chloroform. A 500 ml volumetric flask was filled with 50.0 g of the fish after it had been weighed out. Then chloroform-methanol combination was added and thoroughly mixed. It was kept overnight at room temperature. After three layers were visible. Chloroform's transparent bottom layer held everything the lipids, a thick, paste-like surface, and a colored aqueous layer of methanol that contained only water-soluble material [27]. After discarding the methanol layer, the lower layer was meticulously extracted. The organic layer from each extraction process was carefully evaporated. Additionally, it was mentioned that the sample should be shielded from light by being covered

with dark papers. This resulted from the polymerization and breakdown of some lipids upon exposure to light and air [27].

**b. The second method: Extraction of Fat Fish:**

The Soxhlet method was used to calculate the fish's fat content, and the steps involved were as follows:

After being crushed (ideally dry and passing 40 mesh), the samples were weighed. As 50.0 g and placed in a paper-lead Soxhlet extraction tube. Water for cooling passed via the condenser for five hours, and a test tube filled with 100 ml of diethyl ether solvent was placed on a Soxhlet distillation apparatus. then carried out eight repetitions till the unused sample weight reached 400 g. Each sample repetition received 50 ml of diethyl ether for the second and subsequent times. Fat that has been extracted from the soxhlet and dried in an oven set to 600C for two hours is mixed with diethyl ether. After that, it was chilled for half an hour in a desiccator [26].

**Separation of Phospholipids Crude Extract of Total Lipid by Dissolution:**

Chloroform was used to remove the total lipid that was produced from the extraction process 40 millilitres of chloroform were used to extract 10 grammes of total lipid. Centrifugation was used to separate a non-polar lipid and a chloroform-soluble fraction at 5000.0 rpm for ten minutes. removed from the insoluble portion of a polar lipid. To separate the soluble and insoluble fractions, the soluble fraction was centrifuged after being dried with nitrogen and re-extracted using chloroform[26]. polar lipid's insoluble fraction and combined with the initial extraction's insoluble fraction. Additionally, the polar phospholipid was dissolved in methanol to extract the chloroform insoluble fraction. Centrifugation was used to separate the insoluble fraction for ten minutes at 5000 rpm. Drying the soluble fraction is allowed [28].


The two procedures mentioned above are used to extract lipids from Pangasius.

### 2.2.1 Sample Preparation from *Pangasius Pangasius*

The lipids of *Pangasius Pangasius* can be extracted by solvent extraction process [26,27]. In the solvent extraction process, the collected pangas fish were cut into small pieces and then the small pieces of fish were cleaned with water and DI water. After cleaning the fish sample, it was ground in a hand-grinding machine.

### 2.2.2 Lipid Extraction Process

From the groundfish sample, 20.00 grams of mesh fish sample is taken in a bucket and then the 750.00 ml of chloroform and 250.00 ml of ethanol are taken in that bucket to maintain 3:1 (v/v). Then the chloroform, mesh fish sample and ethanol are mixed well [26-28]. Then the total mixture is divided into four conical flasks in the same equivalent ratio. Then the all-conical flask is shaken well by an orbital shaker (RJH5005, Richard James Hilfiger, UK). The speed of the shaking should be 300.00 rpm maximized.



**Fig. 1.** Preparation route of lipid sample from *Pangasius Pangasius*.

The shaking process is continued for 20.0 hours for complete extraction. Then the well shake is kept quiet for two hours for complete separation of chloroform and ethanol layer [26,27]. The chloroform layer contains the lipid of *Pangasius Pangasius*. Then the two layers are separated by a separating funnel. The ethanol layer is collected in a bucket and another chloroform layer is collected in a beaker. Then the chloroform layer evaporated by nature in the shade. In the evaporation process, the extra heat or sun rays cannot be used because the fat can be broken. After the complete evaporation, the lipids of pangas appear [26-27]. The route is illustrated in Fig. 1.

### **2.2.3 Analytical Techniques**

Gas Chromatography-Mass Spectrometry (GC-MS) and Ultraviolet-Visible Spectrometry Analysis were conducted for this study. The lipid sample is collected in a refrigerator for analysis of chemical composition by GC-MS (Gas Chromatography-Mass Spectrometry) and antioxidant test by UV (Ultraviolet-Visible Spectrometry).

## **3.0 Characterization**

### **3.1 Gas Chromatography Mass Spectrometry (GC-MS) Analysis**

GC-MS analysis was carried out with Clarus ® 690 gas chromatograph (PerkinElmer, CA, USA) using a column (Elite-35, 30m length, 0.25 mm diameter, 0.25 µm thickness of film) and it was equipped with Clarus ® SQ 8 C mass spectrophotometry (PerkinElmer, CA, USA). 1.0 µL sample was injected (split less mode) and pure Helium (He) 99.99 % was used as a carrier gas at a constant flow rate (1.0 mL/min) of 40.0 mins runtime. The sample was analyzed in EI (electron ionization) mode at high energy at 70.0 eV. Though inlet temperature was constant at 280.0 °C, column oven temperature was set at 60.0 °C (for 0.0 min), raised at 5.0 °C per minute to 240.0 °C and held for 4.0 mins. The sample compounds were identified by comparing them to the National Institute of Standards and Technology (NIST) database [27].

### 3.2 Ultraviolet-Visible Spectrometry Analysis

The antioxidant test can be done by the DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity method [28]. Various concentration of the sample was prepared with DPPH such as 25.0  $\mu$ l, 50.0  $\mu$ l, 100.0  $\mu$ l, 150.0  $\mu$ l, 200.0  $\mu$ l, 250.0  $\mu$ l, 300.0  $\mu$ l. The concentration of the blank sample was sorbed in a fixed wavelength at 517.0 nm. The Ultra violet light is absorbed by the lipid sample which estimates the antioxidant activity of the lipid sample. The antioxidant activity can be estimated by the inhibition of free radicals [28].

$$\text{Inhibition rate (\%)} = \{(\text{blank absorbance} - \text{sample absorbance}) / \text{blank absorbance}\} \times 100\%$$

Where blank is the absorbance of the control reaction which contains all reagents except the test compound. Sample absorbance containing test compound.

## 4.0 Result and Discussion

### 4.1. Chemical composition of lipid from *Pangasius Pangasius*


*Pangasius Pangasius* lipids are grey colour solid fat substances. Initially, during the extraction process, it forms a transparent liquid but after evaporation, it forms a solid grey colour. The identified compounds by GC-MS, according to their retention time in the column, the result is shown in Table 1.

**Table 1.** Chemical composition of the extracts from the sample of *Pangasius Pangasius*

Serial No.	Retention time (RT)	Compounds	Molecular Weight	Molecular Formula	Area (%)
1	7.44	propionic acid, 4-hydroxy-3-hexyl ester	174.0	C <sub>9</sub> H <sub>18</sub> O <sub>3</sub>	0.02
2	21.73	benzoic acid,4,4- (1,2-dioxo-1,2-	326.0	C <sub>18</sub> H <sub>14</sub> O <sub>6</sub>	0.05

		ethanediyl) bis-dimethyl ester			
3	21.95	propanoic acid, 2-methyl,2-(hydroxymethyl)-1-propyl butyl ester	216.0	C <sub>12</sub> H <sub>24</sub> O <sub>4</sub>	0.09
4	24.64	diethyl phthalate	222.0	C <sub>12</sub> H <sub>14</sub> O <sub>4</sub>	0.02
5	28.27	Hepta cosanoic acid, 25-methyl, methyl ester	438.0	C <sub>29</sub> H <sub>58</sub> O <sub>2</sub>	0.02
6	29.69	l-(+)-ascorbic acid 2,6-dihexa decanoate	652.0	C <sub>38</sub> H <sub>68</sub> O <sub>8</sub>	0.33
7	33.19	bis(2-ethylhexyl) phthalate	390.0	C <sub>24</sub> H <sub>38</sub> O <sub>4</sub>	12.37
8	37.36	dodecanoic acid,1,2,3-propanetriyl ester	638.0	C <sub>39</sub> H <sub>74</sub> O <sub>6</sub>	86.67

In GC-MS analysis, twenty-four (24.0) compounds were identified from the sample. The bioactive compounds identified from the sample were represented by their retention time (RT), molecular formula, molecular weight and peak area (%) in Table 1 and Fig. 2. represent the distinct GC-MS chromatogram. This chromatogram representation of the vertical line indicates the peak area, where the chemical compound can be identified compared with the National Institute of Standards and Technology (NIST) database. The horizontal line represented the retention time in minutes. The samples were run in the column for 40.0 minutes the sample was analyzed in electron ionization mode at high energy at 70.0 ev. The different peaks are formed at different retention times which represent the different chemical compositions.




**Fig. 2.** Retention time vs area curve from GC-MS.

First times, we observed propionic acid, 4-hydroxy-3-hexyl ester at the 7.44 retention time. It's useful in the food industry for the preservation of foods such as it is used to inhibit bacterial growth and used as a preservative for food [29]. Benzoic acid, 4,4- (1,2-dioxo-1,2-ethanediy) bis-dimethyl ester has an antimicrobial property which is beneficial for food and beverages and strong antibacterial property at pH 2.5 to 4.0 [30]. Propanoic acid, 2-methyl, 2-(hydroxymethyl)-1-propylbutyl ester known as methyl ethyl ketone peroxide acts as a strong oxidizing agent [31]. It's also used in the curing of unsaturated polymer resin as a catalyst. Effect of diethyl phthalate use in plastic and personal care products. It also has a hazardous property [31]. It causes male reproductive, female reproductive, liver and kidney damage and causes cancer [31]. Heptacosanoic acid, 25-methyl, methyl ester a long-chain fatty acid. It is an energy source, structure source for cell membranes and also signalling molecules [31]. L ( $\pm$ ) ascorbic acid 2,6-dihexadecanoate has an antioxidant property, it is also used for the reduction of free radicals from the human body. It also possesses antibacterial, antitumor and wound-healing properties. It is used for various pathogenic diseases [32].

Bis-2-Ethylhexyl is a plasticizer, it is intended use of food contact application. But it also has some side effects that cause cancer [33,25], birth defects or other reproductive harm [33]. It also interferes hormone system. Dodecanoic acid,1,2,3-propanetriyl ester is also known as glyceryl laurate [33]. It is used as an emulsifier, stabilizer, and also flavoring agent. It has also an antimicrobial property which extends the shelf life of products [33]. The major component from the extract is Dodecanoic acid, 1,2,3-Propanetriyl Ester at 86.67%, the second percentage from the extract is Bis-2-Ethylhexyl Phthalate 12.37% and the third is ( $\pm$ ) Ascorbic acid 2,6-Dihexa decanoate at 0.33 %. The GC-MS is used for the identification of various chemical compound and their percentages. In the analysis of the chemical composition of lipids which are extracted from *Pangasius Pangasius* . We use the GC-MS, here we found 24.0 bioactive chemical compounds.

#### **4.2. Antioxidant Properties Analysis**

The degree of hydrolysis could be estimated by DPPH radical which are require for the estimation of antioxidants. The lipids extracted from *Pangasius* fish contain little antioxidant power. DPPH Radical Scavenging Activity is expressed where DPPH is widely acceptable for the determination of free radical scavenging activities of antioxidants [34]. The DPPH radical's reduction capacity can be determined by an absorbance of 517.0 nm induced by antioxidants. The increasing of ultra-violet light with concentration indicates the increasing antioxidant property of this compound. However, during analysis of this compound, the absorption rate had been decreased with concentration. So, the compound has less antioxidant properties. The UV visible spectrum analysis concentration vs absorbance [35-49] of *Pangasius Pangasius* extracts is shown in Fig. 3.



**Fig. 3.** The UV visible spectrum analysis concentration vs absorbance of *Pangasius Pangasius* extracts.

The absorbance capacity of the lipid sample is shown in Fig. 3. It expressed that the increasing the concentration of 25.0 µl to 300.0 µl the absorbance decreased from 1.208 a.u. to 1.168 a.u. respectively. So, the low absorbance indicates that antioxidant properties are decreased expressed in Table 2.

**Table 2.** Absorption of ultra violet light by the extract from *Pangasius Pangasius*.

Observation number	Concentration (µl)	Absorption of UV light (a.u)
1	25.0	1.208
2	50.0	1.202
3	100.0	1.196
4	150.0	1.192
5	200.0	1.186
6	250.0	1.181
7	300.0	1.168
8	Control value	1.297

The exact antioxidant properties are illustrated by the inhibition rate which accumulated in Table 3. So, by calculating the inhibition rate, we can determine of antioxidant activity of the sample.

$$\text{Inhibition rate (I) \%} = \{1 - (A)_{\text{sample}} / (A)_{\text{control}}\} * 100$$

Here, A= absorption and I= inhibition rate

From the above equation, the inhibition rate is determined. If the inhibition rate increases with concentration indicates the absorption rate decreases. If the inhibition rate increases the antioxidant property of the compound also decreases.

**Table 3.** Free radical scavenging activity (DPPH) of lipids from *Pangasius Pangasius*.

Observation Number	Concentration ( $\mu\text{l}$ )	Inhibition rate (%)
1	25.0	0.0686
2	50.0	0.0732
3	100.0	0.0778
4	150.0	0.0809
5	200.0	0.0856
6	250.0	0.0894
7	300.0	0.0995
8	Control value	0.0000

Table 3 shows that the concentration increased from 25.0  $\mu\text{l}$  to 300.0  $\mu\text{l}$ . The inhibition rate increased from 0.0686 to 0.0995 % respectively. So, the high-concentration antioxidant properties of the extracts decreased and with the low-concentration, the low inhibition rate of the antioxidant properties of the extracts increased. The crystallographic details analysis [50-70] were employed its symmetry and orientation. For the purpose of estimation of antioxidant power used UV-spectrometry. Here we use DPPH radical which is a common reagent to determination of the degree of hydrolysis or antioxidant power. Here we found the lipids sample contain small amount of antioxidant properties.

### 4.3 Research Gap Identification and Aim of This Research

The research explored for the behinds of prospect were depicted in below mentioned.

Table 4: Research Gap Identification and aim

Study Status	The outcome of the Research	Gap identification and development study	Reference
Previous research	In the previous research, they worked with <i>Pangasius</i> fish bone and skin collagen with papain enzyme. The result of collagen peptide hydrolysed from fish skin had a greater reduction power activity (0.817) than the collagen peptide hydrolysed from fish bone (0.788). They also demonstrated carbohydrate, protein, cholesterol and lipid contents etc.	Previous studies were not able to show the antioxidant power in <i>Pangasius</i> fish	[12, 26-28]
Present research	In the present study, we worked with the lipids which was extracted from <i>Pangasius</i> . And the result is the lipids also contain small amount of antioxidant power. The maximum inhibition rate is 0.0995 % and the minimum of 0.0686 % by the human body. We also determine the chemical composition of the lipids from <i>Pangasius</i> . The higher percentages of Dodecanoic acid 1,2,3 Propanetriyl ester 86.67 % found in GC-MS. preparations such as makeup, lotions & creams, deodorants, and sunscreen and suntan preparations all include the chemical molecule dodecanoic acid, 1,2,3-propanetriyl ester.	The present study is able to identify the antioxidant power of the studied fish sample. The sample contains a small percentage of 0.33 of L (+) ascorbic acid 2,6 dihexa-decanoate which is an antioxidant agent that reduces free radicals in the human body.	This study

### 5.0 Conclusion

Lipids from *Pangasius Pangasius*, twenty-four (24.0) bioactive chemical compounds were identified by GC-MS. The first major compound is Dodecanoic Acid, 1,2,3-Propanetriyl Ester

86.67 % is a strong antibacterial agent and also a plant metabolite. It is used in soaps and cosmetics. The second major compound is Bis (2-Ethylhexyl) Phthalate 12.37 % which is a plasticizer for resins, in pesticides and as a solvent for ink. The third is L (+) Ascorbic acid 2,6-Dihexa decanoate observed at 0.33 %. Ascorbic acid is helpful for wound healing cells of the human or animal body, to increase the absorption of iron from plant foods and to support the immune system. It works as an antioxidant to protect cells against free radicals. The overall findings of this study could be useful for *Pangasius Pangasius*. Thus, *Pangasius Pangasius* chemical compounds contain little percentage of antioxidants and are more anti-bacterial and pharmaceutical functions. The chemical constituent and antioxidant properties of lipids isolated from the popular freshwater fish named *Pangasius pangasius* are clearly clarified in this study. This outcome highlighted its possibility as an origin of bioactive chemicals and expanded our knowledge of its nutritive and therapeutic properties. This identical investigation is important for scientific researchers because it emphasizes the role of Pangasius lipids in encouraging research on organic antioxidants and the sustainable use of aquatic ecosystems, while also opening up new possibilities for their use in food products and nutraceuticals that improve health.

#### **Disclaimer (Artificial intelligence)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### **Conflict of Interest**

The authors express that no conflict of interest could be influenced in this manuscript.

#### **Data Availability**

The data is available on request.

### **Acknowledgement**

The author's heartiest thanks to Dr. Sharif Md. Al-Reza, Chairman, Department of Applied Chemistry and Chemical Engineering (ACCE), Islamic University, Kushtia-7003, Bangladesh for co-operation this research and availed laboratory facilities. We also expressed thanks to Dr. Shirin Akter Jahan, PSO, BCSIR, Bangladesh for using PC, software and other appliances.

### **References**

- [1] Bechtel, P. J., & Oliveira, A. C. (2006). Chemical characterization of liver lipid and protein from cold-water fish species. *Journal of Food Science*, 71(6), S480-S485.
- [2] Kim, S. K., Kim, Y. T., Byun, H. G., Nam, K. S., Joo, D. S., & Shahidi, F. (2001). Isolation and characterization of antioxidative peptides from gelatin hydrolysate of Alaska pollack skin. *Journal of agricultural and food chemistry*, 49(4), 1984-1989.
- [3] Jun, S. Y., Park, P. J., Jung, W. K., & Kim, S. K. (2004). Purification and characterization of an antioxidative peptide from enzymatic hydrolysate of yellowfin sole (*Limanda aspera*) frame protein. *European Food Research and Technology*, 219, 20-26.
- [4] Islam, M. R., Ahmed, S., Sadia, S. I., Sarkar, A. K., & Alam, M. A. (2023). Comprehensive review of phytochemical content and applications from *cestrum nocturnum*: A Comparative Analysis of Physicochemical Aspects. *Asian Journal of Research in Biochemistry*, 13(4), 43-58.
- [5] Picot, L., Bordenave, S., Didelot, S., Fruitier-Arnaudin, I., Sannier, F., Thorkelsson, G., ... & Piot, J. M. (2006). Antiproliferative activity of fish protein hydrolysates on human breast cancer cell lines. *Process Biochemistry*, 41(5), 1217-1222.
- [6] Bougatef, A., Nedjar-Arroume, N., Ravallec-Plé, R., Leroy, Y., Guillochon, D., Barkia, A., & Nasri, M. (2008). Angiotensin I-converting enzyme (ACE) inhibitory activities of sardinelle (*Sardinella aurita*) by-products protein hydrolysates obtained by treatment with microbial and visceral fish serine proteases. *Food chemistry*, 111(2), 350-356.
- [7] Dong, Y. L., Sheng, G. Y., Fu, J. M., & Wen, K. W. (2005). Chemical characterization and anti-anaemia activity of fish protein hydrolysate from *Saurida elongata*. *Journal of the Science of Food and Agriculture*, 85(12), 2033-2039.
- [8] Baehaki, A., Nopianti, R., & Anggraeni, S. (2015). Antioxidant activity of skin and bone collagen hydrolyzed from striped catfish (*Pangasius pangasius*) with papain enzyme. *J. Chem. Pharm. Res*, 7(11), 131-135.
- [9] Azizah, N., Ochiai, Y., & Nurilmala, M. (2020). Collagen peptides from *Pangasius* fish skin as antioxidants. In *IOP Conference Series: Earth and Environmental Science* (Vol. 404, No. 1, p. 012055). IOP Publishing.
- [10] Ramezani, F., & Khan, M. M. (2024). Effect of *Pangasius Hypophthalmus* Fish Extract on Blood Sugar and Uric Acid Levels in Alloxan-Induced *Rattus Norvegicus*.

- [11] Pradarameswari, K. A., Zaelani, K., Waluyo, E., & Nurdiani, R. (2018, April). The physico-chemical properties of pangas catfish (*Pangasius pangasius*) skin gelatin. In IOP Conference Series: Earth and Environmental Science (Vol. 137, No. 1, p. 012067). IOP Publishing.
- [12] Baehaki, A., Nopianti, R., & Anggraeni, S. (2015). Antioxidant activity of skin and bone collagen hydrolyzed from striped catfish (*Pangasius pangasius*) with papain enzyme. *J. Chem. Pharm. Res.*, 7(11), 131-135.
- [13] Muyonga, J. H., Cole, C. G. B., & Duodu, K. G. (2004). Extraction and physico-chemical characterisation of Nile perch (*Lates niloticus*) skin and bone gelatin. *Food hydrocolloids*, 18(4), 581-592.
- [14] Pradarameswari, K. A., Zaelani, K., Waluyo, E., & Nurdiani, R. (2018, April). The physico-chemical properties of pangas catfish (*Pangasius pangasius*) skin gelatin. In IOP Conference Series: Earth and Environmental Science (Vol. 137, No. 1, p. 012067). IOP Publishing.
- [15] Islam, R., Paul, D. K., Rahman, A., Parvin, T., Islam, D., & Sattar, A. (2012). Comparative characterization of lipids and nutrient contents of *Pangasius pangasius* and *Pangasius sutchi* available in Bangladesh. *J Nutr Food Sci*, 2(2), 1-6.
- [16] Alam, S. M. D., Karim, M. H., Chakraborty, A., Amin, R., & Hasan, S. (2016). Nutritional characterization of the Long-Whiskered catfish *Sperata aor*: A commercially important freshwater fish of Bangladesh. *International Journal of food science and Nutrition Engineering*, 6(1), 1-8.
- [17] Monalisa, K., Islam, M. Z., Khan, T. A., Abdullah, A. T. M., & Hoque, M. M. (2013). Comparative study on nutrient contents of native and hybrid Koi (*Anabas testudineus*) and Pangas (*Pangasius pangasius*, *Pangasius hypophthalmus*) fish in Bangladesh.
- [18] Chakma, S., Rahman, M. A., Siddik, M. A., Hoque, M. S., Islam, S. M., & Vatsos, I. N. (2022). Nutritional Profiling of Wild (*Pangasius pangasius*) and Farmed (*Pangasius hypophthalmus*) *Pangasius* Catfish with Implications to Human Health. *Fishes*, 7(6), 309.
- [19] Karl, H., Lehmann, I., Rehbein, H., & Schubring, R. (2010). Composition and quality attributes of conventionally and organically farmed *Pangasius* filets (*Pangasius hypophthalmus*) on the German market. *International journal of food science & technology*, 45(1), 56-66.
- [20] Orban, E., Nevigato, T., Di Lena, G., Masci, M., Casini, I., Gambelli, L., & Caproni, R. (2008). New trends in the seafood market. *Sutchi* catfish (*Pangasius hypophthalmus*) filets from Vietnam: Nutritional quality and safety aspects. *Food Chemistry*, 110(2), 383-389.
- [21] Kulawik, P., Migdal, W., Tkaczewska, J., Gambus, F., Szczurowska, K., & Özoğul, F. (2016). Nutritional composition of frozen filets from *Pangasius* catfish (*Pangasius hypophthalmus*) and Nile tilapia (*Oreochromis niloticus*) imported to European countries. *Annals of Animal Science*, 16(3).
- [22] Nam, P. V., Van Hoa, N., Anh, T. T. L., & Trung, T. S. (2020). Towards zero-waste recovery of bioactive compounds from catfish (*Pangasius hypophthalmus*) by-products using an enzymatic method. *Waste and Biomass Valorization*, 11, 4195-4206.
- [23] Oktavianawati, I., Andinata, D., Isnaeni, A. N., Hermiastuti, M., Rahmawati, N., Handayani, W., & Winata, I. N. A. (2016). Effects of feeding diets containing *Azolla pinnata* and probiotic on the growth and nutritional content of Patin fish (*Pangasius djambal*). *Agriculture and Agricultural Science Procedia*, 9, 403-410.

- [24] Rahman, M. A., Ullah, M. R., Kabir, M. A., Alam, M. A., Rahman, M., & Hossen, M. F. (2020). Artificial propagation of indigenous yellowtail catfish (*Pangasius pangasius*): experiences and challenges. *Aquaculture*, 523, 735215.
- [25] Sachchu, M. M. H., Hossain, A., Kobir, M. M., Hoda, M. D., Ahamed, M. R., Lima, M. N. J., ... & Alam, M. A. (2024). Heavy Metal Intake by Fishes of Different River Locations in Bangladesh: A Comparative Statistical Review. *Asian Journal of Fisheries and Aquatic Research*, 26(6), 43-67.
- [26] Rahman, M. M., Islam, M. R., Islam, M. R., & Naznin, S. (2018). Extraction and characterization of lipid from Pangus fish (*P. Pangasius*) available in Bangladesh by solvent extraction method. *American Journal of Zoology*, 1(2), 28-34.
- [27] Zilani, M. N. H., Islam, M. A., Biswas, P., Anisuzzman, M., Hossain, H., Shilpi, J. A., ... & Hossain, M. G. (2021). Metabolite profiling, anti-inflammatory, analgesic potentials of edible herb *Colocasia gigantea* and molecular docking study against COX-II enzyme. *Journal of Ethnopharmacology*, 281, 114577.
- [28] Baehaki, A., Nopianti, R., & Anggraeni, S. (2015). Antioxidant activity of skin and bone collagen hydrolyzed from striped catfish (*Pangasius pangasius*) with papain enzyme. *J. Chem. Pharm. Res*, 7(11), 131-135.
- [29] Propionic Acid - an overview | ScienceDirect Topics." Accessed: Mar. 12, 2024. [Online]. Available: <https://www.sciencedirect.com/topics/pharmacology-toxicology-and-pharmaceutical-science/propionic-acid>.
- [30] Benzoic Acid - an overview | ScienceDirect Topics." Accessed: Mar. 12, 2024. [Online]. Available: <https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/benzoic-acid>
- [31] Weaver, J. A., Beverly, B. E., Keshava, N., Mudipalli, A., Arzuaga, X., Cai, C., ... & Yost, E. E. (2020). Hazards of diethyl phthalate (DEP) exposure: A systematic review of animal toxicology studies. *Environment international*, 145, 105848.
- [32] Fig. 2h: L-(+)-Ascorbic acid 2,6-dihexadecanoate," ResearchGate. Accessed: Mar. 12, 2024. [Online]. Available: [https://www.researchgate.net/figure/h-L--Ascorbic-acid-2-6-dihexadecanoate\\_fig9\\_281776251](https://www.researchgate.net/figure/h-L--Ascorbic-acid-2-6-dihexadecanoate_fig9_281776251)
- [33] Green, R., Hauser, R., Calafat, A. M., Weuve, J., Schettler, T., Ringer, S., ... & Hu, H. (2005). Use of di (2-ethylhexyl) phthalate-containing medical products and urinary levels of mono (2-ethylhexyl) phthalate in neonatal intensive care unit infants. *Environmental health perspectives*, 113(9), 1222-1225.
- [34] Sánchez-Moreno, C. (2002). Methods used to evaluate the free radical scavenging activity in foods and biological systems. *Food science and technology international*, 8(3), 121-137.
- [35] Hoque, N. S., Hossain, M. S., Rahaman, M. A., Chowdhury, R. S., Alam, M. A., Hossain, M. I., & Tama, R. T. Molecular Mechanism of Cell Signaling Pathways of Cancer Cells in Forward and Reverse Mutation: A Perspective Review.
- [36] Hasan, M. S., Jahan, R., Alam, M. A., Khatun, M., & Al-Reza, S. M. (2016). Study on physicochemical properties of edible oils available in Bangladeshi local market. *Archives of Current Research International*, 6(1), 1-6.
- [37] Alam, M. A., Munni, S. A., Mostafa, S., Bishwas, R. K., & Jahan, S. A. (2023). An investigation on synthesis of silver nanoparticles. *Asian Journal of Research in Biochemistry*, 12(3), 1-10.

[38] Haque, N. N., Alam, M. A., Roy, C. K., Zenat, M., Akther, E., & Munshi, J. L. (2023). Cyanobacteria mediated CO<sub>2</sub> segregation: A promising alternative method for sustainable bioremediation and biomass production. *Asian Journal of Research in Biochemistry*, 13(3), 28-43.

[39] Ali, M. S., Ahmed, S., Islam, M. R., Ahamed, M. S., Rahaman, M. A., Khatun, M., & Alam, M. A. (2024). Diabetes mellitus control including fruits in diet: Exhaustive review and meta-analysis. *Asian Journal of Food Research and Nutrition*, 3(1), 43-59.

[40] Kobir, M. M., Ali, M. S., Ahmed, S., Sadia, S. I., & Alam, M. A. (2024). Assessment of the physicochemical characteristic of wastewater in Kushtia and Jhenaidah Municipal Areas Bangladesh: A Study of DO, BOD, COD, TDS and MPI. *Asian Journal of Geological Research*, 7(1), 21-30.

[41] Sarkar, A. K., Ahmed, S., Sadia, S. I., Kobir, M. M., Tabassum, S., Islam, M. R., & Alam, M. A. (2024). Overview of the skeleton significance of toothpaste formulation, evaluation and historical perspectives: Insights from Bangladesh's toothpaste industry. *Journal of Materials*

*Science Research and Reviews*, 7(1), 80-101.

[42] Zenat, M., Akther, E., Haque, N. N., Hasan, M. R., Begum, M., Munshi, J. L., ... & Alam, M. A. (2024). Antifungal activity of various plant extracts against aspergillus and penicillium species isolated from Leather-Borne Fungus. *Microbiology Research Journal International*, 34(1), 10-23.

[43] Khatun, M., Kobir, M. M., Miah, M. A. R., Sarkar, A. K., & Alam, M. A. (2024). Technologies for remediation of heavy metals in environment and ecosystem: A critical overview of comparison study. *Asian Journal of Environment & Ecology*, 23(4), 61-80.

[44] Haque, N. N., Alam, M. A., Baidya, A. S., Zenat, E. A., Rahman, M. Z., Roy, C. K., & Munshi, J. L. (2024). Bioremedial capacity of indigenous hydrophytes and microalgae of Bangladesh: A comparative study on their potential in tannery effluent treatment. *Asian Journal of Environment & Ecology*, 23(6), 53-65.

[45] Sachchu, M. M. H., Hossain, A., Kobir, M. M., Hoda, M. D., Ahamed, M. R., Lima, M. N. J., ... & Alam, M. A. (2024). Heavy Metal Intake by Fishes of Different River Locations in Bangladesh: A Comparative Statistical Review. *Asian Journal of Fisheries and Aquatic Research*, 26(6), 43-67.

[46] Haque, N. N., Alam, M. A., Baidya, A. S., Zenat, E. A., Roy, C. K., Hossain, M. K., & Munshi, J. L. (2024). Heavy metal scavenging potential of indigenous microalgae of Bangladesh: A study on its application in textile effluent treatment. *South Asian Journal of Research in Microbiology*, 18(7), 58-75.

[47] Eva, T. N., Nahar, K., Hoda, M. D., Sachchu, M. M. H., Afrin, S., Lima, M. N. J., ... & Alam, M. A. Monitoring of Trace Elements in Groundwater of Municipal Area at Kushtia and Jhenaidah District of Bangladesh.

[48] Khatun, M., Hoque, F., Hoque, N. S., Hossain, M. S., Alam, M. A., Afrin, S., ... & Tama, R. T. (2024). Emerging Role of Probiotics in Advancement of Combating Physical Abnormalities and Diseases: A Systematic Perspective Analysis. *Asian Journal of Biochemistry, Genetics and Molecular Biology*, 16(8), 1-23.

- [49] Tama, R. T., Hossain, M. S., Rahaman, M. A., Alam, M. A., Rahman, M., Parvin, A., ... & Islam, M. S. (2024). Harnessing the Power of Natural Products Against Bacterial Urinary Tract Infections: A Perspective Review for Cultivating Solutions. *Health Sciences Review*, 100199.
- [50] Alam, M. A., Mostafa, S., Bishwas, R. K., Sarkar, D., Tabassum, M., & Jahan, S. A. Low Temperature Synthesis and Characterization of High Crystalline 3c-Ag Nanoparticle. Available at SSRN 4446717.
- [51] Alam, M. A., Mobashsara, M. T., Sabrina, S. M., Bishwas, R. K. B., Debasish, D. S., & Shirin, S. A. J. (2023). One-pot low-temperature synthesis of high crystalline copper nanoparticles. *Malaysian Journal of Science and Advanced Technology*, 122-127.
- [52] Alam, M. A., Tabassum, M., Mostofa, S., Bishwas, R. K., Sarkar, D., & Jahan, S. A. (2023). The effect of precursor concentration on the crystallinity synchronization of synthesized copper nanoparticles. *Journal of Crystal Growth*, 621, 127386.
- [53] Alam, M. A., Bishwas, R. K., Mostofa, S., & Jahan, S. A. (2024). Low-temperature synthesis and crystal growth behavior of nanocrystal anatase-TiO<sub>2</sub>. *Materials Letters*, 354, 135396.
- [54] Tabassum, M., Alam, M. A., Mostofa, S., Bishwas, R. K., Sarkar, D., & Jahan, S. A. (2024). Synthesis and crystallinity integration of copper nanoparticles by reaction medium. *Journal of Crystal Growth*, 626, 127486.
- [55] Kobir, M. M., Tabassum, S., Ahmed, S., Sadia, S. I., & Alam, M. A. (2024). Crystallographic benchmarking on diffraction pattern profiling of Polymorphs-TiO<sub>2</sub> by WPPF for Pigment and Acrylic Paint. *Archives of Current Research International*, 24(1), 62-70.
- [56] Ahamed, M. S., Ali, M. S., Ahmed, S., Sadia, S. I., Islam, M. R., Rahaman, M. A., & Alam, M. A. (2024). Synthesis of silver nanomaterials capping by fruit-mediated extracts and antimicrobial activity: A Critical Review. *International Research Journal of Pure and Applied Chemistry*, 25(1), 45-60.
- [57] Shishir, M. K. H., Sadia, S. I., Ahmed, S., Aidid, A. R., Rana, M. M., Hasan, M. M., ... & Alam, M. A. (2024). Transmission Electron Microscopic and X-ray Diffraction Based Study of Crystallographic Bibliography Demonstrated on Silver, Copper and Titanium Nanocrystals: State of the Art Statical Review. *Asian Journal of Applied Chemistry Research*, 15(3), 1-19.
- [58] Alam, M. A., Bishwas, R. K., Mostofa, S., Sarkar, D., & Jahan, S. A. (2024). X-ray crystallography derived diffraction properties of cuprite crystal as revealed by transmission electron microscopy. *Physics Open*, 20, 100228.
- [59] Alam, M. A., Bishwas, R. K., Mostofa, S., & Jahan, S. A. (2024). Crystallographic phase stability of nanocrystalline polymorphs tio<sub>2</sub> by tailoring hydrolysis pH. *South African Journal of Chemical Engineering*, 49, 73-85.
- [60] Alam, M. A., Bishwas, R. K., Mostofa, S., & Jahan, S. A. (2024). Impact on preferred orientation and crystal strain behavior of nanocrystal anatase-TiO<sub>2</sub> by X-ray diffraction technique. *South African Journal of Chemical Engineering*, 49, 348-352.
- [61] Shishir, M. K. H., Afrin, S., Sachchu, M. M. H., Eva, T. N., Tabassum, S., Ahmed, S., ... & Alam, M. A. Crystalline Copper Nanomaterials for Advanced Ceramic: A Comprehensive Review for Functional Ceramic Coating Approaches.
- [62] Ahmed, S., Shishir, M. K. H., Sadia, S. I., Al-Reza, S. M., Sachchu, M. M. H., Aidid, A. R., ... & Alam, M. A. (2024). Crystallographic phase biographs of copper nanocrystalline material: A statistical perspective review. *Nano-Structures & Nano-Objects*, 39, 101275.

- [63] Sadia, S. I., Shishir, M. K. H., Ahmed, S., Alam, M. A., Al-Reza, S. M., Afrin, S., ... & Jahan, S. A. (2024). Green synthesis of crystalline silver nanoparticle by bio-mediated plant extract: A critical perspective analysis. *Nano-Structures & Nano-Objects*, 39, 101272.
- [64] Al-Mahmud, M. R., Shishir, M. K. H., Ahmed, S., Tabassum, S., Sadia, S. I., Sachchu, M. M. H., ... & Alam, M. A. (2024). Stoichiometry crystallographic phase analysis and crystallinity integration of silver nanoparticles: A Rietveld refinement study. *Journal of Crystal Growth*, 643, 127815.
- [65] Sadia, S. I., Shishir, M. K. H., Ahmed, S., Aidid, A. R., Islam, M. M., Rana, M. M., ... & Alam, M. A. (2024). Crystallographic biography on nanocrystalline phase of polymorphs titanium dioxide (TiO<sub>2</sub>): A perspective static review. *South African Journal of Chemical Engineering*.
- [66] Alam, M. A., Ahmed, S., Bishwas, R. K., Mostofa, S., & Jahan, S. A. (2025). X-ray crystallographic diffraction study by whole powder pattern fitting (WPPF) method: Refinement of crystalline nanostructure polymorphs TiO<sub>2</sub>. *South African Journal of Chemical Engineering*, 51, 68-77.
- [67] Nahar, A., Akbor, M. A., Shawon, M. T. A., Trisha, S. A., Akhtar, U. S., Alam, M. A., ... & Hossain, M. S. (2024). Biotite and ZnO embedded carbon aerogel composite for enhanced photocatalytic activity to degrade ciprofloxacin under sunlight stimulation. *Results in Engineering*, 23, 102524.
- [68] Nahar, A., Hossain, M. S., Akbor, M. A., Dey, S. S., Alam, M. A., Trisha, S. A., ... & Sonia, F. A. (2024). Evaluation of anti-microbial activity of biotite and biotite composite based on crystallographic parameters: Estimation of crystallite size employing X-ray diffraction data. *Results in Engineering*, 23, 102595.
- [69] Hossain, M. I., Rahman, M. M., Ghos, B. C., Gafur, M. A., Alam, M. A., & Rabbi, M. A. (2024). Preparation and characterization of crystalline nanocellulose from keya (*Pandanus tectorius*) L. fiber as potential reinforcement in sustainable bionanocomposite: A waste to wealth scheme. *Carbohydrate Polymer Technologies and Applications*, 8, 100600.
- [70] Rahman, M. M., Hossain, M. I., Ghos, B. C., Gafur, M. A., Alam, M. A., Uddin, M. J., ... & Barmon, J. (2024). Fabrication of CNC-AC bionanosorbents from the residual mass of *Magnolia champaca* l. Bark after methanol extraction for wastewater treatment: Continuous column adsorption study. *Environmental Nanotechnology, Monitoring & Management*, 22, 101015.