

# Impact of Irrigation Regimes and Organic Amendments on Soil Physical Properties, Nutrient Availability, and Productivity in Calcareous Soil

## Abstract

The scarcity of irrigation water in Egypt requires addressing severe water scarcity, such as deficit irrigation and using organic amendments to improve water use efficiency in agriculture. A field experiment was conducted at El-Nubaria Agricultural Research Station in the summer of 2022 for maize and the winter of 2022/2023 for wheat to study the impact of organic amendments and irrigation regimes on calcareous soil properties and productivity. The experiments involved a split plot design with three replicates, irrigation regimes were placed in main plots (50% (I<sub>50</sub>), 75% (I<sub>75</sub>), and 100% (I<sub>100</sub>) of ETC) and organic amendments in subplots included five treatments: 1) 9.6ton ha<sup>-1</sup> (B<sub>1</sub>), 2) 14.4 ton ha<sup>-1</sup> (B<sub>2</sub>) of biochar, 3) potassium humate at the rate of 1.5% (KH<sub>1</sub>), 4) 3% (KH<sub>2</sub>) and 5) compost (12 ton ha<sup>-1</sup>). The obtained results indicated that compost, followed by B<sub>2</sub> (14.4 ton ha<sup>-1</sup>), achieved the highest soil organic matter value. Applied I<sub>100</sub> had the highest soil available NPK and micronutrients, followed by I<sub>75</sub>. Organic amendments significantly affected the soil availability of N, P, K, Fe, Mn and Zn. The maximum potassium content in grains was achieved using I<sub>100</sub>, followed by I<sub>75</sub>. The influence of organic amendments on field capacity and available water took the descending order: B<sub>1</sub>>KH<sub>1</sub>> compost. Water use efficiency was negatively impacted by increasing the amount of applied irrigation water. Applying I<sub>100</sub> resulted in a significant increment in wheat and maize grain yields, followed by I<sub>75</sub>. The influence of organic amendments on wheat grain yield took the descending order: compost > potassium humate > biochar. The effect on maize grain yield took the descending order: potassium humate > compost > biochar.

**Key words:** Irrigation Regime, Compost, Humate, Biochar, Wheat, Maize, Calcareous Soil.

## 27        **1. Introduction**

28            Considering over 80% of Egypt's water sources are used for agriculture, the  
29 availability of water is the greatest obstacle to growing the country's food production. The  
30 increase in irrigated areas, however, is increasing the demand for water sources. Since  
31 irrigation water resources are scarce, great efforts should be made to rationalize the use of  
32 each unit of water. One of the major tactics to save water and improve water use efficiency is  
33 irrigating the crops with a water amount less than the optimal requirement, which refers to  
34 water deficit. However, the application of a deficit water pattern in irrigating crops is  
35 associated with low yields and quality [1]. Physiologically, low water supply, as an abiotic  
36 stress, causes critical issues in cell water integrity, stomata performance, plant pigments, and  
37 photosynthesis, hence a reduction in productivity is anticipated [2]. Therefore, several  
38 practices are required to be applied under water shortage to maintain better plant growth and  
39 productivity [3]. Maize (*Zeamays* L.) is an important food and feed crop that ranks third in  
40 the world, with multiple uses such as human food, animal feed, and industrial raw materials  
41 [4].

42            Wheat (*Triticum aestivum* L.) is a major cereal crop in Egypt, with a planted area of  
43 1.43 million hectares. The total production was 9.3 million tons, with an average of 6.5 tons  
44 per hectare [5]. Another important cereal crop is wheat, which grains contain a range of  
45 protein (6–21%), fats (1.5–2.0%), cellulose (2.0–2.5%), minerals (1.8%), and vitamins  
46 (1.8%), according to [6]. Egypt is the world's largest importer of wheat, providing 40% of the  
47 population's needs, with 10 million tons produced annually from 1.5 million hectares [7].  
48 However, growing wheat in calcareous soils under deficit irrigation resulted in noticeable  
49 decreases in yield and quality. In saline or calcareous soils, crop plants are subjected to  
50 physiological drought, which negatively alters the physiological status and nutrient balance,  
51 suppressing growth. Organic soil amendments enhance soil properties, maintain fertility,

52 increase productivity, preserve the environment, and produce safe organic food [8]. Organic  
53 materials improve the chemical and physical properties of sandy soil by retaining moisture  
54 and recycling nutrients [9]. Organic additions enhance soil quality and increase carbon  
55 storage, potentially aiding in climate change mitigation efforts [10]. Soil amendments, when  
56 properly prepared and applied, can enhance plant productivity, and soil stability [11]. They  
57 cause increased population sizes and proportions of bacteria associated with nutrient cycling  
58 [12].

59 Compost is a fully decomposed, stabilized, homogeneous substance of animal or plant  
60 origin, free from harmful elements that could harm humans, animals, plants, or the  
61 environment [13]. It is a humic-rich amendment that enhances soil properties through the  
62 controlled biological decomposition of organic resources [14, 15]. Compost can enhance the  
63 growth of plants, lower the need for fertilizer, and reduce soil-borne plant  
64 diseases [16]. Composting is an environmentally friendly method of converting organic waste  
65 into organic nutrients [17].

66 Biochar is a porous, high-carbon material produced through pyrolysis in a no-oxygen  
67 environment, which serves as a storage medium for carbon [18, 19] and raises aromatic  
68 structures and essential minerals in clay minerals over time [20]. Biochar is gaining  
69 popularity as a soil conditioner due to its carbon storage, soil fertility enhancement, water  
70 retention, and increased crop yield [21, 22, 23]. Also, the use of biochar in agricultural soils  
71 offers numerous benefits including improved soil health, nutrient retention, efficient transfer,  
72 and climate change mitigation. Its high surface area, porosity, and high levels of COO<sup>-</sup>, OH<sup>-</sup>,  
73 P-O-, and R-OH groups contribute to its benefits [24]. Furthermore, biochar enhances soil  
74 properties, relocates fertilizers, and sequesters carbon, while maintaining stability is crucial  
75 for climate change mitigation [25]. Biochar in alkaline soils was an effective, reasonable, and  
76 environmentally friendly approach [26]. Compost and biochar can be used to improve soil

77 quality and productivity by absorbing nutrients, improving nutrient availability, and  
78 stimulating the microbial. Biochar-blended compost can act as a partial replacement for  
79 mineral fertilizers [27, 28]. Also, Biochar and compost can enhance maize yield and soil  
80 quality, with their combined application potentially enhancing the properties of soil [29, 30].

81 Humic substances (HS) promote plant growth through anatomical and biochemical  
82 changes in the root system via chelation [31]. Humic acid, a plant growth regulator, is broken  
83 down by living microorganisms, allowing plants with strong chemical and biological activity  
84 to easily absorb it [32]. HS comprising 60% of soil organic compounds are crucial for the  
85 agroecosystem and are responsible for various complex chemical reactions [33]. K-humate is  
86 more efficient than K-Si and compost due to its high solubility, smaller dose, and ease of field  
87 application [34]. HS are commonly used in agriculture due to their hormone-like effects,  
88 which promote growth, enhance plant yield, and improve quality [35]. Humates can improve  
89 soil structure and water retention capacity to enhance maize quality [36, 37, 38]. Potassium  
90 humate, 75% N and 100% Reference evapotranspiration ( $ET_0$ ) improved soil chemical  
91 properties, leading to improved wheat yield [39]. Also, foliar applications of potassium  
92 humate can mitigate stress-related damage in soybean by promoting antioxidant growth and  
93 protecting proteins and chlorophyll [40]. The study found that adding 75% NPK, biochar, and  
94 KH to soil improved growth and water efficiency in olive trees under a half-irrigation water  
95 regime [41, 42].

96 The most effective treatment for crop water reduction was 75% full irrigation and 6  
97 tons of biochar, saving 25% of the water needed [43]. Four irrigation treatments had the  
98 longest maturity period, while no irrigation treatments had the shortest [44]. The  
99 physicochemical and fertility properties of sandy soils were enhanced by optimizing water  
100 and nutrients using an 80% water regime with a  $45 \text{ kg ha}^{-1}$  HA rate [45]. An alternate water-  
101 saving method for growing maize is 70%  $ET_C$  [46]. Apply irrigation with a 4200  $\text{m}^2$  fed level

102 and 4 tons of compost to achieve acceptable maize yield and water utilization [47]. This study  
103 aimed to investigate the impact of soil amendments (biochar, potassium humate, and  
104 compost) on soil physical properties, nutrient status, growth, and yield of wheat and maize  
105 cultivated in calcareous soil under varying irrigation regimes

## 106 2. Materials and Methods

### 107 2.1 Field experiments location and design

108 The field experiment was conducted at the EL-Nubaria research farm, located in the  
109 Behaira Governorate in Egypt. The study examined the impact of soil amendments and  
110 irrigation regimes on soil physical properties, nutrient status, and productivity of maize and  
111 wheat during the 2022 summer and 2022/2023 winter seasons, under the Ministry of  
112 Agriculture and Land Reclamation's Agricultural Research Center. The geographical  
113 coordinates of the farm are 30° 90 N, 29° 96 E, with an altitude of 25 m above sea level.

114 The experiment was laid out in a split design with three replicates. The  
115 experimental plot area was 10.5 m<sup>2</sup> (3m x 3.5m). Irrigation regimes were placed in the main  
116 experimental plots, while organic amendments were in the subplots. The treatments were for  
117 irrigation water requirements (100, 75, and 50% of E<sub>Tc</sub>), and organic amendments included  
118 five treatments: 1) biochar 9.6 ton ha<sup>-1</sup>; 2) biochar 14.4 ton ha<sup>-1</sup>; 3) potassium humates 1.5%  
119 (KH<sub>1</sub>); 4) potassium humates 3% (KH<sub>2</sub>) and 5) compost (12 ton ha<sup>-1</sup>).

120 Some physical and chemical characteristics of the surface soil under investigation, as  
121 well as organic amendments, were analyzed before cultivation according to [48], as shown in  
122 Tables 1 and 2.

### 123 2.2 Plant materials and planting

124 Maize hybrid triple white (Giza 310) was sown on June 1<sup>st</sup> and wheat (Misr 1) was  
125 planted on November 18<sup>th</sup> obtained from the Field Crops Institute. Nitrogen fertilizers such as  
126 ammonium sulfate (20.6%) were added to the soil. Phosphorus fertilizer as

127 superphosphate(15.5% P<sub>2</sub>O<sub>5</sub>) and potassium fertilizer as potassium sulfate (48% K<sub>2</sub>O),  
128 micronutrients, weeds, and diseases were added according to the Ministry of Agriculture and  
129 Land Reclamation of Egypt's recommendation. Potassium humate treatments were sprayed to  
130 coat the leaf surface and drenched the soil around plants at a rate of 1440 L/ha. The plants  
131 were sprayed three times at 30, 60 and 90 days after planting.

### 132 **2.3 Plant-water and irrigation water measurements**

133 The E<sub>Tc</sub> value of crops in the Nubaria region was determined using reference  
134 evapotranspiration and crop factor. Table 3 shows the average values of standard  
135 evapotranspiration. The climatic data from the EL-Nubaria Agricultural Research Station,  
136 Ministry of Agriculture and Land Reclamation, Egypt, was used to calculate the E<sub>T</sub> using  
137 CROPWAT 8[49]. The crop evapotranspiration was calculated using the following equation  
138 [50] E<sub>Tc</sub> = E<sub>To</sub> × K<sub>c</sub>, where: E<sub>Tc</sub> is crop evapotranspiration (mm d<sup>-1</sup>), E<sub>To</sub> is the reference  
139 evapotranspiration (mm d<sup>-1</sup>) and K<sub>c</sub>: crop coefficient.

140 Applied irrigation water was calculated according to the following equation: AIW = E<sub>Tc</sub> / E<sub>a</sub>  
141 (1 - LR) where: AIW: applied irrigation water (mm d<sup>-1</sup>), E<sub>Tc</sub> evapotranspiration values (mm d<sup>-1</sup>)  
142 and E<sub>a</sub> is the irrigation application efficiency. LR is the leaching requirement (assuming  
143 that 10% of the calculated applied irrigation water is additionally applied per irrigation during  
144 the growing season for leaching purposes)

145 **Drought Tolerance Efficiency (DTE)**: was calculated using formula [51]:

$$146 \text{ DTE}(\%) = \frac{\text{Yield under stress}}{\text{Yield under non stress}} \times 100$$

147 **Water utilization efficiency (WUE)**: was estimated for each crop in according to [52] as

$$148 \text{ follows: WUE (Kg m}^{-3}\text{)} = \frac{\text{Seed Yield (Kg)}}{\text{Water applied (m}^3\text{)}}$$

### 149 **2.4 Plant and soil sampling and analysis**

150 At harvest time, maize and wheat plants (after 120 and 150 days from planting,  
151 respectively) from each plot were cut and air-dried, and grain yield and straw yield (ton ha<sup>-1</sup>)

152 were estimated. Also, 1000 grain weights for wheat and 100 grains for maize were weighted.  
153 In addition, plant height (cm) and ear weight (gm) of maize were determined as yield  
154 components of maize. Oven-dried plant samples were analyzed for nitrogen (N), phosphorus  
155 (P), potassium (K), iron (Fe), manganese (Mn) and zinc (Zn) as described by [53].

156 Soil samples were analyzed for available N, P, K, Fe, Mn and Zn according to  
157 methods described by [48]. Organic carbon was determined by the modified Walkley-Black  
158 method [54]. Soil hydraulic conductivity (HC) was determined using undisturbed samples  
159 from the cores [55]. Soil moisture retention curves were determined by exposing the  
160 completely saturated samples to constant suction levels of 0.1, 0.33, 1.0 and 15 atmospheres  
161 using a pressure cooker and membrane [56]. The moisture percentages at each suction level  
162 were calculated volumetrically.

## 163 2.5 Statistical Analysis

164 The significant differences among means were tested using the least significant  
165 differences (L.S.D.) [57].

## 166 3. Results and discussions

### 167 3.1 Organic matter

168 Data in Tables 4 and 5 revealed the effect of irrigation treatments and organic  
169 amendments on the soil organic matter (O.M) % in calcareous soil under wheat and maize  
170 crops. It was observed that the maximum value of O.M% was obtained by compost (12 ton ha<sup>-1</sup>),  
171 followed by B<sub>2</sub> (14.4 ton ha<sup>-1</sup>). This increment may be due to the high content of organic  
172 matter in compost as compared to biochar (Table 2). Thus, the soil O.M content resulting  
173 from applying soil amendments can be arranged in the following descending order after  
174 wheat or maize crops: compost > B<sub>2</sub> > KH<sub>2</sub>.

175           These results are consistent with those obtained by [28] who reported that biochar and  
176 compost enhance soil organic matter. Besides, the results achieved by [58] showed that  
177 compost organic amendments markedly increased soil organic matter compared to control  
178 treatments. Additionally, the results achieved by [59], who mentioned that biochar application  
179 markedly intensified the values of soil organic matter in calcareous sandy soil.

### 180 **3.2 Available NPK of calcareous soil**

181 The results in Tables 4 and 5 indicated the effect of irrigation treatments and organic  
182 amendments on the soil availability of NPK. It was observed that applying 100% of Etc (I<sub>100</sub>)  
183 significantly affected the soil available NP, followed by applying 75% of Etc (I<sub>75</sub>). However,  
184 the available K was not significantly affected by irrigation treatments. Applying I<sub>100</sub> increased  
185 soil nitrogen availability possibly due to its effective chelating properties, reducing nutrient  
186 loss [39]. The study revealed that organic amendments significantly impacted the availability  
187 of NPK in calcareous soil, with compost being the most effective, followed by KH<sub>2</sub> and  
188 B<sub>2</sub>. [34] found that compost significantly increased soil nitrogen (104.4%), followed by K-H,  
189 which increased soil available nitrogen by 81.8%. Besides, [60] found that biochar  
190 application significantly increased soil available nitrogen. In this respect, biochar and  
191 compost enhance soil phosphorus [29]. Moreover, the highest available phosphorus value was  
192 found in 100% of irrigation requirements, followed by 80% and 60% WR. [30] achieved  
193 a significant increase in soil macronutrient availability by applying high rates of compost and  
194 biochar to their soil. Concerning the effect of interaction between irrigation regimes and soil  
195 organic amendments on soil macro-nutrient availability, the statistical analysis showed that  
196 there's only interaction between irrigation treatments and soil organic amendments on soil  
197 available N under wheat crop. So, applying compost and I<sub>100</sub> gave the maximum value of soil  
198 available N.

### 199 **3.3 Available micro-nutrients of the calcareous soil**

200 The results obtained in Tables 6 and 7 show that the available Fe, Mn, and Zn  
201 significantly increased as irrigation requirements increased from 50% to 100% of ET<sub>C</sub>.  
202 Consequently, the highest values of available Fe, Mn, and Zn were obtained with I100. This  
203 increase may be attributed to the application of soil organic amendments, which increase the  
204 organic matter content of the soil, improving water retention and plant growth [11, 45]. The  
205 study also found that compost and potassium humate significantly enhance soil micronutrient  
206 availability, with compost providing the highest values of Fe, Mn, and Zn.  
207 Potassium humate enhances soil micronutrient availability by chelating and releasing them  
208 [61]. It was observed that the availability of micronutrients affected by applying soil organic  
209 materials followed this descending order: compost > potassium humate > biochar. Applying  
210 biochar may enhance the soil's capacity to retain nutrients, potentially increasing the  
211 availability of micronutrients [62]. Moreover, biochar's porous structure and large surface  
212 area enhance soil nutrient availability and reduce leaching [13]. Regarding the interaction  
213 between irrigation regimes and soil amendments on the availability of micronutrients in the  
214 soil, highly significant values of available Fe, Mn, and Zn were obtained by applying  
215 compost and I<sub>100</sub> under wheat and maize crops. On the other hand, the least significant values  
216 of soil available Fe, Mn, and Zn were achieved by applying B1 (9.6 ton ha<sup>-1</sup>) and I<sub>50</sub> (50% of  
217 ET<sub>C</sub>).

### 218 **3.4 NPK contents of straw**

219 Data in Tables 4 and 5 revealed that irrigation treatments had a significant effect on  
220 the N and K contents of wheat straw, but there was no significant effect of irrigation regimes  
221 on the P contents of wheat straw. The highest values of N and K in wheat straw were  
222 achieved by applying I<sub>100</sub>, followed by applying I<sub>75</sub> (Table 4). While irrigation regimes  
223 significantly affected the NPK content of maize straw (Table 5), the lowest NPK contents of  
224 maize straw were achieved by applying I<sub>50</sub>. In this respect, organic amendments significantly

225 affected the NP contents of wheat straw. The maximum values of NP were obtained by  
226 compost, followed by KH<sub>2</sub> (Table 4). On the other hand, the NK contents of maize straw were  
227 significantly affected by organic amendments. KH<sub>1</sub> achieved lower values of NK compared  
228 to compost (Table 5). However, the lowest values of NK were achieved by applying 9.6 ton  
229 ha<sup>-1</sup> of biochar (B<sub>1</sub>). These results agree with those reported by [63] who mentioned that  
230 organic materials achieved the highest nitrogen, phosphorus, and potassium contents in wheat  
231 straw compared to the control

### 232 **3.5 Macro-nutrients contents of grains**

233 Data in Tables 4 and 5 reveal that there was no significant effect of irrigation  
234 requirements on the N and P contents of grains. However, the K content of grain is  
235 significantly affected by irrigation regimes. The maximum K content of grain was achieved  
236 by I<sub>100</sub>, followed by I<sub>75</sub>. On the other hand, organic amendments significantly affect the N and  
237 K contents of grain. Although the phosphorus content of grain is not significantly affected by  
238 organic amendments, the highest values of NK contents of grain were under the wheat crop  
239 (Table 4). Additionally, the maximum values of NPK contents of maize grains were achieved  
240 by applying compost, followed by another attained by applying KH. This increase may be  
241 attributed to the high values of available NPK (Table 4). In contrast, the lowest values of  
242 NPK in wheat or maize crops were obtained by applying B<sub>1</sub> in calcareous soil. These results  
243 were in accordance with those obtained by [34], who mentioned that compost significantly  
244 increased the N content of plants cultivated in calcareous soil compared to KH.

245

### 246 **3.6 Micro-nutrients contents of straw**

247 The results in Tables 6 and 7 reveal that the Fe, Mn and Zn contents of straw behaved  
248 similarly to those obtained in wheat and maize grains, as mentioned above. Thus, the  
249 maximum significant increment in Fe, Mn and Zn contents of wheat and maize straw was

250 achieved by applying I<sub>100</sub>. In this respect, applied compost or KH realized a highly significant  
251 increment of Fe, Mn and Zn in wheat and maize crops (Tables 6 and 7). These results were in  
252 agreement with those obtained by [30, 63] who reported that biochar and compost  
253 significantly improved wheat plant straw micronutrient content.

### 254 **3.7 Micro-nutrient contents of grains**

255 Data in Tables 6 and 7 indicated that the Fe, Mn and Zn contents of grains followed  
256 the same trend as soil available Fe, Mn and Zn in the studied calcareous soil. The maximum  
257 increment of Fe, Mn and Zn achieved by I<sub>100</sub>. However, the lowest values of Fe, Mn and Zn  
258 were realized by applying I<sub>50</sub>. Similarly, applied compost attained the highest content of Fe,  
259 Mn and Zn in wheat and maize crops, followed by applying KH<sub>2</sub> (3%). This increment in  
260 micronutrient concentrations of wheat and maize grains may be due to the increase in soil  
261 availability of micronutrients in the studied calcareous soil resulting from the application of  
262 compost or potassium humate as soil conditioners. The results were in accordance with those  
263 obtained by [61, 62, 30]. The interaction between irrigation regimes and organic amendments  
264 on the Fe, Mn and Zn contents of wheat and maize grains, as shown in Tables 6 and 7  
265 indicated a highly significant increase in micronutrients was achieved by applying  
266 compost and I<sub>100</sub> followed by those obtained by applying KH<sub>2</sub> and I<sub>100</sub>. The lowest values of  
267 micro-nutrients contents of wheat and maize grains were realized by applying B<sub>1</sub> and I<sub>50</sub>  
268 (Tables 6 and 7).

269

270

271

### 272 **3.8 Physical properties of calcareous soil.**

273 Data in Table 8 revealed the effect of irrigation regimes on the hydraulic conductivity  
274 (HC) of the studied calcareous soil. The results indicated that the HC (m/day) values  
275 gradually improved as the water requirements increased from I<sub>50</sub> to I<sub>100</sub>. So, the maximum

276 values of HC were obtained by applying I<sub>100</sub>, followed by applying I<sub>75</sub>. This may be attributed  
277 to the increased soil organic matter, which caused an enhancement of soil water retention by  
278 applying I<sub>75</sub> and I<sub>100</sub> (Tables 4 and 5). These results were in agreement with those obtained by  
279 [64]. The obtained results cleared that the effect of organic amendments on HC took the  
280 following descending order: compost>biochar>potassium humate. The beneficial effect  
281 of biochar on soil moisture retention may be due to the porosity of biochar as reported by  
282 [65]. These results were also in accordance with [25, 66].

283 The statistical analysis showed that the interaction of irrigation regimes and soil  
284 organic amendments affected soil HC (m/day) under wheat and maize crops (Table 8). The  
285 highly significant effect of the interaction between irrigation regimes and applied organic  
286 amendments was achieved by applying compost and I<sub>100</sub>, followed by applying B<sub>2</sub> (14.4  
287 ton/ha) and I<sub>100</sub>. However, applying KH<sub>1</sub> (1.5%) and 50% of ETC achieved the least significant  
288 value of HC (m/day) under wheat and maize crops (Table 8). The obtained results in Table 8  
289 revealed the effect of irrigation treatments and organic materials on some soil physical  
290 properties (HC, FC%, WP%, and AW %) under wheat and maize crops. It was found that soil  
291 moisture content at FC% declined gradually with increasing water requirements from 50% to  
292 100% of ETC. Adversely, the soil moisture content of WP% increased with increasing water  
293 requirements, from 50% to 100% of ETC. Consequently, available water decreased gradually  
294 with increasing water requirements, from 50% to 100% of ETC. These results were in  
295 accordance with [43] who mentioned that irrigation deficit using 75% full irrigation and  
296 applying biochar at a rate of 14.4 ton/ha which saved 25% of water requirements.  
297 [45] recorded that utilizing an 80% water regime with 45 Kg ha<sup>-1</sup> HA rate improved the  
298 physicochemical and nutrient status of sandy soils by optimizing water and nutrients. Also,  
299 [46] reported that 70% ETC is an alternative water-saving strategy for maize production.  
300 Regarding the influence of organic amendments on FC% and AW% of the studied calcareous

301 soil, it can be considered in the following descending order: Biochar (B<sub>1</sub>)> potassium humate  
302 (KH<sub>1</sub>) >compost. The increase in soil moisture at FC and AW% resulted from applying  
303 biochar, which may be due to biochar becoming a soil conditioner due to its ability to store  
304 carbon, improve soil fertility, increase water retention, and increase crop yield [21, 22, 23].  
305 Biochar has many advantages for agricultural soils, such as its high chemical and physical  
306 properties, nutrient retention, improved soil health, and the removal of pollutants. It has a  
307 high surface area, porosity, ion exchange, and water-holding capacity [24]. Moreover,  
308 compost and biochar can enhance soil quality and productivity by absorbing nutrients and  
309 improving nutrient cycling and availability [27, 28].

310       Regarding the interaction between irrigation regimes and applied organic amendments  
311 on F.C.% and AW% of the studied calcareous soil, data in Table 8 revealed that at 50% of  
312 ETC, the effect of applied organic amendments on F.C.% and AW% took the following  
313 descending order: KH<sub>1</sub>> B<sub>1</sub>>compost. This increment resulted from the application of  
314 potassium humate, which may have enhanced the physicochemical and fertility properties of  
315 the studied calcareous soil by optimizing water and nutrients [45].Meanwhile, at 75% of ETC,  
316 the effect of applied organic amendments on the available water of the studied calcareous soil  
317 took the following descending order: B<sub>2</sub>> KH<sub>2</sub> >compost (Table 8). However, at 100% ETC,  
318 the obtained results in Table 8 showed that both applied compost (12 ton/ha) and B<sub>1</sub>(9.6  
319 ton/ha) maximized the values of field capacity and available water of the studied calcareous  
320 soil, followed by the first rate of potassium humate.These results were in accordance with the  
321 results obtained by [27] and [28], who reported that compost and biochar can improve soil  
322 quality and productivity by absorbing nutrients.

### 323 **3.9 Drought tolerates efficiency (DTE):**

324       The results in Fig 1 illustrate that increasing applied irrigation water had a positive  
325 effect on the DTE values, which increased from 75.90% at I<sub>50</sub> to 85.0% at I<sub>75</sub>. In

326 addition, increasing the application rate of organic amendments had a positive effect on water  
327 stress in the  $I_{75}$  treatments, where DTE were 88.3, 87.4, 82.3, 85.9, and 82.0% for  $I_{75}B_1$ ,  
328  $I_{75}B_2$ ,  $I_{75}KH_1$ ,  $I_{75}KH_2$ , and  $I_{75}compost$ , respectively. Biochar recorded the highest values, at  
329 88.3% at  $I_{75}B_1$  and 79.0% at  $I_{50}B_1$ . While, the irrigation water regime noted that  $I_{75}$  had high-  
330 value drought tolerance efficiency for wheat in Fig 1. In addition, results in Figure 1 show that  
331 with maize, the highest value of drought tolerance efficiency was 97.3% at  $I_{75}$ , followed by  
332 92.2% at  $I_{50}$  for the irrigation water regime. Soil amendments increased by 98.4, 95.3, 96.4,  
333 97.8, and 97.1% for  $I_{75}B_1$ ,  $I_{75}B_2$ ,  $I_{75}KH_1$ ,  $I_{75}KH_2$ , and  $I_{75}compost$ , respectively. Biochar  
334 exhibited the highest drought tolerance efficiency at 98.4% due to its ability to retain water  
335 and reduce evapotranspiration. These findings align with previous reports [67] that drought  
336 tolerance efficiency is influenced by irrigation treatment and calcium-humate. The treatment  
337 of  $I_{75}$  Ca- $H_{20}$  had the highest drought tolerance efficiency at 96.72%, while the irrigation  
338 water regime had 93.07%. As for the highest drought tolerance efficiency, it was higher in  
339 corn than in wheat under the influence of all treatments (Fig. 1)

### 340 **3.10 Water use efficiency (WUE)**

341 Results in Fig.2 for wheat illustrate that the highest water use efficiency value ( $6.1 \text{ kg/m}^3$ )  
342 was recorded for  $I_{50}$  with compost. Decreasing the amount of applied irrigation water and  
343 increasing soil amendment applications have a highly positive effect on water use efficiency.  
344 Also, increasing the amount of irrigation water used negatively affects the value of the  
345 efficiency of water use, where WUE was better for  $I_{50}$  compared to  $I_{75}$  and  $I_{100}$ , respectively.  
346 The best additions for soil amendments were compost, followed by a high rate of potassium  
347 humate under an irrigation water regime. Also, the data in Fig. 2 for maize illustrates that the  
348 highest value of water use efficiency was  $I_{50}$ , followed by  $I_{75}$ , then  $I_{100}$  of  $ET_C$ . As for soil  
349 amendments, the best additives were compost, followed by potassium humate, and then  
350 biochar under an irrigation water regime. Soil amendments are materials added to soil to

351 improve its physical, chemical, or biological properties. They can have various effects on  
352 water retention, depending on their composition. For example, organic matter such as  
353 compost can improve water retention by increasing soil porosity and enhancing its ability to  
354 hold onto moisture. Irrigation water level at I<sub>50</sub> with compost have significantly impacted  
355 water use efficiency, with a maximum value of 6.1 kg m<sup>-3</sup>. Water use efficiency was higher  
356 with maize than with wheat in all treatments under 50% irrigation water regime (Fig. 2.).  
357 Using compost and increased potassium rates enhances water storage in the root zone, while  
358 potassium humate fertilizer mitigates water stress. Stomata rich in K keep stomata closed,  
359 reducing transpiration rate but allowing plant roots to absorb more water without additional  
360 water.

### 361 **3.11 Wheat and maize yields**

362 Results in Table 9 showed that applying I<sub>100</sub> significantly increased wheat and maize  
363 yields, followed by I<sub>75</sub>. This increase may be attributed to improved soil properties such as  
364 organic matter and macro- and micronutrient availability, as well as increased soil hydraulic  
365 conductivity. These results were obtained by [46], who reported that 70% ETC is an  
366 alternative water-saving strategy for maize production. Concerning the effect of organic  
367 amendments on wheat yield, the data in Table 9 indicates that this effect followed a  
368 descending order: applied compost > potassium humate > biochar. However, the effect of  
369 organic amendments on maize yield as affected by organic amendments followed a  
370 descending order: potassium humate > compost > biochar. The lowest values of wheat and  
371 maize grain yields were achieved by applying the first rate of biochar (Table 9). These  
372 findings align with previous studies by [14] and [15], who suggest that compost as a humic-  
373 rich amendment can enhance soil's chemical, physical, and biological properties. Humic  
374 substances, comprising 60% of soil organic compounds, can be utilized as soil conditioners to  
375 enhance soil structure and water retention capacity, thereby reducing fertilizer usage and

376 enhancing plant growth. Moreover, potassium humate can also improve maize quality [36-  
377 38]. The study revealed that there was no significant difference in wheat grain yield as a  
378 result of the interaction between irrigation regimes and organic materials. However, the  
379 highest maize grain yields were observed when I<sub>100</sub> and compost were applied, followed by  
380 compost and I<sub>75</sub>. The lowest yield was achieved with the first rate of biochar (B<sub>1</sub>) and  
381 I<sub>50</sub>(Table 9).

### 382 **3.12Wheat and maize straw yields**

383 Results in Table 9show that the maximum wheat straw yield was achieved with I<sub>75</sub>  
384 irrigation, followed by I<sub>100</sub>, while maize straw yield was affected in descending order, with  
385 100% irrigation, 50% irrigation, and 75% irrigation resulting in the highest yield.The data in  
386 Table 9 reveals that organic amendments impact wheat straw yield in descending order,  
387 compost and KH<sub>1</sub> above B<sub>2</sub>, and maize straw yield in descending order: compost> potassium  
388 humate> biochar. These results were in accordance with those obtained by [14, 15].

389 Regarding the results in Table 9,the first rate of potassium humate and 75% of Etc  
390 yielded significantly more wheat straw than the second rate of biochar and 75% of ET<sub>C</sub>. The  
391 study found that the highest wheat straw yield was achieved with the second rate of  
392 potassium humate and 50% of ET<sub>C</sub>, but the maximum significant maize straw yield was  
393 achieved with compost and I<sub>100</sub>, followed by potassium humate and I<sub>100</sub>. While the minimum  
394 value of maize straw yield is achieved by applying the first rate of biochar (B<sub>1</sub>) and 50% of  
395 ET<sub>C</sub>.

#### 396 **3.12.1 Weigh 1000 grains of wheat and 100 grains of maize**

397 The obtained results in Table 9 revealed that the effect of irrigation regimes on the  
398 1000 grain weight of wheat and the weight of 100 maize grains caused a highly significant  
399 increment in both the 1000 grain weight of wheat and the weight of 100 maize grains by  
400 increasing irrigation water requirements. The maximum significant increment of both the

401 grain weight of wheat and the weight of 100 maize grains was achieved by applying I100,  
402 followed by applying 75% of ETc. The increase in the grain weight of wheat may be  
403 attributed to increased K, Fe, Mn, and Zn contents in wheat grains (Tables 4 and 6). While the  
404 increase in weight of 100 maize grains may be due to increased N, Fe, Mn, and Zn contents  
405 in maize grains, which may reflect on improving plant growth (Tables 5 and 7). The study  
406 found that increasing irrigation water requirements led to a significant increase in the weight  
407 of wheat and maize grains. Therefore, the application of I<sub>100</sub> and 75% of ETc resulted in the  
408 maximum significant increase in the 1000 grain weight of wheat and 100 maize grains. The  
409 increase in grain weight of wheat may be attributed to increased K, Fe, Mn, and Zn contents  
410 in wheat grains (Tables 4 and 6). While the increase in the weight of 100 maize grains may be  
411 due to increased N, Fe, Mn, and Zn contents in maize grains, which may reflect on improving  
412 plant growth (Tables 5 and 7). Data in Table 9 clarified that the effect of applying organic  
413 amendments on 1000 grain weights of wheat and 100 grain weights of maize grains took the  
414 following descending order: compost > potassium humate > biochar. This increment in  
415 applying compost and potassium humate may be due to an increase in macro- and  
416 micronutrient contents in wheat and maize grains, which resulted in improved plant growth  
417 (Tables 4, 5, 6 and 7). Also, soil amendments can improve soil properties, maintain fertility,  
418 increase productivity, and preserve the environment [8].

419       Regarding the effect of the interaction between irrigation regimes and organic  
420 amendments on the grain weight of wheat, the results obtained in Table 10 revealed that the  
421 highly significant weight of 1000 wheat grains was obtained by applying the second rate of  
422 potassium humate (KH<sub>2</sub>) and I<sub>100</sub>. It was followed by applying compost and I<sub>100</sub>. However, the  
423 lowest weight was achieved by applying the first rate of biochar (B<sub>1</sub>) and 50% of ETc. The  
424 effect of the interaction between irrigation regimes and organic amendments on the weight of  
425 100 maize grains is shown in Table 10. The highly significant value of 100 maize grain

426 weight obtained by compost and I<sub>100</sub> is followed by applying the second rate of potassium  
427 humate and I<sub>100</sub>. However, the least weight of 100 maize grains was achieved by applying B<sub>2</sub>  
428 and 50% of ETc.

### 429 **3.12.2 Plant height of maize**

430 The obtained results in Table 9 revealed that the effect of the irrigation regime on  
431 maize plant height was significantly increased by applying I<sub>75</sub>, followed by applying  
432 I<sub>100</sub>. Concerning the effect of organic amendments on maize plant height, the data in Table  
433 10 showed that a significant increment in plant height was achieved by applying KH<sub>2</sub>,  
434 followed by KH<sub>1</sub>. However, the lowest height was obtained by applying B<sub>1</sub>. These findings  
435 agree with those reported by [36, 29] who, found that combined and single-application  
436 biochar or compost additions increased the plant height of two maize varieties. Higher plant  
437 heights (206.8 cm) were found in plots irrigated five times compared to lower irrigation  
438 regimes.

### 439 **3.12.3 Ear weight of maize**

440 Data in Table 9 indicated that irrigation regimes affected the ear weight of maize. It  
441 was observed that an increase in irrigation water requirements from I<sub>50</sub> to I<sub>100</sub> caused a highly  
442 significant increment in the ear weight of maize. Thus, the maximum value of the ear weight  
443 of maize was achieved by applying I<sub>100</sub>, followed by applying I<sub>75</sub>. This high value of ear  
444 weight in maize may be attributed to improved plant growth under applying I<sub>100</sub>, which  
445 resulted in increasing N, Fe, Mn, and Zn contents in maize grains, as mentioned above  
446 (Tables 5, 7). Concerning the effect of organic application on amendments application on ear  
447 weight of maize, the obtained results in Table 9 revealed that applying compost resulted in  
448 the highest significant increment of ear weight of maize, followed by applying KH<sub>2</sub>.  
449 Whereas, the lowest one achieved B<sub>1</sub>. This increment in ear weight of maize may be due to  
450 improving some soil characteristics and maintaining fertility, resulting in the application of

451 soil amendments that increase the macro- and micronutrient contents of the maize grains  
452 (Tables 5 and 7). These findings agree with those reported by [68] found that reducing water  
453 stress and ensuring moisture during the maize growing season increased ear weight, possibly  
454 due to improved photosynthesis and growth. The study found that the maximum significant  
455 maize ear weight was achieved with compost with I<sub>100</sub>, followed by KH<sub>2</sub> with I<sub>100</sub>, and the  
456 least significant value was achieved with B<sub>1</sub> and 50% Etc (Table 9).

#### 457 **4. Conclusions**

458 In calcareous soil at El-Nubaria Agricultural Research Station, the applied irrigating  
459 water (AIW) of maize was 3677.1 m<sup>3</sup>ha<sup>-1</sup> with 100% ET<sub>C</sub> for the application of compost,  
460 giving the highest yield of 12.84 ton ha<sup>-1</sup> and the highest water use efficiency of 6.1 kg m<sup>-3</sup> of  
461 applied water at 50% ET<sub>C</sub> and compost. Drought tolerance efficiency is affected by irrigation  
462 water regimes and soil amendments. Therefore, I<sub>75</sub>B<sub>1</sub> recorded the highest values, at 98.4%.  
463 Also, the irrigation water regime showed that I<sub>75</sub> had high-value drought tolerance efficiency,  
464 which was 97.3%. While irrigating wheat at 2218 m<sup>3</sup> ha<sup>-1</sup> with 100% ET<sub>C</sub> at the application of  
465 compost gives the highest yield of 8.72 ton ha<sup>-1</sup> and the highest water use efficiency of 6.1  
466 kg m<sup>-3</sup> of applied water were achieved at 50% ET<sub>C</sub> and compost. Drought tolerance efficiency  
467 is affected by irrigation water regimes and irrigation water regime noted that I<sub>75</sub> had high value  
468 drought tolerance efficiency, which was 85.0%. At the end of this study, the recommendation  
469 was to apply irrigation water at 75% of the Etc is saving 25% of the water required for crops  
470 in Egypt. Further research is needed to evaluate the impact of biochar, potassium humate and  
471 compost amendments on enhancing salt and drought resilience in soil and plants. This will  
472 aid in recommending suitable application methods considering soil management and  
473 cropping systems.

#### 474 **Acknowledgments**

475 The authors would like to express their gratitude to the staff of the Nubariya  
476 Agricultural Research Station and the Sandy and Calcareous Soils Research Department for  
477 their invaluable support and outstanding assistance in carrying out this experiment.

#### 478 **Novelty Statement**

479 The study focuses on optimizing calcareous soil conditions using eco-friendly  
480 fertilizers such as compost, K-humate, and biochar under an irrigation water regime, aiming  
481 to differentiate between these fertilizers in terms of crop quality and yield.

#### 482 **Author's Contribution**

483 This study was completed in cooperation between all authors: Dalal H. Sary, Zeinab  
484 M. Abd EL-Rahman and Osama F. EL-Sedfy. Both authors designed the study and designed  
485 the experimentation. The author Dalal H. Sary followed up the field-work, literature survey  
486 and drafted the manuscript. The authors Zeinab M. Abd EL-Rahman and Osama F. EL-Sedfy  
487 managed the laboratory analyses of the study, literature survey and performed the statistical  
488 analysis. The final manuscript was read and approved by all authors.

#### 489 **Compete of interest**

490 The authors declare that they have no conflict of interest.

491

492

#### **References**

493

494

495

496

497

498

499

500

501

502

503

1. E.M.M. Salem, M.K.M. Kenawey, H.S. Saady, and M. Mubarak, "Soil mulching and deficit irrigation effect on sustainability of nutrients availability and uptake, and productivity of maize grown in calcareous soils", *Comm. Soil Sci. Plant Anal*, vol. 52, pp. 1745-1761, 2021. <https://doi.org/10.1080/00103624.2021.1892733>.
2. M.A.A. Ali, M.A. Nasser, A.N. Abdelhamid, I.A.A. Ali, H.S. Saady, and K.M. Hassan, "Melatonin as a key factor for regulating and relieving abiotic stresses in harmony with phytohormones in horticultural plants — a Review", *J Soil Sci Plant Nutr*, vol. 24, pp. 54-73 <https://doi.org/10.1007/s42729-023-01586-9>, 2024.
3. S.M.A. Doklega, H.S. Saady, M.A. El-Sherpiny, A. Abou El-Yazied, H.G. Abd El-Gawad, M.F.M. Ibrahim, M.A.M. Abd El-Hady, M.M.A. Omar, and A.A. Metwally,

- 504 “Rhizospheric addition of hydrogel polymer and zeolite plus glutathione mitigate the  
505 hazard effects of water deficiency on common bean plants through enhancing the  
506 defensive antioxidants” *Gesunde Pflanzen*, vol. 76, pp. 235-249, 2024.
- 507 4. P. Madhavi, V. Sailaja, T.R. Prakash, and S. Hussain, “Characterization of Biochar  
508 and Humic Acid and their Effect on Soil Properties in Maize”. *Int. J. Curr. Microbiol.*  
509 *App. Sci.*, vol. 6, pp. 449-457, 2017.
- 510 5. F. Faostat, “Statistics division. Food and Agriculture Organization of the United  
511 Nations”, 2016.
- 512 6. A.K. Malav, B. Monpara, A Gaur, and S.S. Bhati, “Character association analysis in  
513 yield and yield components in bread wheat (*Triticum aestivum* L.) genotypes”, *J.*  
514 *Plant Develop. Sci.*, vol 9, pp. 77-83, 2017.
- 515 7. S. Asseng, A.M. Kheir, B.T. Kassie, G. Hoogenboom, A.I. Abdelaal, D.Z. Haman  
516 and A.C. Ruane, “Can Egypt become self-sufficient in wheat?” *Environmental*  
517 *Research Letters*, vol. 13, no. 094012, 2018.
- 518 8. K. Singh H.B. Vasava, D. Snoeck, B.S. Das, D. Yinil, D. Field, T. Sanderson, C.  
519 Fidelis, I. Majeed, and N. Panigrahi, “Assessment of cocoa input needs using soil  
520 types and soil spectral analysis”, *Soil Use and Management*, vol. 35, pp. 492-502,  
521 2019.
- 522 9. M. Elshony, I.M. Farid, F. Alkamar, M.H. Abbas, and H. Abbas, “Ameliorating a  
523 sandy soil using biochar and Compost amendments and their implications as slow-  
524 release fertilizers on plant growth”, *Egyptian Journal of Soil Science*, vol. 59, pp.305-  
525 322, 2019.
- 526 10. P. Sisouvanh, V. Trelo-Ges, N.A. Isarangkool, S. Ayutthaya, A. Pierret, N. Nunan, N.  
527 Silvera, K. Xayyathip, and C. Hartmann, “Can organic amendments improve soil  
528 physical characteristics and increase maize performances in contrasting soil water  
529 regimes?”, *Agriculture*, vol. 11, no.132, 2021.
- 530 11. M.W. Kang, M. Yibeltal, Y.H. Kim, S.J. Oh, J.C. Lee, E.E. Kwon, and S.S. Lee,  
531 “Enhancement of soil physical properties and soil water retention with biochar-based  
532 soil amendments”, *Science of the Total Environment*, vol. 836, no.155746, 2022.
- 533 12. L. Hale, D. Curtis, M. Azeem, J. Montgomery, D.E. Crowley, M.E. Mcgiffen Jr,  
534 “Influence of compost and biochar on soil biological properties under turfgrass  
535 supplied deficit irrigation”, *Applied Soil Ecology*, vol. 168, no.104134, 2021.
- 536 13. J. Konz, B. Cohen, and A. van der Merwe,” Assessment of the potential to produce  
537 biochar and its application to South African soils as a mitigation measure”,

- 538 Environmental Affairs Department: Republic of South Africa: Pretoria, South Africa,  
539 2015.
- 540 14. E. Abouhussien, A.M. Elbaalawy, and M. Hamad, “Chemical Properties of Compost  
541 in Relation to Calcareous Soil Properties and Its Productivity of Wheat”, Egyptian  
542 Journal of Soil Science, vol. 59, pp. 85- 97, 2019.
- 543 15. S.M. Mohamed and R.T. Rashad, “Studying some characteristics of sandy soil  
544 amended by water hyacinth, bean straw, and compost”, Egyptian Journal of Soil  
545 Science, vol. 60, pp. 53-65, 2020.
- 546 16. R. Badar, B. Batool, A. Ansari, S. Mustafa, A. Ajmal, and S. Perveen, “Amelioration  
547 of salt affected soils for cowpea growth by application of organic amendments”,  
548 Journal of pharmacognosy and phytochemistry, vol. 3, pp. 87-90, 2015.
- 549 17. J. Barthod, C. Rumpel, and M.F. Dignac, “Composting with additives to improve  
550 organic amendments” A review. *Agronomy for Sustainable Development*, vol. 38, no.  
551 17, 2018.
- 552 18. S. Mansoor, N. Kour, S. Manhas, S. Zahid, O.A. Wani, V. Sharma, L. Wijaya, M.N.  
553 Alyemeni, A.A. Alsahli, and H.A. El-Serehy, “Biochar as a tool for effective  
554 management of drought and heavy metal toxicity”, *Chemosphere*, vol. 271,  
555 no.129458, 2021.
- 556 19. N. Bolan, S.A. Hoang, J. Bei Yuan, S. Gupta, D. Hou, A. Karakoti, S. Joseph, S. Jung,  
557 K.H. Kim, and M. Kirkham, “Multifunctional applications of biochar beyond carbon  
558 storage”, *International Materials Reviews*, vol. 67, pp.150-200, 2022.
- 559 20. F. Jing, Y. Sun, Y. Liu, Z. Wan, J. Chen, and D.C. Tsang, “Interactions between  
560 biochar and clay minerals in changing biochar carbon stability”, *Science of The Total  
561 Environment*, vol. 809, no.151124, 2022.
- 562 21. M. Rodrigues, T.G. Garmus, M. Arrobas, A. Gongalves, E. Silva, L. Rocha, L. Pinto,  
563 C. Brito, B. Martins, and T.D.O. Vargas, “Combined biochar and organic waste have  
564 little effect on chemical soil properties and plant growth”, *Spanish Journal of Soil  
565 Science*, vol. 9, pp. 199-211, 2019.
- 566 22. Y. Xing, J. Wang, S.M. Shaheen, X. Feng, Z. Chen, F.I. Zhang, and J. Rinklebe,  
567 “Mitigation of mercury accumulation in rice using rice hull-derived biochar as soil  
568 amendment”: A field investigation. *Journal of Hazardous Materials*, vol. 388,  
569 no.121747, 2020.

- 570 23. H. Huang, N.G. Reddy, X. Huang, P. Chen, P. Wang, Y. Zhang, Y. Huang, P. Lin, and  
571 A. Garg, “Effects of pyrolysis temperature, feedstock type and compaction on water  
572 retention of biochar amended soil”, *Scientific Reports*, vol. 11, no 7419, 2021.
- 573 24. F.U. Haider, J.A. Coulter, C. Liqun, S. Hussain, S.A. Cheema, W. Jun, and R.  
574 Zhang, “An overview on biochar production, its implications, and mechanisms of  
575 biochar-induced amelioration of soil and plant characteristics”. *Pedosphere*, vol. 32,  
576 pp. 107-130, 2022.
- 577 25. N.K. Yadav, V. Kumar, K. Sharma, R.S. Choudhary, T.S. Butter, G. Singh, M. Kumar,  
578 and R. Kumar, “Biochar and their impacts on soil properties and crop productivity: a  
579 review”, *Journal of Pharmacognosy and Phytochemistry*, vol. 7, pp. 49-54, 2018.
- 580 26. T.M. Salem, K.M. Refaie, A.E.-H.E.-G.Abd, E-L. Sherif, M.A.M. Eid, “Biochar  
581 application in alkaline soil and its effect on soil and plant”, *Acta  
582 AgricultureSlovenica*, vol. 114, pp. 85-96, 2019.
- 583 27. M.A. Sánchez-Monedero, M.L. Cayuela, M. Sanchez-Garcia, B. Vandecasteele, T.  
584 D'Hose, G. Lopez, C. Martinez-Gaitan, P.J. Kuikman, T. Sinicco, and C Mondini,  
585 “Agronomic evaluation of biochar, Compost and biochar-blended Compost across  
586 different cropping systems”: Perspective from the European project FERTIPLUS.  
587 *Agronomy*, vol. 9, pp. 225, 2019.
- 588 28. E. Mahmoud, M. Ibrahim, N. AN, H. AN, “Effect of biochar and Compost  
589 amendments on soil biochemical properties and dry weight of canola plant grown in  
590 soil contaminated with heavy metals”, *Communications in Soil Science and Plant  
591 Analysis*, vol. 51, pp. 1561-1571, 2020.
- 592 29. A.K. Mensah and K.A. Frimpong, “ Biochar and/or Compost applications improve soil  
593 properties, growth, and yield of maize grown in acidic rainforest and coastal savannah  
594 soils in Ghana”, *International / Journal of Agronomy*, vol. 7, pp. 1-8, 2018.
- 595 30. A. Mancy and M. Sheta, “Evaluation of biochar and Compost ability to improve soil  
596 moisture content and nutrients retention”, *Al-Azhar Journal of Agricultural  
597 Research*, vol. 46, pp.153-165, 2021.
- 598 31. K. Jindo, F.L. Olivares, D.J.D.P. Malcher, M.A. Sánchez-Monedero, C. Kempenaar,  
599 and L.P. Canellas, “From lab to field: Role of humic substances under open-field and  
600 greenhouse conditions as biostimulant and biocontrol agent”, *Frontiers in plant  
601 science*, vol.11, no. 426, 2020.

- 602 32. M. Justi, E.G. Morais, and C.A. Silva, "Fulvic acid in foliar spray is more effective  
603 than humic acid via soil in improving coffee seedlings growth", *Archives of*  
604 *Agronomy and Soil Science*, vol. 65, no. 14, pp. 1969-1983, 2019.
- 605 33. J. Gerke, "Concepts and misconceptions of humic substances as the stable part of soil  
606 organic matter": A review. *Agronomy*, vol 8, no.76, 2018.
- 607 34. D.H. Sary and R.T. Rashad, "Nutrient use Efficiency in Calcareous Soil Amended by  
608 the Silicate, Humate and Compost", *Asian Journal of Soil Science and Plant*  
609 *Nutrition*, vol. 7, pp. 1-12, 2021.
- 610 35. Y. Ouni, T. Ghnaya, F. Montemurro, C. Abdelly, and A. Lakhdar, "The role of humic  
611 substances in mitigating the harmful effects of soil salinity and improve plant  
612 productivity", *International Journal of Plant Production*, vol. 8, pp. 353-374, 2014.
- 613 36. M.I Khan and A. Jan, "Impact of Soil Conditioning and Irrigation Regimes on the  
614 Performance of MaizeCrop", *Sarhad Journal of Agriculture*, vol 34, pp.173-187, 2018.
- 615 37. D. Karčauskienė, R. Repšienė, D. Ambrazaitienė, I. Mockevičienė, G. Šiaudinis, and  
616 R. Skuodienė, "A complex assessment of mineral fertilizers with humic substances in  
617 an agroecosystem of acid soil", *Zemdirbyste-Agriculture*, vol. 106, no. 4, pp. 307-314  
618 2019.
- 619 38. R.T. Rashad, "Effect of Soaking Grains in Some Growth Regulators on Wheat Grown  
620 in Sandy Soil", *Egyptian Journal of Soil Science*, vol. 60, pp. 99-108, 2020.
- 621 39. W. Hassan, W. El-Farghal, F. Khalil, and W. EL-Etr, "Effect of some soil amendments  
622 and irrigation treatments on wheat crop productivity in middle Egypt", *Journal of Soil*  
623 *Sciences and Agricultural Engineering*, vol. 8, pp. 553-563, 2017.
- 624 40. S.S. El-Nwehy, D.H. Sary, and R.R.M. Afify, "Effect of potassium humate foliar  
625 application on yield and quality of soybean (*Glycine max L.*) grown on calcareous  
626 soil under irrigation water regime", *Plant Arch*, vol 20, 1495-1502, 2020.
- 627 41. D.Sary and I. Elsokkary, "Effect of irrigation water regime in presence of organic or  
628 biological fertilizer on olive trees", *Egyptian journal of soil science*, vol. 59, pp. 67-  
629 48, 2019.
- 630 42. R.H.Attia and D.H. Sary, "Effect of some soil conditioners on snap bean growth yield  
631 and sandy soil properties", *Journal of Soil Sciences and Agricultural Engineering*, vol.  
632 12, pp. 123-129, 2021.
- 633 43. R.Abdelraouf, E. Essay, M. Saleh, "Sustainable management of deficit irrigation in  
634 sandy soils by producing biochar and adding it as a soil amendment". *Middle East*  
635 *Journal of Agriculture Research*, vol. 6, pp. 1359-1375, 2017.

- 636 44. S. Islam, M. Haque, M. Hasan, A. Khan, and U. Shanta, “Effect of different  
637 irrigation levels on the performance of wheat”, *Progressive Agriculture*, vol. 29, pp.  
638 99-106, 2018.
- 639 45. A.R. Al-Shareef, S.M. Ismail, and F.S. El-Nakhlawy, “Effect of humic acid  
640 application rates on physicochemical and fertility properties of sandy loam soil grown  
641 with mung bean under different irrigation water regimes”, *Current Science*, vol. 115,  
642 pp.1374-1379,2018.
- 643 46. M. Singh, S. Singh, S. Deb, and G. Ritchie, “Root distribution, soil water depletion,  
644 and water productivity of sweet maize under deficit irrigation and biochar  
645 application”, *Agricultural Water Management*, vol 279, no. 108192, 2023.
- 646 47. K.A. Khatab, K.M. Abd El-Latif, E. Osman, and S. Abdou, “Maize productivity and  
647 crop-water relations as affected by irrigation levels and compost rates”, *Journal of*  
648 *Soil Sciences and Agricultural Engineering*, vol. 6, pp.1545-1562, 2015.
- 649 48. A. Page, R. Miller, D. Keeney, D. Baker, J. Roseoc Ellis, and J. Rhodes, “Methods of  
650 soil analysis”, Part 2: chemical and Microbiological Properties, *Agronomy*  
651 *Monograph No. 9. American Society of Agronomy and Soil Science Society America*  
652 *Madison, Wisconsin, USA, 1982.*
- 653 49. J. Doorenbos and A. Kassam, “Yield Response to Water; FAO Irrigation and  
654 Drainage”, Paper No. 33; Food and Agriculture Organization of the United Nations.  
655 *Water Resources and Development Service: Rome, Italy,1986.*
- 656 50. R.G. Allen, L.S. Pereira, D. Raes, and M. Smith, “Crop evapotranspiration –  
657 Guidelines for computing crop water requirements”, *FAO Irrigation and Drainage*  
658 *Paper, No. 56, FAO, Rome, 1998.*
- 659 51. K. Fisher and G. Wood, “Breeding and selection for drought tolerance in tropical  
660 maize”, In: *Proc. Symp. On Principles and Methods in Crop Imprpt. For Drought*  
661 *Resist. Yith Emphasis, 1981.*
- 662 52. A. Iskandar and M. David, “Spatial and temporal variability of water productivity in  
663 the Syr Darya Basin, central Asia”, *Water Resources Research*, vol. 40, 2004.
- 664 53. G. Estefan, R. Sommer, and J. Ryan, “Methods of soil, plant, and water analysis: A  
665 manual for West Asia and North Africa region Beirut”, *Lebanon ICARDA,*  
666 *International Center for Agricultural Research in the Dry Areas 30, 2013.*
- 667 54. D.W. Nelson and L.E. Sommers,” Total carbon, organic carbon, and organic matter.  
668 *Methods of soil analysis”*: Part 3 Chemical methods, vol. 5, pp. 961-1010, 1996.

- 669 55. A. Klute, C. Dirksen, "Hydraulic conductivity and diffusivity: Laboratory methods.  
670 Methods of soil analysis": Part 1 physical and mineralogical methods, vol. 5, pp. 687-  
671 734, 1986.
- 672 56. W.P. Stakman, "Determination of soil waterretention Curves", Sandbox method 11.  
673 Pressuremembrane methods" ICW. Wageningen, theNetherlands, 1966.
- 674 57. G.W. Snedecorand W.G. Cochran,"Statistical Methods" 8th Ed. Iowa State Univ.,  
675 Press, Ames, Iowa, USA, 1990.
- 676 58. D.A. El-Ngar, H.A. Zein El-Abdeen, D.H. Sary, "Effect of Compost and Nano-  
677 Hydroxyapatite on Phosphorus Nutrition and some Properties of Calcareous Soil",  
678 Journal of Soil Sciences and Agricultural Engineering, vol 13, pp.317-324, 2022.
- 679 59. A.E.A.Z.Amin and M.A. Eissa, "Biochar effects on nitrogen and phosphorus use  
680 efficiencies of zucchini plants grown in a calcareous sandy soil", Journal of Soil  
681 Science and plant nutrition, vol. 17, pp. 912-921,2017.
- 682 60. W. Yang, G. Feng, Y. Jia, Y. Yang, X. Gao, L. Gao, and Z. Qu, "Impact of single  
683 biochar application on maize growth and water-fertilizer productivity under different  
684 irrigation regimes", Frontiers in Plant Science, vol 13, 2022.
- 685 61. M. Awwad, K. El-Hedek, M. Bayoumi, and T. Eid, "Effect of potassium humate  
686 application and irrigation water levels on maize yield, crop water productivity and  
687 some soil properties", Journal of Soil Sciences and Agricultural Engineering, vol. 6,  
688 pp. 461-482,2015.
- 689 62. I.C.B.D. Silva, J.J.N. Basilio, L.A. Fernandes, F. Colen, R.A. Sampaio, and L.A.  
690 Frazao, "Biochar from different residues on soil properties and common bean  
691 production", Scientia Agricola, vol. 74, pp. 378-382, 2017.
- 692 63. D.H. Sary and E.N. Hamed, "Effect of Salicylic, Humic and Fulvic Acids Application  
693 on The Growth,Productivity, and Elements Contents of Two Wheat Varieties Grown  
694 Under Salt Stress", Journal of Soil Sciences and Agricultural Engineering, Vol. 12, PP  
695 657 -671, 2021.
- 696 64. P. Sharma, V. Abrol, V. Sharma, S. Chaddha, C.S. Rao, A. Ganie, D.I. Hefft, M.A. El-  
697 Sheikh, and S. Mansoor, "Effectiveness of biochar and Compost on improving soil  
698 hydro-physical properties, crop yield and monetary returns in inceptisol subtropics",  
699 Saudi Journal of Biological Sciences, vol. 28, pp. 7539-7549, 2021.
- 700 65. S. De Jesus Duarte, B. Glaser, and C.E. Pellegrino Cerri, "Effect of biochar particle  
701 size on physical, hydrological and chemical properties of loamy and sandy tropical  
702 soils", Agronomy, vol. 9, no.165, 2019.

- 703 66. M. Ibrahim, E. Mahmoud, L. Gad, and A. Khader, “Effects of biochar and phosphorus  
704 fertilizer rates on soil physical properties and wheat yield on clay textured soil in  
705 middle Nile Delta of Egypt”, *Communications in Soil Science and Plant Analysis*,  
706 vol. 50, pp. 2756-2766, 2019.
- 707 67. D. H. Sary, E.M. Niel, A. Abd El-All, “Effect of Drip Irrigation Water Regime and  
708 Calcium Humate on the Growth Performance, Yield of Faba Bean (*Vicia Faba L.*)  
709 Grown in Sandy Soil”, *American-Eurasian Journal of Scientific Research*, vol. 16, pp.  
710 53-66, 2021.
- 711 68. O.I. Al-Dulaimi, M.H. Al-Ani, A.R. Al-Rawi, S.E. Seadh, “Effect of water stress and  
712 organic fertilization sources on maize growth and yield”, *Int. J. Agricult. Stat. Sci*,  
713 2020.
- 714
- 715

**Table 1. Physical and chemical properties in soil study**

Sand (%)		Silt (%)	Clay (%)	Texture		SP %	O.M (%)	CaCO <sub>3</sub> (%)		
53.5		40.0	6.5	Sandy loam		48	0.75	33.65		
pH (1:2.5)	EC (dS/m) in soil past	Soluble Cations (meq l <sup>-1</sup> )				Soluble Anions (meq l <sup>-1</sup> )				
		Ca <sup>+2</sup>	Mg <sup>+2</sup>	Na <sup>+</sup>	K <sup>+</sup>	HCO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>-</sup>		
7.96	2.4	7.5	4.50	11.45	0.25	1.5	13.5	8.97		
Available Macronutrients (mg kg <sup>-1</sup> )						Available Micronutrients (mg kg <sup>-1</sup> )				
N	P	K				Fe	Mn	Zn		
17.8	10.12	224.38				3.14	1.89	1.37		

**Table 2: Analysis of organic amendments used in the experiment.**

Amendments	Bulk density (kg/m <sup>3</sup> )	Moisture Contents (%)	pH	EC dS/m	O.C %	O.M %	C/N Ratio	Total (%)		
								N	P	K
Biochar	1060	3	7.2	2.7	0.62	1.07	1.19	0.53	0.001	0.38
Compost	580	11	6.6	2.4	26.5	45.7	33.5	0.79	0.350	0.48
k-humate	ND	ND	8.2	10	10.7	18.4	15.89	0.67	0.220	4.5

ND: not determined

**Table 3: The reference evapotranspiration averages for months during the growing seasons at the experimental site with the applied irrigation water (AIW m<sup>3</sup>/ha).**

Planting Seasons	ET <sub>o</sub> mm/day					AIW m <sup>3</sup> /ha		
						100% ET <sub>c</sub>	75% ET <sub>c</sub>	50%ET <sub>c</sub>
<b>Maize 2022</b>	<b>June</b>	<b>July</b>	<b>August</b>	<b>September</b>	<b>October</b>			
	5.9	5.5	4.4	2.93	1.95	3677.1	2757.8	1838.5
<b>Wheat 2022/2023</b>	<b>December</b>	<b>January</b>	<b>February</b>	<b>March</b>	<b>April</b>	<b>100% ET<sub>c</sub></b>	<b>75% ET<sub>c</sub></b>	<b>50% ET<sub>c</sub></b>
	1.05	1.2	1.5	2.6	3.67	2218	1663.5	1109

**Table 4: Effect of irrigation water regime and organic amendments on the organic matter, availability of NPK and their content in wheat under calcareous soil.**

Irrigation water regime	Organic amendments	Soil				Straw			Grain		
		OM (%)	N (mg kg <sup>-1</sup> )	p (mg kg <sup>-1</sup> )	K (mg kg <sup>-1</sup> )	N (%)	P (%)	K (%)	N (%)	P%	K%
<b>I<sub>50</sub></b>	<b>B<sub>1</sub></b>	1.92	18.85	10.27	157.55	0.60	0.08	0.98	1.14	0.27	0.98
	<b>B<sub>2</sub></b>	2.01	21.61	10.74	161.54	0.73	0.10	1.03	1.26	0.31	1.12
	<b>KH<sub>1</sub></b>	1.74	23.72	11.14	169.52	0.90	0.11	1.12	1.54	0.37	1.13
	<b>KH<sub>2</sub></b>	1.82	24.79	12.76	171.51	0.93	0.14	1.16	1.66	0.39	1.15
	<b>Compost</b>	2.17	26.95	13.25	187.46	0.98	0.16	1.23	1.72	0.43	1.18
	<b>Mean</b>	<b>1.93</b>	<b>23.18</b>	<b>11.63</b>	<b>169.52</b>	<b>0.83</b>	<b>0.12</b>	<b>1.10</b>	<b>1.46</b>	<b>0.35</b>	<b>1.11</b>
	<b>I<sub>75</sub></b>	<b>B<sub>1</sub></b>	1.95	21.80	11.59	155.56	0.81	0.14	1.09	1.26	0.29
<b>B<sub>2</sub></b>		1.97	24.92	12.84	171.51	0.82	0.16	1.12	1.33	0.34	1.43
<b>KH<sub>1</sub></b>		1.84	25.93	13.77	191.50	0.88	0.19	1.15	1.47	0.42	1.45
<b>KH<sub>2</sub></b>		1.89	26.46	14.20	195.44	0.92	0.20	1.19	1.82	0.45	1.54
<b>Compost</b>		2.09	27.53	15.05	205.41	0.95	0.20	1.26	1.84	0.47	1.66
<b>Mean</b>		<b>1.95</b>	<b>25.33</b>	<b>13.62</b>	<b>183.88</b>	<b>0.88</b>	<b>0.18</b>	<b>1.16</b>	<b>1.54</b>	<b>0.39</b>	<b>1.47</b>
<b>I<sub>100</sub></b>	<b>B<sub>1</sub></b>	2.13	23.91	15.40	159.52	0.96	0.14	1.14	1.12	0.30	1.67
	<b>B<sub>2</sub></b>	2.35	27.08	16.10	176.33	1.01	0.16	1.17	1.40	0.33	1.68
	<b>KH<sub>1</sub></b>	1.96	29.11	14.20	195.44	1.02	0.18	1.20	1.55	0.44	1.71
	<b>KH<sub>2</sub></b>	2.21	31.10	15.60	205.42	1.04	0.22	1.23	1.70	0.48	1.75
	<b>Compost</b>	2.67	33.81	16.88	216.72	1.18	0.23	1.28	2.18	0.51	1.77
	<b>Mean</b>	<b>2.26</b>	<b>29.01</b>	<b>15.64</b>	<b>190.69</b>	<b>1.04</b>	<b>0.19</b>	<b>1.20</b>	<b>1.59</b>	<b>0.41</b>	<b>1.71</b>
<b>LSD at 0.05</b>	<b>Irrigation</b>	ns	0.97	2.81	ns	0.11	ns	0.07	ns	ns	0.21
	<b>Organic</b>	0.20	0.52	1.13	10.7	0.11	0.04	ns	0.12	ns	0.13
	<b>Interaction</b>	ns	***	ns	ns	ns	ns	ns	ns	ns	ns
<b>Mean of Organic</b>	<b>B<sub>1</sub></b>	1.85	21.52	12.64	157.54	0.79	0.12	1.07	1.17	0.29	1.31

<b>amendments</b>	<b>B<sub>2</sub></b>	1.97	24.25	13.08	169.79	0.85	0.14	1.10	1.33	0.33	1.41
	<b>KH<sub>1</sub></b>	2.00	20.25	13.23	185.49	0.93	0.16	156	1.52	0.41	1.43
	<b>KH<sub>2</sub></b>	2.11	27.45	14.19	190.79	0.96	0.19	1.19	1.72	0.44	1.48
	<b>Compost</b>	2.31	29.43	15.06	203.20	1.04	0.20	1.26	1.91	0.47	1.54

Note: \*\* and \*\*\* refer to significance level of 5% ( $p \leq 0.05$ ) and 1% ( $p \leq 0.01$ ). ns:no-significant.

**Table 5: Effect of irrigation water regime and organic amendments on the organic matter, availability of NPK and their content in maize under calcareous soil.**

Irrigation water regime	Organic amendments	Soil				Straw			Grain		
		OM (%)	N (mg kg <sup>-1</sup> )	p (mg kg <sup>-1</sup> )	K (mg kg <sup>-1</sup> )	N (%)	P (%)	K (%)	N (%)	OM (%)	N (mg kg <sup>-1</sup> )
<b>I<sub>50</sub></b>	<b>B<sub>1</sub></b>	2.24	22.66	11.40	162.11	1.12	0.12	1.04	1.21	0.32	1.51
	<b>B<sub>2</sub></b>	2.39	24.50	12.54	171.65	1.32	0.14	1.05	1.30	0.36	1.64
	<b>KH<sub>1</sub></b>	2.08	26.80	13.49	194.69	1.49	0.17	1.24	1.40	0.43	1.71
	<b>KH<sub>2</sub></b>	2.13	29.77	14.13	209.72	1.55	0.18	1.30	1.59	0.52	1.84
	<b>Compost</b>	2.69	33.70	15.19	223.93	1.74	0.21	1.43	1.68	0.58	1.94
	<b>Mean</b>	<b>2.31</b>	<b>27.49</b>	<b>13.35</b>	<b>192.42</b>	<b>1.44</b>	<b>0.16</b>	<b>1.21</b>	<b>1.44</b>	<b>0.44</b>	<b>1.71</b>
<b>I<sub>75</sub></b>	<b>B<sub>1</sub></b>	2.29	24.50	11.61	183.77	1.18	0.19	1.15	1.33	0.32	1.65
	<b>B<sub>2</sub></b>	2.45	30.71	12.27	202.40	1.36	0.20	1.16	1.41	0.44	1.74
	<b>KH<sub>1</sub></b>	2.15	31.86	13.69	214.03	1.60	0.21	1.34	1.56	0.48	1.82
	<b>KH<sub>2</sub></b>	2.24	33.70	13.95	225.67	1.78	0.22	1.36	1.31	0.53	1.90
	<b>Compost</b>	2.79	35.42	16.12	232.63	2.05	0.25	1.48	1.77	0.66	2.14
	<b>Mean</b>	<b>2.38</b>	<b>32.14</b>	<b>13.53</b>	<b>211.50</b>	<b>1.60</b>	<b>0.21</b>	<b>1.30</b>	<b>1.48</b>	<b>0.49</b>	<b>1.85</b>
<b>I<sub>100</sub></b>	<b>B<sub>1</sub></b>	2.52	28.55	13.25	199.20	1.53	0.23	1.38	2.17	0.35	1.73
	<b>B<sub>2</sub></b>	2.72	30.66	14.46	207.03	1.59	0.28	1.41	2.54	0.45	1.80
	<b>KH<sub>1</sub></b>	2.36	34.31	14.94	226.00	1.79	0.25	1.46	2.84	0.56	1.97
	<b>KH<sub>2</sub></b>	2.40	36.23	15.87	245.41	1.86	0.31	1.82	2.74	0.64	2.15
	<b>Compost</b>	3.31	39.95	16.64	260.62	2.59	0.34	1.90	2.97	0.73	2.27
	<b>Mean</b>	<b>2.66</b>	<b>35.04</b>	<b>15.03</b>	<b>227.65</b>	<b>1.87</b>	<b>0.28</b>	<b>1.59</b>	<b>2.65</b>	<b>0.54</b>	<b>1.98</b>
<b>LSD at 0.05</b>	<b>Irrigation</b>	ns	ns	0.85	17.46	0.15	0.03	0.18	0.11	ns	ns
	<b>Organic</b>	0.25	9.28	0.89	10.61	0.24	ns	0.22	0.12	0.07	0.19

	<b>Interaction</b>	ns	ns	ns	ns	ns	ns	ns	**	ns	ns
<b>Mean of Organic amendments</b>	<b>B<sub>1</sub></b>	2.20	25.42	12.09	181.69	1.28	0.18	1.19	1.57	0.33	1.63
	<b>B<sub>2</sub></b>	2.26	25.62	13.09	193.36	1.43	0.21	1.21	1.75	0.42	1.73
	<b>KH<sub>1</sub></b>	2.35	33.23	14.04	211.57	1.63	0.20	1.35	1.88	0.49	1.83
	<b>KH<sub>2</sub></b>	2.52	36.02	14.65	226.93	1.73	0.24	1.49	1.93	0.56	1.96
	<b>Compost</b>	2.92	37.66	15.98	239.06	2.13	0.27	1.60	2.14	0.66	2.12

Note: \*\* and \*\*\* refer to significance level of 5% ( $p \leq 0.05$ ) and 1% ( $p \leq 0.01$ ). ns:no-significant.

**Table6: Effect of irrigation water regime and organic amendments on the availability of Fe,Zn, Mn and their contentin wheat under calcareous soil.**

Irrigation water regime	Organic amendments	Soil (mg kg <sup>-1</sup> )			Straw (mg kg <sup>-1</sup> )			Grain (mg kg <sup>-1</sup> )		
		Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn	Zn
<b>I<sub>50</sub></b>	<b>B<sub>1</sub></b>	3.22	2.21	1.12	10.60	11.00	10.15	32.40	15.60	15.82
	<b>B<sub>2</sub></b>	3.54	2.30	1.15	14.60	15.16	14.42	33.60	19.17	16.00
	<b>KH<sub>1</sub></b>	3.58	2.38	1.33	15.70	16.37	14.65	34.00	23.80	17.68
	<b>KH<sub>2</sub></b>	3.69	2.47	1.37	16.50	16.92	15.40	35.00	36.56	19.45
	<b>Compost</b>	4.18	2.53	1.52	18.30	18.48	17.80	47.40	36.97	25.20
	<b>Mean</b>	<b>3.64</b>	<b>2.38</b>	<b>1.30</b>	<b>15.14</b>	<b>15.59</b>	<b>14.48</b>	<b>36.48</b>	<b>26.39</b>	<b>18.83</b>
	<b>I<sub>75</sub></b>	<b>B<sub>1</sub></b>	3.26	2.57	1.39	11.20	13.24	13.31	35.60	19.42
<b>B<sub>2</sub></b>		3.54	2.62	1.44	12.90	15.61	15.86	39.60	22.30	16.80
<b>KH<sub>1</sub></b>		3.73	2.66	1.57	16.40	19.34	16.60	41.20	24.35	17.63
<b>KH<sub>2</sub></b>		3.88	2.77	1.95	17.40	21.40	19.26	43.40	29.14	21.48
<b>Compost</b>		4.27	3.25	2.38	19.20	22.21	22.20	54.60	35.83	33.40
<b>Mean</b>		<b>3.74</b>	<b>2.77</b>	<b>1.75</b>	<b>15.42</b>	<b>18.6</b>	<b>17.45</b>	<b>42.88</b>	<b>26.21</b>	<b>20.85</b>
<b>I<sub>100</sub></b>	<b>B<sub>1</sub></b>	3.36	3.13	1.48	13.90	13.52	12.20	38.00	21.60	16.32
	<b>B<sub>2</sub></b>	3.75	3.29	1.77	15.60	14.71	12.94	40.60	28.80	18.40
	<b>KH<sub>1</sub></b>	3.89	3.31	2.19	16.80	19.69	16.62	45.40	29.20	27.75
	<b>KH<sub>2</sub></b>	4.03	3.47	2.53	17.80	23.36	19.80	49.40	33.60	29.84
	<b>Compost</b>	4.63	3.58	2.73	21.20	26.72	27.80	57.80	36.90	35.00
	<b>Mean</b>	<b>3.93</b>	<b>2.36</b>	<b>2.14</b>	<b>17.06</b>	<b>19.60</b>	<b>17.87</b>	<b>46.24</b>	<b>30.02</b>	<b>25.46</b>
<b>LSD at 0.05</b>	<b>Irrigation</b>	0.13	0.11	0.30	0.57	0.74	0.70	1.63	0.95	0.77
	<b>Organic</b>	0.06	0.05	0.04	0.27	0.31	0.29	0.73	0.47	0.40
	<b>Interaction</b>	**	***	***	***	***	***	***	***	***

<b>Mean of Organic amendments</b>	<b>B<sub>1</sub></b>	3.28	2.64	1.33	11.9	12.59	11.89	35.33	18.87	15.69
	<b>B<sub>2</sub></b>	3.61	2.74	1.45	14.37	15.16	14.41	37.93	23.42	1.6
	<b>KH<sub>1</sub></b>	3.73	2.78	1.60'	16.30	18.45	15.96	40.20	25.74	21.02
	<b>KH<sub>2</sub></b>	3.87	2.90	1.95	17.23	20.56	18.15	42.60	33.10	23.59
	<b>Compost</b>	4.36	3.12	2.21	19.57	22.47	22.6	53.27	36.57	31.20

Note: \*\* and \*\*\* refer to significance level of 5% ( $p \leq 0.05$ ) and 1% ( $p \leq 0.01$ ). ns:no-significant.

**Table 7: Effect of irrigation water regime and organic amendments on the availability of Fe,Zn,Mn and their content in maize under calcareous soil.**

Irrigation water regime	Organic amendments	Soil (mg kg <sup>-1</sup> )			Straw (mg kg <sup>-1</sup> )			Grain (mg kg <sup>-1</sup> )		
		Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn	Zn
<b>I<sub>50</sub></b>	<b>B<sub>1</sub></b>	3.30	2.25	1.19	24.32	21.66	19.97	39.24	30.80	31.13
	<b>B<sub>2</sub></b>	3.50	2.41	1.24	25.95	23.22	18.36	40.88	32.20	20.37
	<b>KH<sub>1</sub></b>	3.80	2.48	1.46	27.64	25.47	20.13	42.50	34.75	24.28
	<b>KH<sub>2</sub></b>	3.99	2.74	1.52	33.23	26.97	21.18	43.87	35.27	26.76
	<b>Compost</b>	4.72	3.06	1.72	34.22	29.23	29.84	47.30	36.69	39.26
	<b>Mean</b>	3.86	2.60	1.43	29.07	25.31	21.90	42.76	34.97	28.36
<b>I<sub>75</sub></b>	<b>B<sub>1</sub></b>	3.37	2.85	1.44	25.94	23.80	15.65	44.51	32.33	18.23
	<b>B<sub>2</sub></b>	3.83	2.96	1.52	28.54	26.26	17.29	45.35	33.38	24.22
	<b>KH<sub>1</sub></b>	4.00	3.03	1.68	32.38	27.52	20.74	48.23	34.40	25.36
	<b>KH<sub>2</sub></b>	4.23	3.22	2.13	35.40	28.38	22.59	51.18	35.05	29.04
	<b>Compost</b>	4.78	3.80	2.69	36.90	31.31	34.67	53.33	39.30	41.75
	<b>Mean</b>	4.04	3.24	1.89	31.83	27.45	22.19	48.52	36.28	27.72
<b>I<sub>100</sub></b>	<b>B<sub>1</sub></b>	3.54	3.50	1.53	33.54	24.54	17.19	45.46	33.14	22.99
	<b>B<sub>2</sub></b>	4.06	3.78	1.91	36.66	25.72	19.09	49.26	34.05	27.15
	<b>KH<sub>1</sub></b>	4.24	3.84	2.35	37.12	27.58	28.50	50.90	35.14	36.19
	<b>KH<sub>2</sub></b>	4.47	4.09	2.76	43.83	29.21	30.75	56.49	36.75	40.19
	<b>Compost</b>	5.31	4.37	3.06	44.79	33.05	38.98	60.23	41.34	45.18
	<b>Mean</b>	4.32	3.92	2.32	39.19	28.02	26.90	52.47	33.66	34.34
<b>LSD at 0.05</b>	<b>Irrigation</b>	0.15	0.14	0.33	1.90	0.98	1.65	2.28	1.20	2.96
	<b>Organic</b>	0.07	0.15	0.04	2.16	0.49	1.52	1.22	0.62	2.91
	<b>Interaction</b>	***	ns	***	ns	*	***	**	***	***
<b>Mean of Organic amendments</b>	<b>B<sub>1</sub></b>	3.41	2.87	1.39	27.93	23.33	17.60	43.07	32.09	24.12
	<b>B<sub>2</sub></b>	3.80	3.03	1.56	30.38	25.07	18.25	45.16	33.21	23.91

<b>KH<sub>1</sub></b>	4.01	3.23	1.83	32.38	26.86	23.12	477.21	34.76	28.61
<b>KH<sub>2</sub></b>	4.23	3.35	2.14	37.49	28.19	24.84	50.51	35.67	31.99
<b>Compost</b>	4.94	3.74	2.49	38.64	31.20	34.50	53.62	39.11	42.07

Note: \*, \*\* and \*\*\* refer to significance level of 10 ( $p \leq 0.1$ ), 5% ( $p \leq 0.05$ ) and 1% ( $p \leq 0.01$ ).

**Table 8: Effect of irrigation water regime and organic amendments on some physical soil properties under wheat and maize crops.**


Irrigation water regime	Organic amendments	Maize			Wheat	Maize
		FC %	WP %	AW %	HC (m d <sup>-1</sup> )	HC (m d <sup>-1</sup> )
I <sub>50</sub>	B <sub>1</sub>	32.31	6.10	26.21	1.62	1.31
	B <sub>2</sub>	28.69	5.60	22.79	1.80	1.43
	KH <sub>1</sub>	32.60	5.60	27.00	0.85	1.22
	KH <sub>2</sub>	27.65	5.60	22.05	0.87	1.29
	Compost	30.40	6.60	23.80	2.24	1.47
	Mean	30.33	5.9	24.37	1.46	1.34
I <sub>75</sub>	B <sub>1</sub>	29.82	5.60	24.22	1.96	1.49
	B <sub>2</sub>	30.40	5.60	24.80	2.12	1.57
	KH <sub>1</sub>	30.05	6.60	23.45	1.04	1.38
	KH <sub>2</sub>	30.15	6.30	23.85	1.12	1.45
	Compost	27.44	6.40	21.04	2.26	1.80
	Mean	29.57	6.1	23.47	1.70	1.56
I <sub>100</sub>	B <sub>1</sub>	29.13	6.31	24.23	2.08	1.84
	B <sub>2</sub>	28.46	9.64	18.82	2.19	2.11
	KH <sub>1</sub>	28.74	7.51	21.23	1.74	1.77
	KH <sub>2</sub>	27.80	6.73	21.07	1.44	1.78
	Compost	30.54	6.31	24.23	2.45	2.14
	Mean	28.93	7.29	21.92	1.98	1.93
LSD at 0.05	Irrigation	0.30	0.07	0.30	0.13	0.31
	Organic	0.35	0.09	0.41	0.11	0.04
	Interaction	***	***	***	***	***
Mean of Organic amendments	B <sub>1</sub>	30.38	6.06	24.77	1.92	1.56
	B <sub>2</sub>	29.18	6.95	22.14	2.04	1.70
	KH <sub>1</sub>	30.46	6.56	23.89	1.21	1.46
	KH <sub>2</sub>	28.53	6.21	22.32	1.14	1.51
	Compost	29.39	6.43	22.93	2.33	1.82

Note: \*, \*\* and \*\*\* refer to significance level of 10 (p≤0.1), 5% (p≤0.05) and 1% (p≤0.01).


**Table 9: Effect of irrigation water regime and organic amendments on yield and its components of wheat and maize under calcareous soil.**

Irrigation water regime	Organic amendments	Wheat			Maize				
		1000-grain weight (g)	Grain yield (ton ha <sup>-1</sup> )	Straw yield (ton ha <sup>-1</sup> )	Plant height (cm)	Ear weight (g)	Weight of 100 grains (g)	Grain yield (ton ha <sup>-1</sup> )	Straw yield (ton ha <sup>-1</sup> )
I <sub>50</sub>	B <sub>1</sub>	34.07	5.65	10.85	230.90	334.78	38.33	9.63	14.10
	B <sub>2</sub>	38.30	5.67	10.63	232.90	373.09	40.00	10.05	14.67
	KH <sub>1</sub>	39.80	6.00	9.63	240.97	374.78	42.00	10.59	15.93
	KH <sub>2</sub>	39.93	6.28	8.27	245.47	381.55	42.33	10.86	16.42
	Compost	40.20	6.71	10.51	237.00	416.66	43.00	11.19	18.25
	Mean	38.46	6.06	9.98	237.45	376.16	41.13	10.46	15.87
I <sub>75</sub>	B <sub>1</sub>	39.73	6.31	11.78	239.87	349.22	41.00	10.12	14.57
	B <sub>2</sub>	40.00	6.63	13.56	240.57	374.78	42.33	10.71	14.69
	KH <sub>1</sub>	42.51	6.74	13.70	249.23	410.56	42.67	11.36	15.29
	KH <sub>2</sub>	44.55	7.08	12.77	257.77	420.33	44.33	12.06	15.98
	Compost	53.26	7.15	12.20	245.20	436.01	45.00	12.47	16.45
	Mean	44.01	6.78	12.80	246.53	392.01	43.07	11.03	15.39
I <sub>100</sub>	B <sub>1</sub>	42.53	7.15	10.62	228.37	352.56	41.33	10.28	13.71
	B <sub>2</sub>	42.67	7.60	10.84	246.90	355.22	42.33	11.24	17.79
	KH <sub>1</sub>	55.64	8.19	10.35	249.10	364.00	44.33	11.79	18.01
	KH <sub>2</sub>	58.37	8.24	9.67	253.43	423.89	46.33	12.33	18.12
	Compost	57.55	8.72	10.98	248.33	464.33	52.33	12.84	20.01
	Mean	51.35	7.98	10.49	245.23	398.19	45.33	11.34	17.53
LSD at 0.05	Irrigation	3.25	0.13	0.30	3.72	10.07	1.55	0.329	0.389
	Organic	3.01	0.22	0.50	4.93	14.20	0.730	0.263	0.218
	Interaction	**	ns	***	ns	**	***	***	*
Mean of Organic amendments	B <sub>1</sub>	38.78	6.37	11.09	233.04	345.52	40.22	10.01	14.13
	B <sub>2</sub>	40.32	6.63	11.68	240.12	367.7	41.56	10.67	15.71
	KH <sub>1</sub>	45.98	6.98	11.23	246.43	383.11	43.00	11.24	16.41
	KH <sub>2</sub>	47.62	7.20	10.24	252.22	408.59	44.33	11.75	16.84
	Compost	50.33	7.53	11.23	243.51	439.00	46.78	11.05	18.24

Note: \*\* and \*\*\* refer to significance level of 5% (p≤0.05) and 1% (p≤0.01). \*ns:no-significant.



**Fig.1: Drought tolerance efficiency as affected by irrigation water regime and soil amendment of wheat and maize**



**Fig.2: Water use efficiency as affected by irrigation water regime and soil amendments of wheat and maize**