

Original Research Article

Effect of foliar application of biostimulants and micronutrients on chlorophyll and xanthophylls content of African marigold cv. Pusa Narangi Gainda

ABSTRACT

The present experiment entitled “Effect of foliar application of biostimulants and micronutrients on chlorophyll and xanthophylls content of African marigold cv. Pusa Narangi Gainda” was carried out at Jamuvadi Farm, Department of Horticulture, College of Agriculture, Junagadh Agricultural University, Junagadh, during 2017 to 2019. The experiment was laid out in Randomized Block Design with Factorial concept (FRBD) consisting two factors with three replications. The treatment comprised with four biostimulants and three treatment of micronutrients. The result indicated that foliar application of banana pseudostemsap @ 1% with micronutrient grade-IV @ 1% in addition to recommended dose of fertilizers (200:100:100 kg/ha NPK) produced higher chlorophyll content in leaves and xanthophylls content in flower of African marigold cv. Pusa Narangi Gainda.

Key words: Biostimulants, Micronutrients, African marigold cv. Pusa Narangi Gainda

Introduction

African marigold (*Tagetes erecta* L.) is one of the important commercial flower of India as well as Gujarat and being grown for its spectacular flowers, brilliant colours, delightful appearance, size, shape, forms etc. It belongs to the family Asteraceae (2n=24) and originated from Central to Southern America especially Mexico, from there it reached to Spain and became popular by the name of 'Rose of Indies'. At present, for the increasing flower production, nutrient are supplied through chemical fertilizers. Biostimulants are products of natural and organic origin that stimulates plants to achieve their higher growth, quality of flower and yield potential. Seaweed extracts contain major and micro nutrients, amino acids, vitamins, cytokinins, auxin and abscisic acid like growth promoting substances. The liquid contained macronutrients like P 120 mg/100g, K 4170 mg/100g, Ca 66.98 mg/100g and micronutrients like Fe 147 mg/100g, Mn 5.84 mg/100g, Zn 9.08 mg/100g and Cu 0.36 mg/100g (Yan *et al.*, 2013). Panchagavya is a fermented product made from five ingredients obtained from cow, such as milk, urine, dung, curd and clarified butter (Amalraj *et al.*, 2013). Panchgavya contained macro element like total nitrogen (229 ppm), total phosphorous (209 ppm), total potassium (232 ppm), calcium (25 ppm), IAA (8.5 ppm) and GA (3.5 ppm) (Anon., 2017). While separating fibers from the banana pseudostem, the liquid available is known as sap which contains good amount of essential macro element like 119 ppm N, 50.4 ppm P, 1289 ppm K and micronutrients like Fe-124 ppm, Mn-6.73 ppm, Cu-4.61 ppm and Zn-0.97 ppm (Gundrashiya, 2013) and also growth promoting substance like, cytokinin- 137.8 mg/l and gibberellic acid- 110.2 mg/l present (Desai, 2018).

Micronutrients are needed in very small amounts. Their concentration in plants are generally below the 100 parts per million (ppm) level. Out of 17 essential elements, Fe, Zn, B, Cu, Mn,

Mo although required in very little amount but their importance for the plant is no way less than those of major elements. Due to, deficiency of these minor elements the leaves, branches and flower may not grow properly and they may even affect the flower quality as well as production. These elements also help in development of hormone, enzyme, chlorophyll and in the absorption of the major elements. The need of micronutrients in flower production has long been recognized in India. It is important to keep the need for micronutrient fertilizers in perspective. Application of micronutrients in the hope of increasing crop yields even through there is little evidence to suggest a deficiency exists. The concern about micronutrient deficiencies are encouraged to investigate the need thoroughly and apply the nutrients in test strips if necessary. Considering the above facts, the present study was planned and undertaken with the objective to assess the response of biostimulants and micronutrients on chlorophyll and xanthophylls content of African marigold cv. Pusa Narangi Gaiinda.

Materials and methods

The field experiment was carried out twice during October 2017 to February 2019 at the Jambuvadi Farm, Department of Horticulture, Junagadh Agricultural University, Junagadh (Gujarat). The experiment was laid out in Randomized Block Design with Factorial concept (FRBD) consisting two factors with three replications. The treatment comprised with four biostimulants *viz.*, without spray of biostimulants (B₀), Seaweed extract @ 1% (B₁), Panchgavya @ 3% (B₂) and Banana pseudostem sap @ 1% (B₃) and three treatments of micronutrients *i.e.* without micronutrients (F₀), micronutrients grade-IV @ 0.5% (F₁) and micronutrients grade-IV @ 1% (F₂). Five plants from each treatment plot were randomly selected, labeled and used for recording observation. Chlorophyll content of leaf was analyzed by collecting the healthy fully matured second leaf from the centre of the plant at peak vegetative stage. Chlorophyll-a, chlorophyll-b and total chlorophyll contents of leaf tissue were determined by using Dimethyl sulfoxide (DMSO). Xanthophylls was estimated by AOAC Hot saponification method (Lawarance, 1990).

Table 1: Time of applications

Time of foliar application of biostimulants (both seasons)	Time of foliar application of micronutrinets (both seasons)
1 st 30 days after transplanting	1 st 40 days after transplanting
2 nd 45 days after transplanting	2 nd 55 days after transplanting
3 rd 60 days after transplanting	3 rd 70 days after transplanting

Result and discussion

Table 2: Effect of biostimulants on chlorophyll content in leaves

Significantly maximum chlorophyll-a content in leaves (0.753, 0.790 and 0.772 mg/g) was recorded with application of banana pseudostem sap 1% (B₃) during 2017-18, 2018-19 and in pooled. The maximum chlorophyll-b content was significantly increased in leaves (0.329, 0.343 and 0.336 mg/g) was recorded with application of banana pseudostem sap 1% (B₃) during 2017-18, 2018-19 and in pooled. Which was at par with treatment of panchgavya 3% (B₂) during 2018-19 and in pooled. The total chlorophyll content was significantly increased in

leaves (1.084, 1.103 and 1.093 mg/g) was recorded maximum with application of banana pseudostem sap 1% (B₃) during 2017-18, 2018-19 and in pooled. Significantly maximum chlorophyll-a, b and total chlorophyll content in leaves were recorded with an application of banana pseudostem sap @ 1% (B₃) during the year 2017-18, 2018-19 and in pooled. The organic liquid generated during extracting fiber from banana pseudostem sap used as spraying material in this experiment contained about 124 ppm of Fe (Gundrashiya, 2013). The most well known function of Fe is in enzyme systems in which haem or haemin functions as prosthetic group. Thus Fe plays a somewhat similar role to Mg in the porphyrin structure of chlorophyll. The haem pigments constitutes only about 0.1 % of the total Fe in plant leaves. In green plants, there is good correlation between the levels of Fe supply and the chlorophyll content. Plant well supply with Fe is high in chlorophyll. Along with this iron is not readily mobile between different plant organs. Hence green plants deprived of Fe soon become chlorotic in the younger plant parts. Results are in consonance with the findings of Desai (2018) in tuberose and Gundrashiya, (2013) in okra, cluster bean and cow pea.

Table 2: Effect of micronutrients on chlorophyll content in leaves

The maximum chlorophyll-a content was significantly increased in leaves (0.673, 0.713 and 0.693 mg/g) was recorded in F₂ treatment (micronutrient grade-IV 1%) during the year 2017-18, 2018-19 and in pooled, respectively. This treatment was at par with micronutrient grade-IV 0.5% (F₁) during both the year and in pooled. Significantly maximum chlorophyll-b content in leaves (0.304, 0.319 and 0.312 mg/g) was recorded in F₂ treatment (micronutrient grade-IV 1%) during the year 2017-18, 2018-19 and in pooled, respectively. This treatment was at par with micronutrient grade-IV 0.5% (F₁) during 2017-18 and in pooled. The maximum total chlorophyll content was significantly increased in leaves (0.977, 1.015 and 0.996 mg/g) was recorded in F₂ treatment (micronutrient grade-IV 1%) during the year 2017-18, 2018-19 and in pooled. This treatment was at par with micronutrient grade-IV 0.5% (F₁) during 2017-18, 2018-19 and in pooled. Chlorophyll is a major green pigment found in green leaves and is undoubtedly determining the photosynthetic efficiency and productivity of plants. The micronutrients might have increased the nutrient availability for synthesis and accumulation the chlorophyll content in leaves of marigold plant. The variation in chlorophyll content due to micronutrients may be attributed to decrease in chlorophyll degradation and increased chlorophyll synthesis. Iron is involved in chlorophyll synthesis pathway. The best role of iron is its catalytic function in nutrient absorption and balancing other nutrients and also in biological oxidation, reduction and other metabolic processes in plants. It may also be associated with organic acid metabolism.

Table 2a: Interaction effect of biostimulants and micronutrients on chlorophyll content in leaves

The combined application of banana pseudostem 1% with micronutrient grade-IV 0.5% (B₃F₁) recorded the maximum chlorophyll-a content in leaves (0.776 mg/g) during the year 2017-18. It was at par with B₃F₀ and B₃F₂ during 2017-18. While, banana pseudostem 1% with micronutrient grade-IV 1% (B₃F₂) recorded the maximum chlorophyll-a content in leaves (0.862 and 0.796 mg/g) during the year 2018-19 and in pooled, respectively. However, it was found at par with treatment of B₃F₁, during pooled. The combined application of banana

pseudostem 1% (B_3F_0) recorded the maximum chlorophyll-b content in leaves (0.345 mg/g) during the year 2017-18 and (0.360 and 0.351 mg/g) with combined application of banana pseudostem 1% with micronutrient grade-IV 1% (B_3F_2) during the year 2018-19 and in pooled, respectively. However, it was found at par with treatment of B_3F_1 and B_3F_0 during pooled, respectively. While, found at par with treatment of B_3F_1 , B_3F_2 , B_2F_0 and B_0F_2 during 2017-18 and during 2018-19 it was at par with treatment of B_3F_1 , B_3F_0 , B_2F_2 , B_2F_1 and B_2F_0 . The application of banana pseudostem sap 1% with without micronutrient (B_3F_0) recorded the maximum total chlorophyll (1.098 mg/g) during the year 2017-18, respectively. It was found at par with treatment of B_3F_2 and B_3F_1 during 2017-18, respectively. Application of banana pseudostem 1% with micronutrient grade-IV 1% (B_3F_2) recorded the maximum total chlorophyll (1.134 and 1.102 mg/g) during the year 2018-19 and in pooled. However, it was found at par with treatment of B_3F_1 and B_3F_0 during 2018-19 and in pooled, respectively. In green plants, there is good correlation between the levels of Fe supply and the chlorophyll content. Plant well supply with Fe is high in chlorophyll. Along with this iron is not readily mobile between different plant organs. Hence green plants deprived of Fe soon become chlorotic in the younger plant parts. Results are in consonance with the findings of Desai (2018) in tuberose and Gundrashiya, (2013) in okra, cluster bean and cow pea. The favorable effects chlorophyll content could be due to the positive response of marigold to micronutrient application. Besides, vigorous plant growth, mainly in terms of foliage, resulted in efficient production of photosynthetic products. Another reason could be that the application of Zn and Fe regulate various metabolic activities of plants and also are involved in the auxin production.

Table 3: Effect of biostimulants on xanthophylls content of flower

Significantly maximum xanthophylls content in flower (1.593, 1.543 and 1.567 mg/g) was recorded with application of banana pseudostem sap 1% (B_3). Which was at par with treatment of panchgavya 3% (B_2) during the year 2018-19 and in pooled. It might be due to banana pseudostem sap contains good amount of essential macro and micronutrients as well as growth boosters. Salunkhe, (2010) analysed the samples of banana pseudostem for its elemental composition and found that banana pseudostem contained macro elements in the range of 1.00 to 1.12 % N, 0.50 to 0.71 % P, 2.39 to 20.2 % K and micro nutrients in the range of 259 to 323.2 mg/kg Fe, 47.3 to 241.3 mg/kg Mn, 10.1 to 107.4 mg/kg Zn and 13.4 to 83.6 mg/kg Cu. Beneficial effect of zinc on photosynthetic pigments might be due to its role in increasing the rates of photochemical reduction, enzyme of carbohydrates transformation, photosynthetic electron transfer as well as photosynthesis and iron may be due to indirect role of iron in chlorophyll biosynthesis. Similar result also obtained by Balakrishnan *et al.* (2007) in marigold, EL-Naggar (2009) in carnation and Khalifa *et al.* (2011) in iris.

Table 3: Effect of micronutrients on xanthophylls content of flower

Significantly maximum xanthophylls content in flower (1.350, 1.380 and 1.365 mg/g) was recorded in F_2 treatment (micronutrient grade-IV 1%) during the year 2017-18, 2018-19 and in pooled, respectively. This treatment was at par with micronutrient grade-IV 0.5% (F_1) during the year 2018-19, respectively. Regarding the beneficial effect of iron may be due to indirect role of iron in chlorophyll biosynthesis and beneficial effect of zinc on photosynthetic pigments

might be due to its role in increasing the rates of photochemical reduction, enzyme of carbohydrates transformation, photosynthetic electron transfer as well as photosynthesis. Similar result also obtained by Balakrishnan *et al.* (2007) in marigold, EL-Naggar (2009) in carnation and Khalifa *et al.* (2011) in iris.

Table 3a: Interaction effect of biostimulants and micronutrients on xanthophylls content of flower

The combined application of banana pseudostem 1% with micronutrient grade-IV 1% (B₃F₂) recorded significantly maximum xanthophyll content in flower (1.700, and 1.610 mg/g) during the year 2017-18 and in pooled, respectively. However, it was found at par with treatment of B₃F₁ and B₃F₀ during pooled, respectively. The combined application of banana pseudostem 1% with without micronutrient spray (B₃F₀) recorded maximum xanthophylls content in flower (1.580 mg/g) during the year 2018-19. Which was at par with treatment B₃F₂, B₃F₁, B₂F₂ and B₂F₁ during the year 2018-19. It might be due to banana pseudostem sap contains good amount of essential macro and micronutrients as well as growth boosters. Salunkhe, (2010). This might be due to the indirect role of iron in chlorophyll biosynthesis. Regarding the beneficial effect of zinc on photosynthetic pigment may be due to its role in increasing the rates of photochemical reduction, chloroplast structure, photosynthetic electron transfer as well as photosynthesis. Similar result was also reported by Balakrishnan *et al.* (2007) in marigold and Yadegari (2013) in pot marigold.

Summary and conclusion

On the basis of result obtained in the present investigation it may be concluded that the foliar application of banana pseudostem sap @ 1% at 30, 45 & 60 days after transplanting with micronutrient grade-IV @ 1 % at 40, 55 & 70 days after transplanting in addition to recommended dose of fertilizers (200:100:100 kg/ha NPK) proved to be the best treatment for getting higher chlorophyll content in leaves, vegetative growth and xanthophylls content in flower of African marigold cv. Pusa Narangi Gaiinda.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

- Amalraj, E. L. D.; Praveen, K.G.; Ahmed, S. K. M. H.; Abdul, R. and Kishore, N. 2013. Microbiological analysis of panchagavya, vermicompost, and FYM and their effect on plant growth promotion of pigeon pea (*Cajanus cajan* L.) in Indian. *Organic Agriculture*, **3**: 23–29.
- Anonymous. 2017. The Indian Horticulture Database 2016-17. National Horticulture Board, Gurgaon. Available at [http://nhb.gov.in/area-pro/\(3rdEstimate\).pdf](http://nhb.gov.in/area-pro/(3rdEstimate).pdf) accessed on 5 May, 2017.
- Balakrishnan, V.; Jawaharlal, M.; Senthil Kumar, T. and Ganga, M. 2007. Response of micronutrients on flowering and yield in African marigold. *J. Ornament. Hort.*, **10** (3): 153-156.
- Desai, S. A. 2018. Effect of plant growth enhancers on growth, flowering and yield of tuberose cv. Prajwal. M. Sc. Thesis, Navsari agricultural University, Navsari (Gujarat, India).
- El-Naggar, A. H. 2009. Response of *Dianthus caryophyllus* L. plants to foliar nutrition. *World J. Agri. Sci.*, **5** (5): 622-630.
- Gundrashiya, R. R. 2013. Effect of spraying of banana pseudostem based enriched sap at different concentration on growth and yield of different crops. M. Sc. Thesis, Navsari agricultural University, Navsari (Gujarat, India).
- Khalifa, R.; Shaaban, S and Rawia, A. 2011. Effect of foliar application of zinc sulfate and boric acid on growth, yield and chemical constituents of Iris plant. *Oze. J. Appl. Sci.*, **4** (2): 129-144.
- Lawrence, J. F. 1990. Determination of total xanthophylls and marigold oleoresin. *J. of Assoc. of Official Anal. Chemists.*, **2**: 970-975.
- Salunkhe, J. R. 2010. Feasibility of using banana pseudostem sap as liquid fertilizer in onion under drip irrigation. *M.Sc. Thesis*, Navsari Agricultural University, Navsari.
- Yadegari, M. 2013. Foliar application of Fe, Cu, Mn, and B on growth, yield and essential oil yield of *Calendula officinalis*. *J. Appl. Sci. Agri.*, **8** (5): 559-567.
- *Yan, P.; Enyi, X.; Kai, Z.; Mangaladoss, F.; Xianwen, Y.; Xuefeng, Z.; Yifei, W.; Bin, Y.; Xiuping, L.; Juan, L. and Yonghong, L. 2013. Nutritional and chemical composition

and antiviral activity of cultivated seaweed *Sargassum naozhouense* Tseng et Lu. Mar Drugs., **11** (1): 20-32.

Yan, P.; Enyi, X.; Kai, Z.; Mangaladoss, F.; Xianwen, Y.; Xuefeng, Z.; Yifei, W.; Bin, Y.; Xiuping, L.; Juan, L. and Yonghong, L. 2013. Nutritional and chemical composition and antiviral activity of cultivated seaweed *Sargassum naozhouense* Tseng et Lu. Mar Drugs.,**11** (1): 20-32.

UNDER PEER REVIEW

Table 2: Effect of biostimulants and micronutrients on chlorophyll content in African marigold cv. Pusa Narangi Gainda

Treatments	Chlorophyll-a content in leaves (mg/g)			Chlorophyll-b content in leaves (mg/g)			Total Chlorophyll content in leaves (mg/g)		
	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled
Level of Biostimulants (B)									
B ₀ – Control	0.581	0.607	0.594	0.256	0.246	0.251	0.843	0.857	0.850
B ₁ – Seaweed extract (1%)	0.588	0.619	0.604	0.274	0.280	0.277	0.858	0.906	0.882
B ₂ – Panchgavya (3%)	0.671	0.654	0.663	0.294	0.319	0.306	0.965	1.000	0.982
B ₃ – Banana pseudostem sap (1%)	0.753	0.790	0.772	0.329	0.343	0.336	1.084	1.103	1.093
S.Em.±	0.011	0.012	0.008	0.010	0.009	0.007	0.016	0.020	0.013
C.D. at 5 %	0.033	0.034	0.02	0.030	0.027	0.020	0.047	0.058	0.036
Level of Micronutrients (F)									
F ₀ – Control	0.618	0.611	0.615	0.267	0.282	0.275	0.887	0.912	0.900
F ₁ – Micronutrient Grade-IV (0.5%)	0.654	0.678	0.666	0.293	0.289	0.291	0.948	0.971	0.959
F ₂ – Micronutrient Grade-IV (1%)	0.673	0.713	0.693	0.304	0.319	0.312	0.977	1.015	0.996
S.Em.±	0.010	0.010	0.007	0.009	0.008	0.006	0.014	0.017	0.011
C.D. at 5 %	0.029	0.030	0.020	0.026	0.024	0.017	0.041	0.051	0.032
Interaction (B X F)									
S.Em.±	0.020	0.020	0.014	0.018	0.016	0.012	0.028	0.035	0.022
C.D. at 5 %	0.057	0.060	0.040	0.051	0.047	0.033	0.081	0.101	0.063
CV %	5.21	5.28	5.25	10.51	9.34	9.93	5.1	6.19	5.69

Table 2a: Interaction effect of biostimulants and micronutrients on chlorophyll content in leaves of marigold cv. Pusa Narangi Gainda

Treatments	Chlorophyll-a content in leaves (mg/g)			Chlorophyll-b content in leaves (mg/g)			Total Chlorophyll content in leaves (mg/g)		
	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled	2017-18	2018-19	Pooled
B ₀ F ₀	0.473	0.512	0.493	0.170	0.192	0.181	0.667	0.714	0.690
B ₀ F ₁	0.608	0.628	0.618	0.290	0.242	0.266	0.895	0.874	0.885
B ₀ F ₂	0.661	0.682	0.671	0.309	0.303	0.306	0.969	0.982	0.976
B ₁ F ₀	0.605	0.600	0.603	0.255	0.283	0.269	0.847	0.895	0.871
B ₁ F ₁	0.556	0.636	0.696	0.288	0.263	0.275	0.844	0.902	0.873
B ₁ F ₂	0.602	0.621	0.612	0.279	0.294	0.286	0.883	0.920	0.901
B ₂ F ₀	0.638	0.619	0.629	0.299	0.322	0.311	0.937	0.971	0.954
B ₂ F ₁	0.677	0.655	0.666	0.293	0.315	0.304	0.968	1.004	0.986
B ₂ F ₂	0.698	0.688	0.693	0.288	0.320	0.304	0.988	1.024	1.006
B ₃ F ₀	0.755	0.714	0.735	0.345	0.333	0.339	1.098	1.070	1.084
B ₃ F ₁	0.776	0.794	0.785	0.302	0.336	0.319	1.084	1.103	1.093
B ₃ F ₂	0.729	0.862	0.796	0.341	0.360	0.351	1.070	1.134	1.102
B ₀ F ₀	0.473	0.512	0.493	0.018	0.016	0.012	0.028	0.035	0.022
B ₀ F ₁	0.608	0.628	0.618	0.051	0.047	0.033	0.081	0.101	0.063
S.Em.±	0.020	0.020	0.014	10.51	9.34	9.93	5.1	6.19	5.69
C.D. at 5 %	0.057	0.060	0.040	0.170	0.192	0.181	0.667	0.714	0.690
CV %	5.21	5.28	5.25	0.290	0.242	0.266	0.895	0.874	0.885

Table 3: Effect of biostimulants and micronutrients on xanthophylls content of marigold cv. Pusa Narangi Ganda

Treatments	Xanthophylls content (mg/g)		
	2017-18	2018-19	Pooled
Level of Biostimulants (B)			
B ₀ – Control	1.031	1.062	1.047
B ₁ – Seaweed extract (1%)	1.101	1.199	1.150
B ₂ – Panchgavya (3%)	1.296	1.493	1.394
B ₃ – Banana pseudostem sap (1%)	1.593	1.543	1.567
S.Em.±	0.022	0.023	0.052
C.D. at 5 %	0.065	0.067	0.236
Level of Micronutrients (F)			
F ₀ – Control	1.170	1.268	1.219
F ₁ – Micronutrient Grade-IV (0.5%)	1.246	1.325	1.285
F ₂ - Micronutrient Grade-IV (1%)	1.350	1.380	1.365
S.Em.±	0.019	0.020	0.014
C.D. at 5 %	0.056	0.058	0.039
Interaction (B X F)			
S.Em.±	0.038	0.040	0.027
C.D. at 5 %	0.112	0.116	0.078
CV %	5.25	5.16	5.21

Table 3a: Interaction effect of biostimulants and micronutrients on xanthophylls content of marigold cv. Pusa Narangi Ganda

Treatment combinations	Xanthophylls content (mg/g)		
	2017-18	2018-19	Pooled
B ₀ F ₀	0.860	0.947	0.903
B ₀ F ₁	1.100	1.087	1.093
B ₀ F ₂	1.133	1.153	1.143
B ₁ F ₀	1.030	1.100	1.065
B ₁ F ₁	1.117	1.197	1.157
B ₁ F ₂	1.157	1.300	1.228
B ₂ F ₀	1.260	1.447	1.353
B ₂ F ₁	1.217	1.487	1.352
B ₂ F ₂	1.410	1.547	1.478
B ₃ F ₀	1.530	1.580	1.555
B ₃ F ₁	1.550	1.530	1.540
B ₃ F ₂	1.700	1.520	1.610
S.Em. ±	0.038	0.040	0.027
C.D. at 5%	0.112	0.116	0.078
CV%	5.25	5.16	5.21