

Degradation pattern and residues of atrazine in maize under long term conservation agriculture

Abstract

Field experiment was conducted at Eastern block farm, Tamil Nadu Agricultural University with maize to study the degradation, persistence and residues of atrazine under long term conservation agriculture system. The treatments constituted atrazine 1.0 kg ha⁻¹ on 3 DAS and atrazine 1.0 kg ha⁻¹ on 3DAS +Hand weeding on 45th day under different conservation tillage practices. Degradation, persistence and residue analysis were carried out for atrazine using HPLC. Residues of atrazine was analyzed at different periods viz., 0,15,30,45,60,75 and 90 days after herbicide application and at harvest and the results revealed that degradation of atrazine was found to follow first order reaction kinetics ($R^2 > 0.94$) with the half life of 13.7–16.2 days irrespective of tillage practices and weed management methods. The recommended rate of atrazine @1.0 kgha⁻¹ recorded terminal residues below detectable limit of <0.01 mg kg⁻¹ in soil and plant parts of maize.

Keywords: maize- conservation tillage - atrazine- degradation- persistence- residue.

1.Introduction

Maize (*Zea mays* L.) is one of the most important cereal crops in the world agricultural economy used both as a food and fodder crop. It is a miracle crop with very high yield potential. The productivity of maize largely dependent upon its weed and nutrient management. It is very susceptible to the weed infestation. Atrazine is in a group of human-made systemic herbicides called triazines. It is used for broadleaf weeds both before and after they sprout. It is also used on some grassy weeds. Atrazine interferes with photosynthesis in some broadleaf plants and grasses. Atrazine is almost non-volatile and its half-life in neutral condition is about 200 days but varies from 4-57 weeks (McCormick and Hiltbold, 1966) depending on various environmental factors like pH, moisture content, temperature and microbial activity (Armstrong, et al., 1967; Frank and Sirons, 1985; Nair et al., 1993). Shrinivas *et al.* (2014) reported that combined application of atrazine (50 %) + pendimethalin (50%), showed best control of all the grasses, sedges and broad leaved weeds for long time without any phytotoxic effect. It was well witnessed from the data that, application of herbicides showed lower weeds than hand weeding twice and farmers practice that too from the day of sowing indicating efficiency of the herbicides in controlling weeds. Channabasavanna *et al.* (2015) reported that tank mix application of atrazine 50 WP @ 625 g a.i. ha⁻¹ + pendimethalin 30 EC @ 750 g a.i ha⁻¹ resulted in effective control of grass, broad leaved and sedge weeds.

Conservation agriculture is a sustainable farming that reduces soil disturbance, keeps the soil covered, and uses different crop rotations (Hobbs *et al.*, 2008). It helps to improve soil health and crop productivity and lower environmental impact. Weed management is crucial for obtaining profitable yields in reduced tillage system and achieving satisfactory weed control requires intensive management. Tillage practices were found to modify significantly pesticide degradation in soil, but in contrasted ways according to studies. For example, in no-tillage, vetch residues accelerated the degradation of metolachlor by from 1.5 to 3 times, but had no effect on the degradation of atrazine (Teasdale *et al.*, 2003). The nature and decomposition degree of crop residues both influence interception and retention of pesticides. In combination with other weed management practices, residues mulching prevents weed seeds germination by blocking the light required for weed seeds germination or inhibits weeds growth due to its allelopathic effect (Teasdale and Mohler, 2000).

Once the herbicides are added to the soil, besides under decomposition and physical removal, a portion of them may be taken up by the plant and accumulated in the edible parts. The residue accumulation should not exceed the maximum residue limit (MRL) prescribed by the National and International standards. The information on the degradation of atrazine in maize under conservation agriculture is lacking. Hence, the present study on the degradation and residues of atrazine in maize was carried out.

2. Materials and method

Maize - sunflower cropping system was followed from 2016 as a long term experiment to study the herbicidal weed management in conservation tillage practices. In this study a field experiment was conducted with maize during kharif 2019 at field no.37 Eastern Block farm of Tamil Nadu Agricultural University, Coimbatore. The experimental soil is sandy clay loam with a pH of 7.81 and EC 1.48 dSm⁻¹ respectively. The experiment was conducted in a split plot design with 3 replications. The treatments were as follows.

2.1. Treatment details

Rabi 2018 (sunflower) Previous crop	Kharif 2019 (Maize)
T ₁ Conventional tillage	Conventional tillage
T ₂ Conventional tillage	Zero tillage
T ₃ Zero tillage + Residue	Zero tillage
T ₄ Zero tillage	Zero tillage + Residue

T₅ Zero tillage + Residue

Zero tillage + Residue

Sub plot (Weed management practices)

W₁ Recommended PE Atrazine 1.0 kg/ha

W₂ PE Atrazine 1.0 kg/ha + HW on 45 DAS

W₃ Unweeded check

During Kharif 2019, maize was grown as test crop which received atrazine as pre emergence herbicide to control weeds. Soil and plant samples were collected from the herbicide applied and hand weeding plots on 0,15,30,45,60,75 days after herbicide application and subjected to residue analysis to find out the persistence of herbicides in soil and to study the terminal residues as influenced by the tillage practices, soil, maize grain and straw at harvest were taken and analysed for atrazine residues.

2.2 Residue extraction

The sample was extracted with 100 ml methanol as extracting solution after shaking in orbital shaker for 1 hr. The suspension was filtered through whatman no.42 filter paper and the extract was evaporated in water bath till the suspension reduced to 10 ml. The concentrated extract was transferred to separating funnel and the atrazine residue was portioned with 30 ml of 1% NaCl and 50 ml of hexane. The upper organic layer was collected separately. The lower aqueous layer was re-extracted twice with 25 ml hexane each time. The organic layer was pooled and filtered through whatman no. 42 filter paper containing a pinch (2 gm) of anhydrous sodium sulphate. The filtered residue extract was concentrated to dryness using rotary vacuum evaporator. The dried residues of atrazine compounds were dissolved in 2 ml acetonitrile HPLC grade solvent for HPLC determination.

2.3.HPLC parameters for atrazine detection

- i) Column : Agilent Eclipse C18, 4.6 x 150 mm, 5 µm
- ii) Mobile phase : Acetonitrile :MilliQ-Water (80:20 % v/v)
- iii) Flow rate : 0.8mL min⁻¹
- iv) Detector : Photo Diode Array Detector

Parameters

- v) Wavelength (λ_{\max}) : 221 nm
- vii) Injection volume : 20 µl

- viii) Retention time : 3.29 ± 0.2 min
ix) LOD : 0.01 mg /kg
x) LOQ : 0.05 mg/kg

2.4. Calculation

The amount of atrazine in the sample was calculated as given below:

$$\text{Residue in ppm (mg/kg)} = \frac{A_1 \times C \times V_1}{A_2 \times W} \times R_f$$

Where,

A_1 = Area of compound from sample, in chromatogram

A_2 = Area of compound from standard, in chromatogram

V_1 = Total volume of sample in ml

C = Concentration of analytical standard (96.55%) in ppm

W = Weight of the sample in g

R_f = Recovery factor

Analytical standard of different concentration was used to study the linearity of atrazine. In this study a calibration curve was prepared by taking the areas corresponding to different concentrations of analytical standard (0.01 to 1.0 mg L⁻¹). Before analysing the unknown samples, recovery studies were carried out to establish the reliability of the analytical method employed for the present study. The blank soil and plant samples were fortified with known concentrations of atrazine standard ranged from 0.05 to 0.5 mg L⁻¹ and fortified matrices were subjected to extraction and determination.

3. Results and discussion

The instrumental conditions presented above were followed for the determination of atrazine from different matrices. LOD of atrazine molecule was found to be 0.05 mg/kg and resolved at 3.29 min under the optimized HPLC conditions.

3.1. Standard calibration of atrazine

The chromatogram of atrazine standard as determined by HPLC-DAD was sketched in **(Fig. 1)**. By plotting peak areas and concentration on X and Y-axis, the equation of calibration graph for atrazine was represented in **(Fig. 2)** within range of 0.05 to 0.5 mg L⁻¹.




Fig 1. Chromatograms of atrazine (0.01mg/Lit) standards




Fig 2. Standard graph of atrazine determined by HPLC

3.2.Recovery studies of atrazine (Table 1)

The recovery studies of atrazine carried out in blank soil, maize grain and straw. The range of atrazine fortified were between 0.05 and 1.0 mg kg⁻¹. The average numerical value of recovery percentage of soil was 90.9 per cent among different concentrations. Atrazine recovery detected from maize plant parts and mean recovery from maize grain and straw were 85.7 and 87.8 per cent at spiked concentrations from 0.05 to 0.50 mg kg⁻¹.

Table 1. Recovery of atrazine in field soil and plant parts

Matrix	Concentration (ppm)	Amount recovered	Recovery percentage	Average
Field soil	0.05	0.046	92.0	90.9
	0.10	0.089	89.0	
	0.50	0.458	91.6	
Grain	0.05	0.040	80.0	85.7
	0.10	0.085	85.0	
	0.50	0.461	92.2	
Straw	0.05	0.042	84.0	87.8
	0.10	0.091	91.0	
	0.50	0.442	88.4	

*Average of three replicates

Degradation of atrazine were analyzed at different periods viz., 0, 15, 30, 45, 60, 75 and 90 days after herbicide application and the results are given in Table 2. The dissipation of atrazine was found to follow first order reaction kinetics ($R^2 > 0.94$) irrespective of tillage practices under both the weed control methods with the half life of 13.7 – 16.2 days. Irrespective of tillage practices and weed management methods, >80 % of atrazine dissipated in 45 days. Similar results were reported by Ram Prakash and Madhavi (2014) who found initial concentration of atrazine in soil was 0.541-0.602 mg kg⁻¹ which reduced to 80-88.7% at 45 days. Similarly degradation of pendimethalin and metolachlor in sunflower and soybean followed first order reaction kinetics (Janaki et al., 2015).

Table 2. Influence of conservation tillage and weed management practices on residues of atrazine (mg/kg) in soil with maize (kharif'19) in maize – sunflower system

Treatments	W_1 (Atrazine @ 0.5 kg/ha)						
	0 day	15 day	30 day	45 day	60 day	75 day	90 day
T ₁	0.280	0.156	0.090	0.022	0.018	0.010	BDL
T ₂	0.294	0.163	0.095	0.024	0.020	0.014	BDL
T ₃	0.245	0.134	0.072	0.022	0.017	0.012	BDL
T ₄	0.259	0.112	0.061	0.019	0.014	0.012	BDL
T ₅	0.266	0.108	0.053	0.020	0.012	0.011	BDL

	W₂ (Atrazine 0.5 kg/ha + HW on 45 DAS)						
T ₁	0.272	0.162	0.075	0.024	0.018	0.012	BDL
T ₂	0.283	0.177	0.091	0.028	0.019	0.010	BDL
T ₃	0.259	0.132	0.065	0.027	0.015	0.013	BDL
T ₄	0.281	0.112	0.038	0.021	0.011	0.011	BDL
T ₅	0.278	0.109	0.049	0.029	0.018	0.011	BDL

Residue analysis showed that the terminal residues of atrazine in soil, maize grain and straw from different plots were below 0.01 mg/kg irrespective of the tillage management practices followed for weed control. The findings are in accordance with the results of Tandon et al. [2015] who reported that, when atrazine was sprayed before emergence, the atrazine in corn kernels, soil and straw during harvest was less than 0.005 mg kg⁻¹. Cao *et.al.* (2023) studied a method for simultaneous determination of atrazine in fresh corn, corn kernels, and corn straw was established based on modified QuEChERS pre-treatment and high-performance liquid chromatography–tandem mass spectrometry (HPLC–MS/MS) and the results revealed that the terminal residues of atrazine in corn kernels, and atrazine was below the LOQ (0.01 mg kg⁻¹) and 0.05 mg kg⁻¹ in maize grain and straw (Janaki et al 2012). Babu et al (2015) also reported oxyfluorfen and Imazethapyr residues in groundnut kernel were BDL (0.005 mg kg⁻¹). Janaki et.al. (2012) concluded from the results that the longer persistence of atrazine can be expected at higher dose of application. In addition to this, the bioaccumulation of atrazine in soil was significant in terms of continuous application.

Conclusion

To conclude, persistence studies of atrazine showed that persistence of atrazine in soil was up to 75 days after herbicide application with a half life ranged between 13.7 – 16.2 days. The degradation followed 1st order kinetics for atrazine. Recommended rate of atrazine @ 1.0 kg ha⁻¹ recorded terminal residues below detectable limit of <0.01 mg kg⁻¹ in soil and maize plant parts of grain and straw.

References

Armstrong, D.E., Chesters, G. & Harris, R.F. (1967). Atrazine hydrolysis in soil, Soil Sci. Soc. Am. Proc., 31, 61-66.

Babu,C. Janaki,P, Chinnusamy,C.(2015). Effect of rate of application on degradation of imazethapyr in groundnut and soil under tropical Indian condition, J. Appl Nat Sci. 7(2): 714-718.

Cao, J., Pei, T.,Wang, Y.,Qin, S., Qi, Y., Ren, P.& Li, J.(2023). Terminal residue and dietary risk assessment of atrazine and isoxaflutole in corn using high-performance liquid chromatography–tandem mass spectrometry. *Molecules*, 28, 7225.

Channabasavanna, A.S., Shrinivas, C.S., Rajakumar, H & Kitturmath, M. S. (2015). Efficiency of tank mix application of atrazine 50 WP and pendimethalin 30 EC weedicides in maize. *Karnataka J. Agric. Sci.*, 28(3), 327-330.

Frank, R. & Sirons, G. (1985). Dissipation of atrazine residues from soil, *Bull. Environ. Contam. Toxicol.*,34, 541-548.

Hobbs, Peter R, Ken Sayre, & Raj Gupta. (2008). The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363 (1491),543-555.

Janaki,P, Meena, S. and Chinnusamy,C.(2015a). Dynamics of metolachlor in sandy clay laom soil and its residues in maize and soybean. *Trends Biosci.* 8:131-137.

Janaki,P., Meena, S., Chinnusamy, C., Murali Arthanari, P. & Nalini, K. (2012). Field persistence of repeated use of atrazine in sandy clay loam soil under maize. *Madras Agricultural Journal*,99 (7-9), 533-537.

Mc-Cormick, L.L. & Hiltbold, A.E. (1966). Microbiological decomposition of atrazine and diuron in soil, *Weeds*,14, 77-82.

Nair, D.R., Burken, J.G., Licht, L.A. & Schnoor, J.L. (1993). Mineralization and uptake of triazine pesticide in soil-plant systems, *J. Env. Engrg.*, ASCE, 119, 842-854.

Ramprakash, T & Madhavi, M. (2019). Persistence of herbicides in rice- maize cropping system. In: *Herbicide residue research in India*. Pp 289-303.

Shrinivas, C. S., Channabasavanna. A. S & Mallikarjun. (2014). Evaluation of sequential application of herbicides on nutrient uptake and yield of maize (*Zea mays* L.) under irrigated condition. *Res J Agric Sci.*, 5(5), 924-926.

Tandon, S. & Singh, A. (2015). Field dissipation kinetics of atrazine in soil and post-harvest residues in winter maize crop under subtropical conditions, *Chem. Ecol.* 2015, 31, 273–284.

Teasdale, J.R., Shelton, D.R., Sadeghi, A.M.& Isensee, A.R. (2003). Influence of hairy vetch residue on atrazine and metolachlor soil solution concentration and weed emergence, *Weed Sci.* 51, 628–634.

Teasdale, J.R & Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Sci.* 48(3): 385-392.