

A REVIEW OF ETHNOMEDICINAL USES, PHYTOCHEMISTRY AND PHARMACOLOGY OF NIGERIAN CROTONS

ABSTRACT

Approximately 80% of the world's population relies on traditional plants to produce and synthesize contemporary medications. In the past, the production of plants as sources of pharmaceuticals or medicines for wound healing and treatment was based on superstition and experience that was passed down from one generation to the next. The majority of these plants belong to the families Euphorbiaceae, Leguminosae, Rutaceae, and Piperaceae. The Euphorbiaceae family includes Croton plants. They are found in tropical and sub-tropical areas of both hemispheres and comprise of 1,300 species of trees, shrubs, and plants. This study focused on providing detailed review on chemical components of Croton species found in Nigeria. The term "ethnomedicinal potentials" refers to the therapeutic applications of plants that have positive pharmacological effects on both human and animal bodies. Prior to the development of modern medicine, certain Croton species were used to cure rheumatism, diabetes, diarrhea, cancer, and other illnesses. Active alkaloids are abundant in the Croton genus. In addition to triterpenoids, flavonoids, sesquiterpenoids, phytosterols, N-containing chemicals, cyclohexane derivatives, aliphatic molecules, and ferulic acid ester derivatives, diterpenoids are said to be widely distributed throughout the African Croton. This work is a critical review of ethnomedicinal, phytochemistry and pharmacology of various Nigerian Croton species. The study also discussed the chemical constituents of different Croton species.

Key word: Croton, Nigeria, phytochemistry, euphorbiaceae, African, traditional.

INTRODUCTION

Approximately around 80% of the world's population used traditional plants to produce and synthesize contemporary medications. In the past, production of plants as sources of medications for wound healing and treatment was based on superstition and experience that were passed from generation to generation, essentially through word of mouth (Folkloric).^[1] The majority of these plants belong to the families Euphorbiaceae, Leguminosae, Rutaceae, and Piperaceae. The Euphorbiaceae family comprises of croton plants. They are found in tropical and sub-tropical environments of both hemispheres and comprise of about 1,300 species of trees, shrubs, and plants.^[2] There are approximately 292 species of Croton in Africa, and eleven (11) of them are found in Nigeria. *Crotonhirtus* was recently discovered in Nigeria, Sierra Leone, Cote d'Ivoire, Republic of Congo and Gabon. Around the world, Croton species are applicable to treat a variety of conditions including cancer, diarrhea, intestinal worms, discomfort, ulcers, and weight loss.^{[2];[3];[4]} The term "ethnomedicinal potentials" refers to the therapeutic applications of plants that possess positive pharmacological effects on both human and animals. Prior to the development of modern medicine, certain croton species were used to treats rheumatism, diabetes, cancer, and other illnesses. Active alkaloids are abundant in the Croton genus.^{[5];[6]} In addition to triterpenoids, flavonoids, sesquiterpenoids, phytosterols, N-containing chemicals, cyclohexane derivatives, aliphatic molecules, and ferulic acid ester derivatives, diterpenoids are said to be widely distributed throughout the African Croton.^[4] Antihypertensive, antimalarial,

Comment [AW1]: This word is general name it can't go with the two preceding words 'trees' and 'shrubs'. You better replace it with 'herbs'.

Comment [AW2]: Must come before the last chemical compound 'diterpenoids'.

Comment [AW3]: You better include sentences on the methods you used to conduct your work and conclusion in your abstract.

Comment [AW4]: Replace by medicinal.

Comment [AW5]: Be consistent in capitalization.

Comment [AW6]: Replace with the growth form type preferably 'herbs'

Comment [AW7]: You need citation.

antimicrobial, and myo-relaxant are some other actions of *Croton* isolates that have been determined.^[2]

ETHNOMEDICINAL PROPERTIES AND GEOGRAPHICAL REPRESENTATION OF CROTON GENUS IN NIGERIA

African traditional healers have traditionally used medicinal plants as permanent supply to cure a number of severe diseases.^[7] In Africa, South Asia, and Latin America, *Croton* species have long been used to cure a variety of infections and digestive issues.^{[8];[9];[10]}

CROTONGRATISSIMUSBURCH.

This plant is indigenous to central and tropical West Africa. Senegal, Sudan, Botswana, and South Africa are among its locations.^{[11];[12]} Freestyle and sub-freestyle are its two divisions.^[4] Sub *gratissimus* differs from *gratissimus* in that it has stellate hairs on the upper leaf surface, whereas *gratissimus* does not. They can be found in South Africa's far north, Zimbabwe, and Botswana.^[4] It is used by the indigenous community as essential oil, food flavoring, and medicine.^[12] Malaria is treated with an infusion made from bark slashes. Bleeding gums are treated with the burned and powdered bark.^[12] Soup created from a leaf decoction is used as a wash to cure headaches, fever, and diarrhea.^[13] The shoots are used as a febrifuge, tonic, and to ease menstruation pain. Additionally, the root is employed as an aperient.^[13] Its bark extract has long been used as a styptic, cathartic, and treatment for a variety of ailments, pleurisy, uterine disorders, and intercostal neuralgia.^[14] Because of their fantastic aroma, its young branches are dried and ground into powder for use in perfume industries.^[14]

CROTONGRATISSIMUS VAR. GRATISSIMUS

The Republic of Congo and South Africa are home to this species of *Croton*.^[13] The plant material was gathered from Nigeria, which is said to be the primary location for *Crotonzambesicus*.^[15]

***CROTON HIRTUS* L'Her., Stirp.**

The annual herb *Crotonhirtus*L'Her., Stirp. reaches a height of approximately 60 cm.^[13] It is famous across the West Indies and tropical America, has recently been documented in Nigeria, Sierra Leone, Cote d'ivoire, and Benin, and is now naturalized throughout the tropics.^{[3];[4];[16]}

***CROTONLOBATUS* (Linn.)**

The base of this annual herbaceous plant turns woody. Senegal, Gambia, Guinea Bissau, and Nigeria are among their environments.^[13] It is referred to as "gaasayaa" in Hausa, "ajekofole" in Yoruba, and "Okwe-one" in Igbo. Traditionally, the boiled leaves are injected as a therapy for gynecological infections. When combined with palm oil, the leaves are applied topically to rheumatism and coastal discomfort. As a purgative, the leaf and bark infusion is administered orally or by injection.^[13] In Nigeria, it is used to lessen scorpion sting agony. They are used to treat headaches, skin conditions, and ulcers. In Togo (Prota), the leaf sap is applied as an eye drop. In the event of an abortion or hiccup, the flower and root decoction is given as an antispasmodic (Prota). A leaf infusion is used to alleviate fever, and when combined with honey and palm oil, it can be applied topically on stiff limbs (Prota).

***CROTONMACROSTACHYUS* (Hoscht.)**

Locally it is named as 'Bisana', in Amharic; 'Bakkaniisa', in Afan Oromo and popularly referred to as 'Rush slide' in English.^[17] In the tropics, it is the most prevalent plant.^[85] It is a medium-sized deciduous tree that is indigenous to Ethiopia, Eritria, Kenya and Nigeria.^[18] In addition, it is found in Liberia, Malawi, Zambia, and Zimbabwe. Moreover, Congo (DR) and Angola are home to it infrequently.^{[19];[4]} The bark, fruits, leaves, roots, and seeds of *Crotonmacrostachyus* are said to have a variety of medicinal qualities and are used as herbal remedies for at least 61

Comment [AW8]: Move it before you discuss about the type of chemicals in croton and after the point that you discussed the diseases treated by croton prior to the development of modern medicine.

Comment [AW9]: Elaborate this paragraph. Rearrange you better start the case of the world and come to the case of Africa finally close with the case in Nigeria. To have good flow you better include list of croton species that you discuss in the next paragraphs.

Comment [AW10]: You don't need to capitalize the species name except the begging letter.

Comment [AW11]: Instead of putting this variety under separate subtitle it is better to include the issue raised under the subtitle of the ***CROTONGRATISSIMUSBURCH.*** Just by mentioning the presence of this variety in the species.

Comment [AW12]: What is it? Is it part of the species name? Use symbol on top of Her to use the right phonetics. ***Croton hirtus*L'Hér**

Comment [AW13]: Elaborate ethno medicinal role of the species.

Comment [AW14]: Replace by L. to be consistent.

Comment [AW15]: Replace by dominant or conspicuous.

illnesses and ailments.[17] Blood clots, cancer, constipation, diarrhea, epilepsy, malaria, stomach discomfort, typhoid and wounds are among the conditions for which it has a long medicinal application. [20];[17] Crushed leaves of *Crotonmacrostachyus* are used to treat hemorrhoids, and the shoots are used to treat fever and edema. Its stem bark is macerated and used as a uterotonic to discharge retained placenta and as an abortifacient.[21] A decoction, infusion, or maceration of its leaves, root bark, and stem bark is used as a vermifuge and purgative throughout Madagascar and tropical Africa. [22];[4] To prevent bloating, its powdered roots are consumed with milk.[4] Ethiopian indigenous people compress the roots and fruits to make a water infusion that is used to treat venereal infections. [23];[4] The Nandi and Kikuyu community of Kenya utilized the leaf juice to heal wounds and the root and leaf decoctions to treat malaria. [24];[4] The plant's decoction, which is made from several parts, is used to cure female infertility, constipation, and stomachaches. An infusion of the bark is used to cure rheumatism and chest issues in East Africa. [4];[25]

CROTONMEMBRANACEUS Müll. Arg.

West tropical African countries including Ghana, Cote d'Ivoire, and Nigeria are home to this perennial plant. [12];[4] Urinary retention brought on by measles and an enlarged prostate can be cured with its root bark.[12] The herb is used as a treatment for a number of gastrointestinal issues by the Yoruba people of Nigeria. [26];[4]

CROTONNIGRITANUS

The shrub *Crotonnigritanus* can reach a height of three meters.[13] Senegal, Benin, Guinea, Nigeria, and west tropical Africa are among its locations.[12] In the past, the plant was reportedly collected from the wild and used to make regional medicines. In Sierra Leone's Scarcies River region, the herb is applied as a compress to wounds. [13]

CROTONPENDULIFLORUS Hutch

It resembles a spreading-crowned tree. West tropical Africa, including Gabon, Nigeria, Sierra Leone, and the Central African Republic, are homes to it.[12] In Ghana, the leaf infusion is used to treat fever, while in Cote d'Ivoire, it is used to treat menstruation abnormalities.[12] Its seeds contain oil that is used as a purgative (Proto). The seed extract is used to treat gastrointestinal issues and uterine cancers in Nigeria. [4];[26]

CROTONPSEUDOPULCHELLUS Pax

East Kenya, Cote d'Ivoire, Burkina Fasso, Nigeria, and Ethiopia are the primary locations for this *Croton* genus in tropical Africa. [4];[12] The leaves are traditionally applied to the chest as a medicine for chest ailments and used as a treatment for ulcers.[13] The decoction of the roots is used to treat asthma. To treat a head cold, the powdered root is consumed as snuff.[13] Burning *Crotonpseudopulchellus* and using the smoke to flavor fresh milk is a common condiment in Kenya's coastal regions.[27] Additionally, it is utilized to treat tussive and viral infections. [4];[28] In South Africa's central and southern regions, this *Croton* species is used to treat TB symptoms like fever, coughing, and blood in sputum. [11];[29]

CROTONSYLVATICUS (Hochst.)

Tropical regions of Africa, including, Tanzania, and Kenya, are home to this ornamental tree, which grows quickly.[12] Guinea and southern Nigeria are unusual places to find it.[4] Traditional remedies for TB, fever, digestive issues, and stomach pain in Tanzania and Kenya include decoctions of leaves and root bark.[11] Additionally, it is used as a purgative to reduce inflammation and as a malaria wash.[30] Elephantiasis is treated with *Crotonsyvaticus* wood shavings.[13] Ear infections are treated with the juice of young leaves.[12] Rheumatism, gastrointestinal disorders, dropsy, and uterine issues are all treated with the bark decoction.[12] On swellings, the ground roots are applied as a poultice.[30] To relieve excruciating

Comment [AW16]: But your title mentioned Nigeria. If the species is unusual in Nigeria there is no need to consider this particular species in this article.

stomachaches, the ground bark is soaked in milk.^[4] In Gazaland (present-day Zimbabwe and Mozambique), the bark powder is also used as a fish poison and applied topically on scarifications to alleviate rheumatism, chest pains, and inflammation.^[4] Additionally, cattle gallsickness is treated with the powdered bark.^[32] The root is used to cure pleurisy and as a therapy for indigestion. Malaria is also treated with several plant parts.^{[4];[33]}

CROTONZAMBESICUS Mull. Arg.

Sierra Leone, Mali, Dahomey, Niger, Gambia, and Nigeria are home to these shrubs or tiny trees.^[13] In Sierra Leone and Nigeria, *Crotonzambesicus* is used as a traditional medicine for a variety of purposes. The roots are used as an aperient, and the leaf decoction is used as a fever wash. Convulsions and dysentery are treated internally with it.^[4] In Benin, the leaf decoction is also used to treat urinary tract infections and lower blood pressure.^[34] The Ibibios of Nigeria utilize the root as an antidiabetic and antimalarial, while the Sudanese use it to relieve menstrual pain.^{[4];[34];[35]}

PHYTOCHEMISTRY AND PHARMACOLOGY OF NIGERIAN CROTON

Many research work has been carried out on various species of *Croton* plants which yields the isolation of terpenoid, diterpenoids, triterpenoids, non-terpenoid such as alkaloids, flavonoids, lignans.^[36] Isolation of megastigmane glycosides.^[37] Phytochemical analysis of the leaves and stems of *C. gratissimus* revealed the presence of alkaloids, amino acids, phenolic compounds, flavonoids, carbohydrates, terpenoids, saponins and fixed oils and fats.^[38] *Croton gratissimus* burchwas reported to predominantly yielded cembranoditerpenoids including a new cyclo-cebranoid skeleton (1).^[39] α -Glutinol(2), lupeol (3), eudesm-4(15)-ene-1 β ,6 α -diol (4) and 24-ethylcholesta-4,22-dien-3-one (5) have also been reported from *C. gratissimus*.^{[10];[39]} Isolated flavonoids (6–11), 3-methoxy-4-hydroxybenzoic acid (12), and benzyltetrahydroisoquinoline alkaloids laudanin (13) and laudanin (14) from *C. gratissimus*. [40] isolated fourteen compounds such as caryophyllene oxide (15), 1 β -methoxycaryolan-9 β -ol (16), kaur-16-en-19-oic acid (17), *cis*-ozic acid (18), spathulenol (19), lupeol (3), 7 δ -methoxyopposit-4(15)-en-1 β -ol (20), germacra-4(15),5,10(14)-trien-1 β -ol (21), β -sitosterol (22), *ent*-kaur-16-en-18-ol (23), 15-methoxy-*neo*-clerodan-3,13-dien-16,15-olide-18-oic acid (24), 6 α -methoxyeudesm-4(15)-en-1 β -ol (25), sucrose (26) and N-methyl-*trans*-4-hydroxy-L-proline (27) from *Crotongratissimus*.^[40] [41] Also reported the presence of (-)-(1*R**,4*R**,10*R**)-4-methoxycembra-2*E*,7*E*,11*Z*-trien-20,10-olide (28), (-)-(1*S**,4*R**,10*R**)-1-hydroxy-4-methoxycembra-2*E*,7*E*,11*Z*-trien-20,10-olide (29), (-)-(1*S**,4*S**,10*R**)-1,4-dihydroxycembra-2*E*,7*E*,11*Z*-trien-20,10-olide (30), (-)-(1*S**,4*S**,10*R**)-1,4-dihydroxycembra-2*E*,7*E*,11*Z*-trien-20,10-olide (31), (+)-(10*R**)-cembra-1*E*,3*E*,7*E*,11*Z*,16-pentaen-20,10-olide (32), (+)-(10*R**)-cembra-1*Z*,3*Z*,7*E*,11*Z*,15-pentaen-20,10-olide (33), (+)-(5*R**,10*R**)-5-methoxycembra-1*E*,3*E*,7*E*,11*Z*,15-pentaen-20,10-olide(34), (+)-(1*S**,4*S**,7*R**,10*R**)-1,4,7-trihydroxycembra-2*E*,8(19),11*Z*-trien-20,10-olide (35), (-)-(1*S**,4*S**,7*S**,10*R**)-1,4,7-trihydroxycembra-2*E*,8(19),11*Z*-trien-20,10-olide (36) and (+)-(1*S**,4*R**,8*S**,10*R**)-1,4,8-trihydroxycembra-2*E*,6*E*,11*Z*-trien-20,10-olide (37) in the leaves of *C. gratissimus*. Four flavonoids, kaempferol (38), isovitexin (39), helichryroside-3-methyl ether (40) and tiliroside (41) was found in the leaf extract of *C. gratissimus* var. *gratissimus*.^[42] *Crotongratissimus* var. *subgratissimus* was reported to contain α -phellandrene (42), germacrene D (43), and 1,8-cineole (44) and contains antimicrobial property.^{[43];[44]} In *Crotongratissimus* var. *gratissimus*, the hexane, ethyl acetate, butanol and 20% aqueous methanol extracts showed weak antioxidant capacity and acetyl cholinesterase (AChE) inhibitory effects.^{[4];[15]}

Crotonhirtus L. found in Costa Rica are reported to have bisnor-[15-16]-13 α -hydroxy-2-oxodolabra-1(10)-3-diene (45), 15,16-dihydroxy-2-oxodolabradan-3-ene (46), 16-hydroxy-2,15-

dioxodolabradan-3-ene (47), twelve kaurane diterpenoids; 16 α ,17-dihydroxy-7-oxokaurane (48), 6 β ,16 α ,17-trihydroxy-7-oxokaurane (49), 3 α ,16 α ,17-trihydroxy-7-oxokaurane (50), 3 α ,6 β ,16 α ,17-tetrahydroxy-7-oxokaurane (51), including two cyclopropakauranes, 7 β ,11 β ,16 α ,17-tetrahydroxycyclo-[3,18]-kaurane (52), 16 α ,17-dihydroxy-7-oxocyclo-[3,18]-kaurane (53), the hirtusanes, 16 α ,17-dihydroxy-7-oxohirtusan-3-ene (54), 16 α ,17-dihydroxy-7-oxohirtusan-3,5-diene (55), 3 β ,4 β -epoxy-16 α ,17-dihydroxy-7-oxohirtusan-5-ene (56), 3 α ,4 α -epoxy-16 α ,17-dihydroxy-7-oxohirtusan-5-ene (57), 3 β -H-16 α ,17-dihydroxy-7-oxohirtusan-4(19)-ene (58), 3 β ,16 α ,17-trihydroxy-7-oxohirtusan-4(19)-ene (59), five germacradiene esters, 6 β ,8 α -dihydroxy-8-*O*-benzoylgermacra-1(10)-*Z*,4*E*-dien-14-oic acid (60), 6 β ,8 α -dihydroxy-6-*O*-acetyl-8-*O*-benzoylgermacra-1(10)-*Z*,4*E*-dien-14-oic acid (61), 6 β ,8 α -dihydroxy-8-*O*-benzoylgermacra-1(10)*E*,4*E*-diene (62), 6 β ,8 α -dihydroxy-6-*O*-acetyl-8-*O*-benzoyleleman-1(2),3(4)-dien-14-oic acid (63), 6 β ,8 α -dihydroxy-8-*O*-benzoyleleman-1(2),3(4)-dien-14-carboxyl- γ -lactone (64).^[36] Isolated (-)-5,8-dihydroxyjatrophon-3-one (65) and (+)-14,16,17-trihydroxykauran-1-one (66) from the root of *Croton hirtus* occurring in Malaysia. The main compounds found in the oil of the leaves from *Croton hirtus* collected at Simões were spathulenol (19), E-caryophyllene (67), bicyclogermacrene (68), α -cadinol (69) and cubenol (70).^[45] Dihydro- β -ionol-*O*-[arabinosil(1-6) glucoside] (71), dihydro- β -ionol-*O*-[arabinosil(1 \rightarrow 6) glucoside] (72), β -sitosterol (22) and isorhamnetin-3-*O*-rutinoside (73) was recently isolated.^[46] The essential oil of *Croton hirtus* of Ivory Coast shows the presence of terpene derivatives, monoterpenes and sesquiterpenes.^[47] The toxicity of the essential oils found in *Croton hirtus*, showed 50% lethal concentration. *C. hirtus* extract prevents NO-mediated inflammation by suppressing NF- κ B and inflammatory cytokines. The methanolic extracts presented the highest 1,1-diphenyl-2-picrylhydrazyl (DPPH), 2,20-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS), and ferric reducing antioxidant power (FRAP) values.^{[45],[48],[46]} Its water, methanol and ethyl acetate extracts exhibited inhibitory effects on acetylcholinesterase (AChE) and butyrylcholinesterase (BChE), with a higher activity observed for dichloromethane, while the methanol extract showed the highest impact against tyrosinase.^[46]

Brazilian sample of *Croton lobatus* was reported to contain tertiary and quaternary alkaloids and hemolytic saponins.^{[49],[50]} Compounds such as diterpenes; geranylgeraniol (74), triglyceride lobaricidic acid (75) and triterpenes; betulinic acid (76) have been isolated from the stems and leaves of *Croton lobatus* (Prota). In *Croton lobatus*, betulinic acid (76) is said to be a potent HIV-1 antiviral compound and research have showed that *Croton lobatus* inhibits the growth of plasmodium falciparum and geranylgeraniol (74) induces apoptosis in leukemia cell lines activity against strains that are sensitive to chloroquine as well as resistant ones (Prota), it is said to contain higher amount of anti-oxidant phytochemicals.^[51] Its methanol leaf extracts retards arteriogenic risk factors therefore it can be used as herbal therapy for the treatment of diabetes mellitus and associated cardiovascular complication.^[51] The ethanol leaf extracts of *Croton lobatus* shows the gastro-protective potential in albino rats.^[52] Aqueous leaf extract of *C. lobatus* aids in the prevention of threatened abortion, management of pregnancy and infertility.^[53] Tannins, triterpenoids, and saponin are reported to be responsible for its antimicrobial activity.^[54] *Croton lobatus* leaf extracts could be used in diabetes and gout treatments, based on the antioxidant results of α -amylase inhibition and xanthine oxidase.^[55] Methanol leaf extract of *Croton lobatus* possesses significant analgesic and anti-inflammatory activities.^[56]

Croton macrostachyus is rich in terpenoids (diterpenoids and triterpenoids) and essential oils that contain monoterpenoids, sesquiterpenoids, and some shikimate-derived compounds. Previous studies showed the existence of croton (a chalcone) (77), lupeol (3) (a triterpene), crotopoxide (78) (a cyclohexanediepoxy), proteins, fatty acids, saponins, resins and

alkaloids.^[57] Some of the compounds isolated from *C. macrostachyus* include crotomacrine (**79**), halim-5,10-en-19,6 β ;20,12-diolide (**80**), floridolide (**81**), a labdane; crotomachlin (**82**) and four trachylobane diterpenoids; trachyloban-19-oic acid (**83**), trachyloban-18-oic acid (**84**), 3 α ,18,19-trihydroxytrachylobane (**85**), 3 α ,19-dihydroxytrachylobane (**86**) as well as four triterpenoids; betulin (**87**), a derivative of betulinic acid (**76**), acetyl aleuritolic acid (**88**), and sitosterol palmitate (**89**). *Cis*-clerodane (**90**) and 3 β -Acetoxy tetraer-14-en-28-oic acid (**91**).^{[17];[58]} The essential oil extracted from the leaves confirm the presence of 69.16% terpenoids; Germacrene D (**43**), caryophyllene (**67**), 1-methyl-4-(6-methyhept-5-en-2-yl) cyclohexa-1, 3-diene (**92**), β – Capaene (**93**), β – Pinene (**94**), linalool (**95**) and α – Copaene (**96**) along with ester benzyl benzoate (**97**), hydrocarbons; Naphthalene (**98**) and Cyclododecane (**99**), heterocyclic compounds; Indole (**100**) and Piperidine (**101**), fatty acid; Hexadecanoic acid (**102**), and amine; Phenylephrine (**103**).^[65]

Pharmacological studies shows that *Crotonmacrostachyus* has a wide range of pharmacological effects such as anti-diarrhea, and sedative, antidiabetic, anti-inflammatory, antileishmanic and larvicidal effects. *Crotonmacrostachyus* stem bark extract is active compared to studies in which anti-plasmodial activity has been implicated in a number of classes of secondary plant metabolites, including alkaloids and sesquiterpenes, inonoids, and quassinoids.^{[66];[17]} *CrotonMacrostachyus* is effective against diarrhea.^[67] The chemical constituent in *Crotonmacrostachyus*; terpenoids such as abietic acid (**104**) and steroids such as phytosterols (**105**) have been shown to inhibit the production of prostaglandin E2, which plays a crucial role in stimulating intestinal secretion and hence it has antidiarrheal activities.^{[68];[69];[17]} The anthelmintic activity, the analgesic and anti-inflammatory effects of the aqueous and methylene chloride/methanol stem bark extracts, the antimicrobial and antifungal activities of the methanol and dichloromethane extracts of the leaves and stem, and the antibacterial and antileishmanic activities of the plant's essential oils were all confirmed by numerous pharmacological studies of *C. macrostachyus*, according to the report.^[58] With minimum inhibitory concentration values of 3.75 mg/ml and 7.5 mg/ml, respectively, the methanol extract of *C. macrostachyus* demonstrated a strong effect in inhibiting the growth of tested isolates in both in vitro and in vivo settings. It also demonstrated stronger antibacterial activity against *S. aureus* than *E. coli*.^[59]

The root bark of *Crotonmembranaceus* contains scopoletin (**106**) and julocrotine (**107**) (glutimiride Alkaloid). Traces of calcium oxalate crystals were also reported in its root bark.^[12] Furano-clerodanediterpenoid; crotomembranafuran (**108**), labdanediterpenoid; gomojoside H (**109**), sitosterol; sitosterol 3-*O*- β -*D*-glucoside (**110**) and DL-threitol (**111**) are reported to have been isolated from its root extract.^{[4];[60]} According to reports, *Crotonmembranaceus* exhibits antibacterial activity. Certain phase I metabolizing enzymes are induced and inhibited by *Crotonmembranaceus*, whereas phase II metabolizing enzymes are modestly induced.^[61] According to in vitro studies, *C. membranaceus*'s aqueous and organic stem extracts both exhibit some antioxidant properties.^[62]

The seed extract of *Croton penduliflorus* shows the presence of fatty acid. Isolated from the root bark of West African croton *penduliflorus* is a halimanediterpenoid; Penduliflaworonsin (**112**).^[4] In addition, julocrotine (**107**) and lupeol (**3**) was also reported from the West African *C. penduliflorus*.^[63] Although essential oils derived from *C. penduliflorus* seeds have been shown to be hypocholesterolemic, they may also increase the risk of anemia. Investigation of the gastrointestinal effects and acute toxicity of the essential oils extracted from *C. penduliflorus* seeds, revealed that the oil caused inflammatory reactions in the colon, ileum, and jejunum, as well as hypersecretions in every section of the gastrointestinal tract. Of the visceral organs, the

liver, lung, and myocardium were the most affected.^{[70][71]} It has been shown that the methanolic extract of *C. penduliflorus* seeds increases the concentrations of albumin, total protein, sucrose, and maltase in pregnant rats.^[72] The chicks' foot pad oedema was considerably reduced by the hydro-ethanolic extract of *C. penduliflorus* stem bark.^[73] In the agar-well diffusion method, the stem bark extract of *C. penduliflorus* shown activity against the Gram-negative bacterium *P. aeruginosa* but not *E. coli*, and against the Gram-positive bacteria *S. aureus*, *K. pneumoniae*, *E. faecalis*, and *B. subtilis*.^[73]

The leaves of *Crotonpseudopulchellus* are found to contain taxalbumincrotonin (**113**).^{[13];[74]} Seven kauranediterpenoids, a labdane (**114-120**), three sesquiterpenoids (**36**, **121-122**), triterpenoids (**88,123**) and stigmaterol (**124**).^{[4];[74]} Phytochemical investigation of the root bark extract of *Croton pseudopulchellus* led to the isolation of five secondary metabolites namely, 18-methoxycarbonyl-18-methoxycarbonyl-15,16-epoxy-*ent*-cleroda-3,13(16),14-triene-20,19-olide (megalocaroidolide B) (**125**), 7,8-dehydrocrotonocorylifuran (**126**), vitexin (**127**), lupeol (**3**) and acetyl aleuritic acid (**128**).^[75] The acetone extract of *C. pseudopulchellus*' aerial parts had a minimal inhibitory concentration (IC₅₀) of 0.1 mg/mL against *Mycobacterium tuberculosis*, and a chloroform extract of the stem bark of the plant exhibited 82% minimum inhibitory activity at 50 mg/mL against PfUP1, a strain of *Plasmodium falciparum* that is resistant to chloroquine, and a minimum inhibitory concentration (IC₅₀) of 3.45 mg/mL against the kidney cells of vervet monkeys.^{[4];[28];[74]} Vitexin (**127**) showed antimicrobial activities with minimum inhibitory concentration and minimum microbicidal concentration values ranged between 16 and 32 µg/mL and interesting antioxidant properties very close to those of vitamin C and butyl hydroxyl toluene (BHT).^[75]


The seed of *Croton sylvaticus* is said to contain an oil composed of palmitic acid, stearic acid, linoleic acid and tiglic acid. Other research shows the presence of glutarimide alkaloid; julocrotine (**107**), lupeol (**3**) and penduliflaworonsin (**112**) in the stem bark and tannins in its bark.^[12] The roots are reported to contain the toxalbumincrotonin; a glycoprotein molecule that is attached to crotonin, a dihydrochalcone (**130**) and glycosylated protein.^[32] Hydrodistillation of the leaves is reported to have shown the presence of over fifty-two components including caryophyllene oxide (**15**) and α -humulene-1,2-epoxide (**131**).^[76] β -sitosterol (**22**), stigmaterol (**125**) and a clerodanediterpenoid, hardwickiic acid (**132**) was been reported from the leaves of East African *C. sylvaticus*.^{[4];[76]} Three labdane diterpenoids namely; 18-*nor*-labd-13(*E*)-ene-8 α , 15-diol (**133**), labd-13(*E*)-ene-8 α , 15-diol (**134**) and austroinulin (**135**) were isolated from the stem bark of *Croton sylvaticus*.^[77] Hydrodistillation from *C. sylvaticus* leaves shows the presence of β -caryophyllene oxide (**15**) and α -humulene-1,2-epoxide (**131**) as the major constituents.^{[30];[31]} Three compounds, namely 2-[N-(2-methylbutanoyl)]-N-phenyl-ethylglutarimide (**107**), lup-20(29)-en-3b-ol (**3**) and *ent*-(12*R*)-methyl-15.16-epoxy-9.10-friedolabda-5(10),13(16),14-trien-19-oate-20,12 lactone (**112**) were isolated from the leaves of *C. sylvaticus*.^{[17];[58]} The phytochemical investigation of the leaf extracts of *C. sylvaticus* yielded a clerodanediterpenoids namely sylvaticinol (**136**) and 3-hydroxy-3-((*Z*)-4-hydroxy-but-1-enyl)-2,2,4-trimethyl-cyclohexanone (**137**).^[41] Further phytochemicals analysis of the stem bark and leaves of *C. sylvaticus* yielded *trans*-phytol (**138**), lupenone (**139**), 3 β -acetoxylup-20(29)-ene (**140**), β -amyrin (**141**), lignoceryl *trans*-ferulate (**142**) and (+)-syringaresino (**143**).^{[74][78]} Isolation of 3,3,4,5,7-pentahydroxyflavone (**144**) and 3,4,5,7-tetrahydroxyflavone (**145**) from methanol leaf extracts of *C. sylvaticus* was carried out.^[41]


Previous research on *Croton sylvaticus* shows the presence of antiplasmodial activity, low to high toxicity and the inhibitory of acetyl cholinesterase.^[11] The aqueous methanol extracts showed anti-inflammatory and anti-oxidants activities. Both the water and methanol extracts of

C. sylvaticus exhibited very promising 5-lipoxygenase inhibitory activity.^{[12];[79]} Hardwickii acid (**132**) showed a significant antileishmanial activity on *L. donovani* promastigotes.^[80] The crude extract (1:1 MeOH in CH₂Cl₂) was found to be active at the tested concentration of 10 µg/ml exhibiting cell inhibition of 86 % against drug sensitive leukemia cell.^[77]


In *Croton zambesicus*, five flavone-C-glycosides, vitexin (**128**), orientin (**146**), vicenin-1 (**147**), saponaretin (**148**) and iso-orientin (**149**).^[81] The leaves and bark of Benin and Cameroon *Croton zambesicus* was reported to contain diterpenoids belonging to clerodane (**150-151**) labdane (**152**), kaurane (**153**).^{[63];[64]} Trachylobane (**154-159**).^[82] Pimarane (**160-161**).^[64] Phytol (**162**) classes.^[4] Other compounds reported from *C. zambesicus* include lupeol (**3**), betulinic acid (**76**), betulin (**87**) and sitosterol glycoside (**110**). Monoterpenes, sesquiterpenes, and aliphatic chemicals are present in the essential oils extracted from *Croton zambesicus* leaves.^[4] The antibacterial properties of *C. zambesicus* stem bark have been investigated. Rats subjected to pyrethroid-based insecticides showed improved testicular health after consuming *Croton zambesicus* leaf extracts.^[80] Rats' fecal production increased significantly and dose-dependently when they were given *Croton zambesicus* root extract.^[16] Benin uses *Croton gratissimus* burch. as an anti-malarial, anti-hypertensive, and antimicrobial to treat urinary tract infections.^[65] *Croton gratissimus* leaf extract has positive effects on the immune system, liver, and kidney, making it a potential treatment for hypertension. *Croton gratissimus* extracts have been shown to have potent antibacterial property.^{[4];[7];[83]} *Croton gratissimus* showed strong antioxidant properties without any toxicity.^[84] The methanolic leaf and stem extracts showed significant activity against a variety of bacterial isolates.^[38]

STRUCTURES







- 6** R₁ = OMe, R₂ = OH, R₃ = OMe, R₄ = OMe
7 R₁ = OMe, R₂ = OMe, R₃ = OH, R₄ = OMe
8 R₁ = OMe, R₂ = OMe, R₃ = OMe, R₄ = OMe
9 R₁ = OMe, R₂ = OH, R₃ = OH, R₄ = OMe
10 R₁ = OMe, R₂ = OMe, R₃ = OH, R₄ = OH




11




12




13 OH
14 OMe




15




16




17







18












19




20




28 $R_1 = \text{OCH}_3$, $R_2 = \text{H}$
 29 $R_1 = \text{OCH}_3$, $R_2 = \text{OH}$
 30 $R_1 = \text{OH}$, $R_2 = \text{OH}$







43



44




45



46 R₁ = OH

47 R₁ = O




48 R₁ = R₂ = H

49 R₁ = H, R₂ = OH

53 R₁ = H, R₂ = O

50 R₁ = OH, R₂ = H

51 R₁ = R₂ = OH




52 R₁ = OH, R₂ = β-OH

54

55(5, 6)


56(5,6) 3,4-epoxide

57(5,6) 3,4-epoxide



58 R₁ = H, α-CH₃


59 R₁ = OH, CH₃













60 R₁ = H, R₂ = CO₂H


61 R₁ = Ac, R₂ = CO₂H

62 R₁ = H, R₂ = CH₃




63







80



81



82




83 $R_1 = \text{CH}_3$, $R_2 = \text{CO}_2\text{H}$, $R_3 = \text{H}$

84 $R_1 = \text{CO}_2\text{H}$, $R_2 = \text{CH}_3$, $R_3 = \text{H}$


85 $R_1 = \text{CH}_2\text{OH}$, $R_2 = \text{CH}_2\text{OH}$, $R_3 = \text{OH}$

86 $R_1 = \text{CH}_3$, $R_2 = \text{CH}_2\text{OH}$, $R_3 = \text{OH}$












87













88







89

























111




- 114 $R_1 = \text{CO}_2\text{H}$, $R_2 = R_3 = R_4 = \text{H}$
115 $R_1 = \text{CHO}$, $R_2 = R_3 = R_4 = \text{H}$
116 $R_1 = \text{CO}_2\text{H}$, $R_2 = \text{OH}$, $R_3 = R_4 = \text{H}$
117 $R_1 = \text{CO}_2\text{H}$, $R_2 = \text{OCOCH}_3$, $R_3 = R_4 = \text{H}$
118 $R_1 = \text{CO}_2\text{H}$, $R_2 = \text{H}$, $R_3 = \text{H}$, $R_4 = \text{OH}$
119 $R_1 = \text{CO}_2\text{H}$, $R_2 = \text{H}$, $R_3 = \text{OH}$, $R_4 = \text{H}$












141




142




143














144





145



146





CONCLUSION

The genus *Croton* is a member of the *Euphorbiaceae*. They are found in tropical and sub-tropical areas of both hemispheres and comprise 1,300 species of trees, shrubs, and plants. More than 292 species of the genus *Croton* are known to exist in Africa, whereas 11 species of the genus including *Croton gratissimus*, *Croton gratissimus* var. *gratissimus*, *Lobatus*, *hirtus*, *membranaceus*, *macrostachyus*, *penduliflorus*, *pseudopulchellus*, *sylvaticus*, and *zambesicus* are found in Nigeria. Found all throughout the world, croton species are most commonly used in traditional medicine to cure conditions like cancer, constipation, diabetes, dysentery, external wounds, intestinal worms, discomfort, ulcers, and weight loss. Nigerian *Croton* species have many ethnomedical applications and contain a variety of bioactive chemicals, such as terpenes, alkaloids, and flavonoids, which give them pharmacological properties like antibacterial, antileishmanial, antiplasmodial, antioxidant, and anticancer properties.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

REFERENCES

- [1] Sofowora A. Recent trends in research into African medicinal plants. *Journal of Ethnopharmacology*, 1993; 38(2-3), 197–208.
- [2] Salatino A, Salatino ML, Negri G. Traditional uses, Chemistry and Pharmacology of *Croton* species (Euphorbiaceae). *J. Braz. Chem. Soc. J. Braz. Chem. Soc.*, 2007; 18(1):11-33.
- [3] Koutchiko HS, Attakpa SE, Akotegnon R, Guinnin F, Senou M, Amoussa AM, Baba-Moussa L. Acute toxicity profile of ethanolic extracts of *Croton gratissimus* Burch and *Schrankia leptocarpa* DC in rats: Medicinal plants used in the treatment of arterial hypertension in Beninese traditional medicine. *Journal of Physiology and Pathophysiology*, 2022; 13(1), 17-26.
- [4] Isyaka MS. The chemistry of African *Croton* specie, University of Surrey, 2020.
- [5] Amaral AC, Barnes RA. Clerodane diterpenoids from *Croton hemiargyreus*. *Natural Product Letters*, 1998; 12(1): 41-46.
- [6] Birhanu G. Effect of Seasonal Variations in the Chemical Composition of Essential Oil

- of the Leaves of *CrotonMacrostachyus*. Journal of Indigenous Knowledge and Development Studies. 2021; 3(1):17-24.
- [7] Segla KA, Eugène AS, Félix G, Maximin S, Madjid AA, Latifou L. Antioxidative Effects and Mechanisms of Antihypertensive Potential of *Crotongratissimus* Burch and *Schrankialeptocarpa* DC in Rats. Ann Hypertens. 2022; 2(1):1011.
- [8] Wu X, Zhao Y. Advance on chemical composition and pharmacological action of *Croton* L. Natural product research and development, 2004; 16(5), 467-472.
- [9] Xu WH, Liu WY, Liang Q. Chemical constituents from *Croton* species and their biological activities. Molecules, 2018; 23(9), 2333.
- [10] Mahmoud AB, Danton O, Kaiser M, Khalid S, Hamburger M, Mäser, P. HPLC-based activity profiling for antiprotozoal compounds in *Crotongratissimus* and *Cuscutahyalina*. Frontiers in pharmacology, 2020; 1246.
- [11] Njoya EM, Eloff JN, McGaw LJ. *Crotongratissimus* leaf extracts inhibit cancer cell growth by inducing caspase 3/7 activation with additional anti-inflammatory and antioxidant activities. BMC Complementary and Alternative Medicine. 2018; 18:305
- [12] The Ferns. Tropical.theferns.info/view tropical 2014.
- [13] Burkil HM. Royal botanical gardens; kew. 2004; www.Aluka.org
- [14] Palgrave KC. Trees of Southern Africa; Struik: Cape Town 2002.
- [15] Ndhlala A, Aderogba M, Ncube B, Van Staden J. Anti-oxidative and cholinesterase inhibitory effects of leaf extracts and their isolated compounds from two closely related *Croton* Species. Molecules.2013; 18, 1916–1932.
- [16] Ezeabara CA, Okonkwo EE, Comparison of phytochemical and proximate components of leaf, stem and root of *Crotonhirtus* L'Herit and *Crotonlobatus* Linn. J. Med. Health Res, 2016; 24, 33.
- [17] Abdisa T. Medicinal Value of *Crotonmacrostachyus* and *Solanumincanum* against Causative Agent of Foodborne Diseases. Veterinary Medicine–Open Journal. 2019; 4, 57-68.
- [18] Karmali MA, Gannon V, Sargeant JM. Verocytotoxin-producing *Escherichia coli* (VTEC). Vet Microbiol.2010;140(3-4): 360-370.
- [19] JSTOR. *Croton macrostachyus* Hochst. [family EUPHORBIACEAE]. 2019; https://plants.jstor.org/stable/10.5555/al.ap.upwta.2_97.
- [20] Khameneh B, Diab R, Ghazvini K, Bazzaz B. Breakthroughs in bacterial resistance mechanisms and the potential ways to combat them. MicrobPathog. 2016; 95: 32-42.
- [21] LeJeune JT, Besser TE, Hancock DD. Cattle water troughs as reservoirs of *Escherichia coli* O157. Appl Environ Microbiol. 2001; 67: 3053-3057.
- [22] Mazzanti G, Bolle P, Martinoli L, Piccinelli D, Grgurina I, Animati F, Mugnè Y. *Croton macrostachys*, a plant used in traditional medicine: purgative and inflammatory activity. J. Ethnopharmacol.1987; 19, 213–219.
- [23] Tefera M, Geyid A, Debella A. *In vitro* anti-Neisseria gonorrhoeae activity of *Albizia gummifera* and *Croton macrostachyus*. Pharmacol. Online.2012; 1, 75–83.
- [24] Jeruto P, Mutai C, Ouma G, Lukhoba C. An inventory of medicinal plants that the people of Nandi use to treat malaria. J. Anim. Plant Sci. 2011;3, 1192–1200.
- [25] Tane P, Tatsimo S, Connolly JD. Crotomacrine, a new clerodane diterpene from the fruits of *Croton macrostachyus*. Tetrahedron Lett. 2004; 45, 6997–6998.
- [26] Adesogan EK. The structure of penduliflaworosin, a new furanoid diterpene from *Croton penduliflorus*. J. Chem. Soc. Perkin Trans. 1981; 1(4):1151–1153.
- [27] Pakia M, Cooke JA, VanStaden J. The ethnobotany of the Midzichenda tribes of the

- coastal forest areas in Kenya: General perspective and non-medicinal plant uses. *South African J. Bot.* 2003; 69, 370–381.
- [28]Prozescy EA, Meyer JM, Louw AI. *In vitro* antiplasmodial activity and cytotoxicity of ethnobotanically selected South African plants. *J. Ethnopharmacol.* 2001; 76, 239–245.
- [29]Lall N, Meyer JJ. *In vitro* inhibition of drug-resistant and drug-sensitive strains of *Mycobacterium tuberculosis* by ethnobotanically selected South African plants. *J. Ethnopharmacol.* 1999; 66, 347–354.
- [30]Kapingu MC, Mbwambo ZH, Moshi MJ, Magadula JJ. Brine shrimp lethality of alkaloids from *Croton sylvaticus* Hoechst. *The East and Central African Journal of Pharmaceutical Sciences*, 2012; 15(2), 35-37.
- [31]Kapingu MC, Guillaume D, Mbwambo ZH, Moshi MJ, Uliso FC, Mahunnah RL. Diterpenoids from the roots of *Croton macrostachys*. *Phytochemistry*. 2000; 54(8): 767-770.
- [32]Watt JM, Breyer-Brandwijk MG. *The Medicinal and Poisonous Plants of Eastern Africa*, 2nd ed.; E. & S. Livingstone: Edinburgh 1962.
- [33]Beentje H. *Kenya Trees, Shrubs, and Lianas*; Nairobi, Kenya : National Museums of Kenya 1994
- [34]Okokon J, Ofodum K, Ajibesin K, Danladi B, Gamaniel K. Pharmacological screening and evaluation of antiplasmodial activity of *Croton zambesicus* against *Plasmodium berghei* infection in mice. *Indian J. Pharmacol.* 2005; 37, 243.
- [35]El-hamidi A. Drug Plants of the Sudan Republic in Native Medicine. *Planta Med.* 1970;18, 278–280.
- [36]Rosandy AR, Azman AA, Khalid RM, Othaman R, Lazim AM, Choudary IM, Bakar MA. New diterpenoids from the roots of *Croton hirtus* (*Euphorbiaceae*). *Malaysian Journal of Analytical Sciences*, 2018; 22(1), 64-71.
- [37]Kawakami S, Matsunami K, Otsuka H, Shinzato T, Takeda H. Crotonionosides A–G: Megastigmane glycosides from leaves of *Croton cascarilloides* Rauschel. *Phytochemistry*, 2011; 72: 147–153.
- [38]Naidoo D. Secretory structures of *Croton gratissimus* Burch. Var. *gratissimus* (*Euphorbiaceae*): Micromorphology and Histophytochemistry (Doctoral dissertation) 2018.
- [39]Langat MK, Crouch NR, Smith PJ, Mulholland DA. Cembranolides from the leaves of *Croton gratissimus*. *J. Nat. Prod.* 2011; 74, 2349–2355.
- [40]Canelo LI, Mafuca I, Mata, RS, Mendonça DI. Chemical Constituents of a Population of *Croton gratissimus* (*EUPHORBIACEAE*). *Química Nova*. 2017; 40, 1035-1038.
- [41]Langat M, Mulholland DA, Crouch NR. New diterpenoids from *Croton sylvaticus* and *Croton pseudopulchellus* (*Euphorbiaceae*) and antiplasmodial screening of ent-kaurenoic acid. *Planta Medica*, 2008; 74(09), PB126.
- [42]Ashwell RN, Aderogba MA, Bhekumthetho N, Van-Staden J. Antioxidative and cholinesterase inhibitory effects of leaf extracts and their isolated compounds from two closely related *Croton* species. *Molecules* 2013; 18:1916–1932.
- [43]VanVuuren S. The antimicrobial activity and essential oil composition of medicinal aromatic plants used in African traditional healing. Ph.D. thesis, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg 2007.
- [44]Lawal OA, Ogunwande IA, Osunsanmi FO, Opoku AR, Oyedeji AO.

- Crotongratissimus* leaf essential oil composition, antibacterial, antiplatelet aggregation, and cytotoxic activities. *Journal of Herbs, Spices & Medicinal Plants*, 2017; 23(1), 77-87.
- [45] De Lima SG, Medeirosa LP, Cunha NC, Da Silva D, De Andrade NC, Moita NJ, Lopesa JA, Steffen RA, Araújo BQ, De Reisb AM. Chemical composition of essential oils of *Crotonhirtus* L'Her from Piauí (Brazil). *The Journal of Essential Oil Research*. 2012; 24(4):371–376.
- [46] Dall'Acqua S, Sinan KI, Sut S, Ferrarese I, Etienne OK, Mahomoodally MF, Lobine D, Zengin G. Evaluation of Antioxidant and Enzyme Inhibition Properties of *Croton hirtus* L'Hér. Extracts Obtained with Different Solvents. *Molecules*. 2021; 26, 1902. <https://doi.org/10.3390/molecules26071902>
- [47] Daouda T, Prevost K, Gustave B, Joseph DA, Nathalie G, Raphaël O, Rubens D, Jean CC, Mireille D, Felix T. Terpenes, Antibacterial and Modulatory Antibiotic Activity of Essential Oils from *Croton hirtus* L' Hér. (Euphorbiaceae) from Ivory Coast. *TEOP*. 2014; 17(4):607 – 616.
- [48] Kim MJ, Kim JG, Sydara KM, Lee SW, Jung SK. *Crotonhirtus* L'Hér Extract Prevents Inflammation in RAW264. 7 Macrophages Via Inhibition of NF-κB Signaling Pathway 2020.
- [49] Farnsworth NR, Blomster RN, Messmer WM, King JC, Persinos GJ, Wilkes DA. phytochemical and biological review of the genus *Croton*. *Lloydia*, 1969; 32(1), 1-28.
- [50] Willaman JJ, Li HL. Alkaloid-bearing plants and their contained alkaloids 1970; 1957–1968.
- [51] Fasola TR, Ukwenya B, Oyagbemi AA, Omobowale TO, Ajibade TO. Antidiabetic and antioxidant effects of *Croton lobatus* L. in alloxan-induced diabetic rats. *J Intercult Ethnopharmacol*. 2016; 5(4):364–371.
- [52] Ezugwu HC, Anosike AC, Berinyuy EB, Dasofunjo K, Ezugwu MU, Achu HO. Effect of ethanol leaf extract of *Crotonlobatus* on indomethacin-induced gastric ulceration in albino rats 2018.
- [53] Enohor OO, Oshomoh EO. Uterotonic Evaluation of *CrotonLobatus* Linn On Reproductive Parameters of Wistar Rats. *SAU Science-Tech Journal*. 2020; 5(1), 75-82.
- [54] Kilani MA, Hassan AZ, Fadason ST, Obalowu AM, Aliyu A, Kilani, HB. In-vitro and in-vivo antibacterial effect of *CrotonlobatusLinnaeus* L. on two days post surgical wounds in rats. *Bangladesh Journal of Scientific and Industrial Research*. 2019; 54(2), 139-146.
- [55] Chodaton-Zinsou MD, Assogba FM, Yayi-Ladekan E, Gbaguidi F, Moudachirou M, Gbenou JD. Phytochemical Composition, Biological Activities of *Croton lobatus* L. Leaves, Hydrolysis Effect on Activities and Chemical Composition. *American Journal of Applied Chemistry*. 2020; 8(1):13-22. doi:10.11648/j.jac.20200801.13
- [56] Anafi S, Yaro A, Abbas-Yakubu M, Yakubu A. Evaluation of analgesic and anti-inflammatory activities of methanol leaf extract of *Crotonlobatus* (Euphorbiaceae) in rodents. *Trop J Nat Prod Res*. 2017; 1(6): 255-58.
- [57] Carlet J, Jarlier V, Harbarth S, Voss A, Goossens H, Pittet D. Ready for a world without antibiotics? The pensieres antibiotic resistance calls to action. *Antimicrob Resist Infect Control*. 2012; 1: 11.
- [58] Meresa A. Ethno medicinal uses, phytochemistry and anti-malarial effect of *Croton macrostachyus* (Bisana): A review. *Journal of Medicinal Plants*, 2019; 7(2), 79-88.
- [59] Aylate A, Agize M, Ekeru D, Kiros A, Ayledo G, Gendiche K. *In-Vitro* and *In-Vivo*

Antibacterial Activities of *Croton macrostachyus* Methanol Extract against *E. coli* and *S. aureus*. Adv. Anim. Vet. Sci. 2017;5(3): 107-114.

- [60] Bayor MT, Gbedema S, Annan K. The antimicrobial activity of *Croton membranaceus*, a species used in formulations for measles in Ghana. J. Pharmacogn. Phyther. 2009; 1, 47–51.
- [61] Asare GA, Ongong'a RO, Anang Y, Asmah RH, Rahman H. Effect of a Benign Prostatic Hyperplasia (BPH) Xenobiotic *Croton membranaceus* Müll. Arg. Root Extract on CYP1A2, CYP3A4, CYP2D6, and GSTM1 Drug Metabolizing Enzymes in Rat Model. 2020.
- [62] Afriyie DK, Ofori-Ameyaw E, Acheampong DO, Tuffour I, Appiah-Opong R. In-vitro and in-vivo antioxidant properties of stem extracts of *Croton membranaceus*. 2022.
- [63] Block S, Flamini G, Brkic D, Morelli I, Quetin-Leclercq J. Analysis of the essential oil from leaves of *Croton zambesicus* Muell. Arg. growing in Benin. FlavourFragr. J. 2006; 21, 222–224.
- [64] Block S, Brkic D, Hubert P, Quetin-Leclercq J. A validated method for the quantification of pimarane and trachyloban diterpenes in the leaves of *Croton zambesicus* by capillary gas chromatography. Phytochem. Anal. 2005;16, 342–348.
- [65] Block S, Stévigny C, De Pauw-Gillet MC, de Hoffmann E, Llabrès G, Adjakidjé V, Quetin-Leclercq J. Ent-trachyloban-3 β -ol, a new cytotoxic diterpene from *Croton zambesicus*. Planta Med. 2002;68, 647–649.
- [66] Kiranmayi CB, Krishnaiah N, Mallika EN. Escherichia coli O157:H7- An emerging pathogen in foods of animal origin. Vet World. 2010; 3(8): 382-389.
- [67] Burt S. Essential oils: their antibacterial properties and potential applications in foods: A review. Int J Food Microbiol. 2004; 94: 223-253.
- [68] Linscott A. J. Food-borne illnesses. Clinical Microbiology Newsletter. 2011; 33(6): 41-45.
- [69] Liu LF, Liang CH, Shiu LY, Lin WL, Lin CC, Kuo KW. Action of solamargine on human lung cancer cells—enhancement of the susceptibility of cancer cells to TNFs. FEBS Lett. 2004; 577(1-2): 67-74.
- [70] Ojokuku SA, Odesanmi OS, Magbagbeol OA. The effects of oral administration of *Croton penduliflorus* seed oil and medroxy progesterone acetate on fasting blood sugar, lipid and haematology of pregnant rabbits. Int. J. Trop. Med. 2011; 6, 35–38.
- [71] Asuzu IU, Chineme CN. Acute toxicity and gastrointestinal irritant effect of *Croton penduliflorus* seed oil in mice. Phyther. Res. 1988; 2, 46–50.
- [72] Sarwar M. Methanolic Extract of *Croton Penduliflorus* Affects Intestinal Enzyme Activity and Protein Content in the Late Phase of Pregnancy. International Journal of Pharmacology, 2011; 7(6):670-670.
- [73] Baah KA, Acheampong A, Amponsah IK, Jibira Y, Owusu MO, Tima LA. Evaluation of the Anti-inflammatory, Antimicrobial and Antioxidant activities of the stem bark extract of *Croton penduliflorus* Hutch (Euphorbiaceae). Journal of Pharmacognosy and Phytochemistry, 2017; 6(6): 1705-1710.
- [74] Langat MK. The phytochemistry of three African *Croton* species. University of Surrey (United Kingdom) 2009.
- [75] Tatsimo JS, Tamokou JD, Havyarimana L, Kengne IC, Ekom SE, Lannang AM, Sewald N. Antimicrobial and antioxidant activities of isolated compounds from the root bark of *Croton pseudopulchellus* (Euphorbiaceae). Invest Med Chem Pharmacol, 2020; 3(1), 39.
- [76] Mwangi JW, Thoithi GN, Addae-Mensah I, Achenbach H, Lwande W, Hassanali H.

Aromatic plants of Kenya III: Volatile and some non-volatile constituents of *Croton sylvaticus*. East Cent. African J. Pharm. Sci.1, 1998; 41–43.

- [77] Okerio KN, Kenanda EO, Omosa LK. Antiproliferative Properties of Labdane Diterpenoids from *Croton sylvaticus* Hochst against Drug Sensitive and Resistant Leukemia cell lines 2019.
- [78] Aderogba M A, Ndhala A R, Van-Staden J. Acetylcholinesterase inhibitory activity and mutagenic effects of *Croton penduliflorus* leaf extract constituents. South African Journal of Botany, 2013; 87, 48-51.
- [79] Frum Y, Viljoen AM. *In vitro* 5-lipoxygenase and anti-oxidant activities of South African medicinal plants commonly used topically for skin diseases. Skin Pharmacol. Physiol. 2005; 19, 329–335.
- [80] Crentsil JA, Yamthe LT, Anibea BZ, Broni E, Kwofie SK, Tetteh KA, Osei-Safo D. Leishmanicidal Potential of Hardwickiic Acid Isolated from *Croton sylvaticus*. Frontiers in pharmacology. 2020; 11, 753.
- [81] Wagner H, Hörhammer L, Kiraly IC. Flavon-C-glycoside in *Croton zambezicus*. Phytochemistry. 1970; 9, 897.
- [82] Ngadjui BT, Folefoc GG, Keumedjio F, Dongo E, Sondengam BL, Connolly JD. Crotonadiol, a labdanediterpenoid from the stem bark of *Croton zambezicus*. Phytochemistry. 1999; 51, 171–174.
- [83] Morobe IC, Obi LC, Oyedeji AO, Majinda RT, Hattori T, Idighe JE, Vasaikar SD. Isolation and biological investigation of bioactive compounds from *Croton grattissimus* (Burch) in Mthatha, Eastern Cape, South Africa. Nat. Prod. Chem. Res. 2018; 6, 1–7.
- [84] Ahamed AA, Adam YS, Hussien AM, Hassan ME. Effect of Heat Treatment on Antioxidant and Antimicrobial Activity of *Croton grattissimus* and *Xylopiiaethiopica* Spices 2021.
- [85] Käppeli U, Hächler H, Giezendanner N, Beutin L, Stephan R. Human infections with non-O157 Shiga toxin-producing *Escherichiacoli*, Switzerland, 2000-2009. Emerg Infect Dis. 2011; 17(2): 180-185.