

Overview on Gallstone ileus: An update on management

Abstract

Gallstone ileus is a rare complication of gallstone disease and is seen in less than 5% of cases of intestinal obstruction. The diagnosis is confirmed by a combination of clinical presentation and imaging modalities in the form of abdominal x-ray and computerized tomography which will show gas in the biliary system and dilated bowel. The management of gallstone ileus is by surgical therapy, which can be divided into an enterotomy alone or an enterotomy followed by excision of the cholecystic-enteric fistula and a cholecystectomy. As the choice of surgical therapy is tailor made to the patient, we have conducted this review article to look at the diagnosis and treatment of this condition. We have also looked at the various types of surgical therapy that is done for gallstone ileus.

Keywords- "Gallstone ileus", "Gallstone coelus", "enterotomy", "one-staged procedure", "two staged procedure", "laparoscopy" and "Intestinal obstruction".

Introduction

Gallstone ileus is a rare complication of gallstone disease that is seen in 0.3% to 0.5% of patients with cholelithiasis. It is seen in elderly and predominantly female patients, accounting for symptoms of acute intestinal obstruction. Gallstone ileus occurs when a gallstone is passed into the intestine by a fistulous tract that forms between the inflamed gallbladder and the duodenum, stomach and colon. The average size of the gallstones is 4cm and the most common site of obstruction is at the ileo-cecal valve, followed by the duodenum and colon(1,2). The clinical presentation is depended on the site of obstruction with abdominal pain, vomiting and constipation, as the symptoms progress, they may proceed to acute intestinal obstruction. The diagnosis may be obtained by performing imaging modalities like abdominal Xray which will show a radiopaque stone, air in the biliary system and dilated loops of bowel. This is referred to as the Rigler's triad, but these findings are only seen in 40% of cases, Computerized tomography is the gold standard in the diagnosis of gallstone ileus with a sensitivity of 90% (3).

The management of gallstone ileus can be divided into conservative treatment of intestinal obstruction which involves fluid resuscitation with intravenous fluids, keep the patients fasted,

starting intravenous analgesics and insertion of a Ryles tube. Surgical Intervention is the definitive treatment of choices for gallstone ileus. Surgical therapy can be divided into a one stage procedure which involves an enterotomy, excision of the cholecystic-enteric fistula and a cholecystectomy. This procedure though is associated with a high mortality rate and prolonged anesthesia and is reserved for patients who fit and with no co-morbidities. The two staged procedure involves performing an enterotomy and removal of the gallstone that is causing obstruction and the excision of the fistula and cholecystectomy is performed later as a second procedure once the patient's acute condition has improved. The two-stage procedure is associated with reduced mortality when compared to the one staged procedure(4–7).

Laparoscopy has been attempted in the surgical management of gallstone ileus but there are difficulties encountered with finding the site of obstruction, examining the entire length of the bowel, and performing the surgery in the presence of dilated bowel which prolongs the duration of surgery. Laparoscopy will require an advanced laparoscopic surgeon and in an advanced center that can treat these patients(8). Other methods that have been used include extra-corporeal shock wave lithotripsy, endoscopic shock wave lithotripsy but they are not popular in the management of gallstone ileus(9).

We have conducted this review article to investigate the diagnosis and management options in gallstone ileus. The role of abdominal x-ray and computerized tomography in the diagnosis of gallstone ileus is investigated. We have reviewed the single stage enterotomy and the two stage enterotomy, fistula excision and cholecystectomy. We conducted a literature review using PUBMED, Cochrane database of clinical reviews and Google scholar looking for clinical trial, observational studies, cohort studies systemic reviews, and meta-analysis from 1980 to 2024. We used the following keywords, “Gallstone ileus”, “Gallstone coleus”, “Enterotomy”, “One stage procedure “” Two stage procedure “;” laparoscopy” and “intestinal obstruction”. All articles were in English language only. Further articles were obtained by manual cross referencing of the literature. Case reports and studies with less than 10 patients and editorials were excluded. Adult male and female patients were included in this study. Pregnant patients and pediatric patients were excluded.

Diagnosis of gallstone ileus

Patients with gallstone ileus often present with symptoms of intestinal obstruction and the most common imaging modality that is performed first is an abdominal x-ray. The classical features that are **detected on x-ray include the presence of air in the biliary system, presence of a radio-opaque stone and dilated loops of small bowel. This is known as the Rigler's triad and is diagnostic of gallstone ileus, but it is seen in only 40% of patients.** Ultrasound of the abdomen may be used to detect any pathology in the biliary system like gallstones, but computerized tomography is the gold standard in the diagnosis of gallstone ileus. Computerized tomography can identify the site of obstruction, the presence of air in the biliary system and the site of the

cholecystic -enteric fistula. Computerized tomography has a sensitivity of 90%-93% and a specificity of 100% in the diagnosis of gallstone ileus. Magnetic resonance imaging can also be used in the diagnosis of gallstone ileus with a sensitivity of 97.7% but it is time consuming to performed and not routinely available in most hospitals and hence is not used(10–13).

Management of gallstone ileus

The management of gallstone ileus can be divided into initial conservative treatment with intravenous fluids, analgesics, antibiotics followed by definitive surgical treatment. Surgical treatment can be divided into the single stage procedure which involves performing an enterotomy alone to relieve the obstruction and the two-stage procedure that involves performing an enterotomy, excision and repair of the cholecystic-enteric fistula and a cholecystectomy(14). Enterotomy with stone extraction was compared with enterotomy, excision of the cholecystic-enteric fistula and cholecystectomy and the operative time, morbidity and mortality was reduced in the enterotomy group when compared to the group that underwent enterotomy, excision of the fistula and cholecystectomy(15–20). As most of the patients of gallstone ileus are elderly and have co-morbidities, the enterotomy and removal of the stone is the safest emergency procedure that is associated with better outcomes(21–24).

The single staged procedure which involves performing an enterotomy, excision of the cholecystic-enteric fistula and a cholecystectomy is considered a definitive procedure and should be performed in young patients with an American Society of Anesthesia (ASA) score of less than 3, and with reduced co-morbidities (like hypertension, heart disease and diabetes mellitus(25). There were no significant differences with regards to the morbidity and mortality rates between the single stage and two stage procedure in some centers(26). One of the important complications from the single stage procedure is the risk of leakage from the enterotomy site or from the site of fistula excision, hence the need to provide this procedure for patients who are fit and have a good American Society of Anesthesia(ASA) score(27).

A systemic review and meta-analysis was conducted on the management of gallstone ileus by Vadher et al. 10 studies with 293 patients were in patients this study, and this study showed that there was a reduction in the mortality in patients who underwent a single stage procedure when compared to the two stage procedure, but the morbidity and length of operative time was higher

in the single stage procedure when compared to the two stage procedure. This study than concluded that the single stage procedure had a good risk benefit ratio in terms of decreased mortality but the decision on which procedure to perform will depend on patient factors and the risk of surgery(28).

Laparoscopic management of gallstone ileus has been attempted but some of the problems that can be encountered include the lack of space due to the dilated bowel, the difficulty to identify the site of obstruction, removal of the gallstones and the need to perform bowel resection. The prolonged anesthesia and the effect of gas insufflation may influence the intra-operative management of elderly patients. Laparoscopic surgery may be offered in centers where the experience and services is available(29,30).

Table □ :Table showing the mortality rate between patients with gallstone ileus who underwent enterotomy alone versus enterotomy with fistula excision and cholecystectomy.

Study	Year	Study type	N=numbers	Mortality rate (enterotomy) (%)	Mortality rate (enterotomy +fistula excision cholecystectomy) (%)
Doko et al	2003	Retrospective study	30	9.1%	11.1%
Yakan et al	2010	Retrospective study	12	16.6%	33.3%
Alencastro et al	2013	Retrospective study	12	12.5%	25%
Mallipeddi et al	2013	Retrospective study	127	5.3%	7.1%

Recurrent gallstone ileus

The risk of recurrent gallstone ileus is reported at 5%-8% and is predominantly seen in patients who had undergone an Enterotomy. Most patients experienced recurrence of symptoms of gallstone ileus after six weeks of undergoing treatment. The most common site for impaction is still at the ileum or at the ileocecal valve. The management of recurrent gallstone ileus is with the single staged procedure which involves an enterotomy and removal of the stone followed by excision of the fistula, but the mortality is still high(31).A systemic review and meta-analysis was conducted by Alzerwi et al on recurrent gallstone ileus.50 studies with 56 patients were included in this study and 87.3% of the patients were women, the median time for recurrence was 20.5days,the region where the stone was lodged was similar and the mortality rate was 11.8%(32).

Bouveret's syndrome

Bouveret's syndrome is a variant of gallstone ileus which is characterized by gastric outlet obstruction secondary from an impacted gallstone. It is seen in 2% -3% of all cases of gallstone ileus and is predominantly seen in female patients. The diagnosis is usually confirmed by computerized tomography which has a sensitivity of 93% and a specificity of 100% (33). The management of Bouveret's syndrome can be divided into endoscopic and surgical treatment. Endoscopic treatment which involves the use of esophagogastroduodenoscopy and removal of the stone is difficult and associated with a high failure rate. Surgical treatment can be divided into performing an enterotomy and removal of the stone followed by a gastrojejunostomy if the duodenal wall is friable. This is the most common procedure that is performed. The second type of procedure involves performing an enterotomy, removal of the stone, followed by excision of the fistula and performing a cholecystectomy, but it is associated with a higher morbidity and mortality (34-36).

Endoscopic removal of the stone with a net or basket has been attempted in some centers and for stones that are larger than 2cm, mechanical, extra hydraulic, laser and extracorporeal shockwave lithotripsy can be attempted. Endoscopic therapy requires specialist equipment and training, and these are the drawback of this form of therapy (37,38).

Conclusion

Gallstone ileus is a rare complication of gallstone disease that is predominantly seen in elderly, female patients and the diagnosis is confirmed by performing a computerized tomography. The management of gallstone ileus initially involves fluid resuscitation, antibiotics and analgesics. The definitive therapy is surgical treatment, which can be divided into the single stage repair, which involves performing an enterotomy, excision of the cholecystic-enteric fistula and a cholecystectomy and the two-stage procedure which involves performing an enterotomy alone and the cholecystectomy and fistula excision later. The choice of surgical is often tailor made to the clinical presentation and co-morbidities of the patient. The experience of the surgeon and facilities that are available are also factors that will decide what sort of surgical procedure needs to be performed for this patient.

References

1. Alemi F, Seiser N, Ayloo S. Gallstone Disease: Cholecystitis, Mirizzi Syndrome, Bouveret Syndrome, Gallstone Ileus. *Surg Clin North Am.* 2019 Apr;99(2):231-244. doi: 10.1016/j.suc.2018.12.006. PMID: 30846032.
2. Inukai K. Gallstone ileus: a review. *BMJ Open Gastroenterol.* 2019 Nov 24;6(1):e000344. doi: 10.1136/bmjgast-2019-000344. PMID: 31875141; PMCID: PMC6904169.
3. Ploneda-Valencia CF, Gallo-Morales M, Rinchon C, Navarro-Muñiz E, Bautista-López CA, de la Cerda-Trujillo LF, Rea-Azpeitia LA, López-Lizarraga CR. Gallstone ileus: An overview of the literature. *Rev Gastroenterol Mex.* 2017 Jul-Sep;82(3):248-254. English, Spanish. doi: 10.1016/j.rgm.2016.07.006. Epub 2017 Apr 19. PMID: 28433486.
4. Abou-Saif A, Al-Kawas FH. Complications of gallstone disease: Mirizzi syndrome, cholecystocholedochal fistula, and gallstone ileus. *Am J Gastroenterol.* 2002 Feb;97(2):249-54. doi: 10.1111/j.1572-0241.2002.05451.x. PMID: 11866258.
5. Ayantunde AA, Agrawal A. Gallstone ileus: diagnosis and management. *World J Surg.* 2007 Jun;31(6):1292-7. doi: 10.1007/s00268-007-9011-9. Epub 2007 Apr 15. PMID: 17436117.
6. Dodgion C, de Moya M. Gallstone Ileus. *Emergency General Surgery: A Practical Approach.* 2019:165-73. Available from: http://link.springer.com/10.1007/978-3-319-96286-3_13
7. Saldinger PF, Itskovich A. Gallstone Ileus. *Difficult Decisions in Hepatobiliary and Pancreatic Surgery: An Evidence-Based Approach.* 2016:291-5. Available from: http://link.springer.com/10.1007/978-3-319-27365-5_25
8. Zaliakas J, Munson JL. Complications of gallstones: the Mirizzi syndrome, gallstone ileus, gallstone pancreatitis, complications of "lost" gallstones. *Surgical Clinics of North America.* 2008 Dec 1;88(6):1345-68. <https://doi.org/10.1016/j.suc.2008.07.011>
9. Beuran M, Ivanov I, Venter MD. Gallstone ileus--clinical and therapeutic aspects. *J Med Life.* 2010 Oct-Dec;3(4):365-71. PMID: 21254732; PMCID: PMC3019077.
10. Lassandro F, Gagliardi N, Scuderi M, Pinto A, Gatta G, Mazzeo R. Gallstone ileus analysis of radiological findings in 27 patients. *Eur J Radiol.* 2004 Apr;50(1):23-9. doi: 10.1016/j.ejrad.2003.11.011. PMID: 15093232.

11. Chang L, Chang M, Chang HM, Chang AI, Chang F. Clinical and radiological diagnosis of gallstone ileus: a mini review. *EmergRadiol*. 2018 Apr;25(2):189-196. doi: 10.1007/s10140-017-1568-5. Epub 2017 Nov 16. PMID: 29147883; PMCID: PMC5849656.
12. Ripollés T, Miguel-Dasit A, Errando J, Morote V, Gómez-Abril SA, Richart J. Gallstone ileus: increased diagnostic sensitivity by combining plain film and ultrasound. *Abdom Imaging*. 2001 Jul-Aug;26(4):401-5. doi: 10.1007/s002610000190. PMID: 11441553.
13. Yu CY, Lin CC, Shyu RY, Hsieh CB, Wu HS, Tyan YS, Hwang JI, Liou CH, Chang WC, Chen CY. Value of CT in the diagnosis and management of gallstone ileus. *World J Gastroenterol*. 2005 Apr 14;11(14):2142-7. doi: 10.3748/wjg.v11.i14.2142. PMID: 15810081; PMCID: PMC4305784.
14. Doko M, Zovak M, Kopljar M, Glavan E, Ljubicic N, Hochstädter H. Comparison of surgical treatments of gallstone ileus: preliminary report. *World J Surg*. 2003 Apr;27(4):400-4. doi: 10.1007/s00268-002-6569-0. PMID: 12658481.
15. Requena-López Abner A., Mata-Samperio Brenda K., Solís-Almanza Fernando, Casillas-Vargas Ricardo, Cuadra-Reyes Luis A.. Comparison between surgical techniques in gallstone ileus and outcomes. *Cir. cir. [magazine on the Internet]*. 2020 Jun [cited 2024 Dec 16] ; 88(3): 292-296. Available at: http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S2444-054X2020000300292&lng=es. Epub Nov 15, 2021. <https://doi.org/10.24875/ciru.19001264> .
16. de Alencastro MC, Cardoso KT, Mendes CA, Boteon YL, de Carvalho RB, Fraga GP. Acute intestinal obstruction due to gallstone ileus. *Rev Col Bras Cir*. 2013 Jul-Aug;40(4):275-80. English, Portuguese. doi: 10.1590/s0100-69912013000400004. PMID: 24173476.
17. Martínez Ramos D, Daroca José JM, Escrig Sos J, Paiva Coronel G, Alcalde Sánchez M, Salvador Sanchis JL. Gallstone ileus: management options and results on a series of 40 patients. *Rev EspEnferm Dig*. 2009 Feb;101(2):117-20, 121-4. English, Spanish. doi: 10.4321/s1130-01082009000200005. PMID: 19335047.
18. Muthukumarasamy G, Venkata SP, Shaikh IA, Somani BK, Ravindran R. Gallstone ileus: surgical strategies and clinical outcome. *J Dig Dis*. 2008 Aug;9(3):156-61. doi: 10.1111/j.1751-2980.2008.00338.x. PMID: 18956594.
19. Scuderi V, Adamo V, Naddeo M, Di Natale W, Boglione L, Cavalli S. Gallstone ileus: monocentric experience looking for the adequate approach. *Updates Surg*. 2018 Dec;70(4):503-511. doi: 10.1007/s13304-017-0495-z. Epub 2017 Oct 10. PMID: 29019098.

20. Gungor F, Atalay Y, Acar N, Gur EO, Kokulu I, Acar T, Karasu S, Dilek ON. Clinical outcome of gallstone ileus; a single-centre experience of case series and review of the literature. *ActaChir Belg*. 2022 Feb;122(1):7-14. doi: 10.1080/00015458.2020.1816673. Epub 2020 Sep 8. PMID: 32857668.
21. Halabi WJ, Kang CY, Ketana N, Lafaro KJ, Nguyen VQ, Stamos MJ, Imagawa DK, Demirjian AN. Surgery for gallstone ileus: a nationwide comparison of trends and outcomes. *Ann Surg*. 2014 Feb;259(2):329-35. doi: 10.1097/SLA.0b013e31827eefed. PMID: 23295322.
22. Mallipeddi MK, Pappas TN, Shapiro ML, Scarborough JE. Gallstone ileus: revisiting surgical outcomes using National Surgical Quality Improvement Program data. *J Surg Res*. 2013 Sep;184(1):84-8. doi: 10.1016/j.jss.2013.05.027. Epub 2013 May 31. PMID: 23764312.
23. Vera-Mansilla C, Sanchez-Gollarte A, Matias B, Mendoza-Moreno F, Díez-Alonso M, Garcia-Moreno Nisa F. Surgical Treatment of Gallstone Ileus: Less Is More. *Visc Med*. 2022 Feb;38(1):72-77. doi: 10.1159/000518451. Epub 2021 Oct 5. PMID: 35291697; PMCID: PMC8874244.
24. Vasilescu AM, Tarcoveanu E, Bradea C, Lupascu C, Stagnitti F. Gallstone Ileus. What therapeutic options are there? *Ann ItalChir*. 2022;92:300-306. PMID: 35122424.
25. Riaz N, Khan MR, Tayeb M. Gallstone ileus: retrospective review of a single centre's experience using two surgical procedures. *Singapore Med J*. 2008 Aug;49(8):624-6. PMID: 18756345.
26. Tan YM, Wong WK, Ooi LL. A comparison of two surgical strategies for the emergency treatment of gallstone ileus. *Singapore Med J*. 2004 Feb;45(2):69-72. PMID: 14985844.
27. Kirchmayr W, Mühlmann G, Zitt M, Bodner J, Weiss H, Klaus A. Gallstone ileus: rare and still controversial. *ANZ J Surg*. 2005 Apr;75(4):234-8. doi: 10.1111/j.1445-2197.2005.03368.x. PMID: 15839973.
28. Vadher D, Zacken A, Shah V, Silmi M, Aguilar LS, Patel K, Calcuttawala MA, Rajagopal P. The Rolling Stones: A Systematic Review and Meta-Analysis of the Management of Gallstone Ileus. *Chirurgia (Bucur)*. 2024 Oct;119(5):483-514. doi: 10.21614/chirurgia.3046. PMID: 39520187.
29. Zygomas A, Karamanakos S, Kehagias I. Totally laparoscopic management of gallstone ileus--technical report and review of the literature. *J LaparoendoscAdvSurg Tech A*. 2012 Apr;22(3):265-8. doi: 10.1089/lap.2011.0375. Epub 2012 Feb 3. PMID: 22303929.

30. Owera A, Low J, Ammori BJ. Laparoscopic enterolithotomy for gallstone ileus. *SurgLaparoscEndoscPercutan Tech*. 2008 Oct;18(5):450-2. doi: 10.1097/SLE.0b013e318180570f. PMID: 18936664.
31. Mir SA, Hussain Z, Davey CA, Miller GV, Chintapatla S. Management and outcome of recurrent gallstone ileus: A systematic review. *World J Gastrointest Surg*. 2015 Aug 27;7(8):152-9. doi: 10.4240/wjgs.v7.i8.152. PMID: 26328035; PMCID: PMC4550842.
32. Alzerwi NAN, Idrees B, Alsareii S, Aldebasi Y, Alsultan A. The Regularity of the Site of Impaction in Recurrent Gallstone Ileus: A Systematic Review and Meta-Analysis of Reported Cases. *Can J Gastroenterol Hepatol*. 2021 Dec 2;2021:5539789. doi: 10.1155/2021/5539789. PMID: 34900851; PMCID: PMC8660221.
33. Haddad FG, Mansour W, Deeb L. Bouveret's Syndrome: Literature Review. *Cureus*. 2018 Mar 10;10(3):e2299. doi: 10.7759/cureus.2299. PMID: 29755896; PMCID: PMC5945273.
34. Lowe AS, Stephenson S, Kay CL, May J. Duodenal obstruction by gallstones (Bouveret's syndrome): a review of the literature. *Endoscopy*. 2005 Jan;37(1):82-7. doi: 10.1055/s-2004-826100. PMID: 15657864.
35. Nickel F, Müller-Eschner MM, Chu J, von Tengg-Kobligk H, Müller-Stich BP. Bouveret's syndrome: presentation of two cases with review of the literature and development of a surgical treatment strategy. *BMC Surg*. 2013 Sep 4;13:33. doi: 10.1186/1471-2482-13-33. PMID: 24006869; PMCID: PMC3766223.
36. Dąda P, Pawlik P, Zaroda P, Niewinna P, Żuchowski M, Mańdziuk D, Kołodziej K, Kołodziej W, Wawrzkowicz J, Korga M. Bouveret's Syndrome: Diagnostic Dilemmas and Therapeutic Strategies - A Comprehensive Educational Approach for Healthcare Professionals. *J Educ Health Sport [Internet]*. 2024 May 16 [cited 2024 Dec. 16];69:48190. Available from: <https://apcz.umk.pl/JEHS/article/view/48190>
37. Caldwell KM, Lee SJ, Leggett PL, Bajwa KS, Mehta SS, Shah SK. Bouveret syndrome: current management strategies. *Clin Exp Gastroenterol*. 2018 Feb 15;11:69-75. doi: 10.2147/CEG.S132069. PMID: 29497323; PMCID: PMC5819584.
38. Dumonceau JM, Devière J. Novel treatment options for Bouveret's syndrome: a comprehensive review of 61 cases of successful endoscopic treatment. *Expert Rev Gastroenterol Hepatol*. 2016 Nov;10(11):1245-1255. doi: 10.1080/17474124.2016.1241142. Epub 2016 Oct 5. PMID: 27677937.