

# Chemical composition and antioxidant activity of *Dialiumguineense* fruit (Fabaceae) from Côte d'Ivoire

**Abstract:** This study is devoted to the phytochemical analysis of *Dialiumguineense* fruit, which is particularly popular in urban areas in Côte d'Ivoire. Despite its high consumption, there is still little information about its potential. To assess and identify the levels of various secondary metabolites, hydromethanolic extracts of the pulp, kernel, and shell of *D. guineense* fruit were analyzed via qualitative and quantitative methods (colorimetric and TLC tests) : total phenols ( $40.201 \pm 1.8$  ;  $509.023 \pm 14.5$  and  $194.368 \pm 2.2$   $\mu\text{g}$  EAG/g dry extract), flavonoids ( $58.958 \pm 6.1$  ;  $216.771 \pm 21.7$  and  $238.385 \pm 6.2$   $\mu\text{g}$  EC /g dry extract), condensed tannins ( $264.167 \pm 13.9$  ;  $411.667 \pm 25.8$  and  $282.500 \pm 5.2$   $\mu\text{g}$  EC/ mg dry extract) and hydrolysable tannins ( $0.44 \pm 0.03$  %;  $0.21 \pm 0.02$  % and  $0.30 \pm 0.03$  %). Furthermore, the analysis of antioxidant potential (DPPH and FRAP) showed satisfactory activity overall, with special attention given to the kernel and shell. These results indicate that the fruit of *D. guineense* could be viewed as a potential source of antioxidant compounds that can help prevent or fight various pathologies associated with oxidative stress.

**Key words:** *Dialiumguineense*, fruit, chemical composition, antioxidant activity

In the abstract, it should be added that this study is devoted not only to phytochemical research, but also to the recognition of antioxidant potential of *Dialium guineense*

## 1. Introduction

*Dialiumguineense*, commonly referred to as black tamarind or velvet tamarind, is an endangered plant species that is widely used by people in Côte d'Ivoire [1, 2]. Despite the high level of consumption, this species has not undergone any studies. [3] has conducted research on the shell, pulp, and kernel of this fruit, which is why. The results reveal the presence of various minerals, with a particularly high concentration of calcium (Ca), potassium (K) phosphorus (P) and magnesium (Mg). Despite not being factored into men's eating habits, the kernel is a significant protein source ( $23.58 \pm 1.18\%$ ) in comparison to other parts of the fruit. In terms of energy utilization, the pulp, kernel, and shell can generate  $317.01 \pm 0.14$  kcal/100g,  $364.47 \pm 0.35$  kcal/100g, and  $351.02 \pm 0.32$  kcal/100g, respectively. The kernel contained  $56.79 \pm 1.22$  g/100 g of carbohydrates, while the pulp had  $70.15 \pm 0.9\%$ , and the shell had  $79.44 \pm 0.7\%$ . The hull had a significant amount of total fiber ( $30.14 \pm 0.02\%$ ). The anti-nutritional elements were found to be only in the kernel ( $14.83 \pm 0.07$  mg/100g), and oxalates were found in all parts of the fruit through the evaluation of the anti-nutritional elements. With this information, the understanding of *D. Guineans* in Côte d'Ivoire is enhanced and new opportunities for this fruit's development are opened up [4–6] Incorporating antioxidant-rich food supplements into our diet is essential to combat these conditions [7]. Synthetic antioxidants have been criticized for their potential toxic risks [8]. Many researchers are investigating antioxidant molecules that come from nature for this reason. *D. guineense* has been studied in various regions, particularly in

**Comment [Γ11]:** duplicated (needs to be removed)

**Formatted:** Font color: Red

**Comment [Γ12]:** The author of the classification of the plant species should be indicated

**Formatted:** Font color: Red

**Comment [Γ13]:** the content is not completely clear

**Comment [Γ14]:** *D. guineense*

Nigeria, and it has been found to have phytochemicals with interesting potential [9, 10]. *D. guineense* has the potential to be a promising source of antioxidant substances. The aim of this study is to examine the shell, kernel, and pulp of the fruit of *D. guineense* from Côte d'Ivoire's chemical composition and antioxidant potential.

## 2. Materials and methods

### 2.1. Plant material

*Dialiumguineense* fruits (Figure 1A) were collected in March 2020 in Dimbokro (6°39' north, 4°42' west), a town in the N'Zi region of central Côte d'Ivoire. They were identified by an ethnobotanist from Nangui ABROGOUA University and then certified at the Centre National de Floristique in Abidjan in accordance with existing herbariums (N°UCJ009331). After collection, the fruits were cleaned and shelled. The various resulting parts (pulp, kernel and shell) (Figure 1(B-D)) were dried in an air-conditioned room (18°C) for 21 days, then reduced to powder using an electric grinder.




Figure 1: Unhulled fruit (A), pulp (B), kernels (C) and shells (D) of *Dialiumguineense* (Photo taken by ADOPO in 2020)

### 2.2. Methods

#### 2.2.1. Preparation of samples

100 g of powder from each part of the fruit was macerated in 600 mL of methanol (MeOH, 3 × 24 h) at room temperature (25°C) with continuous stirring. After vacuum filtration, the filtrates were combined and concentrated using a rotary evaporator (Büchi R11) at 40°C. The aqueous extracts obtained were kept in a refrigerator (4°C) for 24 h, in order to precipitate the lipophilic compounds. Once the lipophilic compounds had been removed by decantation, part of the aqueous extracts were oven-dried at 45°C until the dry hydromethanolic crude extracts were obtained, which were used for phytochemical screening using colour reactions, quantitative analysis and antioxidant potential studies. The second part of the aqueous extracts was successively exhausted in solvents of increasing polarity (3×50 mL): hexane (n-C<sub>6</sub>H<sub>14</sub>), chloroform (CHCl<sub>3</sub>), ethyl acetate (AcOEt) and n-butanol (n-BuOH). The selective extracts obtained were used for phytochemical screening by thin-layer chromatography (TLC).

#### 2.2.2. Qualitative analysis

The chemical composition of every part of the *D. guineense* fruit was determined through qualitative analysis. To achieve this, colorimetric tests and TLC were employed [11–18].

#### 2.2.3. Quantitative analysis

The content of certain compounds in each part of the *D. guineense* fruit was evaluated through quantitative analysis using spectrophotometry. The Folin-Ciocalteu colorimetric method was

Comment [F15]: the appearance of water as a solvent is not entirely clear

Comment [F16]: It is not entirely clear whether there is really a solvent in the dry extract?

used to determine the total phenols [12, 19, 20]. Aluminum chloride method was used to estimate the total flavonoid levels [21, 22]. The vanillin hydrochloride assay was used to determinate condensed tannins [22-24] method was used to evaluate the level of hydrolysable tannins.

Formatted: Font color: Red

#### 2.2.3.1. Study of antioxidant potential

DPPH test [25-27] and FRAP method [28, 29] were used to assess the free radical scavenging capacity of the various parts of *D. guineense* fruit.

#### 2.2.4. Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 22 software. Assay results, expressed as the mean of six trials  $\pm$  standard deviation, were subjected to a one-factor ANOVA at the 5% threshold. Tukey's test (verified homoscedasticity) and Tamhane's T2 test (unverified homoscedasticity) were used as post hoc tests, when the ANOVA test led to significant differences between means. These tests make it possible to group the different means into homogeneous subsets. Finally, the Pearson test was applied to establish the correlations between the different levels obtained.

### 3. Results and discussions

#### 3.1. Chemical composition of the different parts of the *D. guineense* fruit

Several secondary metabolites were discovered during phytochemical screening using colorimetric tests (Table I).  $FeCl_3$  solution was used to identify polyphenols in all parts of *D. Guineans* fruit. Kernel and pulp extracts had intense blue-black coloration, while the shell extracts had green-black coloration, suggesting a significant presence of these compounds in these parts. Different classes of flavonoids could be identified by detecting flavonoids in all three-study media using Shinoda, ammonia, and vanillin tests. According to [17], flavones and flavonols can be detected by a yellow color at room temperature or a red-orange color when heated by ammonia test, which can be verified by Shinoda's test. The presence of flavonoids is confirmed by a positive result observed with vanillin. All parts of the fruit contain tannins, coumarins, sterols, and turbanes. Saponins were detected with foam indices of 100, 125 and 200 for the pulp, kernel and shell respectively. The kernel and shell showed positive results from Borntraeger reagent, which suggests that quinones are present, as well as alkaloids detected by the Dragendorff and Wagner reagents. In spite of this, cardiotonic glycosides were not present in all parts of the fruit. According to studies on the species in Nigeria, which [30] observed in ethanolic, methanolic, and aqueous extracts, these results agree with those of other studies.

Comment [Γ17]: a period should be placed between words

Formatted: Font color: Red

Comment [Γ18]: the detector of flavonoids is not temperature but the wavelength of light in which the spots are visible

Comment [Γ19]: should be written: terpenes

Comment [Γ110]: the determination method should be specified

Comment [Γ111]: a period should be placed between words

Formatted: Font color: Red

Table I: Compounds detected in different parts of *D. guineense* fruit

		Pulp	Kernel	Shell
Polyphenols		+	+	+
Flavonoids	Shinoda's test	+	+	+
	Ammonia test	+	+	+
	Vanillin test	+	+	+
Coumarins		+	+	+
Tannins		+	+	+
Alcaloids	Dragendorff's test	-	-	+

Comment [Γ112]: Flavonoids

Comment [Γ113]: Alcaloids

	<i>Wagner's test</i>	-	+	+
	<i>Picric acid test</i>	-	-	-
<b>Sterols and terpenes</b>		+	+	+
<b>Quinones</b>		-	+	+
<b>Saponins</b>		+(Im = 100)	+(Im = 125)	+(Im = 200)
<b>Cardiotonic glycosides</b>		-	-	-

+ : Positive test; - : Negative test

In the end, the fruit of *D. guineense* has phytochemicals that have the potential to be beneficial for the body. The kernel and shell, which are often ignored by the public, could have potential uses in other fields because of their intriguing chemical composition.

### 3.2. TLC profiles of selective extracts

The presence of various phytochemicals, including sterols, terpenes, coumarins, flavonoids, tannins, and anthocyanins, was uncovered through TLC screening of *D. Guineans* shell (Coq), kernel (Noy), and pulp (Pul) (Tables II and III). To study the migration of compounds, present in hexane and chloroform extracts, solvent gradients containing  $n\text{-C}_6\text{H}_{14}/\text{CHCl}_3/\text{AcOEt}$  (4:10:1, V/V/V) and (6:10:6, V/V/V) were used. To determine what compounds were in the ethyl acetate and n-butanol extracts, the  $\text{AcOEt}/\text{CHCl}_3/\text{EtOH}$  (V/V/V) mixture was employed. Different developers were used to identify the compounds. Liebermann-Bürchard reagent was utilized to view sterols, which appeared brown and green to the naked eye and yellow and yellow-green under UV at 365 nm (Figure 2A). So, these compounds were revealed in hexanolic and chloroform extracts from shells ( $R_f = 0.34$ ) and ( $R_f = 0.19; 0.26; 0.4$  and  $0.8$ ), kernels ( $R_f = 0.32; 0.6$  and  $0.92$ ) and ( $R_f = 0.12; 0.23; 0.62$  and  $0.88$ ) and pulps ( $R_f = 0.3$  and  $0.81$ ) and ( $R_f = 0.72; 0.8$  and  $0.92$ ) respectively. The same developer was used to detect terpenes, in particular triterpenegenins [18, 31], which appear as blue and purple spots to the naked eye, and orange-yellow spots under UV light at 365 nm for lupane-type triterpenegenins. Terpenes are revealed in hexanolic extracts: Coq ( $R_f = 0.08; 0.14; 0.2; 0.34; 0.4; 0.5; 0.6; 0.7$  and  $0.9$ ), Noy ( $R_f = 0.08; 0.14; 0.2; 0.3; 0.45; 0.6; 0.7$  and  $0.81$ ) and Pul ( $R_f = 0.08; 0.2; 0.3; 0.41; 0.5; 0.54; 0.56; 0.64; 0.7$  and  $0.9$ ) (Figure 2A) and chloroformic : (Coq ( $R_f = 0.19; 0.22; 0.3; 0.4; 0.54; 0.62; 0.71; 0.86; 0.89$  and  $0.94$ ), Noy ( $R_f = 0.71; 0.88; 0.91$  and  $0.94$ ) and Pul ( $R_f = 0.12; 0.27; 0.35; 0.57; 0.67; 0.72; 0.81; 0.88$  and  $0.92$ )). The sulphuric vanillin solution was employed to show the presence of terpenes, which can be detected by looking at them in violet, pink, and orange and under UV light at 365 nm. In hexane extracts, terpenes were observed at the following  $R_f$  values: 0.08; 0.14; 0.2 and 0.46 (Coq), 0.08; 0.27; 0.5; 0.6; 0.7 and 0.92 (Noy) and 0.08; 0.3; 0.41; 0.5; 0.56; 0.64 and 0.8 (Pul). In chloroform extracts, they are found at  $R_f$  values: 0.08; 0.16; 0.62; 0.71 and 0.86 (Coq) and 0.12 and 0.88 (Pul). Sulphuric vanillin was also used to confirm the existence of sterols [31] in the various parts of the *D. guineense* fruit, visible as blue spots to the naked eye and under UV light at 365 nm. The  $R_f$  values in the hexanic extracts are: 0.21; 0.62 and 0.8 (Coq), 0.27 and 0.8 (Noy) and 0.32; 0.64 and 0.8 (Pul). In chloroform extracts, they are at  $R_f$ : 0.26; 0.37 and 0.88 (Coq), 0.84 and 0.88 (Noy) and 0.8 (Pul). Flavonoids were identified using ammonia vapour and aluminum chloride solution (1 The flavonoids are made yellow by this solution, and it can vary in coloration from blue to brown or yellow-green under UV 365 nm [11, 31]. In chloroform extracts, they are visible at  $R_f$  values of 0.06; 0.17; 0.4; 0.49; 0.54; 0.86 and 0.89 (Coq), at 0.12; 0.23; 0.29; 0.59 and 0.82 (Noy) and at 0.12; 0.35; 0.4; 0.57; 0.85; 0.88 and

**Comment [Γ114]:** it should be indicated: respectively

**Comment [Γ115]:** solvent ratio should be specified

0.9 (Pul (Rf) (Figure 2B), in ethyl acetate extracts, the Rf values are: 0.1; 0.19; 0.27; 0.34; 0.42; 0.56; 0.6; 0.7 and 0.79 (Coq), 0.19; 0.29; 0.4; 0.46; 0.61 and 0.79 (Noy) and 0.08; 0.19; 0.55; 0.66; 0.8 and 0.9 (Pul) and in the n-butanolic extracts the Rf values are 0.06; 0.11; 0.16; 0.2; 0.25; 0.31; 0.45; 0.59; 0.67; 0.74 and 0.8 (Coq), 0.17; 0.31; 0.41 and 0.49 (Noy) and 0.1; 0.1; 0.2; 0.6 and 0.8 (Pul). In the presence of ammonia vapour, flavonoids are visible as yellow and green to the naked eye and under UV 365 nm [32]. This is the case in the chloroform extracts: Coq (Rf = 0.06; 0.21; 0.49 and 0.59), Noy (Rf = 0.12; 0.25; 0.4; 0.87 and 0.91) and Pul (Rf = 0.12; 0.22; 0.29; 0.35; 0.75 and 0.88) (Figure 2B); ethyl acetate: Coq (Rf = 0.07; 0.17; 0.34; 0.49 and 0.62), Noy (Rf = 0.12 and 0.69) and Pul (Rf = 0.1; 0.45; 0.56; 0.71; 0.77 and 0.9) and n-butanol extracts : Coq (Rf = 0.16; 0.31; 0.5 and 0.75), Noy (Rf = 0.19; 0.41; 0.5 and 0.6) and Pul (Rf = 0.75)). In addition, ammonia vapour revealing blue or violet coloration indicates the presence of anthocyanins [11] in chloroform extracts at the following Rf values: 0.87 and 0.91 (Noy) and 0.88 (Pul) (Figure 2B); in ethyl acetate extracts, they are at Rf 0.22 and 0.86 (Coq), 0.19; 0.36 and 0.42 (Noy) and 0.66 (Pul) and in n-butanolic extracts, they are at the following Rf values: 0.07; 0.12 and 0.21 (Coq), 0.06; 0.08; 0.14 and 0.27 (Noy) and 0.12 and 0.89 (Pul (Rf). Yellow spots observed with the naked eye using a 5% methanolic KOH solution indicate the presence of coumarins. This yellow coloration can intensify or diversify under UV/365 nm light. In chloroform extracts, coumarins are visible at the following Rf values: 0.06; 0.12; 0.3; 0.3; 0.47; 0.54 and 0.69 (Coq), 0.06; 0.12; 0.25; 0.5 and 0.82 (Noy), and 0.06; 0.12; 0.35; 0.47; 0.85 and 0.9 (Pul) (Figure 2B). In ethyl acetate extracts, Rf values were: 0.14; 0.19; 0.22; 0.3; 0.41; 0.51 and 0.6 (Coq), 0.5; 0.7 and 0.85 (Noy), and 0.08; 0.12; 0.22; 0.26; 0.82 and 0.9 (Pul). In n-butanol extracts, coumarins appeared at the following Rf values: 0.11; 0.16; 0.2; 0.25; 0.32; 0.45; 0.6; 0.69 and 0.75 (Coq), 0.27; 0.4 and 0.5 (Noy), and 0.07; 0.29; 0.4 and 0.76 (Pul). Tannins were detected with a 2% (w/v) FeCl<sub>3</sub> aqueous solution, appearing as grey or brown stains [31, 33] in ethyl acetate extracts at Rf: 0.08; 0.19; 0.22; 0.31 and 0.71 (Coq), 0.24 (Noy), and 0.71 (Pul) (Figure 2C). In n-butanol extracts, the Rf values for tannins are: 0.16 and 0.25 (Coq), and 0.27 and 0.4 (Noy).

Comment [F116]: information is duplicated in table II

**Table II: Compounds identified in the hexanolic and chloroform extracts of the different parts of the *D. guineense* fruit.**

Solvents	Extracts	Possible compounds : [Rf], Colour, Developer
Hexane	Pul	<b>Sterols</b> : [0.3],Br <sup>e</sup> ; [0.32 ; 0.64],B <sup>a</sup> ; [0.81],Ve <sup>e</sup> , B <sup>a</sup> <b>Terpenes</b> : [0.08],J-Or <sup>e</sup> , Ve <sup>d</sup> , Vi <sup>a</sup> ; [0.2],J-Or <sup>e</sup> , [0.3],J-Or <sup>e</sup> , Vi <sup>a</sup> ; [0.41],J-Or <sup>e</sup> , Or <sup>a</sup> ; [0.5],J-Or <sup>e</sup> , Ve <sup>a</sup> ; [0.54],R <sup>e</sup> ; [0.56],R <sup>e</sup> , Ro <sup>a</sup> ; [0.64],J-Or <sup>e</sup> , Or <sup>a</sup> ; [0.7],J-Or <sup>e</sup> ; [0.8],Ro <sup>a</sup> ; [0.9],J-Or <sup>e</sup>
	Noy	<b>Sterols</b> : [0.32 ; 0.6 ; 0.92],Br <sup>e</sup> ; [0.27 ; 0.8], B <sup>a</sup> <b>Terpenes</b> : [0.08],J-Or <sup>e</sup> , Vi <sup>a</sup> ; [0.14],J-Or <sup>e</sup> ; [0.2],J-Or <sup>e</sup> ; [0.27],Vi <sup>a</sup> ; [0.32 ; 0.45],J-Or <sup>e</sup> ; [0.5],Vi <sup>a</sup> ; [0.6],J-Or <sup>e</sup> , Vi <sup>a</sup> ; [0.7],Vi <sup>e,a</sup> ; [0.81],J-Or <sup>e</sup> ; [0.92],J-Or <sup>a</sup>
	Coq	<b>Sterols</b> : [0.34], Br <sup>e</sup> ; [0.21 ; 0.62 ; 0.8],B <sup>a</sup> <b>Terpenes</b> : [0.08 ; 0.14 ; 0.2],J-Or <sup>e</sup> , Vi <sup>ea</sup> ; [0.34 ; 0.4],J-Or <sup>e</sup> ; [0.46],Vi <sup>a</sup> ; [0.5 ; 0.6 ; 0.7],J-Or <sup>e</sup> ; [0.9],J-Or <sup>e</sup>
Chloroform	pul	<b>Sterols</b> : [0.72 ; 0.8 ; 0.92],Br <sup>e</sup> ; [0.8],Br <sup>e</sup> , B <sup>a</sup> <b>Terpenes</b> : [0.12], J-Or <sup>e</sup> , B <sup>a</sup> ; [0.27 ; 0.35],J-Or <sup>e</sup> ; [0.57 ; 0.67 ; 0.72 ; 0.81 ; 0.92],B <sup>e</sup> ; [0.88],B <sup>e</sup> , B <sup>a</sup> <b>Flavonoïds</b> : [0.12],Ve <sup>b</sup> , J <sup>f</sup> ; [0.22 ; 0.29], J <sup>f</sup> ; [0.35],B <sup>b</sup> , Ve <sup>f</sup> ; [0.47], B <sup>b</sup> ;

		[0.57],J <sup>b</sup> ; [0.75], Ve <sup>f</sup> ; [0.88], B <sup>b</sup> , B <sup>f</sup> ; [0.9],J <sup>b</sup> <b>Anthocyanins</b> : [0.87],B <sup>f</sup> <b>Coumarins</b> : [0.06],Ve <sup>c</sup> ; [0.12 ; 0.35],J <sup>c</sup> ; [0.47],Ve <sup>c</sup> ; [0.54],B <sup>c</sup> ; [0.85 ; 0.9],J <sup>c</sup>
	<b>Noy</b>	<b>Sterols</b> : [0.12],J-Ve <sup>e</sup> ; [0.23; 0.62],Ve <sup>e</sup> ; [0.84],B <sup>a</sup> ; [0.88],J <sup>e</sup> , B <sup>a</sup> <b>Terpenes</b> : [0.71 ; 0.88], J-Or <sup>e</sup> ; [0.91],B <sup>e</sup> ; [0.94], J-Or <sup>e</sup> <b>Flavonoïds</b> : [0.12],Ve <sup>b</sup> ,J <sup>f</sup> ; [0.23],Ve <sup>b</sup> ; [0.25],J <sup>f</sup> ; [0.29],Ve <sup>b</sup> ; [0.4],Ve <sup>f</sup> ; [0.59],J <sup>b</sup> ; [0.82],B <sup>b</sup> ; [0.87],B <sup>f</sup> ; [0.91],B <sup>f</sup> <b>Anthocyanins</b> : [0.87 ; 0.91],B <sup>f</sup> <b>Coumarins</b> : [0.06],Ve <sup>c</sup> ; [0.12 ; 0.25 ; 0.5],J <sup>c</sup> ; [0.4],B <sup>c</sup> ; [0.82],Ve <sup>c</sup> ; [0.91 ],B <sup>c</sup>
	<b>Coq</b>	<b>Sterols</b> : [0.19],Br <sup>e</sup> ; [0.26],Ve <sup>e</sup> , B <sup>a</sup> ; [0.37 ; 0.88],B <sup>a</sup> ; [0.4 ; 0.8],Br <sup>e</sup> <b>Terpenes</b> : [0.08 ; 0.16],B <sup>e</sup> ,Or <sup>a</sup> ; [0.19 ; 0.22 ; 0.3 ; 0.4],B <sup>e</sup> ; [0.62 ; 0.71 ; 0.86], B <sup>e</sup> , Or <sup>a</sup> ; [0.54 ; 0.89 ; 0.94],J-Or <sup>e</sup> <b>Flavonoïds</b> : [0.06],J <sup>b,f</sup> ; [0.17],B <sup>b</sup> ; [0.21],J <sup>f</sup> ; [0.4],B <sup>b</sup> ; [0.49],Ve <sup>b,f</sup> ; [0.54],J <sup>b</sup> ; [0.59],J <sup>f</sup> ; [0.86],J <sup>b</sup> ; [0.89],B <sup>b</sup> <b>Coumarins</b> : [0.06 ; 0.12],Ve <sup>c</sup> ; [0.3 ; 0.35],J <sup>c</sup> ; [0.47 ; 0.54 ; 0.69],J <sup>c</sup> ; [0.88],R <sup>c</sup>

a : compounds developed with sulfuric vanillin; b: compounds developed with aluminum chloride (AlCl<sub>3</sub>); c: compounds developed with 5% potassium hydroxide (KOH); e: compounds developed with 1% Liebermann-Bürchard reagent ; f: compounds developed with ammonia vapour (NH<sub>3</sub>).

B: blue, Br: brown, J: yellow, Or: orange, R: red, Ro: pink, Ve: green, Vi: violet or purplish.


**Table III: Compounds identified in ethyl acetate and n-butanol extracts of different parts of *D. guineense* fruit.**

Solvents	Extracts	Possible compounds : [Rf], Colour, Developer
Ethyl acetate	<b>Pul</b>	<b>Flavonoïds</b> : [0.08 ; 0.19],J <sup>b</sup> ; [0.1],J <sup>f</sup> ; [0.45 ; 0.56],J <sup>f</sup> ; [0.55],Ve <sup>b</sup> ; [0.66],Ve <sup>b</sup> , P <sup>f</sup> ; [0.71 ; 0.77],Ve <sup>f</sup> ; [0.8],Ve <sup>b</sup> ; [0.9],Ve <sup>b,f</sup> <b>Anthocyanins</b> : [0.66],Vi <sup>f</sup> <b>Coumarins</b> : [0.08 ; 0.12 ; 0.22 ; 0.26],J <sup>c</sup> ; [0.71],P <sup>c</sup> ; [0.82],J <sup>c</sup> ; [0.9],Ve <sup>c</sup> <b>Tannins</b> : [0.7],Gr <sup>g</sup>
	<b>Noy</b>	<b>Flavonoïds</b> : [0.12],Ve <sup>f</sup> ; [0.19],J <sup>b</sup> , P <sup>f</sup> ; [0.29],J <sup>b</sup> , P <sup>f</sup> ; [0.36],P <sup>f</sup> ; [0.4],P <sup>b,f</sup> ; [0.46],J <sup>b</sup> ; [0.54],P <sup>f</sup> ; [0.61 ; 0.79],J <sup>b</sup> ; [0.69],Ve <sup>f</sup> <b>Anthocyanins</b> : [0.19 ; 0.36 ; 0.42],Vi <sup>f</sup> <b>Coumarins</b> : [0.06 ; 0.15 ; 0.29 ; 0.4],P <sup>c</sup> ; [0.5],Ve <sup>c</sup> ; [0.6],Gr <sup>c</sup> ; [0.7],Ve <sup>c</sup> ; [0.81],J <sup>c</sup> <b>Tannins</b> : [0.11 ; 0.15],Ro <sup>g</sup> ; [0.24 ; 0.42],Gr <sup>g</sup>
	<b>Coq</b>	<b>Flavonoïds</b> : [0.07 ; 0.17],J <sup>f</sup> ; [0.1],J <sup>b</sup> ; [0.19 ; 0.27],Ve <sup>b</sup> ; [0.22],B <sup>f</sup> ; [0.34],Ve <sup>b,f</sup> ; [0.42],J <sup>b</sup> ; [0.56],B <sup>b</sup> ; [0.49 ; 0.62],J <sup>f</sup> ; [0.6],Ve <sup>b</sup> ; [0.66 ; 0.75],P <sup>f</sup> ; [0.7],J <sup>b</sup> ; [0.79],B <sup>b</sup> ; [0.86],B <sup>f</sup> <b>Coumarins</b> : [0.14 ; 0.19 ; 0.22],J <sup>c</sup> ; [0.31],Ve <sup>c</sup> ; [0.41 ; 0.51 ; 0.6],J <sup>c</sup> ; [0.69 ; 0.71],P <sup>c</sup> ; [0.8 ; 0.9],J <sup>c</sup> <b>Anthocyanins</b> : [0.22 ; 0.86],B <sup>f</sup> <b>Tannins</b> : [0.08 ; 0.19 ; 0.22 ; 0.3 ; 0.71],Gr <sup>g</sup> ; [0.29 ; 0.36],Ro <sup>g</sup> ; [0.4],B <sup>g</sup> ; [0.5 ; 0.6],Ro <sup>g</sup>

n-butanol	Pul	<b>Flavonoids</b> : [0.1],B <sup>b,f</sup> ; [0.15 ; 0.2],Ve <sup>b</sup> ; [0.6],B <sup>b</sup> ; [0.75],J <sup>f</sup> ; [0.89],Vi <sup>f</sup> <b>Anthocyanins</b> : [0.12],B <sup>f</sup> ; [0.89],Vi <sup>f</sup> <b>Coumarins</b> : [0.07 ; 0.29 ; 0.4],Ve <sup>c</sup> ; [0.2 ; 0.47 ; 0.62],Be <sup>c</sup>
	Noy	<b>Flavonoids</b> : [0.06 ; 0.08 ; 0.14],P <sup>f</sup> ; [0.17],J <sup>b</sup> ; [0.19],Ve <sup>f</sup> ; [0.27],P <sup>f</sup> ; [0.3],J <sup>b</sup> ; [0.41],Ve <sup>b</sup> , J <sup>f</sup> ; [0.49],J <sup>b</sup> ; [0.5],Ve <sup>f</sup> ; [0.6],J <sup>f</sup> <b>Anthocyanins</b> : [0.06 ; 0.08 ; 0.14 ; 0.27],Vi <sup>f</sup> <b>Coumarins</b> : [0.08],P <sup>f</sup> ; [0.1],J <sup>c,f</sup> ; [0.14],P <sup>f</sup> ; [0.2],J <sup>c</sup> , Ve <sup>f</sup> ; [0.27],J <sup>c</sup> , P <sup>f</sup> ; [0.4],Ve <sup>f</sup> , J-Ve <sup>f</sup> ; [0.47],J <sup>c</sup> ; [0.61],Ve <sup>c</sup> ; [0.81],J <sup>f</sup> <b>Tannins</b> : [0.27 ; 0.4],Gr <sup>g</sup> ; [0.47],B <sup>g</sup> ; [0.55],Or <sup>g</sup>
	Coq	<b>Flavonoids</b> : [0.06],J <sup>b</sup> , B <sup>f</sup> ; [0.12],B <sup>b,f</sup> ; [0.16],Ve <sup>b,f</sup> ; [0.2],B <sup>b,f</sup> ; [0.31 ; 0.5],J <sup>f</sup> ; [0.25 ; 0.31 ; 0.45 ; 0.59],Ve <sup>b</sup> ; [0.67],J <sup>b</sup> ; [0.75],Ve <sup>b,f</sup> ; [0.8],B <sup>b</sup> <b>Anthocyanins</b> : [0.07 ; 0.12 ; 0.21],B <sup>f</sup> <b>Coumarins</b> : [0.07],Ve <sup>c</sup> ; [0.2],B <sup>c</sup> ; [0.29 ; 0.4],Ve <sup>c</sup> ; [0.47 ; 0.62],B <sup>c</sup> ; [0.76],J <sup>c</sup> <b>Tannins</b> : [0.16 ; 0.25],Gr <sup>g</sup> ; [0.69],Be <sup>g</sup>

b: compounds developed with aluminium chloride (AlCl<sub>3</sub>); c: compounds developed with 5% potassium hydroxide (KOH); f: compounds developed with ammonia vapour (NH<sub>3</sub>); g: compounds developed with 2% iron chloride (FeCl<sub>3</sub>).

B: blue, Gr: grey, J: yellow, Or: orange, P: purple, Ro: pink, Ve: green, Vi: violet or purplish.




**Figure 2: Some chromatographic profiles of the different parts of the *D. guineense* fruit.**


The presence of various phytochemicals was discovered through phytochemical screening, such as sterols, terpenes, flavonoids, anthocyanins, coumarins and tannins, in the various parts of the *D. guineense* fruit. These results show the wide diversity of bioactive compounds present in this fruit. The presence of flavonoids, anthocyanins, coumarins, and tannins has been observed in other *Dialium* species, including *D. gossweileri* and *D. corbisieri*[34].

### 3.3. Quantitative analysis

#### 3.3.1. Total phenol content

The levels of total phenols in the different parts of the *D. guineense* fruit (pulp, kernel and shell) (Figure 3) were measured using a calibration curve with gallic acid, the equation of which is  $y = 0.0232x + 0.0002$ .






**Figure 3: Total phenol content of the different parts of *D. guineense* fruit**

The concentrations of phenols obtained varied from  $40.201 \pm 0.718$  to  $509.022 \pm 5.912$   $\mu\text{g}$  EAG/g in samples. The kernel ( $509.022 \pm 5.912$   $\mu\text{g}$  EAG/g) had the highest content, while the pulp showed the lowest content ( $40.201 \pm 0.718$   $\mu\text{g}$  EAG/g). The total phenol content of the different parts of the *D. Guineense* fruit was significantly different ( $p < 0.05$ ) in the analysis of variance at the 5% level. Due to the variability of the variances, the Tamhane test was used as a post-hoc test. This indicated that the total phenol content of the different parts of the fruit differed significantly from one organ to another. Consequently, they are not comparable. This observation would suggest an uneven distribution of natural compounds in the different organs and physiological tissues of plants. This phenomenon has also been observed by other authors [22, 35, 36]. In addition, the phenol content of the kernel is 12.7 times higher than that of the pulp and around 2.6 times higher than that of the shell. This high concentration of phenols in the kernel suggests that it could be considered a reservoir of phenolic compounds, which could be attributed to the presence of flavonoids, coumarins and tannins detected during phytochemical screening. Similar studies conducted on the different parts of the fruit of *Tamarindus indica* (Fabaceae) revealed levels of  $152 \pm 2.2$   $\mu\text{g}$  EAG/g for the pulp,  $94 \pm 2.1$   $\mu\text{g}$  EAG/g for the shell and  $66 \pm 0.7$   $\mu\text{g}$  EAG/g for the kernel [37]. These results contrast with those obtained for *D. genus*, where the tamarind pulp contains more phenolic compounds than the shell, which is itself richer than the kernel. These observations suggest that there is an unequal distribution of phenolic compounds in the organs of a plant species, despite their belonging to the same family.

### 3.3.2. Total flavonoid content

The figure 4 shows that the flavonoid levels in different parts of the *D. Guineense* fruit varied from  $58.958 \pm 2.476$  to  $238.385 \pm 2.533$  EC/g dry matter. The catechin calibration curve was used to obtain the latter, with the formula  $y = 0.0004x - 0.0026$ .




**Figure 4: Total flavonoid content of different parts of *D. guineense* fruit**

The total flavonoid content of different parts of *D. Guineense* fruit was significantly different ( $p < 0.05$ ) according to analysis of variance at the 5% level. According to the Tamhane test, the flavonoids content in the shell ( $238.385 \pm 2.533 \mu\text{g EC/g}$ ) and kernel ( $217.71 \pm 8.863 \mu\text{g EC/g}$ ) of *D. Guineense* is comparable. *D. Guineense* shell and kernel may have similar pharmacological properties, including antioxidant potential. Flavonoids are recognized in the literature as powerful antioxidants, beneficial for the body [38]. It would therefore be wise to consume all parts of the fruit in order to maximize its many benefits. A Comparison of the proportions of total flavonoids of black tamarind and *Tamarindus indica* reveals that the latter contains fewer flavonoids (pulp ( $24 \pm 1.4 \mu\text{g EC/g}$ ); shell ( $27 \pm 1.0 \mu\text{g EC/g}$ ) and kernel ( $21 \pm 0.7 \mu\text{g EC/g}$ )) [37]. Furthermore, Pearson's test indicated a mean positive correlation ( $r = 0.672$ ) between total phenol and total flavonoid levels. This could indicate an average contribution of flavonoids to the estimation of phenolic compound levels.

### 3.3.3. Condensed and hydrolysable tannin content

The catechin calibration line ( $y = 0.004x + 0.006$ ) was utilized to calculate condensed tannin levels, and hydrolysable tannin levels were measured using a 660 nm spectrometer. The results obtained are shown in figure 5.



**Figure 5: Condensed (A) and hydrolysable (B) tannin content of the various parts of the fruit of *D. guineense***


The percentage of condensed tannins in the different parts of *D. Guineense* fruit was significantly different ( $p < 0.05$ ) in the analysis of variance at the 5% level. The Tamhane test revealed that the shell ( $282.5 \pm 2.141 \mu\text{g EC/g}$ ) and pulp ( $264.167 \pm 5.689 \mu\text{g EC/g}$ ) of *D.*

*guineense* have comparable condensed tannin contents (Figure 5A) but are different from that of the kernel. The kernel ( $411.667 \pm 10.541 \mu\text{g EC/mg}$ ) thus remains the part richest in condensed tannins. In addition, a strong positive correlation ( $r = 0.942$ ) was observed between condensed tannin content and total phenol content, indicating the significant contribution of condensed tannins to the quantification of phenolic compounds. In general, the analysis of variance applied to the hydrolysable tannin content showed a significant difference ( $p < 0.05$ ) between the different averages. The Tukey test showed that the hydrolysable tannin contents of the different parts were not comparable (Figure 5B). The presence of tannins in the fruit of *D. guineense* fruit could explain several properties attributed to it, in particular the astringent flavor felt when the pulp is eaten. This sensation arises from the interaction between the tannins of the consumed matrix and salivary proteins, particularly those rich in proline [39].

### 3.4. Antioxidant activity

#### 3.4.1. Antioxidant capacity by DPPH test

The figure 6 shows the results of the DPPH test.



**Figure 6: Antioxidant profiles of different *D. guineense* extracts**

Overall, the extracts tested had an anti-radical effect on the DPPH. Extracts from the kernel and shell showed significantly greater reduction capacity than those from the pulp. For concentrations of 4 to 0.0312 mg/mL, the reduction percentages (PR) for the kernel range from 92.728 to 26.380% and for the shell, it ranges from 84.873 to 14.756%. They are greater than 50% for concentrations between 0.125 and 4 mg/mL (Figure 6). In addition, for concentrations between 0.5 and 4 mg/mL, PR above 90% were observed for most extracts, except for the shell at 4 mg/mL with a PR of 84.873%. These results suggest that the kernel and shell are potential sources of antioxidant compounds. For the pulp, the highest PR (59.023%) was observed at 4 mg/mL, indicating a reduced capacity to neutralize DPPH compared to other organs. The Table IV presents the different CR<sub>50</sub> values, designating the concentrations that cause the loss of 50% of the DPPH. These values were determined graphically.

**Table IV: CR<sub>50</sub> (mg/mL) of crude extracts of *D. guineense* and vitamin C**

Extract	Pulp	Kernel	Shell	Vitamin C
CR <sub>50</sub>	3.102	0.110	0.193	0.008

According to the literature, the lower the  $CR_{50}$ , the greater the antioxidant activity and vice versa. On the basis of this principle, it can be stated that the kernel extract has a greater antioxidant capacity, followed by the shell. On the other hand, compared to vitamin C, the different parts of *D. guineense* exhibit much lower activity. The antioxidant properties of the kernel and shell of *D. guineense* are much stronger than those of the fruits of *D. gossweileri*. ( $CR_{50} = 500 \mu\text{g/mL}$ ). Conversely, their activity is less than that of *D. corbisieri* ( $CR_{50} = 14.44 \mu\text{g/mL}$ ) [34]. Consumption of these three parts of the fruit could therefore be beneficial for the body.

### 3.4.2. Antioxidant activity by FRAP test

The concentrations of ferric ions ( $\text{Fe}^{3+}$ ) to ferrous ions ( $\text{Fe}^{2+}$ ) were varied from 2115.91 to 51101.52  $\mu\text{g EFS/g DM}$  in the different parts of *D. Guineense* fruit. (Figure 7). These values were obtained after projection onto the calibration line of equation  $y = 0.0197x - 0.0187$  with  $R^2 = 0.9991$ , performed with iron (II) sulfate, prepared under the same conditions as the samples.




Figure 7: Reduction concentrations of  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  in the FRAP test

The shell signed the highest reduction concentration (51101.52  $\mu\text{g EFS/g DM}$ ), while the pulp presented the lowest (2115.91  $\mu\text{g EFS/g DM}$ ). This could be explained by the presence of phytochemicals such as flavonoids in the shell of *D. Guineense*, which are capable of providing electrons to reduce the ferric ion. This hypothesis was confirmed by a strong positive correlation ( $r = 0.999$ ) between the FRAP test and the total flavonoid content ( $238.385 \pm 6.2 \mu\text{g EC/g}$ ). [40] report that most flavonoids possess the ferric ion's reducing properties. However, those which have in their structure, the O-dihydroxy group in the ring B and the 3-hydroxy group and a double bond at the 2,3 positions in the ring C are the most active during the FRAP test [40]. The pulp was the least active part of *D. Guineense* fruit, regardless of the method used to evaluate its antioxidant potential. Consequently, incorporating the other parts of the fruit (shell and kernel) into dietary habits and other uses could be beneficial.

## 4. CONCLUSION

This study focused on improving the value of *D. Guineense* fruit from Côte d'Ivoire by analyzing its chemical composition and evaluating the antioxidant potential of its various components (shell, pulp, and kernel). Phytochemical screening revealed the coexistence of several phytoconstituents (flavonoids, coumarins, tannins, sterols, terpenes, alkaloids and saponins) and

Comment [Γ117]: *D. guineense*

Comment [Γ118]: *D. guineense*

Comment [Γ119]: *D. guineense*

the absence of cardiotoxic glycosides in all parts of the fruit. Furthermore, quinones and alkaloids were not visible in the pulp. Quantitative analysis revealed significant levels of total phenols ( $509.023 \pm 14.5 \mu\text{g EAG/g}$ ) and condensed tannins ( $411.667 \pm 8.7 \mu\text{g EC/ mg}$ ) in the kernel. Flavonoids are most abundant in the shell ( $238.385 \pm 25.8 \mu\text{g EC /g}$ ). The study of antioxidant potential using DPPH and FRAP methods revealed interesting antioxidant compounds in the various parts of the *D. guineense* fruit. These compounds could help to promote *D. guineense* fruit as a health food, offering an effective response to certain chronic diseases. We therefore encourage people to consume the whole fruit of *D. guineense* to reap its full benefits.

## 5. REFERENCES

- [1]. **Ayessou N. C., Ndiaye C., Cissé M., Gueye M., Sakho M. & Dornier M. (2014)**. Nutritional potential of the wild fruit *Dialiumguineense*. *Journal of Food Composition Analysis*, 34(2): 186-191.
- [2]. **Ambé G-A. (2001)**. Wild edible fruits of the Guinean savannahs of Côte d'Ivoire: state of knowledge by a local population, the Malinkés. *Biotechnology, agronomy, society and environment*, 5(1): 43-58.
- [3]. **Adopo, S. S. F., Baguia-Broune, F. D. M., Kabran, G. R. M., Mamyrbékova-Békro, J. A. & Békro, Y-A (2024)**. Chemical characterisation and nutritional potential of the pulp, kernel and shell of *Dialiumguineense* Wild fruit from Côte d'Ivoire. *RARMes Structural and Material Sciences*, 8(1): 120-129.
- [4]. **Fourcade L., Paule P & Mafart B (2007)**. Hypertension in sub-Saharan africa: current situation and outlook. *Revue generale*, 67: 559-567.
- [5]. **Schulze MB, Hu FB (2005)**. Primary prevention of diabetes : what can be done and how much can be prevented. *Annu Rev Public Health*, 26, 445-467.
- [6]. **American Diabetes Association and National Institute of Diabetes, Digestive and Kidney Diseases (2002)**. The prevention or delay of type 2 diabetes. *Diabetes Care*, 25, 742-749.
- [7]. **Pastre J. (2005)**. interest of antioxidant supplementation in the diet of domestic carnivores. thesis for the veterinary doctorate, Paul-Sabatier University of Toulouse. (France), 116p.
- [8]. **Bougandoura, N., and Bendimerad, N. (2012)**. Evaluation of the antioxidant activity of aqueous and methanolic extracts of *Saturejocalamintha* ssp. *Nepeta* (L.) Briq. *Nature and Technology*, (9) 14 - 19.
- [9]. **Onah G. T., Ajaegbu E. E. & Enweani I. B. (2022)**. Proximate, phytochemical and micronutrient compositions of *Dialiumguineense* and *Napoleona imperialis* plant parts. *GSC Biological and Pharmaceutical Sciences*, 18(03), 193–205.
- [10]. **Osanyinlusi R., Awoniyi R., Obajuluwa M. & Ogundare A. (2022)**. Nutritional composition of the fruit, leave, root and bark of Africa black velvet tamarind (*Dialiumguineense*). *International Journal of Academic Research and Development*, 7(4): 50-56.
- [11]. **Ladyguina E., Safronitch L., Otriachenkova V., Bolandina I. & Grinkevitch N. (1983)**. Chemical analysis of medicinal plants. Edition Moska; *VischayaChkola*, 46: 347 p.
- [12]. **Békro Y-A., Mamyrbekova-Békro J. A., Boua B. B., Tra Bi F. H. & Ehilé E. E. (2007)**. Ethnobotanical study and phytochemical screening of *Caesalpinia benthiana* (Baill.) Herend. Et Zarucchi (Caesalpinaceae). *Science and Nature*, 4 (2): 217-225.

- [13]. Boua B. B., Békro Y.A., Mamyrbékova-Békro J. A., Wacothon K. C. &Ehilé E. E. (2008). Evaluation of the sexual stimulant potential of total flavonoids extracted from the leaves of *Palisotahirsuta* Thumb. K. Schum (Commenilaceae). *European Journal of Scientific Research*, 22(4): 533-538.
- [14]. Konan K. (2010). Chemical study and evaluation of the antioxidant activity of four medicinal plants from Côte d'Ivoire. Single doctoral thesis, University of Abobo-Adjamé (Abidjan / Côte d'Ivoire), 112 p.
- [15]. N'Guessan H. A., Dago D. C. E., MamyrbékovaBékro J. A. &Békro Y. A. (2011). TLC of selective extracts of 10 plants used in the traditional treatment of arterial hypertension in Côte d'Ivoire. *European Journal of Scientific Research*, 575-585.
- [16]. Kabran G. R. M., Ambeu N. C., MamyrbékovaBékro J. A. &Békro Y. A. (2011). TLC of selective extracts of 10 plants used in the traditional treatment of breast cancer in Côte d'Ivoire. *European Journal of Scientific Research*, 63(4): 592-603.
- [17]. N'Gaman K. C. C. (2013). Phytochemical study and effect of extracts of *GmelinaarboreaRoxb.* (Verbenaceae) from Côte d'Ivoire on the osmotic stability of erythrocytes. PhD thesis. University NanguiAbrogoua, Abidjan (Côte d'Ivoire), 112 p.
- [18]. Bruneton J. (2016). Pharmacognosy, Phytochemistry, Medicinal plants. 5th edition, Lavoisier Tec. & Doc. Paris (France), 1487 p.
- [19]. Heilerová L., Bučkova M., Tarapčík P., Silhár S. & Labuda J. (2003). Comparison of antioxydative activity data for aqueous extracts of Lemon balm (*Melissa officinalis* L.), Oregano (*Origanum vulgare* L.), Thyme (*Thymus vulgaris* L.), and Agrimony (*Agrimoniaeupatoria*L.) obtained by conventional methods and the DNAbased biosensor. *Czech Journal Food Science*, 21(2): 78-84.
- [20]. Singleton V. L., Ortofer R. &Lamuela-Raventos R. M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of FolinCiocalteu reagent. Packer L (ed) *Methods in enzymology* Orlando Academic Press, 299: 152-178.
- [21]. Swain T and Hillis W., (1959). The phenolicsconstituants of *Prunusdomestica* -I- the quantitative analysis of phenolics constituents. *Journal of the Science of Food and Agriculture*, 10: 63-81.
- [22]. Dif M. M., Benchiha H., Mehdadi Z., Benali-Toumi F., Benyahia M. &Bouterfas K. (2015). Quantitative study of polyphenols in different organs of papaverrhoeas species. L. *Phytothérapie*, 13(5): 314-319.
- [23]. Broadhurst R. and Jones W. (1978). Analysis of condensed tannins using acidified vanillin. *Journal of the science of food and Agriculture*, 29: 788-794.
- [24]. Heilmer D., Vignadini P., Dini M. G., Vincieri F. F. & Romani A. (2006). Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food chemistry*, 99(3): 464-469.
- [25]. Takao T., Kitatami F., Watanabe N., Yagi A. & Sakata K. (1994). A simple screening method for antioxidants and isolation of several antioxidants produced by marine bacteria from fish and shell fish. *Bioscience, Biotechnology and Biochemistry*, 58: 1780-1783.
- [26]. Brand-Williams W., Cuvelier M.E. & Berset C. (1995). Use of a free radical method to evaluate antioxidant activity. *Lebensm.-Wiss. u.-Technologie*. 28: 25-30.

- [27]. Tanoh, S.K., N'Gaman-Kouassi, C.C., Boa, D., Mamyrbekova-Békro, J.A. &Békro, Y.-A. (2019). Antioxidant activity of hydroethanolic and hydroacetone crude extracts of the organs of four medicinal Côte d'Ivoire plants. *Nature and Technology*, 11 (2): 28-34.
- [28]. Patrignani M., Lupano C.E. and Rinaldi G.J. (2016). In vivo effects of Maillard reaction products derived from biscuits, *Food Chemistry*, 196: 204–210.
- [29]. Benzi I. F. and Strain J. J. (1996). The ferric reducing ability of plasma (FRAP) as measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, 239(1) : 70-76.
- [30]. Ajiboye A. E., Babatunde S. K., Adedayo M. R., Ajuwon I. B., Odaibo D. A. &Ihesie I. U. (2018). Antibacterial activity of *Dialiumguineense* seed against selected enteric bacteria. *Covenant Journal of Physical & Life Sciences*, 6(2): 1-10.
- [31]. Lagnika L. (2005). Phytochemical study and biological activity of natural substances isolated from Beninese plants. PhD thesis. University Louis Pasteur, Starsbourg (France)/ University of Abomey, Calavi (Bénin), 280 p.
- [32]. Ramamurthy V. and Schanze K. S. (2003). Semiconductor Photochemistry and Photophysics vol 10. Edition Marcel-Dekker Inc, CRC Press. New-York (USA), 384 p.
- [33]. Ekoumou C. (2003). Phytochemical and pharmacological study of 5 traditional recipes used in the treatment of urinary tract infections and cystitis. University of Bamako, (Mali), 158 p.
- [34]. Maloueki U., Kapetshi J., Kabena N. O., Ndimbo K. S-P., Ngbolua J-P., Mbomba N. B. & Muganza C-D. (2015). Antimicrobial and antioxidant activities of total aqueous extracts of *Dialiumcorbisieri* and *Dialiumgossweileri* fruits (FabaceaeCaesalpinioideae), consumed by bonobos, Bolobo, Democratic Republic of Congo. *International Journal of Biology and Chemistry Science*, 9(2): 633-642.
- [35]. Bouterfas K. (2011). Study of *Marrubiumvulgare* L. from Mount Tessala (western Algeria) autoecology, histology, quantification of some polyphenols and evaluation of the antimicrobial power of flavonoids. Master's thesis, University of Sidi Bel Abbes (Algeria), 244 p.
- [36]. Kouadio A. T., Kabran G. R., Mamyrbekova-Bekro J. A., Virieux D., Pirat JL. &Bekro Y-A. (2020). Total alkaloids and in vitro antioxidant activity of *Crinum jagus* L. (Amaryllidaceae) organs from Côte d'Ivoire. *International Journal of Green and Herbal Chemistry*, Section B: *Herbal Chemistry*, 9(4): 451-463.
- [37]. Atawodi S. E., Liman M. L., Ottu J. O. &Lliemene U. D. (2014). Total polyphenols, flavonoids and antioxidant properties of different parts of *Tamarindus indica* Linn of Nigerian origin. *Annual Research & Review in Biology*, 4(24): 4273-4283.
- [38]. Mbayo M. K., Kalonda E. M., Tshisand P. T., Tatchoua O., Kamulete S., Mbayo G. K., Kihuya E. N., Joseph B. Kahumba J.B. &Lumbu J-B. S. (2015). Chemical screening of some fungi from Katanga (DRC) and evaluation of their biological activity, *International Journal of Innovation and Applied Studies*, 10(1): 435-449.
- [39]. Canon F., Chevrier V. (2013). Wine tannins: from interactions with saliva proteins to the sensation of astringency. *Vitisphere*, 8 p.
- [40]. Firuzi O., Lacanna A., Petrucci R., Marrosu G. & Saso L. (2005). Evaluation of the antioxidant activity of flavonoids by "ferric reducing antioxidant power" assay and cyclic voltammetry. *Biochimica et BiophysicaActa* 1721: 174 – 184.