

Effect of [doses and](#) Storage Time on Acridity of Irradiated Cocoyam Corms

Abstract

Nigeria Colocasia esculentaspp (NCe 001 and NCe 011) and Xanthosoma sagittifolium spp(NXs 001 and NXs 002) were subjected to gamma irradiation of 20, 40, 80, 120 and 150Gy and stored. The acridity content of the controls samples and the irradiated and stored samples were determined. The effect of storage time on the acridity content of the irradiated cocoyam samples was studied. The main and interaction effects of storage time and dosage on acridity content were obtained statistically. The result of the study showed that there was a 50% reduction in the acridity content of the irradiated samples after 28 weeks of storage. A significant impact on the acridity content for all the varieties at 80Gy was observed from the result of the study. Gamma irradiation can with its positive advantage of no contributing to global warming serves as a means of reducing acridity content of cocoyam varieties.

Key words: Acridity, Irradiation, Storage Time

Introduction

Cocoyam (*Xanthosomasagittifolium spp.* and *Colocasia esculenta spp.*), is a member of the *Araceae* family. They are one of the oldest crops grown for its edible corms and leaves (Alabi *et al.*, 2019). Some tropical root and tuber crops, to which cocoyam belongs constitute an under-exploited resource of developing countries. Ranking as the fourteenth most consumed vegetable worldwide; cocoyam leaves are widely grown in tropical and sub-tropical countries (Adeyanjuet *et al.*, 2019). Many of the developing world's poorest farmers and food insecure people are highly dependent on root and tuber crops as a contributing, if not the principal, source of food, nutrition, and cash income (Alabi *et al.*, 2019). About 60% of the world cocoyam production (5.7 million tonnes) is in Africa and most of the remaining 40% in Asia and the Pacific (Owusu-Darko *et al.*, 2014).

Cocoyam forms part of the diet, but to a lesser extent in Nigeria where it is locally known as "Ede" in parts of Igbo land. The principal component of these tropical root and tuber crops is starch, which is increasingly becoming an important raw material for the food and non-food industries worldwide. The current industrial demand for starch is being met by a restricted number of crops mainly corn, potato and wheat. Consequently, the world starch market is dominated by starches from these three crops (Owusu-Darko *et al.*, 2014). Despite being grown on a smaller scale in Nigeria, cocoyam can offer an opportunity as a new source of starch for the Nigerian industry if handled well.

Cocoyam consumption has been affected because of the presence of acridity factors. Acridity in cocoyam causes sharp irritation and burning sensation in the throat and mouth on ingestion (Samaa, 2019). The acridity factor is one of the major problems encountered by those who consume cocoyam and this can be reduced by peeling, grating, soaking and fermentation operations during processing. Other methods of removing acridity include fermentation, baking or extraction with ethanol (Azene and Molla, 2017) and irradiation.

Food irradiation is a postharvest technology used to control insect infestation and the numbers of pathogenic or spoilage microorganisms. It also helps to arrest or reduce ripening in fresh fruits and vegetables; sprouting in tubers and germination that occurs in bulbs (Ravindran and Jaiswal 2019). During radiation processing changes occur in cells leading to the destruction of pest insects and microorganisms (Madu *et al* 2020). These radiations have the ability to remove electrons bonded in an atom and molecule with their high energy making them become electrically charged and are thus known as ionising radiations (Zanardi *et al.*, 2018).

Commented [W1]: Highlight research gap

Materials and methods

The four cocoyam varieties used in this study were NXs 001 and NXs 002 of the *Xanthosoma sagittifolium* species and cultivar; NCe 001 and NCe 011 of the *Colocasia esculenta* species and cultivar. The corms were supplied by the Cocoyam Programme of the National Root Crops Research Institute (NRCRI), Umudike, Abia State, Nigeria.

Forty-two kilogrammes (42kg) each of fresh and healthy corms of the four cocoyam varieties were sorted. The 42kg samples of corms were divided into six different groups, one group served as the control which was analysed immediately after harvest to determine the acidity content. The other five groups were packaged in polyethylene bags of 2 mm thickness and labelled accordingly. Each package containing 7kg of corms was treated with Gamma irradiation using five different doses of 20Gy, 40Gy, 80Gy, 120Gy and 150Gy. The absorbed doses were confirmed using Alanine dosimeters (Bruker Biospin USA (Billerica MA), Lot T030901). The irradiated samples were stored and analysed for acidity. The rapid test method of Association of Official Analytical Chemists (AOAC)(2005) was used to determine the acidity content.

Commented [W2]: Time duration of radiation

Commented [W3]: Unit of acidity

Results and Discussions

The results for the effect of storage on the acidity content for the irradiated NCe 001 cocoyam corms are shown in Table 1. Table 1 indicated that the acidity content of NCe 001 and NCe 011 decreased as the storage period increased. The control samples had higher acidity content than the irradiated corms. The results for the effect of storage on acidity contents for the irradiated NXs 001 and NXs 002 cocoyam corms as presented in Table 1 also indicated that the acidity content decreased as the storage period increased. The control samples also had higher acidity content than the irradiated corms. For all irradiated samples across the four varieties the 80Gy irradiated samples had the least mean acidity content. The results from Table 1 also indicate a 50% reduction in the acidity content of all the irradiated samples.

The results of the statistical analysis of variance for acidity contents of the four irradiated cocoyam corms are presented in Table 2. Table 2 indicates that the irradiated and control samples showed significant differences at $p \leq 0.05$; this is represented by the model term. It also presents that the two-way interaction had no significant difference at $p \leq 0.05$ for the interaction between dosage and storage time (0.191), dosage and variety (0.016), and for storage and variety (0.090).

The regression coefficients for acidity contents for the four irradiated cocoyam corms are presented in Table 3. Table 3 indicates that dosage had a positive effect on NCe 011 (0.065) and NXs 001 (0.516) and a negative effect on NCe 001 (-0.741). Storage had a negative effect on NCe 011 (-0.4181) and a positive effect on NCe 001 (0.0038) and NXs 001 (0.4338). The combined effect of dose and storage showed no significant difference (0.191) at $p \leq 0.05$. The effect of dose on NCe 001 (0.003) and effect of storage on NCe 011 (0.039) and NXs 001 (0.033) showed significant difference (0.191) at $p \leq 0.05$.

Table 1: Acridity content of irradiated cocoyam corms during storage

Commented [W4]: Unit of acridity

Storage Time (weeks)	NCe 001					
	Control	20	40	80	120	150
0	6.5	6.2	5.9	5.7	5.6	5.3
2	5.9	5.9	5.8	5.5	5.4	5.1
6	5.5	4.7	4.2	3.9	4.5	4.8
15	5.3	3.1	3.7	3.1	4.1	3.3
24	5.1	3.5	3.2	2.8	3.8	3.1
28	5.1	3.2	2.3	2.6	3.2	3.1
Storage Time (weeks)	NCe 011					
	Control	20	40	80	120	150
0	7.4	7.1	6.8	6.5	6.3	6.0
2	5.9	5.8	5.6	5.4	5.3	5.1
6	5.7	4.7	4.4	3.7	5.1	4.9
15	5.5	3.0	3.5	3.1	3.5	3.3
24	5.3	3.1	3.0	3.1	3.3	3.1
28	5.1	3.2	3.2	3.5	3.1	3.1
Storage Time (weeks)	NXs 001					
	Control	20	40	80	120	150
0	5.4	4.4	4.2	4.8	4.0	4.2
2	5.2	3.5	3.7	3.5	3.6	3.6
6	5.1	3.3	3.1	2.7	3.1	3.2
15	4.9	2.9	2.4	2.1	2.3	2.6
24	4.7	2.7	2.2	2.1	2.2	2.3
28	4.7	2.4	2.1	1.9	2.1	2.1
Storage Time (weeks)	NXs 002					
	Control	20	40	80	120	150
0	4.2	3.6	3.2	3.4	3.0	3.2
2	4.1	2.8	2.9	2.7	2.6	2.8
6	3.9	2.5	2.3	2.1	2.3	2.7
15	3.8	1.8	1.7	1.7	1.8	2.6
24	3.7	1.6	1.6	1.5	1.5	2.4
28	3.6	1.7	1.5	1.5	1.4	2.1

Table 2: Analysis of variance for acridity content

Source	DF	SS	MS	F-Value	P-Value
--------	----	----	----	---------	---------

Model	14	15.5957	1.1140	22.80	0.000*
Linear	5	13.8537	2.7707	56.71	0.000*
D	1	12.3692	12.3692	253.15	0.000*
ST	1	0.0090	0.0090	0.18	0.673**
V	3	1.4635	0.4878	9.98	0.000*
Square	2	0.1481	0.0740	1.52	0.247**
D*D	1	0.0456	0.0456	0.93	0.347**
ST*ST	1	0.0154	0.0154	0.32	0.581**
2-Way Interaction	7	1.1270	0.1610	3.29	0.019*
D*ST	1	0.0900	0.0900	1.84	0.191**
D*V	3	0.6643	0.2214	4.53	0.016*
ST*V	3	0.3708	0.1236	2.53	0.090**
Error	18	0.8795	0.0489		
Total	32	16.4752			
R ²		0.9466			
S		0.221045			

D = Dose (Gy); ST = Storage (weeks); V= Variety; R² = Coefficient of determination; S = Standard error of the regression; *significant difference (p < 0.05); **no significant difference (p < 0.05); DF (Degree of Freedom) = n - 1 (n = sample size); SS = Sum of Squares; MS (Mean of Squares) = SS ÷ DF

Table 3: Regression Coefficients for acidity content

Term	Effect	Coeff	SE Coeff	T-Value	P-Value
Constant		3.951	0.107	37.02	0.000*
D	2.0641	1.0320	0.0649	15.91	0.000*
ST	-0.0493	-0.0246	0.0575	-0.43	0.673**
V					
NCe 011	-0.7205	-0.3602	0.0677	-5.32	0.000*
NCe 001	0.1636	0.0818	0.0657	1.25	0.229**
NXs 001	0.4070	0.2035	0.0677	3.01	0.008*
NXs 002	0.4073	0.2021	0.0679	3.01	0.008*
D*D	0.257	0.128	0.133	0.97	0.347**
ST*ST	-0.130	-0.065	0.116	-0.56	0.581**
D*ST	0.328	0.164	0.121	1.36	0.191**
D*V					
NCe 011	0.065	0.033	0.108	0.30	0.765**
NCe 001	-0.741	-0.370	0.108	-3.43	0.003*
NXs 001	0.516	0.258	0.108	2.39	0.028*
ST*V					
NCe 011	-0.4181	-0.2090	0.0941	-2.22	0.039*
NCe 001	0.0038	0.0019	0.0885	0.02	0.983**
NXs 001	0.4338	0.2169	0.0941	2.31	0.033*

D = Dose (Gy); ST = Storage (weeks); V= Variety; *significant difference (p < 0.05); **no significant difference (p < 0.05)

Cocoyam is a good base for food preparation for infants because of the high digestibility of its starch, reasonable content of calcium and phosphorus (for bone building), B-complex vitamins and provitamin A (Ihediohanmaet *al.*, 2014). Infants in some developing countries are traditionally weaned solely on starch prepared from pre-cooked, wet-milled and wet-sieved corn (Mahmoud and El Anany, 2014). Forsido *et al.*, (2019) also reported that cocoyam starch can be incorporated in the development of weaning food which is easily digestible and accessible to low-income earners in developing countries. The major limiting factor in the utilisation of cocoyam corms is the presence of needle-like raphids of calcium oxalate crystals compounds (Rozaliet *al.*, 2021). Its oxalate content is responsible for the acrid taste or irritation when the corm is ingested without treatment (Samaa, 2019). The oxalic acid can occur as a soluble or an insoluble acid (Achadu *et al.*, 2019). The effects of these anti-nutrients usually include reduced feed intake, digestibility, nutrient utilisation, and weight gain (Ukomet *al.*, 2018). The mean acidity content of 1.7 – 7.7% obtained in this study for *Colocasia* and *Xanthosoma* spp falls within the range of values (1.8 – 8.2 %) reported by McEwan *et al.* (2014), Acheampong *et al.* (2016), and Amadi *et al.* (2018) in their works with *Colocasia* and *Xanthosoma* spp.

Commented [W5]: Discussion is very poor

Conclusion

Cocoyam with its reported nutritional advantage over other tuber and root crops has its use being limited by its acidity content. The results obtained from the study indicate that there is a direct relationship between dosage and storage on the acidity factor. Irradiation of the cocoyam varieties above 70Gy and storage time of 13 weeks reduced the acidity factor by 50%. Starch from irradiated cocoyam corms therefore can be incorporated in the development and formulation of food and food products.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

References

- Achadu, A. E., Umeh, C.C. and Mohammed, N. (2019). Proximate, phytochemicals and reducing power of leaf extracts of *Colocasia esculenta* and *Ipomoea batatas*. *International Journal of Biochemistry Research and Review*. 24(4): 1 – 11.
- Acheampong, A., Badu, M. and Agyemang A.Y. (2016). Comparative total phenolics and antioxidant activities of *X. colocasia*, *Solanum torvum* and *Allium ascalonicum* L. *International Journal of Chemistry and Biomolecular Science*. 2(4): 73- 79

Commented [W6]: Follow author's guidelines

- Adeyanju, J., Babarinde, G., Abioye, A., Olajire, A., Bolarinwa, D., (2019). Cocoyam processing: food uses and industrial benefits. *International Journal of Scientific and Engineering Research*, Vol 1, 9, 1658-1663
- Alabi, A.O., Awotunde, J.M., Adekunle, A.A., and Adeoye, A.S., (2019). Approaches for Improving Cocoyam Production and Utilization Among Rural Farmers in Ogun and Oyo State, Nigeria. *J of Biol, Agric and Healthcare*, (9)4: 20-27.
- Amadi, G.I., Achonwa, C.C., Ukwu, P.C. and Okoli, I.C. (2018). Physicochemical characteristics of Xanthosoma mafafa tuber meal subjected to palm bunch ash solution treatment. Proc. 43rd Annual Conference of the Nigerian Society for Animal Production, March 18th – 22nd 2018, FUT Owerri. Pp: 1386- 1388.
- AOAC. Official Methods of food analysis (2005). Association of Official Analytical Chemists (18th Edition), Washington DC, USA
- Azene, H., and Molla, H., (2017). Nutritional composition and effects of cultural processing on anti-nutritional factors and mineral bioavailability of Colocasia esculenta (Godere) grown in Wolaita Zone, Ethiopia. *Journal of Food and Nutrition Sciences*, vol. 5, no. 4, pp. 147–154
- Forsido, S. F. Duguma, H. T. Lema, T. B. Sturm, B. and Hensel, O., (2019). Nutritional and sensory quality of composite extruded complementary food. *Food Science & Nutrition*, vol. 7, no. 2, pp. 882–889
- Ihediohanma, N. C., Okafor, D. C., Osuagwu, P. U., and Onuegbu, N. C., (2014). Proximate composition and carotene content of three cultivars of Xanthosoma sagittifolium. *Journal of Environmental Science and Toxicology and Food Technology*, vol. 8, no. 8, pp. 17–22
- Madu, U.O., Chukwu, O., Okolo, C. A., and Haruna. S. A., (2020). Effect of Ionising Radiation and Thermal Processing on Anti-Nutritional Factors of Vegetable Cowpea (*Vigna Sesquipedalis*) Seeds. *Premier Journal of Engineering and Applied Sciences*, Vol. 1, No. 3, 1-8
- Mahmoud, A. H., and El Anany, A. M., (2014). Nutritional and sensory evaluation of a complementary food formulated from rice, faba beans, sweet potato flour, and peanut oil. *Food and Nutrition Bulletin*, vol. 35, no. 4, pp. 403–413,
- McEwan, R., Shangase, F.N., Djarova, T. and Opoku, A.R. (2014). Effect of the three processing methods on some nutrients and antinutritional factor constituents of Colocasia esculenta (Amadumbe). *African Journal of Food Science*, 8(5): 287 – 292.
- Owusu-Darko, P., Paterson, A., and Omenyo, E., (2014) Cocoyam (corms and cormels) – An underexploited food and feed resource. *Journal of Agricultural Chemistry and Environment*, Vol 3, pp 22-29

- Ravindran, R., and Jaiswal, A.K., (2019). Wholesomeness and safety aspects of irradiated foods. *Food Chemistry*, 285, 363–368.
- Rozali, Z.F., Zulmalisa, Z., Sulaiman, I., Lubis, Y.M., Noviasari, S., Eriani, K. and Asrizal, C.W. (2021). Decrease of calcium oxalate levels in the purple taro flour (*Colocasia esculenta*) from Aceh Province, Indonesia using three immersion methods. IOP conference Series: Earth and Environmental Science, 711(2021)012022
- Samaa, M.S. (2019). Reducing the soluble oxalate and phytic acid in taro corm chips by soaking in calcium salt solutions. *Alexandria Journal of Food Science and Technology*, 16(2): 9 – 16.
- Ukom, A.N., Richard, C.P. and Abasiokong, S.K. (2018). Effect of processing on the proximate, functional and anti-nutritional properties of cocoyam *Xanthosoma maffafa* (Schott) flour. *Nigerian Food Journal*. 35(2): 9 – 17.
- Zanardi, E., Caligiani, A., and Novelli, E., (2018). New insights to detect irradiated food: An overview. *Food Analytical Methods*, 11, 224–235.