

## Adoption Of Improved Package Of Practices Of Soyabean Crop And Its Constraint Faced By Soyabean Growers

### ABSTRACT

The Broad bed and furrow system has been mainly developed at the International Crops Research Institute for the Semi-arid Tropics (ICRISAT) in India. The scientist and farmers are concern with the low productivity of Soybean under present agro-situations, keeping the aforementioned perspectives in mind and realizing the importance of Broad Bed Furrow technology for soyabean crops in dissemination of technology the study was carried out to know the effect of demonstration conducted at different locations on the production and profitability of the farmers. The field experiments were conducted at the farmer's fields in the operational village of Krishi Vigyan Kendra, Sangvi (Rly) Yavatmal-II district of Maharashtra. The selection of 50 respondents was randomly drawn from villages a cluster of five villages Dhulapur, Lohi, Tarnoli, Chikhali in Darwha tehsil and Karkheda, in Ner tehsil of Yavatmal. Thus, an attempt was made to examine the adoption of broad bed furrow technology in accordance with all the 21 recommended agronomic practices-wise in order to derive significant findings and recommendations for soyabean growers and extension specialists to update farmers technical knowledge. The majority of soybean growers (60%) had low level of overall adoption of the recommended practices. Out of 21 agronomic practices recommended under BBF technology by the soybean grower; full adoption 52.00 per cent used improve the soybean varieties followed by recommended seed rate by 56.00 per cent, use of herbicide for weed control (76.00%), Majority of respondent partially adopt control insect (56 %) and disease (50%) by improved package of practices by soyabean growers. The majority of soyabean grower encounter obstacles such as the unavailability of BBF machine (96 %) and skill labours (56%), as well as challenges while utilizing BBF technology (92%). Only 50 per cent of large size holding soyabean grower use BBF technology and 44 per cent of those with higher family income have access to it. It was concluded that Improved package of practices of soybean cultivation was found superior in comparison with conventional seed drill.

**Keywords:-** Improved package of practices, Soyabean growers & Constraints

**Commented [CR1]:** Whenever "Broad bed and furrow system" is mentioned in the text follow a uniform format either starting letter of each word is capital or all letters are lower case.

**Commented [CR2]:** Give space before the starting of the sentence.

**Commented [CR3]:** Instead of 'are concern' change it to 'are concerned'.

**Commented [CR4]:** Break the sentence into two sentences. This sentence is too long

**Commented [CR5]:** Add a semicolon or 'namely'

**Commented [CR6]:** "practices-wise" is not required change it to "practices"

**Commented [CR7]:** Reframe the sentence

**Commented [CR8]:** In stead of "out of" give "Among the"

**Commented [CR9]:** Give abbreviation in brackets when it's full form is first mentioned in the text.

**Commented [CR10]:** Remove decimals while representing percentage, follow a uniform format throughout the manuscript.

**Commented [CR11]:** Rewrite the sentence

**Commented [CR12]:** Use lower case

## INTRODUCTION

Soybean (*Glycin max L.*) primarily a kharif season crop is a significant oilseed crop of India, ranking third place next to groundnut and rapeseed & mustard in terms of area and production. This crop possesses a higher potentiality to replace various oilseeds to overcome the shortage of edible oil and protein rich food. Soybean is one of the oldest cultivated crops of the world. In India large portion of the population are vegetarians, under this situation crop like soybean with high protein content and high yield potential became an important crop. Soybean protein is receiving more focus than any other source of protein today. Additionally, it provides numerous vitamins, calcium, phosphorous and iron. These characteristics make them well-suited for human consumption. Food uses of soybean include beverages; fermented products like soya sauce and cheese. Small quantities of soybean.

The scientist and farmers are worried about the low yield of Soybean under present agro situations. In order to overcome these problems, it is needed to educate and bring out the facts of the cultivation practices in the notice of the farmers through demonstration of improved production technology at their field so that they may augment technology based on seeing is believing it will develop the faith among the growers (Singh et al., 2018). In a rainfed eco-system, it is crucial to strategize agriculture by making best use of rainfall potential. For achieve a sustainable crop production system under rainfed condition, it is vital to conserve of rain water and recycle it efficiently. The rainwater can be conserved in either in-situ or ex-situ setting within natural or man-made structure for additional irrigation. In-situ rainwater conservation can be implemented through tillage or management of the land surface (Singh et al., 2000). 'The BBF landform management system primarily lowers the speed of runoff water and thereby enhance the time water to infiltrate and reduces sediment losses. Moreover, during the period of intense rainfall, the furrows facilitate the safe drainage of excess water from the plots and thus preventing water accumulation around the crop' (Kampen, 1982). Thus, keeping the above views in mind and realizing the importance of Broad Bed Furrow technology for soyabean crops in dissemination of technology the study was carried out to know the effect of demonstration conducted at different locations on the production and profitability of the farmers.

## MATERIAL AND METHODS

The field experiments were conducted at the farmer's fields in the operational village of Krishi Vigyan Kendra, Sangvi (Rly) Yavatmal-II district of Maharashtra. The selection of 50 respondents was randomly drawn from villages a cluster of five villages Dhulapur, Lohi, Tamoli, Chikhali in Darwha tehsil and Karkheda, in Ner tehsil of Yavatmal district of Maharashtra and from each village 10 respondent to assess adoption of broad bed furrow (BBF) practices of Soyabean (Var. JS 95-60) was used and constraints faced by them. Therefore, an effort was made to study all the 21 recommended agronomic practices-wise adoption of broad bed furrow technology for drawing the meaningful conclusions and suggestions for farmers and extension professionals for updating the technical know-

**Commented [CR13]:** Give tab before starting of each paragraph

**Commented [CR14]:** Scientific name should be in italics. Check the spelling of the scientific name

**Commented [CR15]:** Check the spelling

**Commented [CR16]:** Replace "Food uses of soybean include " with "Soyabean is also used to prepare"

**Commented [CR17]:** Instead of ; use and

**Commented [CR18]:** This sentence is incomplete

**Commented [CR19]:** The use of quotations '.' should be avoided while citing the text of from literature

**Commented [CR20]:** The use of quotations '.' should be avoided while citing the text of from literature

**Commented [CR21]:** The sentence could be modified by starting with Singh et al. (2008) opined that...

**Commented [CR22]:** The use of quotations '.' should be avoided while citing the text of from literature

**Commented [CR23]:** Replace " For achieve" with " Inorder to achieve"

**Commented [CR24]:** The use of quotations '.' should be avoided while citing the text of from literature

**Commented [CR25]:** Give space after full stop

**Commented [CR26]:** " realizing the importance of Broad Bed Furrow technology for soyabean crops in dissemination of technology" This portion should be replaced as " realizing the importance of Broad Bed Furrow technology for soyabean crop"

**Commented [CR27]:** Give space after full stop

**Commented [CR28]:** Rewrite the sentence.

how of the farmers. Using an interview schedule, the soyabean growers were interviewed in order to get the data. An exploratory design of social research was used for study. For the measurement of extent of adoption, a list of improved package of practices of soyabean crop was prepared and responses of the soyabean growers were collected on it. Extent of adoption was measured on three-point continuum i.e. complete, partial and non-adoption.

**Commented [CR29]:** Include pace after full stop.

**Commented [CR30]:** Change to " the extent"

**Commented [CR31]:** Change to " based on"

## RESULT AND DISCUSSION

**Commented [CR32]:** It is better to start result and discussion with an introductory paragraph rather than table.

**Table 1. Adoption of improved soybean package of practices under BBF technology by soyabean growers.**

Sr. No.	Package of practices	Full adoption	Partial adoption	No adoption
<b>A</b>	<b>Tillage and Land Preparation</b>			
1	Deep summer ploughing once in 3 years	24 (48.00)	22 (44.00)	4 (8.00)
2	Two cross harrowing/ cultivator	10 (20.00)	23 (46.00)	17 (34.00)
3	Apply FYM @ 10 t/ha	10 (20.00)	16 (32.00)	24 (48.00)
<b>B</b>	<b>Variety, Seed treatment, Sowing and plant geometry</b>			
4	Use improved soybean variety	26 (52.00)	16 (32.00)	8 (16.00)
5	Use recommended seed rate	14 (28.00)	28 (56.00)	8 (16.00)
6	Germination test	9 (18.00)	15 (30.00)	26 (52.00)
7	Seed treatment with fungicide	7 (14.00)	15 (30.00)	28 (56.00)
8	Seed treatment with Rhizobium/ PSB culture	7 (14.00)	16 (32.00)	27 (54.00)
9	Timely sowing on BBF	20 (40.00)	25 (50.00)	5 (10.00)
10	Use of intercrop	32 (64.00)	11 (22.00)	7 (14.00)
11	Maintain row to row distance	15 (30.00)	31 (62.00)	4 (8.00)
12	Plant to plant spacing	14 (28.00)	32 (64.00)	4 (8.00)
<b>C</b>	<b>Weed Control</b>			
13	Herbicide use	38 (76.00)	9 (18.00)	3 (6.00)
14	Manual weed management/ inter-culture operation	12 (24.00)	28 (56.00)	10 (20.00)
<b>D</b>	<b>Plant Nutrition</b>			
15	Application of recommended dose of NPK	15 (30.00)	30 (60.00)	5 (10.00)
16	Application of recommended dose of Zinc	11 (22.00)	14 (28.00)	25 (50.00)
17	Application of recommended dose of Sulphur	13 (26.00)	21 (42.00)	16 (32.00)

E	Plant Protection			
18	Insect management	12 (24.00)	28 (56.00)	10 (20.00)
19	Disease management	10 (20.00)	25 (50.00)	15 (30.00)
20	Soil moisture conservation	11 (22.00)	16 (32.00)	23 (46.00)
F	Harvesting			
21	Timely harvesting	18 (36.00)	22 (44.00)	10 (20.00)

Table 1 revealed that component wise adoption of improved package of practices of soyabean crops in soybean growers are as follows;

#### Tillage and land preparation

Out of 21 agronomic practices recommended under BBF technology by the soybean growers; only 48 per cent soyabean growers fully implemented deep summer ploughing once in 3-4 years. Further, followed by 46 percent partial adopted the practice of two criss-cross harrowing for seed bed preparation. Similarly, due to restricted FYM availability, only 32 per cent soyabean growers partially apply the FYM 10 t/ha to soyabean plot.

#### Variety, Seed treatment, Sowing and plant geometry

It is quite encouraging to observe that majority of soyabean grower (52.00%) having full adoption used improve the soybean varieties followed by recommended seed rate by 56.00 per cent along with 50 per cent timely sowing. In case of partial adoption, 50.00 per cent soyabean grower were maintain row to row and plant to plant 64.00 per cent in soyabean crop because lack of knowledge regarding recommended spacing.

#### Weed Control & Plant nutrition

The majority of soyabean growers utilize herbicide for weed control (76.00%) followed by 56 per cent who engage in manual weed control practices/interculture operation for managing weeds in soyabean crop. Whereas, the majority of soyabean growers use and partially apply (60.00%) recommended dose of NPK followed by 42 per cent soyabean growers partially apply the recommended amount of sulphur to soyabean crop.

#### Plant protection and Harvesting

The majority of soyabean growers partially implement for insect control (56 %) and disease (50%) by improved package of practices. Merely 32 per cent of soyabean grower partially implement soil and water conservation practice because of lack of knowledge and unawareness BBF technology. Most of respondent harvest soyabean crop at time recommended time to avoid losses by soyabean growers. The forementioned results are consistent with findings reported by Verma (2008), Singh et al (2011) and Gupta et al (2017).

**Commented [CR33]:** Give space between words

**Commented [CR34]:** Give space between words

**Commented [CR35]:** Remove "followed by"

**Commented [CR36]:** Follow uniform style in heading ie; either sentence case or first letter of each word capital

**Commented [CR37]:** Give space between words

**Commented [CR38]:** Remove decimal while mentioning percentage

**Commented [CR39]:** Replaced this portion "having full adoption used improve the" with "have adopted improved"

**Commented [CR40]:** "followed by recommended seed rate by 56.00 per cent along with 50 per cent for timely sowing" this portion can be replaced by the sentence "The recommended seed rate and timely sowing were adopted by 56 and 50 % farmers, respectively."

**Commented [CR41]:** "In case of partial adoption, 50.00 per cent soyabean grower were maintain row to row and plant to plant 64.00 per cent in soyabean crop because lack of knowledge regarding recommended spacing, ". Rewrite the sentence

**Commented [CR42]:** Add space between words

**Commented [CR43]:** Replace "followed by" by "while"

**Commented [CR44]:** Remove "who"

**Commented [CR45]:** Shift the position of 60%, for example "Whereas, the majority of soyabean growers (60%) growers use and partially apply recommended dose of NPK

**Commented [CR46]:** Add space between words

**Commented [CR47]:** Add space between words

**Commented [CR48]:** Remove "for".

**Commented [CR49]:** Modify "for insect control (56 %) and disease (50%) by improved package of practices" as "insect (56%) and disease (50%) control measures according to improved package of practices"

**Commented [CR50]:** Replace the sentence "Most of respondent harvest soyabean crop at time recommended time to avoid losses by soyabean growers" with "Timely harvest helped to avoid crop loss by majority of the farmers."

**Commented [CR51]:** Check the fond of the sentence

**Table 2. Adoption improved soybean package of practices by soybean grower.**

Adoption level	Number of soybean grower	Percentage
Low	30	60.00
Medium	14	28.00
High	6	12.00
	<b>50</b>	<b>100.00</b>

The data (Table 2) concerning the adoption levels of soybean growers related to broad bed furrow technology revealed that majority of soybean growers (60%) exhibited a low level of adoption of the suggested package of practices. Merely 12 per cent of the soybean growers adopted the technologies at high level, whereas, only 28 per cent soybean growers had medium level adoption, which present a significant issue for the development.

**Table 3. Constraints faced by soybean growers during adoption of improved soybean package of practices**

Sr. No.	Constraints	Respondent N=50	Per-cent
1	Unavailability of BBF Machine	48	96.00
2	Difficulties use of BBF technology	46	92.00
3	Lack of knowledge about package of practices in soybean on BBF technology	40	80.00
4	Lack of knowledge about importance of BBF technology in now days	37	74.00
5	Unavailability of skill labour	35	70.00
6	Lack of knowledge about soil management practices	32	64.00
7	Lack of knowledge about use of seed rate	28	56.00
8	BBF technology adopted only large size holding	25	50.00
9	Less contact with extension worker	24	48.00
10	Higher family income provides access to technology	22	44.00

**Commented [CR52]:** Follow a uniform format throughout the Text

#### Constraints faced by soybean growers during adoption

Further analysis of the data (Table 3) indicated that constraints faced by soybean grower while adopting of improved package of practices i.e. BBF technology in soybean crop. The majority of soybean growers encounters issue such as the lack of access to BBF machines (96 %) and skill labours (56%), they experience challenges when using of BBF technology (92%), are unaware of the package of practice of related BBF technology (80%) and its importance (74%). Only a significant proportion of large size holding soybean grower utilize BBF technology (50%) and those with higher family income have better access to this technology (44%).

**Commented [CR53]:** Change "indicated that" to "indicates the"

**Commented [CR54]:** Give in italics *i.e.*,

**Commented [CR55]:** Give space between words

**Commented [CR56]:** Give space between words

**Commented [CR57]:** Add "some are"

**Commented [CR58]:** Use the term soybean growers uniformly in the text

#### CONCLUSION

Based on the findings from the current study, Soybean cultivation using the tractor operated seed cum fertilizer drill method of sowing, along with harvesting through a combined harvester, is a crucial option for achieving improved crop growth, yield, lower cultivation costs. Additionally, mechanization also helps in avoid the lossescaused by unexpected rainfall during the harvesting periods. In conclusion, it was determined that the method of soybean cultivation utilizing the broad bed furrow seed drill was found superior when compared to the conventional seed drill. The technology is beneficial for increasing soybean productivity and necessitates the execution of demonstrations within the transfer of technology programme by KVK's. Horizontal spread of improved technologies may be accomplished the effective execution of frontline demonstrations and various extensions activities like training program, field day, exposure visit organized in FLDs programs in the farmer's fields

**Commented [CR59]:** Use small letters not capital

**Commented [CR60]:** Replace "through" with "using"

**Commented [CR61]:** Insert "and" after yield

**Commented [CR62]:** Give space between words

**Commented [CR63]:** Mention the abbreviation in brackets when it is first mentioned in text.

**Commented [CR64]:** Add "and " after " field day"

**Commented [CR65]:** Give full form

**Commented [CR66]:** Give full stop

### REFERENCES

- Gupta,R, Gupta B S, Kulmi G S, Somvanshi S P S and Basediya A L. (2017). Influence of broad bed furrow seed drill on growth characters and yield of soybean (*Glycine max L.*) in Mandsaur district of Madhya Pradesh. *Prog Res Int J*12 (S-IV): 2675-2678.
- Kampen, J., 1982. An approach to improved productivity on deep Vertisols. Information Bulletin No. 11, International Crop Research Institute for the Semi-Arid Tropics, Patancheru, A.P., India.
- Katare S., Singh, H.P., Pandey, S.K. and Mustafa M. 2013. Improved soybean production dissemination technology at district Ratlam Madhya Pradesh. *Int. J. of Pl. Sci.* 8(1): 197-200.
- Lakpale, Rajendra and Tripathi, V K. 2012. Broad-bed furrow and ridge and furrow method of sowing under different seed rates of soybean (*Glycine Max L.*) for high rainfall areas of Chhattigarh plains. *Soybean Res*10: 52-59.
- Singh, H.P., Venkateswarlu, B., Vittal, K.P.R., Ramachandran, K., 2000. Management of rainfed agro-ecosystem. In: Yadav, J.P.S., Singh, G.B. (Eds.), Natural Resource Management for Agricultural Production in India. International Conference on Managing Natural Resources for Sustainable Agricultural Production in the 21st Century, New Delhi, February 14-18, 2000, 669-774.
- Singh, D., Vyas, A.K., Gupta, G.K., Ramteke, K.S. and Khan, I.R. 2011. Tractor-drawn broad bed furrow seed drill machine to overcome moisture stress for soybean (*Gylcine Max L.*) in Vertisols. *Indian J. of Agri. Sci.*81(10):941-946.
- Raghuwanshi, S. R. S., O. P. S. Raghuwanshi, R. Umat, G. R. Ambawatia and Bhargav, K. S. 2009. Impact of Improved Technologies on Soybean Productivity. *Haryana J. Agro.* 25(1 & 2): 82-83.
- Verma P D, Parmanand, Jain V and Rajput A S (2018). Evaluation of sowing methods of soybean in Bhatapara District of Chhattisgarh. *J Krishi Vigyan* 6(2) : 109-112.