

EFFECT OF DIFFERENT LEVELS OF POULTRY MANURE ON GROWTH AND YIELD OF A MARANTHUS PLANT (*Amaranthus hybridus*) IN OBIO AKPA, AKWA 'BOM STATE, NIGERIA.

ABSTRACT

The field experiment was carried out at the Teaching and Research farm of Akwa Ibom State University Obio Akpa campus, in 201 B to determine the Optimum Poultry manure rate for Amaranthus,, The experiment was laid out in a randomized complete block design replicated three times. The ~~treatment were~~ four rates of poultry manure and 6 t/ha .Growth and yield parameters were assessed, ~~and Data-data~~ collected were subjected to analysis of variance (ANOVA). Significance difference at 5% probability. Results showed that poultry manure rate had significant (pe 0.05) effects on all the growth and yield parameters assessed except stem girth. The treatment that received 6 t/ ha performed best in all the parameters, while the least was recorded from the control treatment. The application of 6 t/ ha poultry manure, produced 12:19 t/ha fresh foliage yield, followed by ~~the~~ plot that received 4 t/ha poultry (1 1.33 t/ha), The treatment of 2 t/ha produced 7.70 t/ha foliage yield. The ~~least-lowest~~ foliage yield of 3.18 t/ha was harvested from the control plot the application of poultry manure also enhanced dry matter accumulation. The plot of 6 t/ha poultry manure produced 30.71g matter, while the control treatment had 33.14g. Therefore, the study recommended the application of 4 t/ha poultry manure for optimum fresh foliage yield and dry matter accumulation.

Keywords: Poultry manure, different rate, growth yield parameter, dry matter, harvesting interval, Amaranthus plant.

INTRODUCTION

Amaranthus is a perennial vegetable. Amaranthus is a cosmopolitan genus of herbs. The genus Amaranthus, in the family Amaranthaceae has more than 800 species. These include *Amaranthus hybridus*, *Amaranthus cruetus*, and others are consumed as leaf vegetables (Findal, 1975). Amaranthus originated in America and ~~is~~ one of the oldest food crops in the world, with evidence of its cultivation reaching back as far as 6700 BC. At present, Amaranthus is extensively grown as a green, leafy vegetable in many temperate and tropical regions (Khan, 2021). The largest area ever grown was during the height Of the Aztec civilization in Mexico during the 1400s. After the arrival of the Spanish conquistadors in Mexico in the early 1 500s, Amaranthus almost disappeared ~~on from the~~ America as ~~a~~ crop until research began on it in the US in the 1970s. Amaranthus has spread around the world (Including Mexico, Central America, India, and Nepal).

According to Akubugwo (2007), approximate composition Of Amaranthus hybrids L. Leaves ~~are-is~~ moisture content 83.48%, Ash, Content 1 3.80%, crude protein 17.92%, crude oil 4. 65%, crude ~~fbre-fiber~~ 8.61%, available carbohydrate 52.18 and calorific value (Kcal/ 1 00g) 26 8.92 (Khan, 2021). Nigeria has also been claimed as the ~~centre-center~~ of diversity, corroborated by the prevalent use of local names and the enormous genetic diversity available there. Amaranthus plays important roles in nutrition, it provides calcium, iron, Vitamin A (carotene) ~~and~~ Vitamin ~~e-C~~ (ascorbic acid), and vitamin ~~e-C~~ (ascorbic acid) (Palada and Chang, 2003). It is used as a refrigerant, diuretic, and purgative, as an ~~enemafer-enema for~~ stomach troubles, ~~and~~ piles, against cholera, and also as a sudorific. People with hypertension and cardiovascular diseases can use the leaves as an effective alternative to drug therapy (Akubugwo, 2007, Khan, 2021). Their nutrition quality receives world attention, especially as a rich source of minerals and protein (Olufolaji and Tayo, 1980; Ikeh et al., 2023).

Some Amaranthus species are also cultivated for their seeds, which are known to contain about 16% protein and ~~are~~ consequently utilized in confectionaries. Approximately 60 species are presently recognized with inflorescence and foliage ranging from purple and red to gold. The leaf vegetable has a rich profile of essential body metabolites such a-s vitamins A, 8, and C mineral elements such as irons, calcium, ~~and~~ phosphorus, and some measure of protein (Omueti, 1930; Esang et al 2021). It is principally grown for its large fleshly leaves and herbaceous stem Which are succulent palatable and nutritious. It can be boiled and added to soups or stews and ~~is~~ also useful for livestock feed and human consumption. Invariably, the leaves of grain amaranthus could be used for

Formatted: Justified

Commented [BK1]: Tre
treatment then add was

Formatted: Justified

Commented [BK2]: rec

Commented [BK3]: rec

feeding rabbits and other micro livestock at the early stages of growth (Palada and Chang, 2003; Akpaninyang and Opara, 2017).

Amresh et al (2009) reported hepatoprotective and antioxidant activity of 50% ethanol extract of the whole plant of *Amaranthus spinosus* against carbon tetrachloride (CC4) induced. The crop is adapted to a wide range of soil conditions. Sandy soil with slightly acidity is best suited. A temperature range of 20-30°C is required, the land should be ploughed or dug followed by leveling then shallow trenches of width 30-35cm, well rotten farmyard manure is mixed with soil in the trenches (Costea and Demason, 2007). *Amaranthus* is grown during both wet and dry seasons though irrigation is normally required for dry-season crops, since the rate of transpiration by the leaves is fairly high. It prefers soil with a pH above 6. *Amaranthus* is planted either by direct seeding or transplanting. Since *amaranthus* seeds are very small, it is mixed with sand at a ratio of 1g seed and obtain a uniform stand, then cover seed highly with a layer of compost or rice hulls immediately after broadcast (AVROC, 2003; Udounang et al 2022).

In crop production, organic and inorganic materials such as poultry manure are used to supplement chemical fertilizers (Akpan et al. 2024, Khan et al., 2021 and Akata et al., 2024). Poultry manure like other manure is known to improve the soil by enhancing both physical and chemical properties. According to Okoli et al (2015) and Ekwere et al (2022), the use of poultry manure in *amaranthus* production is desirable as it has variable impacts on most parameters assessed, the result of the study showed that the poultry manure rates are capable of improving the yield of *amaranthus*. Nweke and Obasi (2013) observed that higher organic manure rates 15 t/ha and 20 t/ha, and pig manure rates 12 t/ha and 24 t/ha, respectively gave higher yield results than where manure was not applied.

According to Grubben (1976), *Amaranthus* species are endowed with a rapid growth rate which could encourage shoot re-growth following cutting. Furthermore, Ben et al (2024) and Essien et al (2023) viewed that increasing nutrient content is by boosting with organic material; poultry manure. Hence, there is a need to carry out a research on the effect of different levels of poultry manure on the growth and yield of the *amaranthus* plant.

MATERIALS AND METHODS

Experimental Site

This research was carried out between July to October, 2018 at the Teaching and Research Farm, Obio Akpa Campus, Akwa Ibom State University, Nigeria. The research site is situated between latitude 4° 31' 1" and 5° 30' N and longitude 8° 30' E and 8° 40' E, with an annual rainfall of 2500mm. The total area of land for the experiment was 1.5m x 17m (255 m²). Twelve beds were made with the dimensions of 4m X 3m each with pathways of 1m within the land area and 0.05m between other land areas.

EXPERIMENTAL TREATMENTS

Four treatments were replicated three times which includes three different levels of Poultry manure and one in which no poultry manure was added used as control, that is C tons ha⁻¹, 2 tons ha⁻¹, 4 tons ha⁻¹ and 6 tons ha⁻¹. The poultry manures were obtained from the University poultry farm.

NURSERY MANAGEMENT

Amaranthus black seeds were sown in prepared nursery seedbeds on the experiment site. A small quantity of manure was applied in the seedbeds to aid its germination, seeds were sown by broadcasting seeds over the bed. Transplanting was done in the evening so as to reduce temperature stress,

EXPERIMENT DESIGN

The experiment was laid out in a randomized Complete Block Design (RCBD) with three replications.

Land preparation: The land was cleared and debris ~~were~~ was packed

Transplanting: After 30 days in the nursery, the seedlings were transplanted. The transplanting was done at a spacing of 50cm x 50cm.

Weeding: This was done twice at 2 weeks after transplanting (WAT) and at 6 weeks after transplanting (WAT)

Fertilizer Application: Poultry manure was applied at 2 weeks and 6 weeks after transplanting using ring method.

Data Collection: Data were collected on the field from four (4) randomly selected plants and were tagged from each plot.

The growth parameters measured were;

Plant Height (cm): The plant height was measured with the aid of a meter rule from the soil level to the plant apex.

Stem Girth (cm): This was measured by typing a thread around the base area and reading the marks off a ruler.

Leaf Area (cm²): This was determined by measuring the length and width of the leaf of tagged plants. It was later corrected factor.

Thus $LA = 0.8 \times LW$ (Kuet and svobada, 1971)

Where

L=Length Of the leaf

W=Width of leaf

Number Of leaves: This was determined by manual counting of the marketable leaves on the plant.

Number of Branches: ~~Number~~ The number of branches ~~were~~ was carefully counted on each tagged plant. The yield components were:

Dry Matter Accumulation (g): This was determined by removing the moisture content of fresh foliage either through Sun drying or oven drying.

Foliage Weight (t/ha): This can be determined by cutting the leaves of the amaranthus plant and weighed

Germination Percentage (%): The germination percentage was determined by dividing the unsuccessful transplants from the successful transplants and multiplying the total by the percentage mark (100). The number of unsuccessful plants ~~were~~ was out of 576 plant-s, given a germination percentage of 83.33% was recorded.

1 STATISTICAL ANALYSIS

Data collected were subjected to analysis of variance (ANOVA) to determine the level of significance of treatment and means were significantly separated by the use of Least Significant Difference (LSD) at a 5% probability level.

RESULT AND DISCUSSION

Result of soil physico-chemical properties at the Experimental Site before planting (Table 1), The result showed that the soil was sandy ~~soil~~. Organic carbon and total N content were low (Essien et ~~at~~ al., 2023). The soil was acidic (Essien et al. 2024) Therefore, the soil growth yield required the application of manure for proper yield

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Soil Properties

Sand	88.70
Silt	6.30
Clay	5.00
PH (H2O)	5.10
Electrical conductivity (ds/m)	0.08
Toatal Nitrogen (%)	0.11
Available phosphorus	87.55
Organic Carbon (%)	1.42
Exchangeable Ca	3.01
Exchangeable Mg	1.59g
Exchangeable K	0.11
Exchangeable Na	0.07
Exchangeable Acidity	1.95
ECEC	6.73
Base Saturation (%)	71.03

ECEC- Effective cation exchangeable capacity

Number ~~Of aof~~ Amaranthus leaves Per Plant as Influenced by Poultry Manure Rates.

~~Number~~ The number of leaves per plant of Amaranthus was significantly influenced by poultry manure application at 4, 6, 8, 10, and 12 weeks after transplanting (WAT), shown in Table 2. The amaranthus that received the higher dose of poultry manure (4 and 6 t/ha) was significantly taller than those that received 2t/ha manure, which were also taller than the control. The treatment of 6 t/ha had 38.40, 70.33, 97.52, 103.42, and 119.55 leaves per plant at 4, 6, 8, 10, and 12 WAT, respectively. The control treatment had 14.60, 27.40, 33.70, 38.30 and 46.40 leaves per plant, respectively.

Table 2, Number Of Leaves Per Plant as influenced by poultry manure

Rates of Poultry Manure (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	780	14.6	27.40	33.72	38.30	46.40

Formatted: Justified

4	9.40	34.60	68.09	91.30	101.01	12.40
6	9.42	38.40	70.33	97.52	103.42	119.55
LSD (PSO.05)	NS	3.71	5.00	5.39	6.50	6.80

* NS Not Significant

Amaranthus Height (cm) as Influenced by Poultry Manure Rates

Plant height as an influence by poultry manure rates indicated a significant difference in all the weeks under investigation except at 2 WAT- The increase in poultry manure rate from 2 to 4 t/ha indicated a significant increase in plant height (Table 3), whereas statistically there was no significant difference observed between the plant height in 4 t/ha and 6 t/ha rates, at 12 WAT- The plant that received 6 t/ha was significantly different from the height in 2 t/ha.

Formatted: Justified

Table 3: Amaranthus Height as Influenced by Poultry Manure Rate

Poultry manure Rate (t/ ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	16.75	24.46	26.31	29.40	31.01	35.73
2	15.88	46.67	49.25	53.11	53.14	61.06
4	16.42	57.75	52.36	56.91	61.33	63.9B
6	16.45	58.33	52.54	57.20	64.02	64.51
CSO (pso.05)	NS	2.62	2.70	3.01	3.18	3.25

NS = Not Significant

Leaf Area of Amaranthus as Influenced by Poultry Manure Rates

recorded in the treatment of 6 t, ~~the~~ the poultry manure rate was not statistically significant, when compared to 91.30, 98.30, 110.66, 9, 3.75, and 3.70, respectively, recorded from the treatment of 4 t/ha with the leaf area of 3 5.02, 291 5, 2560, 2048 and 20.11 cm², respectively.

Table 4: Leaf Area (cm²) Of Amaranthus as influenced by poultry manure rates

Poultry manure Rate (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	13.93	35.02	29.15	25.60	20.48	20.11
2	13.99	50.85	53.01	39.65	39.64	39.25
4	14.01	91.30	98.32	110.66	93.75	93.70
6	13.97	91.33	100.09	113.70	95.01	94.33
LSD (P<0.05)	NS	2.17	3.70	3.92	3.42	3.20

NS: Not Significant

Stem Girth Of Amaranthus as Influenced by Poultry Manure Rates

Stern girth as affected by poultry manure ~~were~~ was not significantly different (P<0.05) in all the sample weeks (Table 5). However, in the stem girth, the highest dose of poultry manure treatment recorded the biggest stem girth and was statistically significant when compared to the mean values recorded in the control treatment

Table 5: Stem girth Of Amaranthus as influenced by poultry manure rates.

Poultry manure Rate (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	1.71	2.84	3.73	3.78	3.79	3.81
2	1.75	3.71	4.81	4.88	4.89	5.03
4	1.76	3.75	4.98	5.06	5.08	5.10
6	1.73	3.76	5.10	5.11	5.15	5.17
LSD (P<0.05)	NS	NS	NS	NS	NS	NS

NS Not significant

Number Of branches per plant as influenced by poultry manure rates

~~Number~~ The number of branches per plant differed significantly (P<0.05) at 4, 6, 8, 10, and 12 WAT (T

Formatted: Right: 0.43

Formatted: Justified

Formatted: Justified

able 6). The result showed that the Amaranthus plants that received the higher dose of poultry manure (4 and 6t/ha) branched profusely compared to the number of branches in 2t/ha plots and control treatment.

Formatted: Justified

The result further indicated a significant increase in the number of branches per plant when 2t/ha rate was compared to the control plot. At 12 WAT, the plot that received 6 t/ha poultry manure had 31.75 branches, while 28.60 and 21.77 were recorded in 4 t/ha and 2t/ha poultry manure rates, respectively. The least number of branches per plant (14.06) was observed from the control treatment.

Commented [BK4]: rec

UNDER PEER REVIEW

Table 6: Number of Branches Per Plant as influenced by Poultry Manure Rates

Poultry manure Rate (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	0.33	2.89	5.75	972	1241	14.06
2	0.33	4.55	8.06	14.98	19.34	21.77
4	0.32	6.33	10.31	19.11	24.67	28.60 31
6	0.36	6.81	12.41	22.40	26.40	75
LSD (P<0.05)	NS	1.60	1.71	2.18	2.36	3.23

NS = Not significant

Total Fresh Foliage Yield and Dry Matter Accumulation

Total foliage yield as influenced by poultry manure rate varied significantly ($P < 0.05$) when compared to 7.70 t/ha and 3.18 t/ha from 2 t/ha poultry manure and control plots respectively. The total fresh foliage yield obtained from the treatment that received 6 t/ha poultry manure (12.19 t/ha) was not significantly different ($+0.05$). When compared to 1.33 t/ha total fresh foliage yield harvested from the treatment of 4 t/ha poultry manure.

Formatted: Justified

Table 7: Foliage Yield and dry Matter Accumulation (g) of Amaranthus as Influenced by Poultry Manure Rates.

Poultry manure rate (t/ha)	Total fresh foliage (t/ha)	Dry matter accumulation (g)
Control	3.1 B	23.14
2	7.70	29.25
4	11.33	30.71
6	12.19	30.71
LSD (0.05)	1.88	1.51

DISCUSSION

The significant number of leaves per plant, leaf area, plant height, total fresh foliage yield, and dry matter accumulation in the plots with poultry manure rates could be attributed to the improvement of soil fertility which in turn increased the vigor of the plants. The findings of this investigation are in consonance with the report of Ogedegbe (2013); and Akata (2015) that an increase in the dose of poultry manure significantly increases the growth and fresh foliage yield of *Amaranthus crenatus*.

Formatted: Justified

Apparently, the poultry droppings obtained from battery cage houses, which contain higher nitrogen than those from deep litter houses, where sawdust and wood shaving constitute a significant part of total manure, thereby having a high CN ratio and a lower rate of mineralization (Akata, 2006).

Ogedegbe (2013) and Akata (2015) reported higher *Amaranthus* yield in plots treated with manure obtained from battery cages. The increase in growth and yield of *Amaranthus* could be due to higher nutrient content in the manure (Akpaninyang et al., 2015; Ekwere et al., 2019).

The increase in vegetative growth with an increase in manure rate could be that the nitrogen content of the poultry manure promotes vegetative growth of the plants, this observation is in conformity with Akata et al., (2016); Ikeh et al., (2013) that N enhanced the photosynthesis process in turn resulted in high vegetative yield.

CONCLUSION

The application of poultry manure enhanced the growth and yield of Amaranthus. The fresh foliage yield and dry matter accumulation obtained from the treatment of 6t/ha ~~was~~ were significantly different when compared to 2t/ha and control treatments. Furthermore, the fresh foliage yield and dry matter accumulation recorded from the treatment that received 6t/ha and 4 t/ha poultry manure were not significantly different. It is recommended that the application of 4t/ha poultry manure should be appropriate for optimum production of Amaranthus in Obio Akpa soils of Akwa Ibom State.

REFERENCES

- Akata, O. R., Essien, O. A., Akpan, E. A. and Akpan E. G. (2024). Effect of Inorganic Fertilizer and Lime Rate on Chemical Properties of Soil and Yield of Cassava (*Manihot Esculenta* Crantz) in Obio Akpa, Akwa Ibom State. *Journal of Agriculture Environmental resources and Management* 6 (5): 48-60.
- Akata, O. R., Uko, A.E., Nwagwu, F.A., Ndaeyo, N. U., Ikeh, A.O., and Essang, D.M. (2016). Weed dynamics, growth and yield of cassava (*Manihot esculenta* Crantz) as influenced organic fertilizer sources and rates in an Ultisol. *Journal of Basic and Applied Research International*, 18(2), 68-76.
- Akpan, E. A., Akpan, A.A., Elijah, Essien, O. A. (2021). Response of Carrot (*Daucus Carota* L.) to Different Organic Fertilizers in Obio Akpa, Akwa Ibom State. *Journal of Agricultural Technology*. 1 3:46-52.
- Akpan, E. Essien, O. A., Akata, O. R. and Opeolu, E. M. (2024). Effect of Oil Palm Bush Ash and Harvesting intervals on Okra (*Abelmoschus esculentus* L.) Cultivars in Obio Akpa, Akwa Ibom State, Nigeria. *Nigerian Journal of Agriculture, Food and Environment* 20(1); 21-30.
- Akpaninyang, F. E., & Opara, E. U. (2017). The influence of toxins in disease symptom initiation in plants: a review. *Journal of Agriculture and Sustainability*, 10(1).
- Akpaninyang, F., Okpara, D. A., Njoku, J. C., Ogbologwung, P., Okpara, B., Akpan, A. E., & Jan, L. (2015). Combinations of organic and inorganic fertilizers for vine, carotene and root yields of orange-fleshed sweetpotato at Umudike, southeastern Nigeria. *Nigeria Agricultural Journal*, 46(1), 87-94.
- Akubugwo, I. E., Obasi, N. A., Chinyere, G. C. and Igbogu, A. E. (2007). Nutritional and chemical value of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. *African Journal of Biotechnology*. 6 (24), 2833-2839.
- Amresh, G., Zeashan, He, Singh, S. and Raw, C. (2009). Hepatoprotective and Antioxidant Activity of *Amaranthus spinosus* Against CCl₄ Induced Toxicity. *Journal of Ethnopharmacology*. 125: 364-6.
- AVROC (2003). Progress Report 2003. Asian Vegetable Research and Development Centre, Shanghai, Taiwan. pp. 80-9046.
- Ben, F. E., Okpara, D. A., Akpan, E. A., Udounang, P. I., Akata, O. R., & Inyang, P. (2024). Response of orange-fleshed sweetpotato (*Ipomea batatas* (L) Lam) to ratoon cropping and nutrient management in Southern Nigeria. *International Journal of Agricultural Policy and Research*, 12(2), 23.
- Costea, M. and Demason, D. (2007). "Stem Morphology and Anatomy in *Amaranthus* L. (Amaranthaceae). Taxonomic Significance". *Journal of the Torrey Botanical Society*. 128 254 - 281.
- Ekwere, O. J., Akpan, E. A., & Akata, O. R. (2019). Performance and yield of fluted pumpkin (*Telfairia occidentalis* (L.) hook) on an ultisol amended with soybean meal. *J. Agric. Environ. Resour. Manag. ISSN2245-1800 (paper) ISSN, 2245-2943*.
- Ekwere, O. J., Ilimoh, F. O. and Essien, O. A. (2022). Effect of Poultry Manure and Lime on Growth and Yield of Okra. *Mgenan Journal of Crop Science*. 8(1): 15-22.
- Essang, D. M., Akata, O. R., & Ikeh, A. O. (2021). Adaptability of Seven Maize (*Zea mays* L.) Varieties in High Humid Rainforest Zone of Nigeria. *American Journal of Agricultural Science, Engineering, and Technology*, 5(2), 92-101.

- Essien O. A., Akpan, E. A. and Akata, OR. (2023). Comparative Methods of Land Use Practices Effect on Clay Dispersion in Coastal Plain Sand Soils in Southeastern Nigeria. *Journal of Agriculture Environmental Resources and Management* 5(6) 10221320.
- Essien, O A, Inyang, E. A. and Simeon, S. O (2024). Aggregate-size Distribution: Impact on Bulk Density and Hydraulic Conductivity of Soil Derived from Two Parent Materials in South South Nigeria. *Nigerian Journal of Agriculture, Food and Environment* 20(1):78-89.
- Findall, H. D. (1975). *Commercial Vegetable Growing Oxford Tropical Handbook*. Oxford University Press.
- Grubben, G. J. H, (1976). Cultivation of Amaranthus as a Tropical Leaf vegetable With Special Reference to Southern Oahomey communication Department Of Agriculture research Royal tropical Institute (Netherland) 67: 65-80.
- Ikeh AO, Orji JO, Sampson JO, HU. Akata, OR. (2023). Effect of oil palm bunch refuse ash in sustainable production of Egusi-Melon (*Colocynthis citrullus*) in an Ultisol. *Asian Journal Agricultural and Horticultural Research*. 10 (3)138- 148.
- Ikeh, A. O, Ndaeyo, N. Ll., Akpan, E. m, Lidoh, E. I and Akata, O. R. (201 3). Evaluation of Complementary use of Organic Manure for Sustainable Water Yam Production in Oyo, Southeastern Nigeria. *American Journal Of Research Communication*. 1 (2), 33-48.
- Khan, A.U., Ema, I.J., Afsana, A.S. and Khan, A.U. (2021). Home Garden and Women Participation: A Mini Review. *Current Research in Agriculture and Farming*, 2(4): 46-52. Doi: <http://dx.doi.org/10.18782/2582-7146.152>.
- Khan A. U. (2021). Diseases Management in The Agriculture Plants and Crops. *Insect Pests, Diseases and Weeds Management*. 6376: 1-46. Available at: <https://easychair.org/publications/preprint/zWpF>.
- Khan A. U. (2021). Insect Pests Management in The Agriculture Plants and Crops. *Insect Pests, Diseases and Weeds Management*. 6375: 1-50. Available at: <https://easychair.org/publications/preprint/bsnk>.
- Khan A. U. (2021). Weeds Management in The Agriculture Plants and Crops. *Insect Pests, Diseases and Weeds Management*. 6377: 1-35. Available at: <https://easychair.org/publications/preprint/7xDW>.
- Nweke, I. A. and Obasi, M. N. (2013). Effect of Different Levels of Pig Manure on the Growth and Yield of Okra Proceedings of the 47th Annual Conference of the Agricultural Society of Nigeria, Ibadan, 201 3, Pp 23-26.
- Ogedegbe, S. A, Ogah, J. J and Ajaja, 8. A. (2013) Response of Amarath (*Amaranthus Sp p. L.*) Varieties Of Organic Fertilizer in Vom- Nigeria in University Of Nigeria Nsukka pp. 9086.
- Okoli, P.S.O. and Nweke I. A (2015). Effect of Different rate of Poultry Manure on Growth and Yield of Amarath us (*Amaranthu s Cruentus*). /OSR Journal of Agriculture and veterinary Science (IOSR JAYS) eISSN: 2319 - 2380. ISSN: 2319 - 2372. www.io srjournals.org.
- Olufolaji, A. O. and Tayo. T.O. 1980. Growth, Development and Mineral Contents of Three Cultivals of Amaranth. *Scientia Horticulture*, 13: 1 31-189.
- Omueti, O. (1930). Effects of Age of Celosia Cultivars. *Experimental Agriculture*. 1 6: 279286. 49
- Palada M.C. and Chang L. C. (2003). AVRDC International Cooperators Guide. Suggested Cultural Practices for Vegetable Amaranth <http://www.aardc.org>
- Udounang, P. I., Ekwere, O. J., & Akpainyang, F. E. (2022). Comparative effect of fertilization on cocoyam and component crops on Cocoyam-based intercropping systems in Uyo, Southeastern Nigeria. *International Journal of Agricultural Policy and Research*, 10(1), 16.